

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Vegetation influence on soil formation rate in a proglacial chronosequence (Lys Glacier, NW Italian Alps)**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/154334> since 2016-10-13T14:58:38Z

*Published version:*

DOI:10.1016/j.catena.2013.10.001

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted here by agreement between Elsevier and the University of Turin. Changes resulting from the publishing process - such as editing, corrections, structural formatting, and other quality control mechanisms - may not be reflected in this version of the text. The definitive version of the text was subsequently published in CATENA, 113, 2014, 10.1016/j.catena.2013.10.001.

You may download, copy and otherwise use the AAM for non-commercial purposes provided that your license is limited by the following restrictions:

- (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-ND license.
- (2) The integrity of the work and identification of the author, copyright owner, and publisher must be preserved in any copy.
- (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), 10.1016/j.catena.2013.10.001

The publisher's version is available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0341816213002452>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/2318/154334>

1 **Vegetation influence on soil formation rate in a proglacial chronosequence (Lys Glacier, NW**  
2 **Italian Alps)**

3 Michele E. D'Amico\*, Michele Freppaz, Gianluca Filippa, Ermanno Zanini

4 DISAFA, Università degli Studi di Torino, Via Leonardo da Vinci 44, Grugliasco (TO), Italy and

5 NatRisk

6 \*: corresponding author. tel: +39 3490611313; e-mail: ecomike77@gmail.com

7

8 **Abstract**

9 Climate change has huge impacts on alpine ecosystems . One of the most visible effects in the Alps  
10 is glacier retreat since the end of the Little Ice Age (LIA), which caused the exposure of previously  
11 glaciated surfaces. These surfaces are open-air laboratories, verifying theories regarding ecosystem  
12 and soil development.

13 In order to increase our knowledge on the effects of time and vegetation primary succession on soil  
14 development in proglacial areas, we sampled soils and surveyed plant communities on stable points  
15 in the proglacial area of the Lys glacier, in the Italian north-western Alps (Valle d'Aosta Region).

16 The sampling points were located on dated sites (based on literature and/or historical photographs).

17 Glacial till is attacked by weathering processes immediately after deposition and stabilization, with  
18 a consequent loss of soluble compounds, decrease of pH and primary mineral weathering. The  
19 speed of these processes was largely increased after the establishment of a continuous vegetation  
20 cover, thanks to the organic matter accumulation caused by litter input and root decomposition  
21 below the soil surface.

22 On sialic glacial tills, below timberline and under a quasi-climax Larch - Rhododendron forest, a  
23 fast and steady decrease in pH values, increase in organic matter content and horizon differentiation  
24 was observed. In particular, genetic eluvial horizons formed in approximately 60 years, while  
25 diagnostic albic horizons were developed after ca. 90 years, pointing to an early start of the  
26 podzolization processes. Cheluviation of Fe and, secondarily, Al were analytically verified.

27 However, illuviation of Fe, Al and organic matter in incipient B horizons were not sufficient to  
28 obtain diagnostic spodic horizons on LIA materials.

29 Under grazed grassland below timberline and alpine prairie above timberline, acidification and  
30 weathering were slightly slower, and no redistribution with depth of Fe and Al oxi-hydroxides was  
31 observed. A cambic Bw horizon developed on the oldest LIA moraines.

32 Therefore it seemed that this fast onset of the podzolization process in comparison to other  
33 proglacial chronosequences in the Alps was mainly driven by vegetation properties rather than by  
34 specific climatic conditions.

35

36 **Keywords**

37 Soil formation; proglacial area; soil chronosequence; podzolization; subalpine soils; alpine soils

38

39 **Highlights**

- 40 • Plant succession and soil development were slower above than below timberline;
- 41 • Under grassland below and above timberline, Cambisols were found on old moraines;
- 42 • Under typical subalpine forest-shrub vegetation, podzolization was very fast;
- 43 • E horizons and “real” Podzols appeared faster than in other forefields in the Alps;

- Vegetation seemed very important in the early podzolization onset.

45  
46

## 1. Introduction

47 Climate change is having huge impacts on mountain ecosystems. One of the most visible effect is  
48 the glacier retreat, which continued with few interruptions since the end of the Little Ice Age (LIA,  
49 between ca. years 1300 and 1821/1861, Ivy-Ochs et al., 2009). The bare surfaces left by retreating  
50 glaciers offer the opportunity to observe the early stages of soil development and the primary  
51 ecogenetic succession, thus validating existing theories about ecosystem evolution and allowing to  
52 determine the speed of soil forming processes. A large number of soil chronosequences in proglacial  
53 areas has been studied on mountain ranges worldwide and in arctic and antarctic environments (e.g.  
54 Ugolini, 1966, Burt and Alexander, 1996, Egli et al., 2001, Dümig et al., 2011, Mahaney et al.,  
55 2009, Hodkinson et al., 2003). According to most of these studies, soil chronosequences under  
56 boreal/subalpine forests frequently end up with Podzols, but the rate of development of a podzolic  
57 morphology and chemistry widely differ in several studies considering different environments. For  
58 example, in superhumid basins in New Zealand, with an average annual rainfall higher than 10000  
59 mm (Tonkin and Basher, 2001), Podzols were formed in about 1000 years, despite the extremely  
60 fast weathering of the gravel and sand fractions and the consequent increase in silt and clay contents  
61 in surface horizons. In Alaska, high rainfall and humidity leads to the formation of weakly  
62 developed E and Bs horizons after only 70 years, and “real” Podzols after 230 years (Ugolini, 1966,  
63 Burt and Alexander, 1996). In these areas, E horizons immediately appeared after the establishment  
64 of spruce in the forest succession. This happened despite the presence of abundant carbonatic  
65 materials in the parent till. Similar fast rates of Podzol formation (less than 230 years) were  
66 observed on marine terraces in Finland (Mokma et al., 2004).

67 Focusing on the European Alps, many studies have been performed on soil and ecosystem  
68 development in proglacial areas in subalpine forest habitats (e.g. Egli et al., 2003, Egli et al., 2006,  
69 Dümig et al., 2011), while a smaller number was devoted to alpine (i.e. above timberline) habitats,  
70 with most of them considering mainly primary plant succession (e.g., Andreis et al., 2001,  
71 Caccianiga et al., 2006, Raffl et al., 2006).

72 In the fifties some works described Podzol-like soils on recently deglaciated surfaces also in the  
73 Alps: for example, Jenny (1958) showed a 315 years old soil which had a 11 cm thick light gray  
74 horizon overlying a 14 cm thick brownish one, under subalpine ericaceous shrubs in the Rhone  
75 Gletscher forefield. However, in the alpine range, much longer periods since deglaciation were  
76 normally required for the establishment of the podzolization process than in other boreal habitats.  
77 Under subalpine vegetation, the youngest soils were usually Lithic and Skeletic Leptosols; after 120  
78 years soil evolution led to the development of Humi-Skeletic Leptosols or, locally, to weakly  
79 developed Dystric Cambisols (Righi et al., 1999, Egli et al., 2001, Egli et al., 2006). A continuous

80 humus layer appeared in the oldest soils, together with weak signs of chemical weathering  
81 (formation of Fe and Al oxyhydroxides). Oxalate extractable Fe and Al decreased with depth, and  
82 no cheluviation was visible (Egli et al., 2006). In the Alps, Dystric Cambisols are normally found  
83 on 250-300 years old surfaces, while Podzols appear after 600 years in very humid sites (1800  
84 mm/y, Zech and Wilcke, 1977), but usually more than 1300 years are needed for the development of  
85 E horizons (Egli et al., 2003). In central Switzerland, in areas characterized by mean annual  
86 precipitation equal to ~2000 mm/y, Podzols are found in 3300 years old sites (Egli et al., 2001,  
87 Dümig et al., 2011), but they are normally weakly developed (Righi et al., 1999).

88 The establishment of vegetation on the bare deglaciated areas initiates gradients in many soil  
89 properties. The primary succession of vegetation in several glacier forefields in the European Alps  
90 include pioneer stages dominated by *Oxyrietum digynae* and *Epilobietum fleischeri* (Braun-  
91 Blanquet 1948), lasting around 30 years, followed by the establishment of an open forest. *Larix*  
92 *decidua*, *Rhododendron ferrugineum* and other Ericaceae start colonizing regolith surfaces after  
93 about 20-30 years since deglaciation, but *Rhododendron* generally shows low cover values in most  
94 Swiss chronosequences, where the LIA primary succession was normally dominated by *Alnus*  
95 *viridis* (e.g., Burga, 1999, Föllmi et al., 2009, Burga et al., 2010) (except in the Damma glacier  
96 forefield, Bernasconi et al., 2011).

97 Despite these well-established patterns of soil and plant community chronosequences, in many  
98 glacier forefields of the Aosta Valley (North-western Italian Alps), both plant community succession  
99 (particularly at the subalpine belt) and pedogenesis seem faster than in most other proglacial areas  
100 in the Alps. For example, in the proglacial area considered in this study, *Rhododendron ferrugineum*  
101 dominates the understory vegetation already after few decades from glacial retreat (Treter et al.,  
102 2002). During many soil survey campaigns in Aosta Valley, a well-defined bleached E horizon was  
103 commonly recognized in soils developed on 130 years old moraines, associated with weakly  
104 developed BCs horizons, slightly enriched in pedogenic oxyhydroxides; this pattern was recorded  
105 also in the Miage glacier forefield in the Mont Blanc Massif and in the Verra Grande glacier  
106 forefield in the Monte Rosa Group. In these areas, above the treeline, ecosystem and soil  
107 development had a similar rate as in other similar alpine environments (Andreis et al., 2001), and  
108 “climax” soils and plant communities are reached in longer periods than at lower elevation.

109 Thanks to the well-known chronology of deglaciation of the Lys glacier (e.g., Monterin, 1932), we  
110 chose its proglacial area to investigate the rates of the soil forming processes in the Alps. The Lys  
111 forefield includes habitats both below (subalpine) and above timberline (alpine), thus evidencing  
112 multidirectional trends in pedogenesis and vegetation succession (Huggett, 1998), hence  
113 contributing to the detection of the main driving factors of these processes.

114 The comparison between contrasting subalpine and alpine primary vegetation successions and soil  
115 chronosequences provides useful information to evaluate the importance of pedoclimatic and  
116 vegetation spatial variability in driving the direction of pedogenic processes.  
117

## 118 **2 Material and Methods**

### 119 **2.1 Study area** 120

121 The proglacial area of the Lys Glacier is located in the upper Lys Valley (Aosta Valley, Pennine  
122 Alps, Italy, Figure 1). The morainic systems left by several glacier fluctuations during the Holocene  
123 were usually erased by the larger advance of the Little Ice Age (LIA), between ca. years 1350 and  
124 1850 (Joerin et al., 2006). The maximum LIA glacier advance was reached in 1821; a secondary  
125 advance ended in 1861, when the glacier approached the moraine deposited in 1821, leaving no  
126 frontal morainic arc because of river erosion. Since 1861, approximately 1.8 km were left free of  
127 ice. A minor advance (1915-1921) left a small morainic arc about 800 m from present day glacier  
128 terminus.

129 We sampled a recent soil (P1, ca. 7-10 years old) as representative of fresh, raw till (starting point of  
130 soil development and vegetation succession for both alpine and subalpine chronosequences). Above  
131 timberline (“alpine” chronosequence), we sampled soils formed at 4 different sites under common  
132 vegetation covers (Table 1). Below timberline (subalpine chronosequence), we sampled soils at 5  
133 different sites (Table 1); if two different vegetation types occurred on same-age surfaces, we opened  
134 a soil pit under each type (i.e. S3, S4, S6 were observed under larch forest with *Rhododendron*  
135 *ferrugineum*, SG2, SG5, SG7 below grazed grassland). S10 and A6 are on late-glacial (Younger  
136 Dryas) till (more than 10000-11000 years old, according to Pelfini et al., 1997). This age is  
137 attributed thanks to similarities with most other proglacial areas in the Alps (Ivy-Ochs et al., 2009).  
138 We excluded areas visibly disturbed by erosion or deposition processes. All soil profiles were  
139 chosen as representative among a much larger number of soil observations by soil coring. Since  
140 small scale topographic differences have strong effects on soil development and vegetation  
141 succession dynamics (Burga et al. 2010), sites with similar surface rockiness and slope were  
142 chosen.

143 Unfortunately, below the present-day glacier front the slope is steep and eroded (“Rocce di  
144 Salzen”), and it becomes reasonably stable only 200 m below, where the material was left by the  
145 glacier around year 1945. Therefore, no soil has been sampled and analyzed with ages ranging  
146 between 5/10 and 60/70 years.

147 The altitude of the proglacial area ranges from 1990 m a.s.l. to about 2480 m a.s.l.. The highest  
148 morainic ridge (2480 m a.s.l.) was deposited in 1755 (Strada, 1988). The present-day glacier tongue  
149 is almost 100 m above the natural timberline in the Lys valley, even though young individuals of  
150 larch (*Larix decidua*) are found up to 2400 m a.s.l., evidencing an ongoing increase in timberline  
151 associated with recent climate change and/or reduction in cattle grazing pressure. The whole Lys  
152 proglacial area is roughly exposed to the south and only the most ancient subalpine LIA soil profiles  
153 are located on the northward slopes of the 1821 morainic arc.

154 The parent glacial till is made of granitic gneiss and paragneiss belonging to the Monte Rosa nappe,  
155 with minor (ca. 10%) mafic and ultramafic inclusions derived from ophiolitic outcrops in the  
156 southernmost portions of the glaciated part of the massif, belonging to the Piedmontese Ophiolitic  
157 Units (Mattirolo et al., 1951). Based on the observation of the stone fraction, the lithological  
158 composition of the glacial till was similar in every stage of the soil chronosequences. Only the pre-  
159 LIA alpine site (A6) had a higher mafic-ultramafic content (ca. 30% in weight of rock fragments)  
160 than the other sites.

161 The soil moisture regime is udic (Soil Survey Staff, 2010), with a mean yearly rainfall around 1200  
162 mm (Figure 2) and no dry season (alpine subatlantic climate). The south-north direction of the Lys  
163 Valley increases the advection of warm, moist Mediterranean air masses from the south, increasing  
164 summer rainfall, while the proximity to the main Alpine divide allows some spillover of  
165 precipitation also from the north during strong foehn wind events. The mean annual air temperature  
166 ranges between ca. 2°C at 1900 m a.s.l. and -1°C at 2400 m a.s.l., while winter average temperature  
167 is below -4°C (Mercalli, 2003).

168

## 169 **2.2 Soil description vegetation survey and numerical analysis**

170 Floristic relevées were performed on 4X4 m square surfaces around the soil pits. The plant species  
171 were determined according to Pignatti (1992) and single species cover (%) was visually estimated.  
172 Total vegetation cover (%), proxy for plant productivity, was estimated as well.

173 Field description of soil profiles was done according to FAO (2006). Approximately 0.5-1 kg of soil  
174 material was collected from every horizon in the soil pits. In the field we were not able to obtain  
175 samples for the calculation of bulk densities because of excessive stoniness and/or the abundant  
176 presence of medium and/or large roots. The soil chemical and physical analyses were performed  
177 according to standard methods (Ministero delle Politiche Agricole e Forestali, 2000).

178 All samples were air-dried and sieved to separate the fine earth (below 2 mm) from the coarse  
179 fraction. pH was measured in water (soil:water=1:2.5); total carbon (corresponding to total organic  
180 carbon, TOC, thanks to the absence of carbonates) and nitrogen (TN) were analyzed by dry



181 combustion with a CN elemental analyzer (CE Instruments NA2100, Rodano, Italy). The cation  
182 exchange capacity (CEC) was measured with the ammonium acetate extraction (pH 7) method, in  
183 order to classify soils according to the IUSS Working Group (2006). Exchangeable base content and  
184 saturation, on the ammonium acetate extracts, were measured by AAS (flame atomic absorption  
185 spectrometer, Analyst 400, Perkin Elmer, Waltham, MS, USA). The particle size distribution was  
186 determined by the pipette method. In order to detect the spodic properties, the oxalate and  
187 dithionite-extractable fractions of Fe and Al (Fe<sub>o</sub>, Al<sub>o</sub>) were measured.

188 Chronofunctions of TOC concentration changes in surface horizons, of the Spodic Index (later  
189 on, IS, equation 1, Soil Survey Staff, 2010) in CB-BC-Bs horizons and the ratio between the IS in  
190 subsurface CB-BC-Bs horizons and in surface A-AE-E ones (later on, IS<sub>ratio</sub>, equation 2) in  
191 subalpine soils were calculated with the *lm* function, included in the R software (R Development  
192 Core Team, 2000).

$$193 \text{ IS} = 0.5 * \text{Fe}_o + \text{Al}_o \quad (1)$$

$$194 \text{ IS}_{\text{ratio}} = \text{IS}_{(\text{B-BC})} / \text{IS}_{(\text{E-AE})} \quad (2)$$

195 As reported above, we could not calculate the soil carbon stock because no bulk density data have  
196 been measured. However, we believe that the changes in concentration of the different soil  
197 compounds can effectively show pedogenic trends in the studied soils, considering that the skeletal  
198 fraction resulted quite constant, particularly in the first few hundred years of soil development. The  
199 best variable transformation (logarithmic or power) was chosen according to the R<sup>2</sup> and the  
200 significance of the regression coefficients. The chronofunctions were only descriptive, as the  
201 sampling site number was excessively small to obtain statistically significant data. Moreover no  
202 data were available for the 260-11000 years BP time span, and the precise ages of pre-LIA S10 and  
203 A6 were not available. Significant differences in many edaphic parameters between different plant  
204 covers were also checked and displayed as boxplots, using the *multcomp* R package (Hothorn et al.,  
205 2008).

206

## 207 **3. Results**

208

### 209 **3.1. Vegetation primary succession**

#### 210 **3.1.1 Alpine primary succession**

211

212 Above timberline (Table 2, Figure 3), two different grassland communities were normally found  
213 outside the LIA glacial till, roughly corresponding to the lower alpine belt: flat, humid areas were

214 dominated by the *Carex curvula* association (*Caricetum curvulae*, Braun-Blanquet 1948), while  
215 steeper and drier sites were characterized by *Festuca varia* (*Festucetum variae*, Braun-Blanquet,  
216 1948); the observed pre-LIA site was colonized by a rather xerophilous community dominated by  
217 *Festuca varia* (A5, Table 2).  
218 Immediately after moraine stabilization (5-7 years after glacier retreat, P1), the pioneer species of  
219 the *Epilobietum fleischeri* (quite similar to the pioneer community described by Burga et al., 2010  
220 in the Morteratsch forefield) began the colonization of the raw till. Some of these species, often  
221 typical of base-rich soils (Pignatti 1992), were still present on 60 years old moraines (A2, A3). Mid  
222 successional species (Table 1) were common in 60-190 years old sites (A2, A3, A4); these species  
223 were characteristic of disturbed, rocky and eroded soils, only weakly acidified in the surface  
224 horizon. Species typical of later stages of succession appeared on 60 years old moraines (e.g., *Carex*  
225 *curvula*, *Festuca varia*, *Festuca halleri*), but became more common in later stages (A4, A5). The  
226 vegetation growing on 260 years old materials (A5) and the pre-LIA site (A6) was almost  
227 completely devoid of early and mid-successional species. The A5 site, in particular, was covered by  
228 a hygrophilous facies of the *Caricetum curvulae* (Braun-Blanquet, 1948), rich in dwarf Ericaceae  
229 such as *Loiseleuria procumbens* and *Vaccinium uliginosum* subsp. *gaulterioides*, probably because  
230 of microclimatic conditions favoring a long-lasting snow cover (north-west aspect).

231

### 232 3.1.2 Subalpine primary succession

233 Below timberline (i.e., subalpine primary succession, Table 3, Figure 3), the steep surfaces  
234 deglaciaded between ca. 1950 and 1987 (not sampled) were mostly colonized by *Salix* spp.,  
235 accompanied by *Rhododendron ferrugineum* and young and scattered larch trees (Treter et al.,  
236 2002). An extreme vegetation patchiness existed on surfaces deglaciaded between 1922 and 1950  
237 (excluding stony mounds, not considered in this study). Under an open *Larix decidua* forest,  
238 surfaces covered mostly by *Rhododendron ferrugineum* (S3, S4) were intermixed with more open,  
239 grazed, grass-dominated ones (SG2, SG5), which still included a few species characteristic of  
240 pioneer *Epilobietum fleischeri*. Mid-successional species, such as *Salix* spp., were locally common  
241 but never dominating. Species typical of subalpine forest-heath communities already showed high  
242 covering rate in grassland patches, such as *Avenella flexuosa* and *Calamagrostis villosa*. Other  
243 common species were typical of subalpine prairie (e.g., *Festuca varia*, *Nardus stricta*,  
244 *Anthoxanthum odoratum*).

245 Surfaces deglaciaded between 1860 and ca. 1922 (S6) were covered by a quasi-climax subalpine  
246 *Larix decidua* open forest, with *Rhododendron ferrugineum* dominating the understory, together  
247 with *Vaccinium* spp; grasses were already characteristic of a well-developed subalpine forest

248 (*Avenella flexuosa*, *Calamagrostis villosa*, *Homogyne alpina*). The easternmost part of this area has  
249 always been used for intensive cattle grazing and only scattered trees and shrubs were present  
250 (SG7). Here, species typical of subalpine acidophilous grazed grassland were common, such as  
251 *Nardus stricta*, *Festuca varia* and *Rhinanthus alectorolophus* (Table 3).

252 The 1821 moraine is steep and north-facing; a thick, long-lasting and creeping snow cover  
253 characterizes the microclimate of this site, which is colonized mainly by green alder (*Alnus viridis*),  
254 which partly inhibits high covering values by *Vaccinium myrtillus* and *Rhododendron ferrugineum*,  
255 which are only locally dominant (S9).

256 The older surfaces, originated from glacial till abandoned during late glacial maxima (around 10000  
257 years B.P., S10) were colonized by a climax open Larch forest, with an understory dominated by  
258 *Rhododendron ferrugineum* and *Vaccinium* spp. The specific composition and covering values were  
259 similar to those characterizing most of the stable forest sites in the area older than 90 years.

260 The plant cover was already around 100% in 65 years old soils, while above timberline it reached  
261 the same value after 90 years.

## 262 **3.2 Soil properties along the chronosequences**

263 Soil development trends were well correlated with the primary succession of the vegetation cover  
264 and land use; three different pathways were observed under alpine prairie, subalpine grassland and  
265 typical subalpine forest-heath communities (Figure 4).

266

### 267 **3.2.1. Alpine soil chronosequence**

268 Above timberline (Table 4, Figure 4a), organic matter accumulation and acidification in the soil  
269 surface horizons were the main pedogenetic processes in soils younger than 200 years. These  
270 processes led to the formation of A horizons, with a maximum thickness and organic carbon content  
271 in 260 years old soils (A5). The C/N ratio in upper alpine mineral soil horizons ranged between ca.  
272 11 (A2) and 17 (A5, influenced by Ericaceae), while during the first stages of soil development  
273 subsurface horizons showed much lower values. The difference between surface and subsurface  
274 horizons decreased with increasing soil age, suggesting a more efficient organic matter  
275 incorporation in the soil profile.

276 Weathering in subsurface horizons, releasing free Fe and Al oxi-hydroxides, led to the formation of  
277 weakly developed, brownish AB and BC horizons in 190 years old soils (A4). Younger soils had  
278 only A, AC and C horizons while a “true” brown, structured Bw appeared in 260 years (A5). Well-  
279 developed pre-LIA soils were characterized by thick and well developed brown Bw horizons with  
280 strongly acidic pH values, particularly in the A horizon.

281 Amorphous and crystalline Fe and Al oxi-hydroxides were weakly redistributed with depth  
282 (increasing contents in subsurface horizon associated with a depletion in surface ones) in the 260  
283 years old and in the “late glacial” soils (A5 and A6, Table 5).  
284 Initial soils (P1) had near-neutral pH values and high Base Status, thanks to the abundance of  
285 freshly ground, highly reactive primary minerals. Acidification and desaturation proceeded quite  
286 fast, particularly in surface A horizons, together with the strong increase in organic matter. Base  
287 status below 50% appeared after more than 65 years.  
288 According to the WRB soil classification (IUSS Working Group, 2006) the soils up to 65 years in  
289 the forefield (P1, A2, A3) were classified as Haplic Regosol (Eutric, Skeletic). After 190 years (A4),  
290 weakly developed but already acidified soils were classified as Haplic Regosol (Dystric, Skeletic).  
291 On the oldest LIA morainic arc and on late glacial materials (A5 and A6) Haplic Cambisol (Dystric,  
292 Skeletic) were found, thanks to the presence of well developed Bw horizons (Table 4).  
293 RhizoMODER humus forms characterized most of the soils older than 60 years, while A5, with an  
294 ericaceous cover, had a hemiMODER (Zanella et al., 2011).  
295

### 296 **3.2.2. Subalpine soil chronosequences**

297 Considering only stable sites below timberline, two different soil chronosequences were observed,  
298 associated with subalpine forest-shrub vegetation or with subalpine grazed grassland.  
299 Under subalpine grazed grassland (Figure 4b), soil processes and horizon formation were similar,  
300 but faster, to the ones observed above timberline. TOC accumulation in A horizons increased with  
301 age and pH values decreased to ~5.5 (SG7); in the meantime, weakly structured, brownish Bw  
302 horizons formed (in 130 years old prairie sites, SG7). Where the vegetation was dominated by  
303 grassland species (SG5, SG7) or by *Alnus viridis* (S8), the C/N values were below 14 (Figure 5a).  
304 Under grassland vegetation, the albic horizon did not form, and the Fe and Al oxi-hydroxides did  
305 not redistribute with depth.  
306 Under subalpine larch forest with Rhododendron understory (Figure 4c), great morphological and  
307 chemical changes characterized soils in the first 60-65 years since deglaciation (S3). Up to 6 cm of  
308 litter accumulated on the soil surface, below which 3-6 cm thick, dark grayish A horizons developed  
309 and weathering created yellowish BC ones (Table 6). These A horizons were characterized by the  
310 absence of structure and by the juxtaposition of mineral and organic particles, creating a “salt and  
311 pepper” appearance. pH values dropped from 6.5 to 5.0 in the upper horizons, while the C/N values  
312 were already close to 20. The C/N values were significantly higher than under other vegetation  
313 types (Figure 5a) while the pH values were lower (Figure 5b). Also the thickness of all O horizons  
314 was significantly higher under forest than under grassland (Figure 5c).

315 In the following 30 years (S4, on the 1921 moraine), the A horizons were substituted by thin, visibly  
316 bleached E horizons, characterized by a decrease in pedogenic Fe oxi-hydroxides (Table 7). A  
317 further decrease in pH values (4.7 in the E horizon) was measured in ~130 years old soils (S6) and  
318 the E horizons met the morphological requirements for the diagnostic albic horizons (Soil Survey  
319 Staff, 2010 and IUSS Working Group, 2006). Under these albic horizons, weak Bs horizons formed  
320 (here called CBs or BCs due to the lack of structure and the light yellowish colors; the  
321 macromorphological requirements for the spodic horizon were not met, according to Soil Survey  
322 Staff 2010, and IUSS Working Group, 2006). Fe redistribution was evidenced by yellowish colors  
323 and by Fe<sub>o</sub> and Fe<sub>d</sub> depth trends (Figure 6). The E and BCs horizons further developed in 190 years  
324 old soils (S8, S9), when the redistribution of Al (Al<sub>o</sub>) became measurable. Higher Fe<sub>o</sub>  
325 concentrations in subsurface horizons was verified by the ratio between Fe<sub>o</sub> concentrations in B-  
326 BC-CB and in surface E-AE-A horizons (Figure 5d), which was significantly higher under forest-  
327 shrub vegetation also in young soils. The same increasing concentration in subsurface horizons was  
328 observed for Fe<sub>d</sub> (Figure 5e). Al<sub>o</sub> was extremely low in all LIA soils, and its redistribution with  
329 depth was not significantly different under the considered vegetation types. Thus, the chemical  
330 requirements (TOC > 0.5%, IS > 0.5%, ISratio >= 2) for the diagnostic spodic horizons were not  
331 met. The ISratio resulted significantly higher under forest-shrub vegetation than under other land  
332 covers (Figure 5f).

333 Pre-LIA climax soils were Podzols with an extremely well developed morphology, both under forest  
334 and under anthropogenic grassland (not shown). These Podzols were characterized by a strong  
335 illuviation of organic matter and pedogenic Fe, Al and Si oxides in the spodic horizons. Deep  
336 cemented horizons (ortstein) were generally developed below the Bs (S10).

337 According to the WRB soil classification (IUSS Working Group, 2006), the soils under subalpine  
338 forest/shrubs on the LIA materials, and soils under grassland up to 90 years old, were classified as  
339 Haplic Regosol (Dystric, Skeletic). Subalpine “climax” soils were Ortsteinic Podzols (Skeletic)  
340 (S10). Dystric Cambisols were found on 130 years old surfaces under grazed prairie (SG7).

341 Under subalpine grazed grassland, rhizoMODER humus forms were identified, characterized by the  
342 presence of OF and, sometimes, OH horizons, overlying root-rich, single grain A ones. Under  
343 Rhododendron-larch forest, the humus form was dysMODER in 60 years old soils (S3). It quickly  
344 evolved towards MOR forms (hemiMOR in S4, S6 and S9, euMOR in S8); pre-LIA soils had  
345 humiMOR forms (Zanella et al. 2011).

346

## 347 4. Discussion

### 348 4.1 Vegetation primary successions along the chronosequences

349

350 Microtopography governs the establishment of pioneer species and their turnover in the first few  
351 hundred years during the primary succession (Burga et al., 2010). Considering only stable, fine  
352 earth-enriched sites (as in this study), the trends in the vegetation succession are more linear.  
353 As in other proglacial areas worldwide, the early stages of soil development and vegetation  
354 succession were similar above and below timberline, as almost the same plant species colonized the  
355 “young” morainic till, mostly belonging to the “*Epilobietum fleischeri*” (Braun Blanquet, 1948).  
356 Above the treeline, the development of soils and the vegetation succession was slightly faster than  
357 in other proglacial areas described in the Italian Alps (Andreis et al. 2001). The considered time  
358 span (260 years) was enough for the establishment of a quasi-climax vegetation (*Curvuletum* or  
359 *Festucetum variae*) while 500 years is the minimum time normally required for the establishment of  
360 the climax *Curvuletum* in the Alps (Andreis et al., 2001).

361 The primary plant succession above timberline was strongly influenced by topography and  
362 microclimate: for example, *Festuca varia* was common on 60 and 190 years old moraines (A3 and  
363 A4) and in the climax, pre-LIA site (A6), while it was absent from the cooler and more humid 260  
364 years old moraine (A5). It seems thus plausible that the further development of the vegetation cover  
365 in the A3 and A4 sites will lead to the development of *Festucetum variae*, similar to “climax” A6,  
366 and not towards a *Curvuletum*, similar to the one growing on the A5 site.

367 The similarity between primary vegetation successions in areas above and below timberline ends  
368 soon, and already after about 60 years, larch trees (*Larix decidua*) have largely invaded the areas at  
369 lower elevations, accompanied by ericaceous shrubs (e.g. *Vaccinium* spp., *Calluna vulgaris* and  
370 *Rhododendron ferrugineum*) (similar to the situation described in Alaska by Boggs et al., 2010).

371 The establishment of a *Rhododendron* understory, accompanied by *Salix helvetica* and *Salix*  
372 *appendiculata*, was observed earlier than in other proglacial areas in the Alps (e.g., Bernasconi et  
373 al., 2011). A stabilization of the species composition below timberline was observed after the  
374 establishment of ericaceous shrubs, particularly *Rhododendron ferrugineum*, on ca. 90 years old  
375 surfaces. *Rhododendron ferrugineum* is the main limiting factor to the survival of pioneer species,  
376 because of soil shadowing and because of the thick litter layer created by this species (Pornon and  
377 Doche, 1996). A quasi-climax vegetation structure and composition was generally reached after  
378 100-120 years: a much faster vegetation succession is thus observed in this work, if compared to  
379 other well studied chronosequences on the Alps (e.g., Burga et al., 2010, Dümig et al., 2011).

380 The early establishment of a quasi-climax vegetation below the timberline differs from what  
381 reported by Burga (1999), Burga et al. (2010) and Föllmi et al. (2009) in the Morteratsch and the  
382 Rhöne proglacial areas respectively, where the most common shrub was Green alder (*Alnus viridis*),  
383 with associated “megaphorbiae”, while Ericaceae (*Rhododendron ferrugineum* and *Vaccinium* spp.)  
384 never reached covering values higher than 5%. The dominance by *Rhododendron ferrugineum* in  
385 the Lys proglacial area was observed in sites stabilized before 1945 by Treter et al. (2002), who  
386 associated this dominance to light grazing. The ecosystem mosaic, related with microtopography,  
387 was particularly visible in the 1921-1945 time span, where stable sites covered by a larch-  
388 *Rhododendron* forest were mixed with grazed open grassland patches and stony mounds; quasi-  
389 climax subalpine shrubs (*Rhododendron ferrugineum* in particular) colonized most of the older  
390 surfaces, except where grazing pressure was higher. Stony mounds (not considered in this study)  
391 were common in the area younger than 91 years, and were colonized by pioneer species (Treter et  
392 al., 2002).

393 Grazed sites, devoid of ericaceous shrubs and Larch trees, were colonized by different herbaceous  
394 species compared to alpine grasslands, and the oldest sites had a species assemblage close to the  
395 typical subalpine acidophilous *Nardus stricta* prairie. Grazing locally inhibits shrub and tree growth  
396 also on “old” surfaces (SG7), but a low-pressure grazing influence on the abundance of *Ericaceae*  
397 and scarcity of *Alnus viridis* and associated species can be hypothesized.

398

#### 399 **4.2 Soil properties along the chronosequences**

400 Together with the primary vegetation succession, the speed of diagnostic horizon development and  
401 taxonomic reference group change (IUSS Working Group, 2006) was faster below timberline than  
402 at higher elevation. Under subalpine grazed grassland, in fact, a Haplic Cambisol (Dystric) formed  
403 in 130 years (SG7), while above timberline the same taxonomic level was reached in 260 years  
404 (A5). 60 years old soils below subalpine grassland were morphologically similarly developed as  
405 190 years old ones above timberline (A4).

406 This higher soil development rate was probably caused by the higher productivity of ecosystems  
407 below timberline, thanks to less harsh climatic conditions and by hypothetically weaker  
408 cryoturbation phenomena. However, cryogenic features were not significantly more developed  
409 above than below timberline, as demonstrated by the ubiquitous presence of thin and weak silt caps  
410 (table 1, 3) in shallow subsurface horizons particularly in young soils (process well described by  
411 Forman and Miller, 1994). Above timberline, silt caps were best developed in 60 years old soils,  
412 and became less visible with increasing ages, until they disappeared in 260 years old soils. Below  
413 timberline, silt caps were not evident under forest/shrub vegetation, while thin silt caps were visible

414 in shallow subsurface horizons in 60 and 90 years old soils and only in deeper C horizons of 130  
415 years old soils under grazed grassland. Here they were not visible in soils older than 130 years. Silt  
416 caps disappearance was probably related to the bioturbation caused by roots, which was only  
417 slightly faster below than above timberline.

418 A higher ecosystem productivity below than above timberline was verified by the higher TOC  
419 concentration in the fine earth of upper mineral horizons in the 130 years old soil (SG7, table 4, 6);  
420 however, its trend with time was disturbed by a large variability. Above timberline, the rate of TOC  
421 concentration increase declined from  $0.28 \text{ g*kg}^{-1}\text{y}^{-1}$  in 70 years old soils to  $0.16 \text{ g*kg}^{-1}\text{y}^{-1}$  on 260  
422 years old moraines; at this point, the balance between organic carbon inputs and loss via  
423 decomposition in surface horizons (steady state) was reached. Subsurface accumulation continued,  
424 in quantity and depth, as shown by the increase in TOC concentration and in thickness of Bw and  
425 BC-CB horizons (Table 6). Overall, under grassland the TOC concentration in surface horizons  
426 tended to increase for the first few hundreds years until reaching a steady state between inputs (via  
427 root and litter decay) and outputs (mineralization, erosion and leaching), according to the indicative  
428 chronofunction (Figure 7a,  $R^2 = 0.284$ , p-value < 0.05 for all coefficients):

$$429 \quad \text{TOC} = -0.287 + 0.616 * \ln(\text{age}) \quad (3)$$

430  
431  
432  
433 Lower TOC concentrations characterized surface mineral horizons in forest-shrub soils (Figure 7b),  
434 which also showed a completely different TOC variation with time (Figure 7a). The highest TOC  
435 concentration was reached in 60 years old soils. After this time, TOC concentrations started to  
436 decrease, according to the descriptive chronofunction ( $R^2 = 0.201$ , p-value < 0.1):

$$437 \quad \text{TOC} = 1.723 - 0.09 * \ln(\text{age}) \quad (4)$$

438 The slowdown of the organic matter accumulation started quite early compared to other  
439 chronosequences (e.g. Dümig et al., 2011), because of the quick establishment of the podzolization  
440 process under the subalpine vegetation (He and Tang, 2008) and the development of an albic  
441 horizon below the litter layer. This trend was not observed above the treeline or under subalpine  
442 grazed grassland, because of limited vertical migration of soluble organic molecules and of a lower  
443 organic matter production caused by the less productive vegetation. The increasing trend of TOC  
444 concentration in the subsurface horizons of forest soils was faster than in the grassland ones (Figure  
445 7c). Forest soils had higher TOC concentrations in subsurface horizons (Figure 7d) than grassland  
446 soils, probably thanks to a vertical migration of soluble organic compounds (possibly associated to  
447 a higher organic matter produced by root decay).

448 The C/N ratio was significantly related to the vegetation cover, with the highest values observed in  
449 the organic horizons composed of ericaceous shrubs and Larch leaves (e.g. Boettner and Kalisz,  
450 1990). Below the forest-shrub vegetation, the lowest C/N value was found under an understory



451 vegetation dominated by the N-fixing *Alnus viridis* (S8). Where the alpine prairie was enriched in  
452 dwarf Ericaceae, such as *Loiseleuria procumbens* and *Vaccinium uliginosum* subsp. *gaulterioides*  
453 (A5), the C/N ratio was higher. This ratio is characteristic of slowly decomposing organic matter,  
454 rich in fulvic and low-molecular weight acids, and is associated to low pH values. Indeed the fastest  
455 and strongest pH decrease was observed under subalpine forest/shrub vegetation, where values  
456 below 5 characterized the surface mineral horizon already after 60 years since deglaciation. Under  
457 grazed grassland and alpine prairie the pH reached comparable values after 130 and 260 years,  
458 respectively. Since the leaching of organic acids derived from organic matter degradation is the  
459 main acidifying factor in proglacial soils (Bernasconi et al. 2011), the slower pH decrease under  
460 alpine vegetation could be attributed to the lower biomass production and the stronger disturbances  
461 characterizing alpine habitats. However, a weak acidification was visible also in the youngest soils,  
462 extremely poor in organic matter, thanks to incipient mineral weathering and leaching caused by  
463 rain and snowmelt.

464

#### 465 **4.3 Evidences of podzolization processes**

466 Both Fe<sub>o</sub> and Al<sub>o</sub> in the initial soil (P1) were higher in the top C1 horizon than in the underlying  
467 C2. This distribution could be attributed to an initial mineral weathering caused by the organic  
468 matter accumulation produced by the few plants which colonize these sites, and by oxidative  
469 reactions involving fresh, reactive particles. The already detectable Fe and Al oxi-hydroxides in the  
470 fresh material of the C2 horizons in the 6 years old soils could be attributed to subglacial  
471 weathering processes, a well-known source of already weathered material to initial soil formation  
472 processes (Anderson et al., 2000 and Dümig et al., 2011).

473 This depth trend was soon reversed: an early redistribution of pedogenic Fe oxi-hydroxides to  
474 subsurface horizons was observed under subalpine larch forest with *Rhododendron* understory.  
475 Higher values of Fe<sub>o</sub> and Fe<sub>d</sub> in subsurface BC horizons compared to the overlying A ones were  
476 already observed in the ca. 60 years old soils. Despite the weaker Al<sub>o</sub> redistribution with depth in  
477 young soils, the ISratio tended to increase with time, according to the (qualitative) chronofunction  
478 ( $R^2 = 0.801$ , p-value < 0.01 for the age coefficient, Figure 8a):

$$479 \text{ISratio} = 0.676 * \text{age}^{0.185} \quad (5).$$

480 Thus, around 350 years are needed in order to meet the ISratio requirements for the diagnostic  
481 spodic horizon according to Soil Survey Staff (2010) and IUSS Working Group (2006).

482 Observing the IS and Fe<sub>d</sub> depth trends (Figure 5 and Figure 6), the cheluviation seems to have  
483 actively translocated Fe and Al oxi-hydroxides into subsurface BC and CB horizons. The IS, in

484 subsurface horizons of soils under subalpine forest-shrubs (*Rhododendron* and *Larch*), increased  
485 according to the chronofunction (Figure 8b):

$$486 \text{ IS} = 0.045 + 0.003 * \text{age}^{0.8} \quad (6)$$

487 therefore, the time needed was ca. 530 years.

488 Based on the few data available, all the regressions between time since deglaciation and the spodic  
489 properties had similar results: the time required to obtain a “chemical” Podzol, according to the Soil  
490 Survey Staff (2010) and IUSS Working Group (2006) is around 300-600 years, which is a much  
491 shorter period than the 3000 years previously calculated on the Alps (Egli et al. 2001).

492 However, these findings apply only to sites located below the timberline and covered by larch forest  
493 with ericaceous understory. Below subalpine grazed grassland and below alpine prairie, Fe and Al  
494 oxyhydroxides were more abundant in surface A horizons compared to subsurface BC or Bw  
495 horizons. An exception was the 60 years old grassland soil (SG3), influenced by nearby larch trees,  
496 and A5, thanks to abundant ericaceous dwarf shrubs (*Vaccinium uliginosum* subsp. *gaulterioides*  
497 and *Loiseleuria procumbens*).

498 Similarly to other chronosequences (e.g., Burt and Alexander 1996, Egli et al. 2001), the slower  
499 release and mobility of Al from the parent material, compared with Fe, could be related to the  
500 presence of easily weatherable Fe-rich minerals included in the small amounts of mafic rocks in the  
501 parent material. The silic minerals weathered more slowly in the considered soils, as demonstrated  
502 by the  $\text{Al}_0/\text{Fe}_0$  ratio increase with time, both in surface (not shown) and in subsurface horizons  
503 (Table 7). Conversely, in more acidic parent materials, the onset of the podzolization process tended  
504 to remove Al faster than Fe from the young and weakly developed E horizons (Mokma et al., 2004).  
505 The E horizon formation in alpine soils has sometimes been interpreted as a stagnogley feature,  
506 caused by seasonal waterlogging and alternation of reductive and oxidative conditions, common in  
507 high altitude soils thanks to the abundant snow cover and high water input during the snowmelt  
508 (e.g. Gensac, 1990). However, the E horizon formation under subalpine vegetation in the Lys  
509 proglacial area cannot be ascribed specifically to seasonal waterlogging, as stronger seasonal  
510 reductive conditions should be met above timberline, thanks to deeper winter and spring snow  
511 covers. Here, E and CBs/Bs horizons were never found, and Fe/Al redistribution to subsurface  
512 horizons normally was not observed. Moreover the coarse texture of the soil material should  
513 facilitate drainage, decreasing the waterlogging potential in the topsoil.

514 The  $\text{Fe}_0/\text{Fe}_d$  ratio, called “activity ratio” and considered as a measure of the crystallinity of  
515 pedogenic Fe oxides (Burt and Alexander, 1996), did not show any significant time trends but had  
516 significantly lower values under subalpine forest-shrub vegetation than under grassland (Figure 9),  
517 particularly in subsurface horizons. Alternation of reductive and oxidative conditions are known to

518 favor the crystallization of pedogenic Fe-oxi-hydroxides; however, the lower values found under  
519 subalpine vegetation, particularly in subsurface BC and Bs horizons, are not explainable according  
520 to surface seasonal redox conditions, for the same reason explained above. Nor the higher  
521 crystallinity found in soils under forest vegetation, compared to the soils developed under even  
522 more humid climates on the Alps (Dümig et al., 2011), could be easily explainable. Low Feo/Fed  
523 values, however, are typical of many well developed podzolic soils in the Valle d’Aosta Region  
524 (e.g. D’Amico et al., 2008).

525

526 On the Alps, Podzols are known to develop from silica-rich glacial till in late stages of pedogenesis  
527 (3300 and 10000 years), as reported by Egli et al. (2001). Faster rates of E horizon formation have  
528 often been measured in other boreal or mountain areas (Sauer et al., 2007). Faster Podzol  
529 development in Alaska (Alexander and Burt, 1996, Ugolini, 1966), Norway (Mellor, 1986) or China  
530 (He and Tang, 2008) compared to the European Alps was attributed to the maritime climate (Egli et  
531 al., 2006, Dümig et al., 2011), which accelerates plant growth, soil organic matter accumulation  
532 and, consequently, the rate of soil development, thanks to longer growing season and much higher  
533 winter temperature. However, the establishment of mixed coniferous forests caused an immediate  
534 onset of the podzolization process also in continental climates (ca. 225 years, Lichter, 1998). An  
535 even faster redistribution of Fe oxi-hydroxides was observed on sandy parent materials under boreal  
536 Scots pine in north-western Russia (Abakumov et al., 2010), where higher Feo contents in BCs than  
537 in AE horizons were measured in 10 years old soils, and E horizons appeared in 20 years; in fact,  
538 sandy parent materials are known to increase the speed of the podzolization process thanks to the  
539 already weathered grain surfaces (Schaetzl and Anderson, 2005) and higher water mobility that  
540 promotes vertical fluxes of elements.

541 Thus, on a global scale, the “fast” podzolization process characterizing subalpine soils developed on  
542 the gneissic materials of our study area is in the “normal” ranges, but much faster than in other  
543 alpine chronosequences. Similarly developed soils have been observed on the LIA morainic arcs of  
544 the Miage glacier, Val Veny, in the Mont Blanc Massif and in the Verra Grande forefield, Monte  
545 Rosa Group (D’Amico 2011, unpublished data). These proglacial areas in Valle d’Aosta are  
546 characterized by temperature and precipitation regimes similar to other proglacial areas, in  
547 particular in the northern parts of the Alps, where the podzolization processes start later and seem  
548 slower than in our study area. Thus, climatic differences cannot be the cause of such a slower  
549 pedogenesis on the northern side of the Alps, while the early establishment of ericaceous shrubs  
550 below the treeline appeared the main driver of the early appearance of the bleached E horizon in 90  
551 years old soils.

552 Plant communities on Swiss glacier forefields were dominated by alder (*Alnus viridis*), normally  
553 associated with N-fixing bacterial communities (Egli et al., 2001, Burga et al., 2011), and by  
554 herbaceous plant species, while *Rhododendron ferrugineum* appeared in later stages of succession  
555 (Dümig et al., 2011). This vegetation creates a litter which has a weak complexing capacity, while  
556 the ericaceous shrubs are able to begin a quick podzolization process in previously non-podzolic  
557 soils (Bernier and Gillet 2012, 1993, Boettcher and Kalisz, 1990). This is caused by the slow  
558 decomposition rates of the litter of ericaceous shrubs, due to their high amount of lignin, cellulose  
559 and other recalcitrant substances, such as phenolic compounds, which reduce the soil biological  
560 activity (Pornon and Doche, 1995). The litter of coniferous trees and Ericaceae produces large  
561 quantities of low molecular weight and fulvic acids, which cause intense mineral weathering  
562 (Schaetzl and Anderson, 2005). As humus forms are considered a good indicator of forest  
563 ecosystem functioning (Michalet et al., 2001), the fast development of Mor humus forms under  
564 subalpine typical *Rhododendron*-larch vegetation confirms the slow mineralization of the soil  
565 organic matter, typically associated with the onset of podzolization.

566 The strong vegetation effect on pedogenesis could be enhanced by mycorrhizal fungi associated  
567 with different plant species. In fact, ectomycorrhizal and ericoid (associated to Ericaceae) fungi are  
568 known to increase the weathering rate in surface mineral horizons, particularly under coniferous or  
569 ericaceous species, where they form mat-like structure at the boundary between the organic layer  
570 and the upper mineral horizon, and extending down to the E horizons (Koele et al., 2011). These  
571 fungi are able to dissolve mineral grains, extracting and chelating metals and nutrients via the  
572 release of phenolic compounds, low weight organic acids, oxalate, citrate and malate (Landeweert  
573 et al., 2001), which have a stronger acidifying and weathering capacity than humic molecules  
574 (Ochs, 1996). Ericoid mycorrhizal fungi, in particular, produce siderophores molecules, able to  
575 efficiently extract and bind Fe and other metals from primary metals (Hoffland et al., 2004).  
576 Substances produced and released by ectomycorrhizal fungi, thus, increase the podzolization rate  
577 under coniferous trees (Lundström et al., 2000, van Breemen et al., 2000), and even more under  
578 Ericaceae, thanks to siderophore substances. Hence, also this process may contribute to the faster  
579 podzolization rates found in this study.

580

## 581 **5. Conclusions**

582

583 In this study we characterize the main pedogenetic processes occurring in recently deglaciated areas  
584 under different vegetation covers. We furthermore provide evidence of fast rates of podzolization

585 under a subalpine larch-Rhododendron forest, previously undocumented for the Alps: bleached E  
586 horizons are visible in 60 years old soils, and diagnostic albic horizons appear in 120 years. In this  
587 time span the cheluviation of organo-metal compounds in the underlying BC horizons was not  
588 sufficient to create diagnostic spodic features, but 300-500 years seemed enough for the formation  
589 of a diagnostic spodic Bs horizon and, consequently, of a “real” Podzol.  
590 We suggest this relatively fast rate of podzolization be due to the specific plant community  
591 succession rather than to the climatic conditions in the study area, characterized by cold  
592 temperatures and significant but not exceptional precipitation amounts. The appearance of a larch-  
593 Rhododendron forest could significantly influence the soil organic matter characteristics, driving  
594 the soil development and the onset of soil forming processes in this proglacial area.

595

596

### 597 **Acknowledgments**

598 This study was accomplished thanks to the agreement with the “Centro Funzionale” of the Aosta  
599 Valley Region within the project “Optimization of the regional alert system regarding  
600 hydrogeological hazard, towards citizen protection: soil vulnerability map, 1:100000”, and thanks  
601 to the Italian MIUR Project (PRIN 2010-11): “Response of morphoclimatic system dynamics to  
602 global changes and related geomorphological hazards” (national coordinator C. Baroni). We also  
603 thank three anonymous reviewers for the great suggestions on how to improve the quality of the  
604 paper.

605

### 606 **References**

607

608 Abakumov, E., Trubetskoj, O., Demin, D., Celi, L., Cerli, C., Trubetzksaya, O., 2010. Humic acid  
609 characteristics in a Podzol soil chronosequence. *Chem. Ecol.* 26:S2, 59-66

610

611 Alexander, E.B., and Burt, R., 1996. Soil development on moraines of the Mendenhall Glacier,  
612 southeast Alaska. 1. The moraines and soil morphology. *Geoderma* 72, 1-17

613

614 Anderson, S.P., Drever, J.I., Frost, C.D., Holden P., 2000. Chemical weathering in the foreland of a  
615 retreating glacier. *Geochim. Cosmochim. Acta* 64, 1173-1189.

616

617 Andreis, C., Caccianiga, M., Cerabolini, B., 2001. Vegetation and environmental factors during  
618 primary succession on glacier forelands: some outlines from the Italian Alps. *Plant Biosystems*  
619 135(3), 295-310.

620

621 Bernasconi, S.M., Bauder, A., Bourdon, A., Brunner, I., Bünemann, E., Christl, I., Derungs, N.,  
622 Edwards, P., Farinotti, D., Frey, B., Frossard, E., Furrer, G., Gierga, M., Göransson, H., Gülland, K.,  
623 Hagedorn, F., Hajdas, I., Hindshaw, R., Ivy-Ochs, S., Jansa, J., Kiczka, M., Kretschmar, R.,  
624 Lemarchand, E., Luster, J., Magnusson, J., Mitchell, E.A.D., Venterink, H., O., Plötze, M.,

625 Reynolds, B., Smittenberg, R., H., Stähli, M., Tamburini F., Tipper, E.T., Wacker, L., Welc, M.,  
626 Wiederhold, J.G., Zeyer, J., Zimmermann, S., and Zumsteg, A., 2011. Chemical and biological  
627 gradients along the Damma Glacier soil chronosequence, Switzerland. *Vadose Zone J.* 10, 867-883.  
628

629 Bernier, N., Gillet, F., 2012. Structural relationships among vegetation, soil fauna and humus form  
630 in a subalpine forest ecosystem: a Hierarchical Multiple Factor Analysis (HMFA). *Pedobiologia* 55,  
631 321-334  
632

633 Boettcher, S.E., Kalisz, P.J., 1990. Single-Tree Influence on Soil Properties in the Mountains of  
634 Eastern Kentucky. *Ecology* 71 (4), 1365-1372  
635

636 Boggs, K., Klein, S.C., Grunblatt, J., Boucher, T., Koltun, B., Sturdy, M., Streveler, G.P., 2010.  
637 Alpine and subalpine vegetation chronosequences following deglaciation in coastal Alaska. *Arctic,*  
638 *Antarctic and Alpine Res.* 42(4), 385-395.  
639

640 Braun-Blanquet, J., 1948. Übersicht der pflanzengesellschaften rhaetiens (IV). *Vegetatio* 2(1), 20-  
641 37.  
642

643 Burga, C., 1999. Vegetation development on the glacier forefield Morteratsch (Switzerland).  
644 *Applied Veg. Sci.* 2, 17-24.  
645

646 Burga, C.A., Krüsi, B., Egli, M., Wernli, M., Elsener, S., Ziefle, M., Fischer, T., Mavris, C., 2010.  
647 Plant succession and soil development on the foreland of the Morteratsch glacier (Pontresina,  
648 Switzerland): straight forward or chaotic? *Flora* 205, 561-576.  
649

650 Burt, R., and Alexander, E. B., 1996. Soil development on moraines of Mendenhall Glacier,  
651 southeast Alaska. 2. Chemical transformations and soil micromorphology. *Geoderma* 72,19-36.  
652

653 Caccianiga, M., Luzzaro, A., Pierce, S., Ceriani, R. M, Cerabolini, B., 2006. The functional basis of  
654 a primary succession resolved by CSR classification. *Oikos* 112, 10-20  
655

656 Dümig, A., Smittenberg, R., Kögel-Knaber, I., 2011. Concurrent evolution of organic and mineral  
657 components after retreat of the Damma glacier, Switzerland. *Geoderma* 163, 83-94  
658

659 Egli, M., Fitze, P., Mirabella, A., 2001. Weathering and evolution of soils formed on granitic, glacial  
660 deposits: results from chronosequences of Swiss alpine environments. *Catena* 45, 19-47.  
661

662 Egli, M., Mirabella, A, Fitze, P., 2003. Formation rates of smectite derived from two Holocene  
663 chronosequences in the Swiss Alps. *Geoderma* 117, 81-98  
664

665 Egli, M., Wernli, M., Kneisel, C., Haeberli, W., 2006. Melting glaciers and soil development in the  
666 proglacial area Morteratsch (Swiss Alps): I. Soil type chronosequences. *Arctic, Antarctic and Alpine*  
667 *Res.* 38(4), 499-509  
668

669 FAO, 2006. Guidelines for Soil Description. 4th ed. FAO, Rome, 97 pp.  
670

671 Föllmi, K.B., Arn, K., Hosein, R., Adatte, T., Steinmann, 2009. Biogeochemical weathering in  
672 sedimentary chronosequences of the Rhone and Oberaar glaciers (Swiss Alps): rates and mechanisms  
673 of biotite weathering. *Geoderma* 151, 270-281  
674

675 Forman, S.L., Miller, G.H., 1984. Time-dependent soil morphologies and pedogenic processes on  
676 raised beaches, Bröggerhalvöya, Spitsbergen, Svalbard archipelago. *Arctic and Alpine Res.* 16, 4,  
677 381-394  
678

679 He, L., Tang, Y., 2008. Soil development along primary succession sequences on moraines of  
680 Hailuoguo Glacier, Gongga Mountain, Sichuan, China. *Catena* 72, 259-269  
681

682 Hoffland, E., Kuiper, T.W., Wallander, H., Passard, C., Gorbushina, A.A., Haselwandter, K.,  
683 Holmström, S., Landeweert, R., Lundström, U.S., Rosling, A., Sen, R., Smits, M.m., van Hees,  
684 P.A.W., van Breemen, N., 2004. The role of fungi in weathering. *Frontiers in Ecology and the*  
685 *Environment* 2, 5, 258-264  
686

687 Hodkinson, I.D., Coulson, S.J., Webb, N.R., 2003. Community assembly along proglacial  
688 chronosequences in the high Arctic: vegetation and soil development in north-west Svalbard. *J.*  
689 *Ecol.*, 91, 651-663  
690

691 Hothorn, T., Bretz, F., and Westfall, P. (2008). Simultaneous Inference in General Parametric  
692 Models. *Biometr. J.* 50(3), 346—363.  
693

694 Huggett, R.J., 1998. Soil chronosequences, soil development, and soil evolution: a critical review.  
695 *Catena* 32, 155-172  
696

697 IUSS Working Group WRB, 2006. World reference base for soil resources 2006. World Soil  
698 Resources Reports No. 103. FAO, Rome.  
699

700 Ivy-Ochs, S., Kerschner, H., Maisch, M., Christl, M., Kubik, P.W., and Schluechter, C., 2009. Latest  
701 Pleistocene and Holocene glacier variations in the European Alps: *Quat. Sci. Rev.* 28, 2137–2149  
702

703 Jeanroy, E., Guillet, B., Ortiz, R., 1984. Applications pédogénétiques de l'étude des forms du fer par  
704 les réactifs d'extraction: cas des sols brunifiés et podzolisés sur roches cristallines. *Sci. Sol* 3, 199–  
705 211.  
706

707 Jenny, H., 1958. Role of the plant factor in the pedogenic function. *Ecology* 39 (1), 5-46  
708

709 Joerin, U.E., Stocker, T.F., Schlüchter, C. 2006. Multicentury glacier fluctuations in the Swiss Alps  
710 during the Holocene. *The Holocene* 16 (5), 697-704  
711

712 Koele, N., Storch, F., Hildebrand, E.E., 2011. The coarse-soil fraction is the main living space of  
713 fungal hyphae in the BhBs horizon of a Podzol. *J. Plant Nutr. Soil Sci.* 174, 750–753  
714

715 Landeweert, R., Hoffland, E., Finlay, R.D., Kuiper, T.W., van Breemen, N, 2001. Linking plants to  
716 rocks: ectomycorrhizal fungi mobilize nutrients from minerals. *Trends in Ecol. & Evol.* 16, 5, 248-  
717 254  
718

719 Landolt, E., (1977): *Ökologische Zeigerwerte zur Schweizer Flora*. Veröffentlichungen des  
720 Geobotanischen Institutes der Eidgenössischen Technischen Hochschule, Stiftung Rübel, Zürich 64:  
721 1–208.  
722

723 Lichter, J., 1998. Rates of weathering and chemical depletion in soils across a chronosequence of  
724 Lake Michigan sand. *Geoderma* 85, 255-282  
725

726 Lundström, U.S., van Breemen, N., Bain, D.C., van Hees, P.A.W., Giesler, R., Gustafsson, J.P.,  
727 Ilvesniemi, H., Karlton, E., Melkerud, P.A., Olsson, M., Riise, G., Wahlberg, O., Bergelin, A.,  
728 Bishop, K., Finlay, R., Jongmans, A.G., Magnusson, T., Mannerkoski, H., Nordgren, A., Nyberg, I.,  
729 Starr, M., Tau Strand L., 2000. Advances in understanding the podzolization process resulting  
730 from a multidisciplinary study of three coniferous forest soils in the Nordic Countries. *Geoderma*  
731 94, 335-353  
732  
733 Mahaney, W.C., Kalm, V., Milner M.W., Hancock, R.G.V., 2009. A soil chronosequence in late  
734 glacial and neoglacial moraines, Humboldt Glacier, northwestern Venezuela Andes.  
735 *Geomorphology* 109, 236-245.  
736  
737 Mattiolo, E., Novarese, V., Franchi, S., Stella, A., 1951. Carta Geologica d'Italia 1:100000, foglio  
738 29. Istituto Geografico Militare (Firenze, Italy)  
739  
740 Mellor, A., 1987. A pedogenic investigation of some soil chronosequences on neoglacial moraine  
741 ridges, Southern Norway: examination of soil chemical data using principal component analysis.  
742 *Catena* 14, 369-381  
743  
744 Mercalli L., 2003. Atlante climatico della Valle d'Aosta. Società Meteorologica Italiana (Ed),  
745 Torino, p. 405.  
746  
747 Michalet, R., Gandoy, C., Cadel, G., Girard, G., Grossi, J., Joud, D., Pache, G., 2001. Modes de  
748 fonctionnement d'humus des forêts sempervirentes des Alpes internes françaises. *Humus*  
749 functioning types in evergreen coniferous forests of the French Inner Alps. 324 (1), 59-70  
750  
751 Ministero delle Politiche Agricole e Forestali, 2000. Metodi di Analisi Chimica dei Suoli. In:  
752 Violante P, Sequi P (eds), *Collana di Metodi Analitici per l'Agricoltura*. Franco Angeli (Ed.),  
753 Milano, pp 1-474  
754  
755 Mokma, D.L., Yli-Halla, M., Lindqvist, K., 2004. Podzol formation in sandy soils of Finland.  
756 *Geoderma* 120, 259-272  
757  
758 Monterin U., 1932. Le variazioni secolari del clima del Gran S. Bernardo: 1918-1931 e le  
759 oscillazioni del ghiacciaio del Lys al Monte Rosa: 1789-1931. *Boll. Com. Glac. It., ser. I, 12.* 59-  
760 189  
761  
762 Ochs M., 1996. Influence of humified and non-humified natural organic compounds on mineral  
763 dissolution. *Chem. Geol.* 132, 119-124  
764  
765 Pelfini, M., Belloni, S., Rossi, G., Struma, G., 1997. *Geogr. Fis. Din. Quat.* 20, 329-338  
766  
767 Pignatti, S., 1992. *Flora d'Italia*, vol 1-3. Edagricole, Bologna  
768  
769 Pornon, A., Doche, B., 1996. Age Structure and Dynamics of *Rhododendron ferrugineum* L.  
770 populations in theNorthwestern French Alps. *J. Veg. Sci.* 7(2), 265-272  
771  
772 Raffl, C., Mallaun, M., Mayer, R., Erschbamer, B., 2006. Vegetation succession pattern and  
773 diversity changes in a glacier valley, central Alps, Austria. *Arctic, Antarctic and Alpine Res.* 38(3),  
774 421-428.  
775



776 Righi, D., Huber, K., Keller, C., 1999. Clay formation and Podzol development from postglacial  
777 moraines in Switzerland. *Clay Min.* 34, 319-322  
778

779 Sauer, D., Sponagel, H., Sommer, M., Giani, L., Jahn, R., Stahr, K., 2007. Podzol: soil of the year  
780 2007; a review of its genesis, occurrence, and functions. *J. Plant Nutr. Soil Sci.* 170, 581-597  
781

782 Schaetzl, R., Anderson, S., 2005, *Soils. Genesis and geomorphology*. Cambridge University Press  
783 (Ed), Cambridge (UK)  
784

785 Soil Survey Staff, 2010. *Keys to Soil Taxonomy*. United States Department of Agriculture, Natural  
786 Resources Conservation Service, Eleventh Edition, 2010  
787

788 Strada, E., 1988. Le variazioni del ghiacciaio del Lys dalla “Piccola Glaciazione” ai nostri giorni.  
789 *Natura bresciana, Ann. Mus. Civ. Sc. Nat.* 24: 275-188  
790

791 Tonkin, P.J., Basher, L.R., 2001. Soil chronosequences in subalpine superhumid Cropp Basin,  
792 western Southern Alps, New Zealand. *New Zealand J. Geol. Geophys.* 44 (1), 37-45  
793

794 Treter, U., Ramsbeck-Ullmann, M., Böhmer, H.J., Bösche, H., 2002. Vegetationsdynamic im  
795 vorfeld des Lys-gletschers (Valle di Gressoney/Region Aosta/Italien). *Erdkunde* 56, 253-267  
796

797 Ugolini, F.C. 1966. Part 3. Soils. In: A. Mirskey, (Ed.), *Soil development and ecological succession*  
798 *in a deglaciated area of Muir Inlet, southeast Alaska*. Institute of Polar Studies report Number 20,  
799 Ohio State University, Columbus, Ohio, USA.  
800

801 Ugolini FC, Corti G, Certini G. (2006). Pedogenesis in the sorted patterned ground of Devon  
802 Plateau, Devon Island, Nunavut, Canada. *Geoderma* 136, 87-106  
803

804 Van Breemen, N., Lundström, U.S., Jongmans, A.G., 2000. Do plants drive podzolization via rock-  
805 eating mycorrhizal fungi? *Geoderma* 94. 163–171  
806

807 Zanella A, Jabiol B, Ponge JF, Sartori G, De Waal R, Van Delft B, Graefe U, Cools N, Katzensteiner  
808 K, Hager H, Englisch M, Brethes A, Broll G, Gobatl JM, Brun JJ, Milbert G, Kolb E, Wolf U,  
809 Frizzera L, Galvan P, Kolli R, Baritz R, Kemmerse R, Vacca A, Serra G, Banas D, Garlato A,  
810 Chersich S, Klimo E, Langohr R (2011a) *European Humus Forms Reference Base*.  
811

812 Zech, W., Wilcke, B.M., 1977. Vorlauge ergebnisse einer Bodenchronosequenzstudie im Zillertal.  
813 *Mitteilungen der Deutschen Bodenkundlichen Gesellschaft* 25, 571-586  
814

815 **Figure captions**

816 Fig. 1: the Lys proglacial area in the Aosta Valley Region, North-western Italy. The sampling sites  
817 and the main phases of glacial retreat since 1821 are also shown.

818  
819 Fig. 2: 1961-1990 average daily temperature, monthly rainfall or Snow Water Equivalent (SWE,  
820 white columns) and snow on the ground (dark columns) in the Gressoney d'Eyola weather station,  
821 only few km far from the studied proglacial area.

822  
823 Fig. 3: conceptual diagram of vegetation development in the Lys proglacial area; the plant  
824 successions observed below timberline are shown on the left, above timberline (i.e. alpine  
825 successions) on the right.

826  
827 Fig. 4: soil chronosequences under alpine (a), subalpine grassland (b) and typical subalpine forest-  
828 heath (c).

829  
830 Fig. 5: comparison between C/N ratio in surface mineral horizons (a); surface pH values (b);  
831 cumulative thickness of O horizons (OL+OF+OH, c); ratio between Feo (d) and Fed (e) in  
832 subsurface CB, BC, Bs and Bw horizons and in surface A, AE and E ones; ISratio (Spodicity Index  
833 ratio between subsurface and surface horizons,  $IS_{(B-BC)}/IS_{(E-AE)}$ , f) in grassland and forest-shrub soils  
834 (p-value < 0.05). In a), the forest soils low-values outlier is caused by the presence of N-fixing  
835 *Alnus viridis* (S8)

836  
837 Fig. 6: Fed depth trends in the LIA sites analyzed in the Lys proglacial area. Different depth trends  
838 are visible for the soils under different primary successions.

839  
840 Fig. 7: (a), chronofunctions of TOC concentration in surface A, AE and E horizons under grassland  
841 (empty circles, continuous line) and forest-shrub vegetation (dashed line, filled squares); (b),  
842 surface TOC concentration (%) in subalpine forest-shrub soils and grassland; (c), TOC  
843 concentration increase with time in grassland (filled squares, continuous line) and in forest soils  
844 (dashed line, empty circles); (d), almost significantly different (p-value = 0.08) TOC concentration  
845 in forest-shrub compared to grassland subsoil (CB, BC, BCs, Bs and Bw horizons).

846  
847 Fig. 8: (a) descriptive chronofunctions of the ISratio in forest (straight line) and grassland soils  
848 (dashed line); . (b) chronofunction of IS increase in subsurface forest soil horizons.

849  
850 Fig. 9: activity ratio in surface and subsurface horizons of grassland and forest soils (p-value <  
851 0.05).

Table 1: Main environmental properties of the study sites; phytoclimatic level (alpine/subalpine), years of deposition of the parent material, altitude, aspect, slope steepness, land cover type and total plant cover (%) are shown.

|     | Phytoclimatic belt | Year of moraine deposition | Altitude | Aspect | Slope | Land cover, total plant cover (%) |
|-----|--------------------|----------------------------|----------|--------|-------|-----------------------------------|
| P1  | Alpine/subalpine   | 2000                       | 2385     | /      | 5°    | Pioneer veg. (5)                  |
| A2  | Alpine             | 1945                       | 2420     | 140°   | 15°   | Grassland (25)                    |
| A3  | Alpine             | 1945                       | 2415     | 180°   | 5°    | Grassland (80)                    |
| A4  | Alpine             | 1820                       | 2460     | 140°   | 20°   | Grassland (100)                   |
| A5  | Alpine             | 1755                       | 2480     | 280°   | 30°   | Grassland / dwarf shrubs (100)    |
| A6  | Alpine             | ca. 11500 BP               | 2510     | 220°   | 30°   | Grassland (100)                   |
| SG2 | Subalpine          | 1945                       | 2180     | 200°   | 5°    | Grazed grassland (100)            |
| S3  | Subalpine          | 1945                       | 2180     | 220°   | 5°    | Larch-rhododendron Forest (100)   |
| S4  | Subalpine          | 1920                       | 2155     | 340°   | 2°    | Larch-rhododendron Forest (100)   |
| SG5 | Subalpine          | 1920                       | 2155     | 160°   | 2°    | Grazed grassland (80)             |
| S6  | Subalpine          | 1880                       | 2105     | 180°   | 2°    | Larch-rhododendron Forest (100)   |
| SG7 | Subalpine          | 1880                       | 2138     | 160°   | 2°    | Grazed grassland (100)            |
| S8  | Subalpine          | 1821                       | 2005     | 20°    | 30°   | Larch-rhododendron Forest (100)   |
| S9  | Subalpine          | 1821                       | 2005     | 0°     | 30°   | Larch-rhododendron Forest (100)   |
| S10 | Subalpine          | ca. 11500 BP               | 1995     | 90°    | 20°   | Larch-rhododendron Forest (100)   |

Table 2: Alpine primary succession; increasing age of the site is from left to right; the % cover of each species (based on visual estimation) is shown in each site. The differentiation between early successional, mid successional, late successional and ubiquitous species is based on Pignatti (1992) and on the Landolt indices (Landolt, 1977): on acidic parent materials, basophilous species should be restricted on weakly developed, initial soils, not yet acidified and desaturated.

|   | P1 | A2 | A3 | A4 | A5 | A6 |
|---|----|----|----|----|----|----|
| Early successional                                  |    |    |    |    |    |    |
| <i>Cerastium uniflorum</i>                          | 1  | 3  | 0  | 0  | 0  | 0  |
| <i>Epilobium fleischeri</i>                         | 1  | 5  | 2  | 0  | 0  | 0  |
| <i>Linaria alpina</i>                               | 1  | 1  |    |    |    |    |
| <i>Oxyria digyna</i>                                | 1  | 0  | 0  | 0  | 0  | 0  |
| <i>Saxifraga oppositifolia</i>                      | 0  | 0  | 1  | 0  | 0  | 0  |
| <i>Trisetum distichophyllum</i>                     | 1  | 3  | 0  | 0  | 0  | 0  |
| Mid-successional                                    |    |    |    |    |    |    |
| <i>Achillea nana</i>                                | 0  | 5  | 0  | 5  | 0  | 0  |
| <i>Agrostis schraderiana</i>                        | 1  | 0  | 0  | 0  | 0  | 0  |
| <i>Anthyllis vulneraria</i> subsp. <i>valesiaca</i> | 0  | 2  | 2  | 0  | 0  | 0  |
| <i>Aster alpinus</i>                                | 0  | 1  | 1  | 0  | 0  | 0  |
| <i>Campanula cochleariifolia</i>                    | 0  | 0  | 0  | 1  | 0  | 0  |
| <i>Carex norvegica</i>                              | 0  | 0  | 1  | 0  | 0  | 0  |
| <i>Carex ornithopodioides</i>                       | 0  | 0  | 1  | 0  | 0  | 0  |
| <i>Erigeron alpinus</i>                             | 0  | 1  | 1  | 0  | 0  | 0  |
| <i>Gentiana nivalis</i>                             | 0  | 0  | 1  | 0  | 0  | 0  |
| <i>Luzula spicata</i>                               | 0  | 0  | 2  | 3  | 0  | 0  |
| <i>Myosotis alpestris</i>                           | 0  | 0  | 10 | 10 | 0  | 0  |
| <i>Saxifraga exarata</i>                            | 1  | 0  | 2  | 0  | 0  | 0  |
| <i>Saxifraga paniculata</i>                         | 0  | 0  | 2  | 0  | 0  | 0  |
| <i>Silene excapa</i>                                | 1  | 0  | 5  | 1  | 0  | 0  |
| <i>Silene rupestris</i>                             | 0  | 0  | 0  | 1  | 0  | 0  |
| <i>Trifolium pallescens</i>                         | 0  | 30 | 10 | 5  | 1  | 0  |
| Late successional                                   |    |    |    |    |    |    |
| <i>Antennaria dioica</i>                            | 0  | 0  | 0  | 0  | 1  | 0  |
| <i>Anthoxanthum alpinum</i>                         | 0  | 0  | 1  | 0  | 0  | 0  |
| <i>Bellis perennis</i>                              | 0  | 0  | 0  | 0  | 1  | 0  |
| <i>Botrychium lunaria</i>                           | 0  | 0  | 0  | 1  | 0  | 0  |
| <i>Carex curvula</i>                                | 0  | 0  | 20 | 5  | 5  | 0  |
| <i>Carex sempervirens</i>                           | 0  | 0  | 0  | 0  | 0  | 5  |
| <i>Coeloglossum viride</i>                          | 0  | 0  | 0  | 0  | 1  | 0  |
| <i>Euphrasia rohoskoviana</i>                       | 0  | 0  | 3  | 0  | 0  | 0  |
| <i>Festuca halleri</i>                              | 0  | 1  | 3  | 5  | 2  | 2  |

|   |   |    |    |    |    |    |
|---|---|----|----|----|----|----|
| <i>Festuca varia</i>                                    | 0 | 1  | 10 | 20 | 0  | 50 |
| <i>Galium anysophyllon</i>                              | 0 | 0  | 0  | 5  | 0  | 0  |
| <i>Gentiana acaulis</i> subsp. <i>Koch</i>              | 0 | 0  | 0  | 1  | 0  | 1  |
| <i>Geum montanum</i>                                    | 0 | 0  | 0  | 5  | 0  | 10 |
| <i>Hieracium piloselloides</i>                          | 0 | 0  | 0  | 0  | 1  | 0  |
| <i>Homogyne alpina</i>                                  | 0 | 0  | 0  | 0  | 1  | 0  |
| <i>Hupertia selago</i>                                  | 0 | 0  | 0  | 1  | 1  | 0  |
| <i>Juncus trifidus</i>                                  | 0 | 0  | 10 | 0  | 0  | 0  |
| <i>Juniperus communis</i>                               | 0 | 0  | 5  | 0  | 0  | 0  |
| <i>Loiseleuria procumbens</i>                           | 0 | 0  | 0  | 0  | 40 | 0  |
| <i>Lotus corniculatus</i> subsp. <i>alpinus</i>         | 0 | 0  | 5  | 3  | 0  | 0  |
| <i>Luzula alpinopilosa</i>                              | 0 | 0  | 0  | 0  | 2  | 1  |
| <i>Luzula lutea</i>                                     | 0 | 0  | 0  | 0  | 1  | 0  |
| <i>Minuartia recurva</i>                                | 0 | 5  | 1  | 0  | 0  | 0  |
| <i>Nardus stricta</i>                                   | 0 | 0  | 0  | 0  | 0  | 5  |
| <i>Pedicularis kernerii</i>                             | 0 | 0  | 0  | 1  | 1  | 0  |
| <i>Pedicularis verticillata</i>                         | 0 | 0  | 0  | 1  | 0  | 0  |
| <i>Poa nemoralis</i>                                    | 0 | 0  | 2  | 0  | 0  | 0  |
| <i>Pulsatilla alpina</i>                                | 0 | 0  | 0  | 1  | 0  | 1  |
| <i>Rhinanthus alectorolophus</i>                        | 0 | 0  | 1  | 0  | 0  | 0  |
| <i>Soldanella verna</i>                                 | 0 | 0  | 0  | 0  | 0  | 10 |
| <i>Trichophorum caespitosum</i>                         | 1 | 0  | 0  | 0  | 3  | 15 |
| <i>Trifolium alpinum</i>                                | 0 | 0  | 5  | 1  | 5  | 10 |
| <i>Vaccinium uliginosum</i> subsp. <i>gaulterioider</i> | 0 | 0  | 20 | 0  | 60 | 0  |
| <i>Valeriana celtica</i>                                | 0 | 0  | 0  | 0  | 2  | 0  |
| <i>Veronica aphylla</i>                                 | 0 | 0  | 1  | 0  | 0  | 2  |
| Ubiquitous  |   |    |    |    |    |    |
| <i>Agrostis rupestris</i>                               | 0 | 0  | 5  | 0  | 1  | 2  |
| <i>Bartsia alpina</i>                                   | 0 | 0  | 1  | 0  | 2  | 0  |
| <i>Cerastium cerastioides</i>                           | 0 | 0  | 1  | 0  | 0  | 10 |
| <i>Cirsium spinosissimum</i>                            | 0 | 0  | 1  | 1  | 0  | 1  |
| <i>Festuca rubra</i>                                    | 0 | 1  | 0  | 0  | 0  | 0  |
| <i>Festuca violacea</i>                                 | 0 | 0  | 5  | 0  | 0  | 0  |
| <i>Leontodon helveticus</i>                             | 0 | 0  | 1  | 2  | 0  | 0  |
| <i>Leucanthemopsis alpina</i>                           | 0 | 0  | 0  | 5  | 0  | 0  |
| <i>Minuartia sedoides</i>                               | 0 | 0  | 2  | 0  | 0  | 0  |
| <i>Poa alpina</i>                                       | 0 | 2  | 5  | 5  | 2  | 0  |
| <i>Polygonum viviparum</i>                              | 0 | 0  | 2  | 0  | 5  | 0  |
| <i>Potentilla frigida</i>                               | 0 | 0  | 0  | 3  | 0  | 0  |
| <i>Primula hirsuta</i>                                  | 0 | 0  | 0  | 0  | 1  | 0  |
| <i>Rhododendron ferrugineum</i>                         | 0 | 0  | 0  | 0  | 5  | 0  |
| <i>Salix helvetica</i>                                  | 0 | 0  | 5  | 0  | 5  | 0  |
| <i>Saxifraga moschata</i>                               | 0 | 0  | 1  | 0  | 0  | 0  |
| <i>Sempervivum arachnoideum</i>                         | 0 | 10 | 2  | 1  | 0  | 0  |

|                              |   |    |   |   |   |   |
|------------------------------|---|----|---|---|---|---|
| <i>Sempervivum montanum</i>  | 0 | 3  | 3 | 2 | 0 | 0 |
| <i>Veronica bellidioides</i> | 0 | 0  | 1 | 0 | 0 | 0 |
| <i>Veronica fruticans</i>    | 0 | 10 | 0 | 0 | 0 | 0 |



|  |   |    |    |   |    |    |    |   |    |    |
|--|---|----|----|---|----|----|----|---|----|----|
| <i>Cirsium spinosissimum</i>                               | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 0 | 1  | 0  |
| <i>Erigeron alpinus</i>                                    | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 0 | 1  | 0  |
| <i>Festuca varia</i>                                       | 0 | 20 | 15 | 0 | 10 | 0  | 0  | 0 | 30 | 0  |
| <i>Festuca violacea</i>                                    | 0 | 5  | 0  | 0 | 0  | 0  | 0  | 0 | 15 | 0  |
| <i>Galium album</i>  | 0 | 10 | 0  | 0 | 0  | 0  | 0  | 0 | 0  | 0  |
| <i>Galium anisophyllum</i>                                 | 0 | 5  | 0  | 0 | 0  | 0  | 0  | 0 | 0  | 0  |
| <i>Geum montanum</i>                                       | 0 | 1  | 0  | 0 | 0  | 0  | 0  | 1 | 5  | 0  |
| <i>Helianthemum nummularium</i> subsp. <i>grandiflorum</i> | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 0 | 1  | 0  |
| <i>Hieracium pilosella</i>                                 | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 0 | 5  | 0  |
| <i>Hieracium prenanthes</i>                                | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 1 | 0  | 0  |
| <i>Knautia arvensis</i>                                    | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 0 | 1  | 0  |
| <i>Leucanthemum vulgare</i>                                | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 0 | 1  | 0  |
| <i>Lotus corniculatus</i> aggr.                            | 0 | 0  | 5  | 0 | 1  | 0  | 0  | 1 | 5  | 0  |
| <i>Nardus stricta</i>                                      | 0 | 0  | 0  | 0 | 5  | 20 | 0  | 0 | 20 | 0  |
| <i>Nigritella nigra</i>                                    | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 0 | 1  | 0  |
| <i>Phyteuma orbicularis</i>                                | 0 | 1  | 0  | 0 | 0  | 0  | 0  | 0 | 0  | 0  |
| <i>Plantago alpina</i>                                     | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 0 | 1  | 0  |
| <i>Poa supina</i>  | 0 | 2  | 0  | 0 | 0  | 0  | 0  | 0 | 0  | 0  |
| <i>Polygonum viviparum</i>                                 | 0 | 1  | 0  | 0 | 0  | 0  | 0  | 1 | 0  | 0  |
| <i>Rhinanthus alectorolophus</i>                           | 0 | 0  | 5  | 0 | 0  | 0  | 0  | 0 | 5  | 0  |
| <i>Schoenus nigricans</i>                                  | 0 | 2  | 0  | 0 | 0  | 0  | 0  | 0 | 0  | 0  |
| <i>Sempervivum montanum</i>                                | 0 | 0  | 2  | 0 | 0  | 0  | 0  | 0 | 0  | 0  |
| <i>Solidago virgaurea</i> subsp. <i>minor</i>              | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 1 | 0  | 0  |
| <i>Thymus serpyllum</i>                                    | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 0 | 5  | 0  |
| <i>Trichophorum caespitosum</i>                            | 1 | 5  | 0  | 0 | 0  | 0  | 0  | 0 | 3  | 0  |
| <i>Trifolium pratense</i> aggr.                            | 0 | 1  | 10 | 0 | 0  | 0  | 0  | 0 | 3  | 0  |
| <i>Trollius europaeus</i>                                  | 0 | 0  | 0  | 0 | 0  | 0  | 0  | 1 | 0  | 0  |
| Subalpine forest– climax (selection)                       |   |    |    |   |    |    |    |   |    |    |
| <i>Alnus viridis</i>                                       | 0 | 30 | 0  | 0 | 0  | 0  | 30 | 1 | 0  | 10 |





Tab. 4: Macromorphological properties of the soils in the alpine chronosequence. Structure: GR = granular; PL = platy; PS = subangular polyhedral; MA = massive; RS = rock structure; AB = absent; M = matted (O horizons). The number indicates the size class of the soil structure: 1 = very fine, 2 = fine, 3 = medium, 4 = coarse; Consistence: LO loose; FR friable; FI firm; VFI very firm; The second number symbolizes the strength of the aggregates: 1 = very weak, 2 = weak, 3 = moderate, 4 = strong, 5 = very strong. Silt caps and their thickness are shown as well (1= up to 1 mm thick, visible on few rock fragments, and 2 = up to 2 mm thick and visible on many rock fragments).

| Profile | Year of deposition | Horizon | Depth (cm) | Munsell Color (moist) | Sand (%) | Clay (%) | Structure (or cementation) | Consistence (moist) | Rock fragments (volume, %) | Siltcaps |
|---------|--------------------|---------|------------|-----------------------|----------|----------|----------------------------|---------------------|----------------------------|----------|
| P1      | 2004               | C1      | 0-8        | 5Y5.5/3               | 69.1     | 2.1      | PL3/AB                     | LO                  | 60                         |          |
|         |                    | C2      | 8-47+      | 5Y6/1                 | 80.3     | 0.8      | PL3/AB                     | LO                  | 60                         | 1        |
| A2      | c. 1950            | AC      | 0-7        | 2.5Y4/3               | 82.3     | 1.1      | AB                         | LO                  | 50                         |          |
|         |                    | C1      | 7-16       | 2.5Y4/4               | 83.4     | 1.1      | AB                         | LO                  | 60                         | 1        |
| A3      | c. 1950            | C2      | 16-35+     | 2.5Y5/3               | 81.5     | 1.0      | AB                         | LO                  | 60                         | 2        |
|         |                    | OL      | 0-0.5      |                       |          |          |                            |                     |                            |          |
| A4      | 1821               | A       | 0-8        | 10YR4/3               | 81.2     | 1.8      | GR2                        | LO1                 | 50                         |          |
|         |                    | AC      | 8-25       | 5Y4/2                 | 81.8     | 1.5      | AB                         | LO                  | 70                         |          |
|         |                    | C       | 25-48+     | 5Y4/1                 | 82.9     | 1.5      | AB                         | LO                  | 70                         | 2        |
|         |                    | OF      | 0-3        |                       |          |          |                            |                     |                            |          |
| A5      | c. 1755            | A       | 3-14       | 10YR3/2               | 78.5     | 2.1      | M                          | LO2                 | 40                         |          |
|         |                    | AB      | 14-17      | 10YR4/4.5             | 79.2     | 2.0      | SP1                        | LO2                 | 60                         |          |
|         |                    | BC      | 17-37      | 10YR4/3               | 81.2     | 1.6      | AB                         | LO                  | 50                         |          |
|         |                    | C       | 37-45+     | 2.5Y5/4               | 80.6     | 1.4      | AB                         | LO                  | 50                         | 1        |
|         |                    | OL      | 0-1        |                       |          |          |                            |                     |                            |          |
| A6      | c. 11000 BP        | OF      | 1-3        | 7.5YR2/1              |          |          |                            |                     |                            |          |
|         |                    | A1      | 3-10       | 7.5YR2.5/1            | 79.5     | 2.6      | M, PL3                     | FR3                 | 20                         |          |
|         |                    | A2      | 10-22      | 7.5YR3/2              | 81.3     | 1.9      | SP2                        | FR2                 | 20                         |          |
|         |                    | Bw      | 22-40      | 10YR5/4               | 84.6     | 1.1      | SP2                        | LO1                 | 30                         |          |
|         |                    | C       | 40-50+     | 2.5Y5/4               | 94.2     | 0.5      | AB                         | LO                  | 30                         |          |
|         |                    | OL-OF   | 0-6        |                       |          |          |                            |                     |                            |          |
| A6      | c. 11000 BP        | A       | 6-12/16    | 7.5YR2.5/2            | 69.2     | 5.2      | GR/GM2                     | LO2                 | 5                          |          |
|         |                    | BA      | 12/16-48   | 7.5YR3/3              | 73.0     | 4.7      | GR2                        | LO2                 | 40                         |          |
|         |                    | Bw      | 48-72      | 8YR3/6                | 84.1     | 3.5      | GR2                        | FR2                 | 30                         |          |
|         |                    | IIBC    | 72-106+    | 10YR3/6               | 82.1     | 1.7      | GR1                        | FR1                 | 10                         |          |

Tab. 5: Chemical properties of the alpine soil chronosequences.

|    | Horizon | pH  | C     | Feo  | Alo  | Fed   | Feo/Fed |
|----|---------|-----|-------|------|------|-------|---------|
|    |         |     | %     | g/kg | g/kg | g/kg  |         |
| P1 | C1      | 6.6 | 0.04  | 0.87 | 0.21 | 1.87  | 0.47    |
|    | C2      | 6.8 | 0.00  | 0.80 | 0.18 | 1.50  | 0.53    |
| A2 | AC      | 5.9 | 1.71  | 1.00 | 0.22 | 1.80  | 0.56    |
|    | C1      | 5.2 | 0.51  | 1.20 | 0.19 | 1.90  | 0.63    |
| A3 | C2      | 5.6 | 0.17  | 0.80 | 0.18 | 1.40  | 0.57    |
|    | A       | 6.1 | 1.46  | 1.35 | 0.44 | 3.15  | 0.44    |
|    | AC      | 6.4 | 0.33  | 1.23 | 0.26 | 2.60  | 0.46    |
| A4 | C       | 6.5 | 0.07  | 1.73 | 0.21 | 2.15  | 0.77    |
|    | O       | 5.1 | 28.00 |      |      |       |         |
|    | A       | 4.7 | 1.90  | 1.80 | 0.70 | 3.40  | 0.53    |
|    | AB      | 5   | 0.93  | 1.60 | 0.70 | 3.50  | 0.46    |
|    | BC      | 5   | 0.50  | 1.30 | 0.50 | 2.80  | 0.46    |
| A5 | C       | 5.1 | 0.31  | 1.30 | 0.40 | 2.80  | 0.46    |
|    | A1      | 4.5 | 4.14  | 1.80 | 0.90 | 4.00  | 0.45    |
|    | A2      | 4.9 | 2.53  | 3.00 | 1.60 | 8.20  | 0.37    |
|    | Bw      | 5.4 | 0.55  | 1.70 | 1.10 | 8.90  | 0.19    |
| A6 | OL      | 4.6 | 23.24 |      |      |       |         |
|    | A       | 4.6 | 4.83  | 7.25 | 2.46 | 16.20 | 0.45    |
|    | BA      | 5.4 | 2.19  | 7.70 | 5.68 | 14.80 | 0.52    |
|    | Bw      | 5.6 | 1.45  | 4.65 | 4.11 | 17.10 | 0.27    |
|    | BC      | 5.7 | 0.77  | 4.28 | 3.58 | 16.40 | 0.26    |

Tab. 6: Macromorphological properties of the soils in the subalpine chronosequence. Structure: GR = granular; PL = platy; PS = subangular polyhedral; MA = massive; RS = rock structure; AB = absent; M = matted (O horizons). The number indicates the size class of the soil structure: 1 = very fine, 2 = fine, 3 = medium, 4 = coarse; Consistence: LO loose; FR friable; FI firm; VFI very firm; The second number symbolizes the strength of the aggregates: 1 = very weak, 2 = weak, 3 = moderate, 4 = strong, 5 = very strong. Silt caps and their thickness are shown as well (1= up to 1 mm thick, visible on few rock fragments, and 2 = up to 2 mm thick and visible on many rock fragments, 3 = up to 2 mm thick and visible on most rock fragments).

| Profile | Year of deposition | Hor.  | Depth (cm)  | Munsell Color (moist) | Sand (%) | Clay (%) | Structure (or cementation) | Consistence (moist) | Rock fragments (volume, %) | Silt caps |
|---------|--------------------|-------|-------------|-----------------------|----------|----------|----------------------------|---------------------|----------------------------|-----------|
| P1      | 2004               | C1    | 0-8         | 5Y5.5/3               | 69.1     | 2.1      | PL3/AB                     | LO                  | 60                         |           |
|         |                    | C2    | 8-47+       | 5Y6/1                 | 80.3     | 0.8      | PL3/AB                     | LO                  | 60                         | 1         |
| SG2     | c. 1950            | OL    | 0-1         |                       |          |          |                            |                     |                            |           |
|         |                    | OH    | 1-3         | 7.5YR2/2              |          |          |                            |                     |                            |           |
|         |                    | A     | 3-6.5       | 10YR4/2               | 81.3     | 1.1      | GR1                        | LO                  | 40                         |           |
|         |                    | CB    | 6.5-18      | 10YR5/4               | 82.3     | 1.1      | GR1                        | LO                  | 50                         |           |
| S3      | c. 1950            | C     | 18-40+      | 2.5Y5/3               | 86.4     | 0.8      | M                          | M                   | 50                         | 3         |
|         |                    | OL    | 0-1         |                       |          |          |                            |                     |                            |           |
| S4      | 1921               | OF    | 1-3         | 7.5YR2/2              |          |          |                            |                     |                            |           |
|         |                    | OH    | 3-6         | 7.5YR2/2              |          |          | M                          | 3                   |                            |           |
|         |                    | A     | 6-12        | 2.5Y4/1.5             | 78.6     | 1.9      | GR1                        | LO                  | 30                         |           |
|         |                    | CB    | 12-25+      | 1Y5/4                 | 81.3     | 1.1      | AB                         | LO                  | 80                         |           |
|         |                    | OL    | 0-1         |                       |          |          |                            |                     |                            |           |
| SG5     | 1921               | OF    | 1-2         | 5YR2/2                |          |          |                            |                     |                            |           |
|         |                    | OH    | 2-6         | 5YR2/2                |          |          |                            |                     |                            |           |
|         |                    | E     | 6-9         | 5YR5/1                | 78.3     | 2.1      | GR1                        | LO                  | 40                         |           |
|         |                    | CBs/C | 9-37+       | 10YR5/4,<br>10YR6/3   | 81.7     | 1.6      | AB                         | LO                  | 60                         |           |
| SG6     | c. 1880            | OF    | 0-4         | 10YR2/1               |          |          | GR1                        | LO                  |                            |           |
|         |                    | A     | 4-17        | 10YR4/3               | 80.5     | 1.7      | GR1                        | LO                  | 60                         |           |
|         |                    | C     | 17-30+      | 10YR5/3               | 81.7     | 0.8      | AB                         | LO                  | 80                         | 1         |
| SG7     | c. 1880            | OL    | 0-1         |                       |          |          |                            |                     |                            |           |
|         |                    | OF-OH | 1-5         | 5YR2.5/2              |          |          | M                          | 3                   |                            |           |
|         |                    | E     | 5-7.5       | 10YR6/2               | 68.7     | 2.2      | GR1                        | M                   | 50                         |           |
|         |                    | BCs   | 7.5-21      | 10YR4.5/4             | 74.5     | 1.1      | PS2                        | FR1                 | 70                         |           |
|         |                    | C     | 21-27+      | 10YR5/3               | 75.6     | 0.9      | AB                         | LO                  | 70                         |           |
| S8      | 1821               | OH    | 0-1         |                       |          |          |                            |                     |                            |           |
|         |                    | A     | 1-3         | 7.5YR2/1              | 76.5     | 3.1      | M                          | 3                   | 20                         |           |
|         |                    | Bw    | 3-20/30     | 10YR4/5               | 78.3     | 2.2      | PS2                        | FR1                 | 70                         |           |
|         |                    | CB    | 20/30-30/38 | 10YR5/3               | 80.2     | 1.1      | PS1                        | LO                  | 50                         | 1         |
|         |                    | C     | 30/38-45    | 2.5Y5/3               | 81.6     | 1.0      | AB                         | Lo                  | 60                         | 3         |

|     |             |       |                 |                       |      |     |  |                      |       |    |
|-----|-------------|-------|-----------------|-----------------------|------|-----|--|----------------------|-------|----|
|     |             | OF    | 2-4             | 7.5YR2.5/2            |      |     |  | M1                   | LO    |    |
|     |             | E     | 4-7             | 10YR5.5/2,<br>10YR3/1 | 67.5 | 2.1 |  | GR1                  | LO1   | 50 |
|     |             | BCs   | 7-21            | 10YR5/4               | 81.9 | 1.0 |  | GR1                  | LO1   | 60 |
|     |             | C     | 21-42           | 10YR5/3               | 82.6 | 0.8 |  | AB (PS1)             | LO1   | 80 |
| S9  | 1821        | OL    | 0-1             |                       |      |     |  |                      |       |    |
|     |             | OF    | 1-5             | 5YR2/2                |      |     |  |                      |       |    |
|     |             | OH    | 5-10/7          | 5YR2/2                |      |     |  |                      |       |    |
|     |             | AE    | 10/7-12         | 10YR4/3               | 68.2 | 1.9 |  | GR0                  | M     | 30 |
|     |             | E     | 12-18           | 10YR5/2               | 66.5 | 2.1 |  | GR1                  | M     | 40 |
|     |             | BCs   | 18-30           | 10YR4/4               | 79.0 | 2.0 |  | PS2                  | FR1   | 40 |
|     |             | C     | 30-40+          | 10YR5/3               | 84.2 | 0.9 |  | AB                   | LO    | 50 |
| S10 | c. 11000 BP | OL-OF | 0-3             |                       |      |     |  |                      |       |    |
|     |             | OH    | 3-21/12         | 7.5YR2/1              |      |     |  | GR1                  | LO    |    |
|     |             | AE/OH | 21/12-<br>20/30 | 7.5YR2/2              | 59.2 | 8.6 |  | AB                   | MA3   | 0  |
|     |             | Eh    | 20/30-<br>34/46 | 10YR5/3               | 64.4 | 7.6 |  | AB/PL2               | LO    | 60 |
|     |             | EBh   | 34/46-54        | 7.5YR4/4              | 66.3 | 7.5 |  | PL2                  | LO    | 60 |
|     |             | Bhs   | 54-68           | 5YR4/4                | 77.8 | 6.1 |  | PS2                  | LO2   | 60 |
|     |             | Bsm1  | 68-87           | 5YR4/4                | 79.2 | 5.9 |  | Cemented             | M     | 80 |
|     |             | Bsm2  | 87-120          | 6YR5/8                | 81.3 | 5.1 |  | Cemented             | M     | 80 |
|     |             | CBm   | 120-135         | 10YR5/8               | 83.4 | 4.9 |  | Strongly<br>cemented | M     | 80 |
|     |             | C(m)  | 135-150+        | 2.5Y5/6               | 79.6 | 4.8 |  | Partly cemented      | M2/LO | 80 |

Tab. 7: Chemical properties of the subalpine soils along the Lys forefield chronosequences.

|      | Horizon | pH   | C     | Feo  | Alo  | Fed  | Feo/Fed | Alo/Feo |
|------|---------|------|-------|------|------|------|---------|---------|
|      |         |      | %     | %    | %    | %    |         |         |
| P1   | C1      | 6.4  | 0.04  | 0.09 | 0.20 | 0.19 | 0.47    | 0.22    |
|      | C2      | 6.6  | 0     | 0.08 | 0.18 | 0.15 | 0.53    | 0.25    |
| SG2  | OH      | 5.9  | 26.58 |      |      |      |         |         |
|      | A       | 5.8  | 1.5   | 0.11 | 0.27 | 0.19 | 0.58    | 0.27    |
|      | CB      | 5.8  | 0.44  | 0.14 | 0.36 | 0.25 | 0.56    | 0.29    |
| S3   | C       | 5.9  | 0.15  | 0.10 | 0.31 | 0.21 | 0.48    | 0.30    |
|      | OL/OF   | 5.6  | 26.89 |      |      |      |         |         |
|      | A       | 5    | 1.92  | 0.07 | 0.19 | 0.18 | 0.39    | 0.29    |
| S4   | CB      | 5.2  | 0.68  | 0.11 | 0.31 | 0.23 | 0.48    | 0.27    |
|      | OL/OF   | 5.1  | 41.2  |      |      |      |         |         |
|      | OH      | 4.9  | 18.44 |      |      |      |         |         |
| SG5  | E       | 5.1  | 1.24  | 0.07 | 0.18 | 0.32 | 0.22    | 0.29    |
|      | CBs     | 5.2  | 0.34  | 0.14 | 0.22 | 0.42 | 0.33    | 0.14    |
|      | OF      | 5.8  | 14.32 |      |      |      |         |         |
| S6   | A       | 5.5  | 0.55  | 0.17 | 0.44 | 0.38 | 0.45    | 0.24    |
|      | C       | 5.8  | 0.19  | 0.12 | 0.51 | 0.16 | 0.75    | 0.50    |
|      | OF-OH   | 4.4  | 35.24 |      |      |      |         |         |
| SG7  | E       | 4.7  | 1.12  | 0.07 | 0.49 | 0.16 | 0.44    | 0.71    |
|      | BCs     | 5.1  | 0.859 | 0.17 | 0.58 | 0.40 | 0.43    | 0.35    |
|      | C       | 5.3  | 0.51  | 0.15 | 0.71 | 0.35 | 0.43    | 0.47    |
| S8   | OH      | 5.6  | 11.78 |      |      |      |         |         |
|      | A       | 5.6  | 8.18  | 0.15 | 0.61 | 0.28 | 0.54    | 0.40    |
|      | Bw      | 5    | 0.74  | 0.15 | 0.43 | 0.27 | 0.56    | 0.27    |
| S9   | CB      | 5.7  | 0.33  | 0.13 | 0.34 | 0.24 | 0.54    | 0.23    |
|      | C       | 5.4  | 0.14  | 0.09 | 0.22 | 0.18 | 0.5     | 0.22    |
|      | OL-OF   | 4.6  | 16.32 |      |      |      |         |         |
| S10  | E       | 5    | 1.19  | 0.11 | 0.47 | 0.17 | 0.65    | 0.45    |
|      | BCs     | 5.1  | 0.75  | 0.17 | 0.80 | 0.67 | 0.26    | 0.47    |
|      | C       | 5.2  | 0.31  | 0.15 | 0.48 | 0.29 | 0.52    | 0.33    |
| S9   | OH/OF   | 5    | 18.72 |      |      |      |         |         |
|      | E       | 4.5  | 0.8   | 0.11 | 0.38 | 0.24 | 0.46    | 0.36    |
|      | Bs      | 4.7  | 0.76  | 0.20 | 0.74 | 0.41 | 0.49    | 0.35    |
| S10  | OL-OF   | 3.6  | 45.21 |      |      |      |         |         |
|      | OH      | 3.5  | 38.53 |      |      |      |         |         |
|      | AE/OH   | 3.6  | 5.4   | 0.12 | 0.15 | 0.38 | 0.32    | 1.17    |
|      | E       | 4.1  | 1.02  | 0.09 | 0.09 | 0.23 | 0.41    | 1.00    |
|      | EBh     | 4.2  | 1.1   | 0.45 | 0.13 | 0.77 | 0.58    | 0.29    |
|      | Bs      | 4.9  | 2.63  | 1.13 | 0.77 | 4.06 | 0.28    | 0.68    |
|      | Bsm1    | 5.3  | 2.31  | 0.76 | 0.82 | 4.02 | 0.19    | 1.08    |
|      | Bsm2    | 5.4  | 0.86  | 0.29 | 0.73 | 1.56 | 0.19    | 2.52    |
|      | CBm     | 5.4  | 0.3   | 0.11 | 0.53 | 0.81 | 0.14    | 4.82    |
| C(m) | 6       | 0.11 | 0.11  | 0.34 | 0.44 | 0.25 | 3.00    |         |



















