

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**A point mutation in the splice donor site of intron 7 in the as2-casein encoding gene of the Mediterranean River buffalo results in an allele-specific exon skipping**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1507122> since 2018-03-18T00:00:22Z

*Published version:*

DOI:10.1111/j.1365-2052.2009.01897.x

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**This is the author's final version of the contribution published as:**

G. Cosenza, A. Pauciullo, M. Feligini, A. Coletta, L. Colimoro, D. Di Bernardino, L. Ramunno. A point mutation in the splice donor site of intron 7 in the as2-casein encoding gene of the Mediterranean River buffalo results in an allele-specific exon skipping. *Animal Genetics* 40, 788–794 (2009).

doi:10.1111/j.1365-2052.2009.01897.x

**The publisher's version is available at:**

<https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2052.2009.01897.x>

**When citing, please refer to the published version.**

**Link to this full text:**

<https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2052.2009.01897.x>

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

1 **A point mutation in the splice donor site of intron 7 in the *as2*-casein encoding gene of the**  
2 **Mediterranean River buffalo results in an allele-specific exon skipping**

3

4 G. Cosenza\*, A. Pauciullo\*, M. Feligini†, A. Coletta‡, L. Colimoro\*, D. Di Bernardino\* and L.  
5 Ramunno\*

6

7 \*Dipartimento di Scienze del Suolo, della Pianta, dell'Ambiente e delle Produzioni Animali,  
8 Universita' degli Studi di Napoli "Federico II", Portici, Italy.

9 †Istituto Sperimentale Italiano Lazzaro Spallanzani, Lodi, Italy.

10 ‡Associazione Nazionale Allevatori Specie Bufalina (ANASB), Caserta, Italy

11

12 Correspondence: L. Ramunno ([ramunno@unina.it](mailto:ramunno@unina.it))

13

14 *Source/description:* The CSN1S2 cDNA of 10 unrelated Mediterranean River Buffaloes reared in  
15 Southern Italy was amplified by RT-PCR, while the region from the 6th to the 8th exon of the  
16 CSN1S2 gene was amplified from genomic template. All amplicons were sequenced twice and in  
17 both directions. Fiftythree individuals randomly chosen from four breeding herds were genotyped for  
18 an AluI-restriction fragment length polymorphism (RFLP). Primer sequences and PCR conditions are  
19 given in Tables S1 & S2. Three individual milk samples from buffaloes with different genotypes at  
20 CSN1S2 were analysed by reverse-phase-high pressure liquid chromatography (RP-HPLC).

21

22 *Polymorphism detection:* cDNA sequence comparisons showed that five individuals had a normal  
23 transcript only (lodged on EMBL, accession FM865618, named CSN1S2A), one had a deleted  
24 transcript only (lodged on EMBL, accession FM865619, named CSN1S2B), because of the splicing  
25 out of the 27-bp of exon 7, and the remaining four had a heterozygous pattern. Analysis of the  
26 genomic sequences revealed a FM865620:g.773G>C transversion that caused inactivation of the

1 intron 7 splice donor site and, consequently, the allele-specific exon skipping characteristic of the  
2 CSN1S2B allele. The g.773G>C mutation creates a new AluI restriction site enabling a PCR–RFLP  
3 rapid genotyping assay (Fig. S1). PCR–RFLP genotypes for the AluI site were consistent with the  
4 cDNA sequence data for all 10 animals. The cDNA sequences showed three additional exonic  
5 mutations forming an extended haplotype with the g.773G>C polymorphism: FM865618: c.459C>T,  
6 c.484A>T and c.568A>G homozygous and heterozygous respectively in the CSN1S2BB and  
7 CSN1S2AB buffaloes. The first is silent, while the remaining two are non-conservative (p.Ile162Phe  
8 and p.Thp200Ala respectively). Chromatographic analysis of three individual samples with CSN1S2  
9 AA, AB and BB genotypes showed the same retention time for the as2 casein fraction (Fig. S2), but  
10 the hydrophobic characteristics of each allele do not allow their chromatographic separation. Allelic  
11 frequencies: The genotype frequencies (37 CSN1S2A/A, 15 CSN1S2A/B and one CSN1S2B/B) are  
12 in agreement with Hardy–Weinberg equilibrium ( $\chi^2 = 0.13$ , d.f. = 1), with the frequency of the deleted  
13 B allele being 0.16.

14

15 *Comments:* The results indicate that buffalo, similar to goats<sup>1,2</sup> and cattle<sup>3,4</sup> have a CSN1S2 allele  
16 resulting from a non constitutive splicing event. The predicted bubaline as2B protein is 198 aa long  
17 instead of 207 aa and would also be characterized by the presence of Phe at position 147 and Ala at  
18 185.

19

20 *Acknowledgement:* This research was financially supported by the Italian Ministry for Agriculture  
21 and Forestry Policy-MiPAF (SELMOL project).

22

### 23 *References*

24 1 Ramunno L. et al. (2001) Anim Genet 32, 264–8.

25 2 Martin P. et al. (1999) Int Dairy J 9, 163–71.

26 3 Bouniol C. et al. (1993) Gene 128, 289–93.

1 4 Mohr U. et al. (1994) Gene 143, 187–92.

2

3 *Supporting information*

4 Additional supporting information may be found in the online version of this article.

5 Table S1 Oligonucleotide primer sequences and positions.

6 Table S2 Thermal amplification programmes for (a) RT-PCR, (b) PCR and (c) AluI PCR-RFLP.


7 Figure S1 Observed genotypes after AluI digestion of fragments obtained by PCR of the DNA region  
8 spanning the 7th exon and flanking regions of the Mediterranean river buffalo CSN1S2 gene.

9 Figure S2 RP-HPLC chromatogram of individual buffalo milk samples.


**Table S1.** Primer sequences and amplification conditions for PCR.

<b>CSN2-amplified fragment</b>	<b>Position<sup>1</sup></b>	<b>Primer sequence (5'-3')</b>	<b>Use</b>	<b>Annealing temperature</b>
Promoter	1292-1311	F: TCCATTATAGCTTAAGCACT	AS-PCR	58.5 °C
		F: TCCATTATAGCTTAAGCACC		
	Complementary to 1457-1475	R: TGGGATGCACGGAAGTTT		
Exon 9	10323-10340	F: GGGGGTGAGATGAAGAGT	Internal control	
	Complementary to 10663-10682	R: AATGACTGGTTAGGAAATAG		


<sup>1</sup>Numbering of primers was according to the goat *CSN2* gene (AJ011018).




**Figure S1.** Observed genotypes after AluI digestion of fragments obtained by PCR of the DNA region spanning the 7<sup>th</sup> exon and flanking regions of the Mediterranean river buffalo *CSN1S2* gene. M=100 bp DNA ladder (Promega); lane 1: *CSN1S2* A/ A; lane 2: *CSN1S2* A/ B; lane 3 *CSN1S2* B/ B.



**CSNIS2AA**



**CSNIS2AB**



**CSNIS2BB**



**Figure S2.** RP-HPLC chromatogram of individual buffalo milk samples. Subjects with different genotypes at the *CSNIS2 locus* (*CSNIS2AA*, AB and BB) showed the same retention time for the  $\alpha_2$  casein fraction. Peaks 1, 3 and 4:  $\kappa$  casein, peaks 5 and 6:  $\alpha_1$  casein, peak 7:  $\beta$  casein, peak 2:  $\alpha_2$  casein.

**Table S1** Oligonucleotide primer sequences and positions.

	Position nt	Primers sequence (5'-3')*		EMBL	Amplicon size
RT-PCR**	1-18	Forward	ATATTCCATTGCCTGGAC	DQ133467	796 bp 769 bp
	Complementary to: 775-796	Reverse	ATGTGAAACTGTAGAAGATAGA		
PCR ex 6-8	175-190	Forward	GAGAACCTTTGCTCCA	FM865618	~1750 bp
	Complementary to: 235-251	Reverse	CTAGATGAGCCGATAGA		
AluI PCR-RFLP***	668-687	Forward	TTCCTTTTATATTCAGGAGA	FM865620	288 bp
	Complementary to: 938-955	Reverse	GATTTAGGTGGACATTAC		
PCR Mix composition	The 100- $\mu$ l PCR reaction mix comprised: 20 $\mu$ l of RT reaction product or 100 ng of genomic DNA, 1x PCR Buffer, 10 pmol of each primer, dNTPs each at 0.2 mM, 5 U of Taq DNA Polymerase (Promega), 3 mM MgCl <sub>2</sub>				

\* Primers were designed using OLIGO 5.0 software (National Biosciences Inc., Plymouth, MN)

\*\*The RT was performed using Improm-IIk Reverse Transcriptase (Promega)

\*\*\*Digestion with 10 U of AluI endonuclease of 17 $\mu$ l of PCR product was carried out for 5 h at 37 °C following the buffer manufacturer's instructions (Promega).

**Table S2** Thermal amplification programs for a) RT-PCR b) PCR and c) AluI PCR-RFLP

a)

Cycle	Denaturation	Annealing	Extension
1	97 °C – 2 min	57.5 °C – 45 sec	72 °C – 1.5 sec
29	94 °C – 45 sec	57.5 °C – 45 sec	72 °C – 1.5 sec
1	94 °C – 45 sec	57.5 °C – 45 sec	72 °C – 10 min

b)

Cycle	Denaturation	Annealing	Extension
1	97 °C – 2 min	54 °C – 45 sec	72 °C – 2 min
29	94 °C – 45 sec	54 °C – 45 sec	72 °C – 2 min
1	94 °C – 45 sec	54 °C – 45 sec	72 °C – 10 min

c)

Cycle	Denaturation	Annealing	Extension
1	97 °C – 2 min	54 °C – 45 sec	72 °C – 1.5 sec
29	94 °C – 45 sec	54 °C – 45 sec	72 °C – 1.5 sec
1	94 °C – 45 sec	54 °C – 45 sec	72 °C – 10 min