

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Relationship between European Corn Borer injury, *Fusarium proliferatum* and *F. subglutinans* infection and moniliformin contamination in maize

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1526393> since 2015-10-21T12:50:29Z

Published version:

DOI:10.1016/j.fcr.2015.07.014

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



UNIVERSITÀ DEGLI STUDI DI TORINO

1

2

3 This Accepted Author Manuscript (AAM) is copyrighted and published by Elsevier. It is posted
4 here by agreement between Elsevier and the University of Turin. Changes resulting from the
5 publishing process - such as editing, corrections, structural formatting, and other quality control
6 mechanisms - may not be reflected in this version of the text. The definitive version of the text was
7 subsequently published in “Relationship between European Corn Borer injury, *Fusarium*
8 *proliferatum* and *F. subglutinans* infection and moniliformin contamination in maize.”, *Field Crops*
9 *Research*, 183: 69-78. DOI:10.1016/j.fcr.2015.07.014

10 You may download, copy and otherwise use the AAM for non-commercial purposes provided that
11 your license is limited by the following restrictions:

12

13 (1) You may use this AAM for non-commercial purposes only under the terms of the CC-BY-NC-
14 ND license.

15 (2) The integrity of the work and identification of the author, copyright owner, and publisher must
16 be preserved in any copy.

17 (3) You must attribute this AAM in the following format: Creative Commons BY-NC-ND license
18 (<http://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>), [+DOI: 10.1016/j.fcr.2015.07.014]

19

20

21 **FIELD CROP RESEARCH**

22

23 **Relationship between European Corn Borer injury, *Fusarium***
24 ***proliferatum* and *F. subglutinans* infection and moniliformin**
25 **contamination in maize.**

26

27 Authors:

28 Valentina Scarpino¹, Amedeo Reyneri¹, Francesca Vanara¹, Cristina Scopel², Roberto
29 Causin², Massimo Blandino^{1*}

30

31 Affiliation:

32 ¹*University of Torino, Department of Agricultural, Forest and Food Sciences, Largo Paolo*
33 *Braccini 2, 10095 Grugliasco (TO), Italy.*

34 ²*University of Padova, Department of Land, Environment, Agriculture and Forestry -*
35 *AGRIPOLIS - Viale dell'Università 16, 35020 Legnaro (PD), Italy.*

36

37 * Corresponding author: Tel: +39-011-6708895; fax +39-011-6708798.

38 E-mail address: massimo.blandino@unito.it

39

40 **Acknowledgements**

41 The authors would like to thank Fabio Carnaroglio, Mattia Ciro Mancini, Alessandro Peila,
42 Federico Marinaccio and Valentina Sovrani for their precious help and cooperation in the
43 laboratory and field work.

44 The research has been conducted thanks to the financial support of the Italian Ministry of
45 Agricultural, Food and Forestry Policies (MIPAAF), as a part of the MICOPRINCEM project
46 (Coordinator Dr. Maria Grazia D'Egidio, CRA – QCE).

47

48 **Abstract**

49 The European Corn Borer (ECB), *Ostrinia nubilalis* (Hübner), plays an important role in
50 promoting *Fusarium* infection and fumonisin production in maize kernels. Moniliformin
51 (MON) is a mycotoxin that is usually found in association with fumonisins (FB), and as a
52 consequence ECB is also expected to play a role in MON contamination.

53 The aim of the current study was to investigate the influence of ECB activity on MON
54 contamination and on the fungi responsible for its contamination.

55 A comparison has been made between maize infected naturally by insect larvae and
56 protected maize, from 2008 to 2010 in North-West Italy; the latter was obtained by
57 positioning an entomological net at the end of maize flowering.

58 The *F. proliferatum* infection of the maize grain was higher in each growing season than
59 that caused by *F. subglutinans*, although both species were significantly increased due to
60 the ECB damage to the maize ears. The ECB activity significantly increased the MON
61 content 26-fold in 2008 from 43 to 1137 $\mu\text{g kg}^{-1}$, 25-fold in 2009 from < LOQ to 77 $\mu\text{g kg}^{-1}$
62 and 94-fold in 2010 from 6 to 564 $\mu\text{g kg}^{-1}$. The relationship between the MON content and
63 the *Fusarium* species producers of MON was closer for *F. proliferatum* ($R^2 = 0.93$, $P <$
64 0.001) than for *F. subglutinans* ($R^2 = 0.68$, $P < 0.001$). An *in vitro* assay was carried on in
65 order to assess the toxigenic capacity of different *F. proliferatum* and *F. subglutinans*
66 strains isolated from a field experiment and artificially inoculated on a maize substrate. On
67 average, the *F. proliferatum* isolated strains showed a significantly higher ($P < 0.001$)
68 toxigenic capacity than the *F. subglutinans* strains.

69 In conclusion, the combination of the high toxigenic capacity of *F. proliferatum*, and its
70 more frequent occurrence and greater intensity in the field make it possible to state that
71 the production of MON in maize temperate areas, such as North Italy, is mainly due to *F.*
72 *proliferatum* infections and is closely linked to the injury caused by ECB larvae.

73 **Keywords:** maize, Moniliformin, *Fusarium proliferatum*, *Fusarium subglutinans*, European

74 Corn Borer

75

76 **Abbreviations**

77 ANOVA, Analysis of variance; EC, European Commission; ECB, European Corn Borer;
78 EFSA, European Food Safety Authority; ESI, electrospray ionization; FB, fumonisins,
79 GDD, Accumulated growing degree days; HILIC, hydrophilic interaction chromatography;
80 HPLC, high-performance liquid chromatography; LC-MS/MS, Liquid chromatography
81 coupled to tandem mass spectrometry detection; LOD, limit of detection; LOQ, limit of
82 quantification; MON, moniliformin; MS, mass spectrometry detection; RSD, relative
83 standard deviation.

84

85 **1. Introduction**

86 Maize ear rot is a common disease throughout the world that is caused by several
87 *Fusarium* species which can co-occur to give rise to the production of different mycotoxins
88 in infected kernels, with a considerable impact on human and animal health (Bottalico,
89 1998). In temperate areas, maize pink ear rot, which is mainly caused by *F. verticillioides*,
90 leads to an extensive occurrence of FB, a group of toxins among which fumonisin B₁ (FB₁)
91 has been evaluated, by the International Agency for Research on Cancer (IARC), as a
92 Group 2B carcinogen, i.e., possibly carcinogenic to humans (IARC, 2002). Regulatory
93 limits have been set for maize grain within the European Union (European Commission
94 No. 1881/2006 and EC No. 1126/2007), the USA (FDA - Food and Drug Administration,
95 2000) and other countries.

96 Moniliformin (MON) is a mycotoxin that is usually found in association with FB (D'Mello et
97 al., 1999; Sanhueza and Degrossi, 2004). This mycotoxin is produced by *Fusarium*
98 *subglutinans* (Logrieco et al., 2002) *Fusarium temperatum* (Scauflaire et al 2011;
99 Scauflaire et al 2012) and *F. proliferatum*, which is also a FB producer. These *Fusarium*
100 species, which refer to the *Liseola* section, could infect maize kernels in temperate areas
101 and co-participate with *F. verticillioides* in maize ear rot (Srobarova et al., 2002).

102 No regulatory limits have been established about the presence of MON in food or feeds.
103 Although its toxic effects have not yet been fully established, Jonsson et al. (2013) have
104 recently reported a high acute toxicity of MON in rats, with the LD₅₀ value being at the
105 same level as the most toxic *Fusarium* mycotoxins, that is, T-2 and HT-2.

106 The European Food Safety Authority (EFSA) is currently working on establishing a
107 scientific opinion on the risks to public health related to the presence of MON in feeds and
108 food (EFSA 2010). In addition to studies conducted to clarify the occurrence and
109 toxicological effects, it is also of extreme importance to individuate the conditions which

110 could lead to a higher contamination of this mycotoxin, in order to understand the Best
111 Management Practices that can be used to limit contamination.

112 The European Corn Borer (ECB), *Ostrinia nubilalis* (Hübner), is a maize pest that has an
113 economic impact on several producing areas (Bode and Calvin, 1990; Szóke et al., 2002).
114 Two generations of ECB larvae usually occur per year in North Italy: the first generation
115 attacks plants during the mid to late vegetative stages and the second generation attacks
116 during the reproductive stages (from early milk stage to maturity) (Blandino et al., 2009).
117 Moreover, second generation larvae play a very important role in promoting *Fusarium*
118 infection and mycotoxin production in maize kernels (Sobek and Munkvold, 1999).

119 Other *Fusarium* infection pathways are known, through seed and seedlings (systemic) or
120 silks at flowering, although these seem to play a minor role in the final mycotoxin
121 contamination compared to the kernel damage caused by insects (Munkvold et al., 1997).
122 In temperate areas, *F. verticillioides* is generally more favoured by ECB larva feeding than
123 other *Fusarium* species (Lew et al., 1991; Sobek and Munkvold, 1999) and the direct and
124 indirect control of this pest is one of the main strategies adopted to minimize FB
125 contamination (Blandino et al., 2009).

126 Moreover, Papst et al. (2005) and Magg et al. (2003), in comparative studies on Bt maize
127 and its isogenic counterparts conducted in Central Europe, have also reported a significant
128 correlation between the percentage of ECB damaged ears and the MON concentration.
129 Unfortunately, at the moment, no data are available on the effect of ECB on MON contents
130 in warmer maize cultivation areas, where the pressure and damage caused to ears and
131 kernels by this insect are higher. Moreover, so far, it has not been clearly reported, in the
132 literature, which *Fusarium* species is the most responsible for MON contamination in
133 temperate areas or what relationship exists between these species and ECB activity.

134 The aim of the current study was to investigate the influence of ECB activity on MON
135 contamination and the fungi responsible for its contamination, in order to evaluate the

136 effectiveness of the insect control in managing the MON contamination. Moreover, another
137 objective of this work was to assess which of the two *Fusarium* species, *F. proliferatum*
138 and *F. subglutinans*, is the more responsible for MON contamination in maize cultivated in
139 temperate areas, subjected to ECB damage.

140

141

142 **2. Materials and Methods**

143 *2.1. Chemicals*

144 All the chemicals and analytical standards used for the chemical measurements were
145 purchased from Sigma Aldrich (St. Louis, MO), or VWR (Milan, Italy). The solvents were all
146 gradient grade or LC-MS grade.

147 The MON standard was purchased as sodium salt and a 93 mg L⁻¹ stock solution of MON
148 was prepared in acetonitrile/water (84:16, v/v) and stored at 4 °C. This stock solution was
149 used to prepare standard solutions through dilution with acetonitrile/water (84:16, v/v).

150 The FB₁ and FB₂ standards were purchased as a 50 mg L⁻¹ stock solutions in
151 acetonitrile/water (50:50, v/v) and stored at -20 °C. These stock solutions were used to
152 prepare standard solutions containing both FB₁ and FB₂, through dilution with
153 acetonitrile/water (50:50, v/v).

154 All the chemicals used for the mycological measurements were purchased from Sigma (St.
155 Louis, MO), Eppendorf-Germany (Hamburg, Germany), Applied Biosystems by Life
156 Technologies (Life Technologies Italia, Monza, Italy), or PRIMM (PRIMM srl, Milan, Italy).

157

158

159

160

161 2.2. *Field Experimental Design and Samples*

162 The effect of ECB larva feeding activity on MON contamination in maize kernels was
163 studied from 2008 to 2010 in North-West Italy under naturally-infected field conditions in
164 the experimental farm of the Department of Agricultural, Forest and Food Science,
165 University of Turin located in Carmagnola (44° 50' N, 7° 40' E; altitude 245 m), in a sandy-
166 medium textured soil (Typic Udifluvents).

167 The natural maize ear infestation, due to the insect larvae, was compared each year with
168 the protected maize (artificial control of ECB larvae), which was obtained by positioning an
169 entomological net at the end of maize flowering [Growth stage (GS) 69, Lancashire et al.
170 1991] in order to avoid ECB ovideposition.

171 The ECB natural infestation and artificial protection treatments were assigned to
172 experimental units using a completely randomized block design with 4 replicates. Each
173 plot consisted of 4 rows 0.75 m apart and 4 m long. The plot alleys, orthogonal to the
174 maize rows, were one meter wide.

175 The entomological net had a mesh size of 1 mm, and it was placed on a 4.20 m long, 4.20
176 m wide and 3.80 m high, steel structure, in order to minimize any significant microclimatic
177 changes. The position of the net in the field changed every growing seasons. The edge of
178 the net was buried, to prevent adult insects from entering, while the plants within the net
179 were carefully checked for possible first generation attack. If the plants presented any
180 typical leaf injuries caused by first generation ECB larvae, they were cut at the bottom and
181 removed from the plots.

182 No foliar insecticides were applied to the experimental field or to an approximately 20 ha
183 area around the field to control ECB or other insects during the entire growing period.

184 The normal agronomic techniques was adopted. Briefly, the previous crop was maize, and
185 the field was ploughed each year. Studies were carried out each year on the commercial

186 dent corn hybrid Syngenta NX7444 (FAO maturity class 600; 130 days relative to
187 maturity). The crop density was approximately 75.000 plants per hectare and the
188 experiment field received 250, 90 and 100 kg ha⁻¹ of N, P and K, respectively, each year.
189 The field was cultivated under irrigation with full length maturity hybrids, planted in each
190 growing season in the period between the last decade of March and the first decade of
191 April. Metolachlor and terbutylazine were applied to control weeds in pre-emergence while
192 sulcotrione and nicosulfuron were applied in post-emergence.

193 Thirty randomly selected ears (included the ears used for the evaluation of ECB incidence
194 and severity at harvest) were collected by hand from each plot at the end of maturity at a
195 grain moisture content of between 23 and 27%, and shelled using an electric sheller. The
196 kernels from each plot were mixed thoroughly to obtain a random distribution; 4 kg
197 samples were then taken and dried at 60°C for 3 days to analyze the MON and FB
198 contents and to quantify *F. verticillioides*, *F. subglutinans* and *F. proliferatum* using Real
199 Time PCR. All the amount of grains collected from each plot was completely ground using
200 a ZM 200 Ultra Centrifugal Mill (Retsch GmbH, Haan, Germany) and reduced to 1 kg
201 subsample by dynamically sampling the flowing flour during the grinding. This subsample
202 was re-milled and reduced to 500 g with the same method and subsequently an aliquot of
203 100 g, randomly sampled, was used in the molecular analysis, while the remaining 400 g
204 were employed in the chemical analysis.

205 During all the investigated period, rainfall and temperature data have been recorded daily
206 from a weather station, located next to the experimental field.

207

208 **2.3. Entomological measurements**

209 ECB damage incidence was calculated as the percentage of ears per plot with kernel
210 injury or apical and basal tunnels in the cob due to larva activity. ECB damage severity

211 was calculated as the percentage of kernels per ear with injuries due to larva activity. A 7-
212 class rating scale ~~scale of 1 to 7~~ was used in which each numerical value corresponded to
213 a percentage interval of surfaces exhibiting visible kernel damage due to larva activity,
214 according to the following schedule: 1 = no injuries, 2 = 1-5%, 3 = 6-10%; 4 = 11-20 %, 5 =
215 21-35%, 6 = 35-60%, 7 > 60% (Blandino et al., 2009). The ECB damage severity scores
216 were converted to percentages of ears exhibiting symptoms, by using the mid-point of
217 each class interval.

218

219 *2.4. Mycological measurements*

220

221 *2.4.1 Fungal ear rot incidence and severity*

222 Fungal ear rot incidence was calculated as the percentage of ears per plot with symptoms,
223 while fungal ear rot severity was calculated as the percentage of kernels per ear with
224 symptoms. A scale of 1 to 7 was used in which each numerical value corresponds to a
225 percentage interval of surfaces exhibiting visible symptoms of the disease according to the
226 following schedule: 1 = no symptoms, 2 = 1-3 %, 3 = 4-10%; 4 = 11-25 %, 5 = 26-50%, 6 =
227 51-75%, 7 > 75% (Blandino et al., 2009). The ear rot severity scores were converted to
228 percentages of ears exhibiting symptoms and each score was replaced with the mid-point
229 of the interval.

230

231 *2.4.2. Fusarium kernel infection*

232 The evaluation of the total *Fusarium* spp (section *Liseola*) infections was carried out using
233 100 kernels randomly sampled from the untreated control at the dough stage (GS 85). The
234 kernels were surface disinfested for 3 min in a 2% solution of sodium hypochlorite, then
235 rinsed 3 times with sterile water. The kernels were placed in Petri dishes containing

236 dicloran cloramfenicol peptone (DCPA) and incubated at 20 °C. The *Fusarium* colonies
237 were identified after 7 to 10 days through colony and conidial morphology, as reported by
238 Nelson et al. (1983).

239

240 *2.4.3. DNA extraction from the maize samples*

241 The total DNA from the fungal reference cultures was extracted from approximately 200-
242 300 mg (wet weight) of fresh mycelium, according Henrion et al. (1992), with slight
243 modifications. The samples were crushed in liquid N₂ and immediately suspended in one
244 mL of lysis buffer (100 mM Tris-HCl, pH 8, 20 mM EDTA, pH 8, 1.4 mM NaCl, 2%
245 cetyltrimethylammonium bromide (CTAB), 1% polyvinylpyrrolidone (PVP), 1%
246 mercaptoethanol) for 60 minutes at 65°C. The proteins were denatured and removed
247 through one or more extractions by gently shaking the solution for 1 hour in ice with 1 mL
248 chloroform/isoamyl alcohol (24:1, v/v), followed by centrifugation at 17300 × g for 10
249 minutes, to separate the phases. The nucleic acids were precipitated from the aqueous
250 phase by adding 2/3 volume of isopropanol and 1/10 volume of sodium acetate 3 M (pH
251 5.2) and maintaining the sample at -20°C for 20 min. The DNA was then pelleted by
252 centrifugation at 17300 × g for 10 minutes, washed with 70% (v/v) ethanol, vacuum dried
253 briefly, resuspended in 50-150 µL of TE buffer (10 mM Tris-HCl pH 8 + 1 mM EDTA) and
254 stored at -20°C until use.

255 The flour aliquot of 100 g, obtained as previously described, was milled again in a knife
256 mill Grindomix GM 200 (Retsch Technology GmbH, Haan, Germany). Then 10 g of re-
257 milled flour were further ground in liquid N₂ and the DNA was extracted from 1 g, using the
258 DNeasy Plant Maxi Kit, according to the manufacturer's instruction.

259 The concentration of nucleic acids was determined using a Biophotometer (Eppendorf AG,
260 Hamburg-Germany) and DNA purity was evaluated by comparing the A260/A280 and
261 A260/A230 absorbance ratios.

262

263 2.4.4. Preliminary species-specific PCR

264 End-Point PCR was carried out using the SUB1 (CTGTGCGCTAACCTCTTTATCCA) and
265 SUB2 (CAGTATGGACGTTGGTATTATATCTAA) primer for the detection of *F.*
266 subglutinans; the PRO1 (CTTCCGCCAAGTTTCTTC) and PRO2
267 (TGTCAGTAACTCGACGTTGTTG) primer for the detection of *F. proliferatum*; the VER1
268 (CTTCCTGCGATGTTTCTCC) and VER2 (AATTGGCCATTGGTATTATATATCTA) primer
269 for the detection of *F. verticillioides*, all designed by Mulè et al., 2004 on the *calmodulin*
270 gene. These primers amplify respectively 631 bp for *F. subglutinans*, 585 bp for *F.*
271 *proliferatum* and 578 bp for *F. verticillioides*.

272 Moreover, the primers subsequently used in the Real Time PCR were tested to verify high
273 specificity of the investigated *Fusarium* species.

274 The PCR assays were conducted in 50 µL of reaction mixtures containing 1.25 U of Taq
275 (HotMaster™ Taq DNA Polymerase, Eppendorf-Germany), a 1X HotMaster Taq Buffer
276 with Mg²⁺, 400 nM of each primer (PRIMM srl, Milan, Italy), 150 µM of each dNTP
277 (Eppendorf-Germany), and 50 ng of total DNA (from maize). Reactions were performed
278 using the following PCR conditions: denaturation at 95 °C for 5 min; 35 cycles of
279 denaturation at 94 °C for 50 s, annealing at 56 °C for 50 s, extension at 72 °C for 1 min;
280 final extension at 72 °C for 7 min, followed by cooling at 4°C until the samples were
281 recovered. Amplification products were assessed on 1,2% agarose gel stained with Green
282 Gel Plus™ (Fisher Molecular Biology-USA).

283

284 2.4.5. Quantification of *F. verticillioides*, *F. proliferatum* and *F. subglutinans* using Real
285 Time PCR

286 The primers used for the detection of *F. subglutinans* were Fsub565 F
287 (TCATTGGTATGTTGTCGCTCATG) and Fsub622A R
288 (GTGATATGTTAGTACGAATAAAGGGAGAAC), designed on the *EF1 α* gene (Nicolaisen
289 et al., 2009; Boutigny et al. 2011). The primers Fp3-F (CGGCCACCAGAGGATGTG) and
290 Fp4-R (CAACACGAATCGCT TCCTGAC), designed by Jurado et al. (2006) on the
291 intergenic sequence of the ribosomal RNA gene cluster, for the detection of *F. proliferatum*
292 and the primers VER1 and VER2, designed by Mulè et al., 2004 on the *calmodulin* gene,
293 for the detection of *F. verticillioides*, were used in Real Time as described by Nutz et al.
294 (2011); all were synthesized by PRIMM (PRIMM srl, Milan, Italy).

295 Real-time PCR assays were carried out in a total volume of 25 μ L, consisting of 12.5 μ L
296 2X SYBR Green PCR Master Mix (Applied Biosystems), 600nM concentration of each
297 primer for *F. proliferatum*, 300 nM for *F. subglutinans* and *F. verticillioides* and 2 μ L
298 template DNA (approximately 10 - 20 ng); only for *F. subglutinans*. the reaction mixture
299 contain also 0.5 μ g μ L⁻¹ of bovine serum albumin (BSA).

300 The Real-time PCR reactions were performed in triplicate on all the samples and were
301 performed on an ABI 7500 Real-Time PCR System (Applied Biosystems) using three
302 different cycling protocols. The first, for *F. proliferatum*, was 2 min at 50 °C; 10 min at 95
303 °C; 40 cycles of 15 s at 95 °C and 1 min at 62 °C. The second, for *F. subglutinans*, was: 2
304 min at 50 °C; 4 min at 95 °C; 35 cycles of 35 s at 95 °C, 32 s at 64 °C, 32 s at 72°C and 5
305 min at 72°C. The third, for *F. verticillioides*, was 2 min at 50 °C; 10 min at 95 °C; 40 cycles
306 of 50 s at 94 °C, 50 s at 60 °C, 1 min at 72°C and 5 min at 72°C.

307 All cycling protocols were followed by a dissociation curve analysis at 60 to 95 °C; the
308 fluorescence was measured in the first and the second protocol during the annealing
309 phase; in the third during the extension phase.

310 In order to test for linearity and the presence of inhibitors in the pathogen DNA, standard
311 curves were prepared by diluting the DNA of each pathogen (approximately 121 ng μL^{-1}
312 for *F. subglutinans*, 256 ng μL^{-1} for *F. proliferatum* and 97 ng μL^{-1} for *F. verticillioides*) 5-,
313 25-, 125-, 625-, 3125-, 15625-, 78125-fold in maize DNA that was free of fungal
314 contamination (approximately 105 ng μL^{-1}). Standard curves were generated by plotting
315 threshold cycle values (Ct values) against the logarithm of the starting DNA quantities. The
316 standard curves slopes were used to calculate the reaction efficiency (E) of the PCR
317 assays. The amounts of fungal DNA obtained were normalized to the total DNA amount
318 extracted from the meal samples.

319 Real-time PCR reactions were performed in triplicate for each sample.

320

321 *2.5. In vitro MON production by means of different monoconidial strains of* 322 *F. verticillioides, F. proliferatum and F. subglutinans*

323 An *in vitro* assay was performed in order to analyze the toxigenic capacity of the fungal
324 monoconidial strains isolated from the field experiment. Five *F. proliferatum*, five *F.*
325 *subglutinans* and five *F. verticillioides* strains were randomly chosen from all the isolated
326 strains and they were grown for a week on potato dextrose agar (PDA) at 25°C in Petri
327 dishes in the dark.

328 Reference Cultures (*Fusarium proliferatum*, KSU D-4854, Kansas State University;
329 *Fusarium subglutinans*, KSU E-0990, Kansas State University and *F. verticillioides*, KSU A-
330 0999, Kansas State University) were instead grown for DNA extraction on potato dextrose
331 broth (PDB) in 100 mL Erlenmeyer flasks at room temperature on a rotary shaker. The

332 mycelium was harvested after 14 days by means of filtration and then rinsed in distilled
333 water.

334 A commercial maize lot was ground using a ZM 200 Ultra Centrifugal Mill (Retsch GmbH,
335 Haan, Germany) and the flour was used as a substrate for the *Fusarium proliferatum*,
336 *Fusarium subglutinans* and *F. verticillioides* strains. The grain was preliminary tested by
337 means of LC-MS/MS in order to correct the results and to conduct the assays using a
338 meal in which the MON content was as low as possible.

339 The meal was autoclave sterilized and moistened with 25 g of sterile water in 35 g of
340 milled kernels; the Petri dishes (Ø 9 cm) were then inoculated with the selected *Fusarium*
341 strains, using 5 mycelium plugs of 6 mm², taken from one-week-old colonies actively
342 growing on PDA at 25°C in the dark. Three replicates were performed for each fungal
343 strain and 6 Petri dishes were prepared for each replicate. Six Petri dishes without
344 mycelium plugs were also produced as a control.

345 All the Petri dishes were incubated at 25°C in the dark for 20 days and the flour was then
346 exsiccated in a dry kiln for 36 hours at 65°C and for 12 hours at 50°C before being ground
347 again in order to homogenize the mycelium with the meal. These samples were tested for
348 the MON content by means of LC-MS/MS analysis, and *F. subglutinans*, *F. proliferatum*
349 and *F. verticillioides* were quantified using Real Time PCR.

350 The toxigenic capacity of the different fungal strains was expressed as the ratio between
351 the MON content and the fungal DNA, normalized for the total DNA.

352

353 2.6. Chemical Analyses

354 2.6.1. Sample Preparation and Extraction

355 For MON extraction 25 g of maize flour was extracted by mechanical shaking at 100 rpm
356 for 1 h (shaker mod. M102-OS, MPM Instruments, Milan, Italy) with 100 mL

357 acetonitrile/water (84:16, v/v). The extracts were filtered through Whatman[®] no. 1 filters
358 (Brentford, UK) and subjected to clean-up and purification.

359 For FB₁ and FB₂ extraction 50 g of maize flour was extracted by mechanical shaking at
360 100 rpm for 20 min (shaker mod. M102-OS, MPM Instruments, Milan, Italy) with 100 mL
361 methanol/water (80:20, v/v). The extracts were filtered through Whatman[®] no. 1 filters
362 (Brentford, UK), diluted 1:5 with Dulbecco's Phosphate Buffer Saline (PBS, 1X solution,
363 Sigma Aldrich, St. Louis, MO), filtered again through Whatman[®] glass microfiber filters,
364 Grade 934-AH[®] (Brentford, UK) and subjected to clean-up and purification.

365

366 2.6.2. *Clean-up*

367 The clean-up method for MON was performed with MycoSep[®] 240 Mon clean-up columns
368 (Romer Labs[®], Tulln, Austria). The clean-up procedure (Scarpino et al., 2013) was
369 adapted from the Romer Labs[®] procedure. The cleanup MycoSep[®] columns were pushed
370 into test tubes containing 5 mL of the sample extracts, and the extracts were allowed to
371 filter upwards through the packing material of the columns. The interferences adhered to
372 the chemical packing in the columns and the purified extracts, containing MON, passed
373 through the columns. The purified extracts (1.5 mL) were transferred to HPLC vials and
374 analyzed by means of LC-MS/MS, according to the method described below.

375 The clean-up method for FB (FB₁ and FB₂) was performed with immunoaffinity columns
376 (IA) FumoniTest[™] WB Columns (VICAM[®], USA). Five mL of the sample extracts were
377 loaded and passed through the columns and after a washing step with 2.5 mL of PBS 1 X
378 solution FB were recovered and eluted with 2 mL of methanol. The purified extracts were
379 transferred to HPLC vials and analyzed by means of LC-MS/MS, according to the method
380 described below.

381

382 2.6.3. LC-MS/MS Analysis

383 LC-MS/MS analysis was carried out on a Varian 310 triple quadrupole mass spectrometer
384 (Agilent, Italy) equipped with an electrospray ionization ESI source, a 212 LC pump, a
385 ProStar 410 AutoSampler and dedicated software.

386 For MON analysis LC separation was performed on a 100 mm × 2.1 mm i.d., 3.5 μm, 100
387 Å ZIC[®]-HILIC (Merck, SeQuant, Milan, Italy) column. The mobile phase consisted of water
388 buffered with 100 mM ammonium formate (pH 6.4) (A) and acetonitrile (B) delivered at 200
389 μL min⁻¹. The chromatographic and mass spectrometric parameters of the investigated
390 analytes were described by Scarpino et al. in 2013. The deprotonated molecule ($m/z =$
391 97.0) was fragmented to its product ion ($m/z = 41.0$) and used for quantification and
392 identification purposes.

393 The average percentage of recovery for MON was: 81% (Relative Standard Deviation,
394 RSD%: 9%). The results of the MON concentrations were corrected for the recovery rate.

395 The limit of detection (LOD) and the limit of quantification (LOQ) were 1 μg kg⁻¹ and 4 μg
396 kg⁻¹, respectively.

397 For FB analysis LC separation was performed on a 50 mm × 2.1 mm i.d., 5 μm, Varian
398 Pursuit 5 (Agilent, Italy) C18 column. The mobile phase consisted of water acidified with
399 0,1% CH₃COOH (A) and acetonitrile acidified with 0,1% CH₃COOH (B) delivered at 200
400 μL min⁻¹ for 15 min. Mass spectrometric analyses were performed in the positive-ion
401 mode. For FB₁ MS detection the protonated molecule ($m/z = 722.5$) was fragmented to its
402 product ions ($m/z = 334.2$ and $m/z = 352.3$) and used for quantification and identification
403 purposes.

404 For FB₂ MS detection the protonated molecule ($m/z = 706.5$) was fragmented to its
405 product ions ($m/z = 318.0$ and $m/z = 336.0$) and used for quantification and identification
406 purposes.

407 The average percentage of recovery for FB was: 75% (Relative Standard Deviation,
408 RSD%: 8%). The results of the FB concentrations were corrected for the recovery rate.

409 The limit of detection (LOD) and the limit of quantification (LOQ) were 2 $\mu\text{g kg}^{-1}$ and 8 μg
410 kg^{-1} , respectively.

411

412 2.6.4. Calibration

413 Ten different MON concentrations between 0.93 and 930 $\mu\text{g L}^{-1}$, were prepared for
414 calibration in acetonitrile/water (84:16, v/v). A linear regression was used to obtain the
415 regression curve.

416 Ten different FB concentrations between 1 and 10000 $\mu\text{g L}^{-1}$, were prepared for calibration
417 in acetonitrile/water (50:50, v/v). A linear regression was used to obtain the regression
418 curve.

419

420 2.7. Statistics

421 The normal distribution and homogeneity of variances were verified by performing the
422 Kolmogorov–Smirnov normality test and the Levene test, respectively.

423 An analysis of variance (ANOVA) was utilized to compare the fungal ear rot incidence and
424 severity, the *Fusarium* kernel infection, the *F. verticillioides*, the *F. proliferatum* and
425 *subglutinans* DNA quantification and the MON and FB contamination separately for each
426 year, using a completely randomized block design, in which the natural presence of ECB
427 larva feeding injuries was the independent variable. The incidence and the severity values
428 of fungal ear rot, the *Fusarium* kernel infection, and the *F. verticillioides*, the *F. proliferatum*
429 and *F. subglutinans* DNA quantification were previously transformed using
430 $y' = \arcsin(\sqrt{x} \cdot 180/\pi)$. The concentration of MON and FB was transformed using the
431 $y' = \ln(x+1)$ equation to normalize the residuals.

432 Linear regression analysis was performed for the relationship between the *F. proliferatum*
433 DNA, the *F. subglutinans* DNA, the sum of *F. proliferatum* and *F. subglutinans* DNA and
434 the MON concentration in 24 maize field samples over the three year growing season
435 period (2008-2010). Since the x-values were free to vary and subject to error, the Reduced
436 Major Axis Model II linear regression analysis method was applied to better fit the data
437 (Ludbrook, 2012).

438 Pearson correlation coefficients were obtained for MON and FB, relative to each another
439 and to ECB incidence and severity, fungal ear rot incidence and severity, *Fusarium* kernel
440 infection, *F. verticillioides*, *F. proliferatum* and *F. subglutinans* by joining the data sets that
441 referred to the three growing seasons.

442 An analysis of variance (ANOVA) was used to compare the toxigenic capacity of isolated
443 *F. proliferatum* and *F. subglutinans* strains. In order to apply ANOVA, the toxigenic
444 capacity values were subjected to a rank transformation, a valid alternative bridging
445 parametric and non-parametric analysis proposed by Conover and Iman (1981) when
446 parametric model assumptions do not hold. An REGWQ test was subsequently used as a
447 Post-hoc test.

448 The SPSS Version 21.0 for Windows statistical package, (SPSS Inc., 2012) was used for
449 the statistical analysis.

450

451 **3. Results**

452 **3.1. Meteorological data**

453 The three growing seasons showed remarkably different meteorological trends as far as
454 both rainfall and temperature (expressed as growing degree days, GDDs) from flowering
455 to harvesting are concerned (Table 1). The years 2008 and 2010 were characterized by
456 low GDDs and high rainfall in May and June, and also close to flowering. Heavy rainfall
457 occurred, in particular during the 2010 growing season, from the early milk stage to the
458 harvest. The year 2009 instead had less rainfall during the spring, although it was more
459 concentrated in July, after maize flowering. The average GDDs from June to September
460 were higher in 2009 than those in 2008 and 2010, and this led to an early harvest at the
461 beginning of September (Table 1).

462

463 **3.2. ECB damage**

464 Table 2 summarizes the mean ECB incidence and severity recorded for each growing
465 season for the natural infestation of ears collected for each sampling year. The ears
466 collected in the plots protected with entomological nets were free from ECB attack, in each
467 growing season, while those collected in the plots subject to natural insect attacks showed
468 a variable damage severity that depended on the insect pressure in each growing season.
469 The samples collected in the 2008 and 2010 growing seasons showed more higher ECB
470 damage, while the ECB severity recorded in 2009 was more than 3 times lower than in
471 2008 and 2010 (Table 2).

472

473

474

475 3.3. *Mycological measurements in the field samples*

476 The presence of ECB larvae significantly affected the fungal ear rot incidence and severity
477 in each growing season (Table 3). The ECB attack increased the fungal ear rot severity
478 22-fold and 153-fold in the 2008 and 2010 growing seasons, respectively. Conversely, in
479 2009, the year characterized by the lowest ECB pressure, the fungal ear rot severity only
480 increased 3-fold. The *Fusarium* kernel infection was significantly higher in the natural
481 infected ears in each growing season, according to the ECB attack (Table 3).

482 As far as DNA quantification is concerned, the *F. proliferatum* occurrence significantly
483 increased in the natural infection conditions of 29-fold, 15-fold and 21-fold in the 2008,
484 2009 and 2010 growing seasons, respectively, compared to the artificial control. The
485 insect activity on the ears increased the infection of *F. subglutinans* of 23-fold, 19-fold and
486 25-fold in the 2008, 2009 and 2010 growing seasons, respectively, although its effect was
487 only significant in the first two years (Table 3). The ECB activity significantly increase *F.*
488 *verticillioides* infection in all the growing seasons.

489

490 3.4. *FB and MON contamination in the field samples*

491 The MON contamination was affected significantly by the ECB larva feeding activity on the
492 maize ears in all the considered growing seasons ($P = 0.002$, 2008; $P = 0.002$, 2009; $P <$
493 0.001 , 2010) (Table 3). The presence of ECB damage increased the MON content 26-fold
494 in 2008 from 43 to 1137 $\mu\text{g kg}^{-1}$, 25-fold in 2009 from $\exists < \text{LOQ}$ to 77 $\mu\text{g kg}^{-1}$ and 94-fold in
495 2010 from 6 to 564 $\mu\text{g kg}^{-1}$. Moreover, as reported in Table 3 the ECB larva feeding
496 activity on the maize ears significantly affected the FB contamination in the years with the
497 higher ECB pressure, 2008 and 2010.

498 As far as the relationships between the major *Fusarium* species producers of MON and
499 mycotoxin contamination are concerned, the amount of *F. proliferatum* DNA (Fig. 1), the

500 amount of *F. subglutinans* DNA (Fig. 2) and the sum of the amounts of the 2 previous
501 *Fusarium* species DNA (Fig. 3) showed a significant linear relationship with the MON
502 content.

503 Moreover, as reported in Table 4, MON and FB contamination, fungal ear rot incidence
504 and severity, *Fusarium* kernel infection, *F. verticillioides*, *F. proliferatum* and *F.*
505 *subglutinans* were significantly correlated to each other and to ECB damage parameters,
506 incidence and severity.

507 As can be seen in Fig. 1 and Fig. 2, the relationship between the MON content and the
508 DNA of the *Fusarium* species producers of MON was closer to *F. proliferatum* ($r = 0.96$, P
509 < 0.001) than to *F. subglutinans* ($r = 0.83$, $P < 0.001$). Moreover, as shown in Fig. 3, the
510 sum of the DNA of the 2 previous *Fusarium* species did not significantly improve the
511 relationship with the MON content ($r = 0.97$, $P < 0.001$), but respect to the relationship
512 between the MON content and the DNA of the only *F. subglutinans* r value was slightly
513 improved.

514

515 3.5. *In vitro* MON production by monoconidial strains of *F. verticillioides*, *F.* 516 *proliferatum* and *F. subglutinans*.

517 The results related to the toxigenic capacity of the different strains of *F. proliferatum* and *F.*
518 *subglutinans* tested during the *in vitro* assay are summarized in ~~Table 4~~ Table 5. As can
519 be seen in this table, the different strains have significantly different toxigenic capacity ($P <$
520 0.001).

521 On average, the *F. proliferatum* strains showed a significantly higher ($P < 0.001$) toxigenic
522 capacity than the strains of *F. subglutinans*. Two strains of *F. proliferatum* (n° 1 and 2)
523 showed the highest toxigenic capacity, which was significantly higher than all of the
524 analyzed *F. subglutinans* strains. Only *F. subglutinans* strains 6 and 7 showed a

525 comparable toxigenic capacity with the other *F. proliferatum* strains, while two *F.*
526 *subglutinans* strains (9 and 10) showed an extremely low toxigenic capacity towards this
527 mycotoxin. On the other hand, the 5 *F. verticillioides* strains tested in this assay have
528 never produced MON (the contamination was always under the LOQ).

529

530 **4. Discussions and Conclusions**

531 The results related to the MON content in maize fields sampled over 3 growing seasons
532 show that the contamination of this mycotoxin is promoted by a complex and integrated
533 system in which the meteorological conditions, the ECB activity and the *Fusarium* species
534 all play important roles.

535 The collected data underline a clear increase, due to ECB feeding activity on the maize
536 ears, not only of the FB content, as widely reported in literature, but also of the MON
537 content, a mycotoxin, which like FB, is produced by *Fusarium* spp. of the *Liseola* section.
538 Thus, the relationship between MON contamination and ECB injuries is extremely high in
539 temperate areas.

540 The results of the present work point out a significant effect of ECB larva activity on the
541 occurrence of both *F. proliferatum* and *F. subglutinans* species. Munkvold et al. (1997) and
542 Gatch and Munkvold (2002) reported an important increase in maize ear and stalk rot
543 caused by *F. verticillioides*, *F. proliferatum* and *F. subglutinans*, all of which are species of
544 the *Liseola* section, following ECB larva feeding damage. In a field experiment conducted
545 in Austria, Lew et al. (1991) reported that the increase in MON contamination in maize
546 ears damaged by ECB was related to a higher colonization of the grains by *F.*
547 *subglutinans*.

548 Although MON contamination has been shown to be closely linked to ECB activity, the
549 meteorological and climatic conditions are also contributing factors that have the effect of
550 changing the intensity of the MON contamination. The average MON content was higher in
551 the 2008 and 2010 growing seasons, years characterized by lower GDD from the
552 flowering to the end of ripening than 2009. Furthermore, in the 2010 growing season, the
553 MON occurrence in maize grain was increased remarkably by ECB (93-fold), as this
554 mycotoxin was only found in traces in the insect protected plot. In 2008, the insect

555 protected plot showed an average contamination of 43 $\mu\text{g kg}^{-1}$, which increased 26-fold in
556 the ears naturally infected by the insect. On the other hand, in the 2009 experiment, with a
557 lower natural ECB pressure and warmer and drier climatic conditions from flowering to
558 harvest, the occurrence of MON was the lowest, in both the damaged and undamaged
559 ears.

560 As reported by Marin et al. (2001) and by Samapundo et al. (2005), this mycotoxin
561 synthesis is affected to a great extent by the temperature and by rainfall, which could
562 influence both the growth rates of the fungi and the mycotoxin production. Lew et al.
563 (2001) observed that a significant increase in *F. proliferatum* infection occurred in the
564 nineties in Austria, due to changed climatic conditions, with milder and more humid winters
565 followed by drier and warmer summers, which favored the progress of *F. proliferatum*.

566 Both of the *Fusarium* species responsible for MON contamination can be found throughout
567 the world, but as reported by Doohan et al. (2003), the optimum temperature for the
568 growth of *F. subglutinans* in laboratory conditions is lower (Temperature = 15-25°C,
569 Castellá et al., 1999) than that of *F. proliferatum* (Temperature = 30°C, Marin et al., 1995).

570 As far as the European distribution and prevalence of the two different *Fusarium* species
571 that cause MON production is concerned, *F. proliferatum* is more common in southern
572 European areas, while it is substituted by *F. subglutinans* in central areas (Logrieco et al.,
573 2002). As reported in literature, *F. proliferatum* is widely present in Italy (Logrieco et
574 al., 1995) and in nearby European countries (Lević et al., 1997; Srobárová, 1997), in
575 association with *F. verticillioides*, but the occurrence of *F. proliferatum* has rarely been
576 recorded in Austria (Krüger, 1989; Krska et al., 1997), Croatia (Jurjević et al., 1997),
577 Slovakia (Piecková and Jesenská, 1997), Hungary (Szécsi, 1994) and Poland (Kostecki et
578 al., 1995) to the advantage of *F. subglutinans*. A remarkable presence of *F. temperatum*,
579 which is able to produce MON, has recently been reported in Belgium by Scauflaire et al.
580 (2012); but, because of the low number of strains examined, the authors underlined the

581 necessity of extending their screening before assessing the importance of this new
582 species among the MON producer *Fusaria* species.

583 Until now, it has been assumed that the MON contamination of maize in Europe is mainly
584 associated with an *F. subglutinans* infection (Lew et al., 1991; Kostecki et al., 1995; 1997);
585 no other works that have investigated in field conditions the relationship between *F.*
586 *proliferatum* infection and MON production are present in the literature.

587 The results obtained in the current study underline that *F. proliferatum* could be the main
588 agent responsible for MON production in maize grain in the temperate maize growing
589 areas, such as in North-West Italy, where there is a high ECB pressure. In the *in vitro*
590 study on *F. proliferatum*, the collected strains showed a lower infection growth rate on the
591 maize substrate compared to those of *F. subglutinans*, but the toxigenicity capacity of the
592 *F. proliferatum* strains was about 3600 times higher. This finding appears to be in contrast
593 with what Logrieco et al. (2002) reported, that is, that the toxigenicity of *F. proliferatum* in
594 Europe seemed comparable with that of *F. subglutinans*, considering the *Fusarium*
595 species from maize ear rot. However, Logrieco et al. (1995) reported that some strains of
596 *F. proliferatum* from pre-harvest maize ear rot in Italy were able to produce very large
597 quantities of MON on autoclaved maize. In fact, as underlined by our results the *F.*
598 *proliferatum* strain 1 was able to produce a very large amount of MON compared to the
599 others and this behavior could be due to the natural variability among strains of this
600 species. Several authors report the presence in the natural population of *F. proliferatum* of
601 strains with different capabilities in MON synthesis (Logrieco and Bottalico 1988; Logrieco
602 et al., 1995; Vesonder et al., 2000).

603 The *F. proliferatum* infection of maize grain in the field experiments was higher than that
604 caused by *F. subglutinans* in each growing season. Moreover, the relationship between
605 the MON content and the DNA quantification of the *Fusarium* species producers of MON
606 was closer for the *F. proliferatum* species than for *F. subglutinans*.

607 In conclusion, the combination of the high toxigenic capacity of *F. proliferatum*, in the
608 considered experimental conditions, and its more frequent occurrence and greater
609 intensity in the field make it possible to state that the production of MON in the temperate
610 maize areas, such as the North Italian maize areas investigated during this work, is mainly
611 due to *F. proliferatum* infections. Thus, owing to the close link between MON
612 contamination in kernels and injuries caused by ECB larvae on maize ears, any of the
613 strategies that can control this insect could also reduce the contamination of this
614 mycotoxin. Further studies are required to evaluate whether an the integrated field
615 program that is able to minimize the FB content in maize could also lead to a comparable
616 control of MON.

617

618 **References**

- 619 Blandino, M., Reyneri, A., Vanara, F., Pascale, M., Haidukowski, M., Campagna, C., 2009.
620 Management of fumonisin contamination in maize kernels through the timing of
621 insecticide application against European corn borer. Food Addit. Contam. A. 26(11),
622 1501-1514.
- 623 Bode, W. M., Calvin, D. D., 1990. Yield-Loss Relationships and Economic Injury Levels for
624 European Corn Borer (Lepidoptera : Pyralidae) Populations Infesting Pennsylvania
625 Field Corn. J. Econ. Entomol. 83(4), 1595-1603.
- 626 Bottalico A., 1998. *Fusarium* diseases of cereals: species complex and related mycotoxin
627 profiles. J. Plant Pathol. 80, 85-103.
- 628 Boutignya, A.-L., Beukes, I., Smalla, I., Zuñhke, S., Spiteller, M., Van Rensburg, B. J.,
629 Flett, B., Viljoen, A., 2012. Quantitative detection of *Fusarium* pathogens and their
630 mycotoxins in South African maize. Plant Pathol. 61, 522-531.
- 631 Castellá, G., Munkvold, G.P., Imerman, P., Hyde, W.G., 1999. Effects of temperature,
632 incubation period and substrate on the production of fusaproliferin by *Fusarium*
633 subglutinans ITEM 404. Nat. Toxins 4, 129-132.
- 634 EFSA Request for a scientific opinion on the risks for public health related to the presence
635 of moniliformin in feed and food, Mandate M-2010-0312, Reception Date 21-07-
636 2010, Acceptance Date 09-09-2010.
- 637 D'Mello, J. P. F., Placinta, C. M., Macdonald, A. M. C., 1999. *Fusarium* mycotoxins: A
638 review of global implications for animal health, welfare and productivity. Anim. Feed
639 Sci. Tech. 80, 183-205.
- 640 Doohan, F. M., Brennan, J., Cooke, B. M., 2003. Influence of climatic factors on *Fusarium*
641 species pathogenic to cereals. Eur. J. Plant Pathol. 109, 755-768.

642 European Commission, 2006. Commission regulation No. 1881/2006, of 10 December
643 2006 setting maximum levels for certain contaminants in food stuff. Official Journal
644 of the European Union L 364, pp. 5-24.

645 European Commission, 2007. Commission regulation No. 1126/2007, of 28 September
646 2007 amending Regulation (EC) No 1881/2006 setting maximum levels for certain
647 contaminants in foodstuffs as regards *Fusarium* toxins in maize and maize products
648 L 255, pp. 14-17.

649 FDA - Food and Drug Administration, 2000, Guidance for Industry-Fumonisin Levels in
650 Human Foods and Animal Feeds. Draft Guidance, June 6th, pp.1-3.

651 Gatch, E. W., Munkvold, G. P., 2002. Fungal Species Composition in Maize Stalks in
652 Relation to European Corn Borer Injury and Transgenic Insect Protection. Plant
653 Dis. 86(10), 1156-1162.

654 Henrion, B., Le Tacon, F., Martin, F., 1992. Rapid Identification of Genetic Variation of
655 Ectomycorrhizal Fungi by Amplification of Ribosomal RNA Genes. New Phytol.,
656 122, 289-298.

657 International Agency for Research on Cancer (IARC), 2002. Fumonisin B1. IARC
658 Monographs on the Evaluation of Carcinogenic Risks to Humans, Some Traditional
659 Herbal Medicines, Some Mycotoxins, Naphthalene and Styrene, Vol. 82 (Lyon:
660 IARC), pp. 301-366.

661 Jonsson, M., Jestoi, M., Nathanail, A. V., Kokkonen, U.-M., Anttila, M., Koivisto, P.,
662 Karhunen, P., Peltonen, K., 2013. Application of OECD Guideline 423 in assessing
663 the acute oral toxicity of moniliformin. Food Chem. Toxicol. 53, 27-32.

664 Jurado, M., Vazquez, C., Marin, S., Sanchis, V., Gonzalez-Jaen, M. T., 2006. PCR-based
665 strategy to detect contamination with mycotoxigenic *Fusarium* species in maize.
666 Syst. Appl. Microbiol. 29, 681-689.

667 Jurjević, Z., Cvjetković, B., Jurjević, V., Ceović, S., 1997. Occurrence of *Fusarium* spp. on
668 corn in the region of endemic nephropathy in Croatia. *Cereal Res. Commun.* 25,
669 455-456.

670 Kostecki, M., Szczesna, J., Chelkowski, J., Wisniewska, H., 1995. Beauvericin and
671 moniliformin production by Polish isolates of *Fusarium subglutinans* and natural co-
672 occurrence of both mycotoxins in cereal grain samples. *Microbiologie, Aliments,*
673 *Nutrition* 13, 67-70.

674 Kostecki, M., Grabarkiewicz-Szczesna, J., Golinski, P., 1997. Simultaneous analysis of
675 beauvericin and moniliformin in fungal cultures and in cereal grain samples.
676 *Mycotoxin Res.* 13, 17-22.

677 Krska, R., Schuhmacher, R., Grasserbauer, M., Lemmens, M., Lemmens-Gruber, R.,
678 Adler, A., Lew, H., 1997. Effects of beauvericin to mammalian tissue and its
679 production by Austrian isolates of *Fusarium proliferatum* and *Fusarium*
680 *subglutinans*. *Mycotoxin Res.* 13, 11-16.

681 Krüger, W., 1989. Maize diseases caused by Fusaria: Involved species and mycotoxins.
682 In: Chelkowski J. (ed.) *Fusarium – Mycotoxins, Taxonomy and Pathogenicity.*
683 Elsevier, Amsterdam pp. 297-317.

684 Lancashire, P.D., Bleiholder, H., Van Den Boom, T., Langeluddeke, P., Stauss, R., Weber,
685 E., Witzemberger, A., 1991. A uniform decimal code for growth stages of crops and
686 weeds. *Ann. Appl. Biol.* 119, 561-601.

687 Levíc, J., Tamburić-Ilićić, L., Petrović, T., 1997. Maize kernel infection by Fusaria in the
688 period 1994-1996. *Cereal Res. Commun.* 25, 773-775.

689 Lew, H., Adler, A., Edinger, W., 1991. Moniliformin and the European maize borer
690 (*Ostrinia nubilalis*). *Mycotoxin Res.* 7A, 71-76.

691 Lew, H., Adler, A., Edinger, W., Brodacz, W., Kiendler, E., Hinterholzer, J., Oberforster, M.,
692 2001. Shifting of *Fusarium* species and their toxins in Austrian cereals. *Mycotoxin*
693 *Res.* 17(1), 1-4.

694 Logrieco, A., Bottalico, A., 1988. *Fusarium* species of the *Liseola* section associated with
695 stalk and ear rot maize in Southern Italy, and their ability to produce moniliformin.
696 *Trans. Br. Mycol. Soc.* 90, 215-219.

697 Logrieco, A., Moretti, A., Ritieni, A., Bottalico, A., Corda, P., 1995. Occurrence and
698 toxigenicity of *Fusarium proliferatum* from pre-harvest maize ear rot, and associated
699 mycotoxins, in Italy. *Plant Dis.* 79, 727-731.

700 Logrieco, A., Mulè, G., Moretti, A., Bottalico, A., 2002. Toxigenic *Fusarium* Species and
701 Mycotoxins Associated with Maize Ear Rot in Europe. *Eur. J. Plant Pathol.* 108,
702 597-609.

703 Lozzia, G. C., Manachini, B., 2003. Susceptibility of *Ostrinia nubilalis* to *Bacillus*
704 *thuringiensis* var . *kurstaki*. *B. Insectol.* 56(2), 215-219.

705 Ludbrook, J., 2012. A primer for biomedical scientists on how to execute Modell II linear
706 regression analysis. *Clin. Exp. Pharmacol. Physiol.* 39, 329-335.

707 Magg, T., Bohn, M., Klein, D., Merditaj, V., Melchinger, A. E., 2003. Concentration of
708 moniliformin produced by *Fusarium* species in grains of transgenic *Bt* maize hybrids
709 compared to their isogenic counterparts and commercial varieties under European
710 maize borer pressure. *Plant Breeding* 122, 322-327.

711 Marín, S., Sanchis, V., Magan, N., 1995. Water activity, temperature and pH effects on
712 growth of *Fusarium moniliforme* and *Fusarium proliferatum* isolates from maize.
713 *Can. J. Microbiol.* 41, 1063-1070

714 Marín, S., Albareda, X., Ramos, A. J., Sanchis V., 2001. Impact of environment and
715 interactions of *Fusarium verticillioides* and *Fusarium proliferatum* with *Aspergillus*

716 parasiticus on fumonisin B1 and aflatoxins on maize grain. J. Sci. Food Agric.
717 81(11), 1060-1068.

718 Mulè, G., Susca, A., Stea, G., Moretti, A., 2004. A species-specific PCR assay based on
719 the calmodulin partial gene for identification of *Fusarium verticillioides*, *F.*
720 *proliferatum* and *F. subglutinans*. Eur. J. Plant Pathol. 110, 495-502.

721 Munkvold, G.P., McGee, D.C., Carlton, W.M., 1997. Importance of different pathways for
722 maize kernel infection by *Fusarium moniliforme*. Phytopathology 87, 209-217.

723 Munkvold, G.P., Hellmich, R.L., Showers, W.B., 1997. Reduced fusarium ear rot and
724 symptomless infection in kernels of maize genetically engineered for European corn
725 borer resistance. Phytopathology 87, 1071-1077.

726 Nelson, P.E., Toussoun, T.A., Marasas, W.F.O., 1983. *Fusarium* species: an illustrated
727 manual for identification. Pennsylvania State University, University Park.

728 Nicolaisen, M., Supronienė, S., Nielsen, L. K., Lazzaro, I., Spliid, N. H., Justesen, A. F.,
729 2009. Real-time PCR for quantification of eleven individual *Fusarium* species in
730 cereals. J. Microbiol. Methods 76 234-240.

731 Nutz, S., Döll, K., Karlovsky, P., 2011. Determination of the LOQ in real-time PCR by
732 receiver operating characteristic curve analysis: application to qPCR assays for
733 *Fusarium verticillioides* and *F. proliferatum*. Anal. Bioanal. Chem. 401, 717-726.

734 Papst, C., Utz, H. F., Melchinger, A. E., Eder, J., Magg T., Klein, D., Bohn, M., 2005.
735 Mycotoxins produced by *Fusarium* spp. in Isogenic Bt vs. non-Bt maize hybrids
736 under European maize borer pressure. Agron. J. 97, 219-224.

737 Piecková, E., Jesenká, Z., 1997. *Fusarium moniliforme* and *F. subglutinans* in maize-
738 based foodstuffs in the Slovak Republic. Cereal Res. Commun. 25, 609-610.

739 Samapundo, S., Devlieghere, F., De Meulenaer, B., Debevere, J., 2005. Effect of Water
740 Activity and Temperature on Growth and the Relationship between Fumonisin

741 Production and the Radial Growth of *Fusarium verticillioides* and *Fusarium*
742 *proliferatum* on Corn. J. Food Prot. 68(5), 1054-1059.

743 Sanhueza, C. E. P., Degrossi, M. C., 2004. Moniliformin, a *Fusarium* mycotoxin. Revista
744 Mexicana de Micologia 19, 103-112.

745 Scarpino, V., Blandino, M., Negre, M., Reyneri, A., Vanara, F., 2013. Moniliformin analysis
746 in maize samples from North-West Italy using multifunctional clean-up columns and
747 the LC-MS/MS detection method. Food Addit. Contam. A 30(5), 876-884.

748 Scauflaire, J., Gourgue, M., Munaut, F., 2011. *Fusarium temperatum* sp. nov. from maize,
749 an emergent species closely related to *Fusarium subglutinans*. Mycologia 103, 586-
750 597.

751 Scauflaire, J., Gourgue, M., Callebaut, A., Munaut, F., 2012. *Fusarium temperatum*, a
752 mycotoxin-producing pathogen of maize. Eur. J. Plant. Pathol. 133, 911-922.

753 Sobek, E.A., Munkvold, G.P., 1999. European Corn Borer (Lepidoptera: Pyralidae) Larvae
754 as Vectors of *Fusarium moniliforme*, Causing Kernel Rot and Symptomless
755 Infection of Maize Kernels. J. Econ. Entomol. 92, 503-509.

756 Srobárová, A., 1997. Occurrence of *Fusarium* spp. from Slovakia maize kernels. Cereal
757 Res. Commun. 25, 617-618.

758 Srobárová, A., Moretti, A., Ferracane, R., Ritieni, A., Logrieco, A., 2002. Toxigenic
759 *Fusarium* species of *Liseola* section in pre-harvest maize ear rot , and associated
760 mycotoxins in Slovakia. Eur. J. Plant Pathol. 108, 299-306.

761 Szécsi, A., 1994. Occurrence of *Fusarium* species in section *Liseola* isolated from
762 Hungarian corn samples in 1991 and 1992. Növényvédelem 30, 313-318.

763 Szóke, C., Zsubori, Z., Pók, I., Rácz, F., Illés, O., Szegedi, I., 2002. Significance of the
764 european corn borer (*Ostrinia nubilalis* Hübn.) in maize production. Acta
765 Agronomica Hungarica 50, 447-461.

766 Vesonder, R. F., Wu, W., Weisleder, D., Gordon, S. H., Krick, T., Xie, W., Abbas H. K.,
767 McAlpin, C. E., 2000 . Toxigenic strains of *Fusarium moniliforme* and *Fusarium*
768 *proliferatum* isolated from dairy cattle feed produce fumonisins, moniliformin and a
769 new C₂₁H₃₈N₂O₆ metabolite phytotoxic to *Lemna minor* L. J. Nat. Toxins 9(2), 103-
770 112.
771

772 **TABLES**

773

774 **Table. 1.** Total rainfall, rainy days, relative humidity and growing degree days (GDD) from

775 June to October 2008-2010 at the research site.

Year	Month	Rainfall (mm)	Rainy days	GDD ^a (°C d ⁻¹)
2008	May	121	16	204
	June	95	17	304
	July	63	8	382
	August	52	6	372
	September	57	8	228
	October	30	5	151
	May-October	418	60	1641
2009	May	30	10	292
	June	26	7	341
	July	121	8	391
	August	56	11	404
	September	62	8	273
	October	54	6	163
	May-October	349	50	1864
2010	May	117	12	214
	June	192	11	332
	July	37	8	420
	August	116	11	354
	September	51	12	240
	October	105	9	120
	May-October	618	63	1680

776 ^a Accumulated growing degree days for each month using a 10°C base.777 Source: weather station in the experimental farm of the Department of Agricultural, Forest and Food
778 Science, University of Turin located in Carmagnola.

779

780

781 **Table 2.** Natural ECB infestation recorded per year, for the field experiments conducted at
782 Carmagnola (TO) in the 2008 - 2010 period.

Year	ECB incidence ^a	ECB severity ^b
2008	100.0	25.7
2009	60.0	5.6
2010	88.9	20.8

783

784 ^a ECB incidence was calculated as the percentage of ears with symptoms, based on 4 replications of 30 ears
785 each.

786 ^b ECB severity was calculated as the mean percentage of kernels with symptoms per ear, based on 4
787 replications of 30 ears each.

788

789 **Table 3.** Effect of ECB infestation on fungal ear rot incidence and severity, *Fusarium* kernel infection, *F. verticillioides*, *F. proliferatum*
 790 and *F. subglutinans* occurrence and FB (FB₁ + FB₂) and MON contaminations; field experiments conducted at Carmagnola (TO) in the
 791 2008 - 2010 period.

Year	ECB infestation	Fungal ear rot incidence ^c		Fungal ear rot severity ^d		<i>Fusarium</i> kernel infection		<i>F. verticillioides</i> infection ^e		<i>F. proliferatum</i> infection ^e		<i>F. subglutinans</i> infection ^e		FB		MON	
		T	N (%)	T	N (%)	T	N (%)	T	N (Normalized DNA)	T	N (Normalized DNA)	T	N (Normalized DNA)	T	N (µg kg ⁻¹)	T	N (µg kg ⁻¹)
2008	Natural	86.2	98.3	27.9	22.0	65.4	82.4	15.6	75.0	14.6	68.3	6.4	12.6	8.5	4817	7.0	1137
	Artificial control	32.9	29.8	5.6	1.0	41.6	44.1	5.96	12.76	2.29	2.32	1.20	0.54	7.6	2036	3.3	43
	<i>P</i> (F) ^a	< 0.001***		< 0.001***		< 0.001***		0.006**		0.004**		< 0.001***		0.001**		0.002**	
	sem ^b	5.4		2.1		4.3		4.06		4.64		0.82		0.3		0.7	
2009	Natural	48.0	55.0	8.8	2.4	59.7	71.2	3.76	4.61	3.70	4.76	2.26	1.71	7.2	1745	4.1	77
	Artificial control	21.1	13.3	4.5	0.7	25.6	20.3	0.83	0.25	0.87	0.31	0.51	0.09	5.5	298	1.4	<LOQ ^f
	<i>P</i> (F) ^a	< 0.001***		0.014*		0.012*		0.013*		0.043*		0.019*		0.090		0.002**	
	sem ^b	5.4		1.5		11.8		0.80		1.12		0.53		0.9		0.5	
2010	Natural	73.4	91.7	22.9	15.3	56.3	68.7	15.55	84.58	9.48	31.85	1.99	2.54	8.8	7024	6.2	564
	Artificial control	7.5	3.3	1.2	0.1	32.3	29.3	3.44	4.54	2.00	1.54	0.56	0.10	5.7	644	1.6	6
	<i>P</i> (F) ^a	< 0.001***		< 0.001***		0.009**		0.037*		0.031*		0.298		0.016*		< 0.001***	
	sem ^b	5.6		1.4		7.9		6.29		3.67		1.81		1.3		0.6	

792 The reported data are the averages of 4 replications.

793 ^a *P* (F) = ANOVA level of significance, * *P* < 0.05, ** *P* < 0.01, *** *P* < 0.001.

794 ^b sem: standard error of mean.

795 ^c Fungal ear rot incidence was calculated as the percentage of ears with symptoms, based on 4 replications of 20 ears each.

796 ^d Fungal ear rot severity was calculated as the mean percentage of kernels with symptoms per ear, based on 4 replications of 20 ears each.

797 ^e *F. verticillioides*, *F. proliferatum* and *F. subglutinans* infections were calculated as the pg Fungal DNA / ng Total DNA ratio. The amounts of fungal DNA obtained
798 were normalized to the total DNA amount extracted from the meal samples.

799 ^f LOQ = Limit of Quantification of the LC-MS/MS analytical method = 4 µg kg⁻¹.

800 The fungal ear rot incidence and severity, the *Fusarium* kernel infection, the *F. verticillioides*, *F. proliferatum* and *F. subglutinans* occurrence means reported are
801 transformed (T; $y' = \arcsin \sqrt{x} * 180/\pi$) and not transformed (N) values. The FB and MON contamination means reported are transformed [T; $y' = \ln(x + 1)$] and not
802 transformed (N) values.

803

804

805

806

807

808

809

810

811

812 **Table 4.** Correlation matrix between ECB incidence and severity, fungal ear rot incidence and severity, *Fusarium* kernel infection, *F.*
 813 *verticillioides*, *F. proliferatum* and *F. subglutinans* infections and FB and MON contaminations in maize kernels in the 2008 - 2010
 814 period.

Correlation	ECB severity	Fungal ear rot incidence	Fungal ear rot severity	<i>Fusarium</i> kernel infection	<i>F. verticillioides</i> infection	<i>F. proliferatum</i> infection	<i>F. subglutinans</i> infection	FB	MON
ECB incidence	0.970**	0.965**	0.922**	0.803**	0.798**	0.813**	0.825**	0.822**	0.849**
ECB severity		0.956**	0.967**	0.774**	0.845**	0.848**	0.863**	0.841**	0.892**
Fungal ear rot incidence			0.926**	0.838**	0.816**	0.800**	0.811**	0.849**	0.837**
Fungal ear rot severity				0.759**	0.802**	0.877**	0.893**	0.746**	0.888**
<i>Fusarium</i> kernel infection					0.634**	0.660**	0.699**	0.754**	0.700**
<i>F. verticillioides</i> infection						0.584**	0.621**	0.751**	0.604**
<i>F. proliferatum</i> infection							0.761**	0.749**	0.965**
<i>F. subglutinans</i> infection								0.587**	0.829**
FB									0.767**

815 (**) correlation significant at $P \leq 0.01$. The data reported in the table are Pearson product-moment correlation coefficients.

816 **Table 5.** In vitro MON production by different fungal strains of *Fusarium proliferatum* and
 817 *F. subglutinans* on maize substrates.

Fungal species	Fungal strain	Normalized DNA ^c	MON (µg kg ⁻¹)	MON / Normalized DNA ^d
		Average ± SD ^a	Average ± SD ^a	Average
<i>Fusarium proliferatum</i>	1	0.015 ± 0.001	699 ± 1	47383.5 a
	2	3.635 ± 0.895	2605 ± 338	738.4 a
	3	0.735 ± 0.071	154 ± 20	214.5 ab
	4	5.113 ± 1.547	411 ± 156	98.7 abc
	5	0.578 ± 0.067	< LOD ^b	1.6 cd
<i>Fusarium subglutinans</i>	6	139.041 ± 3.000	1360 ± 1030	9.9 bcd
	7	64.068 ± 1.612	110 ± 53	1.7 cd
	8	77.760 ± 39.253	61 ± 11	1.2 de
	9	70.399 ± 7.425	24 ± 3	0.3 ef
	10	60.826 ± 19.564	11 ± 4	0.2 f

818

819 The reported data are the averages of 3 replications.

820 ^a SD = standard deviation.

821 ^b LOD = Limit of Detection of the LC-MS/MS analytical method = 1 µg kg⁻¹.

822 ^c Normalized DNA = pg Fungal DNA / ng Total DNA.


823 ^d MON / DNA = Ratio of the MON concentration and Fungal DNA normalized for the Total DNA.

824 Means followed by different letters are significantly different (P < 0.001). The REGWQ test was used as a
 825 Post-hoc test.

826

827 **FIGURES**

828



829

830 **Figure 1.** Relationship between the *F. proliferatum* DNA and the MON concentration in 24
831 maize field samples over three growing season periods (2008-2010).

832 The amount of *F. proliferatum* DNA was determined by means of real-time PCR. The MON concentration
833 was determined using an LC-MS/MS method.

834 Reduced Major Axis linear regression analysis (Model II linear regression analysis) was applied to fit the
835 data; r: Pearson product-moment correlation coefficient; P: level of significance, * P < 0.05, ** P < 0.01, *** P
836 < 0.001.

837

838

839

840

841

842

843


844

845

846

847

848



849

850 **Figure 2.** Relationship between the *F. subglutinans* DNA and the MON concentration in 24
851 maize field samples over three growing season periods (2008-2010).

852 The amount of *F. subglutinans* DNA was determined by means of real-time PCR. The MON concentration
853 was determined using an LC-MS/MS method.

854 Reduced Major Axis linear regression analysis (Model II linear regression analysis) was applied to fit the
855 data; r: Pearson product-moment correlation coefficient; P: level of significance, * P < 0.05, ** P < 0.01, *** P
856 < 0.001.

857

858

859

860

861

862

863

864

865


866

867

868

869

870



871

872 **Figure 3.** Relationship between the sum of *F. proliferatum* and *F. subglutinans* DNA and
873 the MON concentration in 24 maize field samples over three growing season periods
874 (2008-2010).

875 The amount of *F. proliferatum* and *F. subglutinans* DNA was determined by means of real-time PCR. The
876 MON concentration was determined using an LC-MS/MS method.

877 Reduced Major Axis linear regression analysis (Model II linear regression analysis) was applied to fit the
878 data; r: Pearson product-moment correlation coefficient; P: level of significance, * P < 0.05, ** P < 0.01, *** P
879 < 0.001.

880

881

882

883