

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Risk, prevalence, and impact of hospital malnutrition in a Tertiary Care Referral University Hospital: a cross-sectional study

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1689896> since 2019-02-04T22:55:20Z

Published version:

DOI:10.1007/s11739-018-1987-7

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

Risk, prevalence, and impact of hospital malnutrition in a Tertiary Care Referral University

Hospital: a cross-sectional study

Concetta Finocchiaro¹, Giovanni Fanni², Simona Bo²

1. Unit of Clinical Nutrition, Hospital of Turin, Città della Salute e della Scienza, Turin, Italy
2. Department of Medical Sciences, University of Turin, Turin, Italy

Corresponding author: Simona Bo, c.so AM Dogliotti 14, 10126, Turin, Italy

Tel. 0039 11 6336036; e-mail: simona.bo@unito.it

Conflict of interest: The Authors declare that they have no conflict of interest

Malnutrition is a high-impacting condition in terms of public health. Hospital malnutrition (i.e. malnutrition in a healthcare setting) has been demonstrated to affect several major clinical outcomes, such as immune competence, wound healing, quality of life, risk of hospital infections, length of stay, risk of readmission, overall survival, and to be a striking economic burden for the health care system [1–6]. Still now, hospital malnutrition has a high prevalence rate all over the world that varies from 5% to 60% according to patients' clinical condition, geographical region, healthcare setting or hospital department and, remarkably, diagnosis criteria [1–7]. A gap between the number of patients screened and diagnosed with malnutrition and receiving nutritional recommendations and those effectively malnourished remains, being this latter number much higher [8]. Indeed, scarce awareness and a low attention to this issue are present among healthcare professionals and governments [9, 10]. Providing nutritional care through early and appropriate nutritional assessment and intervention has been shown to improve outcomes, particularly among high-risk elderly hospitalized patients. However, recent meta-analyses concluded that there is low-quality evidence for the effects of nutritional support on mortality and serious adverse events [11], though a reduction in healthcare costs indeed [12].

According to the European Society for Parental and Enteral Nutrition (ESPEN), malnutrition consists in both over- and under-nutrition; as regards undernutrition, the differences between cachexia

(severe body weight, fat and muscle loss with increased protein catabolism due to underlying diseases), sarcopenia (loss of muscle mass and function) and malnutrition have been recently highlighted [13]. Malnutrition in hospitalized patients is often a combination of cachexia (disease-related chronic malnutrition with inflammation), acute disease- or injury-related malnutrition and inadequate nutrient consumption. Contributors to this complex interplay might be disease-related acute or chronic metabolic alterations, the reduced availability of nutrients (reduced intake and/or impaired absorption and/or increased losses) and increased energy requirements [1, 13]. Therefore, the nutritional status often worsens during hospitalization.

The impact of in-hospital stay on the individual nutritional status is extremely difficult to be evaluated, since the specific patient clinical status might act as a major confounder.

An interesting insight into the problem has recently been provided by Rinninella, who studied whether the hospitalization itself could be detrimental in malnourished patients [14]. This work is a sub-group analysis nested on an Italian cross-sectional study aimed at estimating risk and prevalence as well as the impact of malnutrition in a cohort of patients from a large Italian hospital [14]. A within-patient comparison of anthropometrical measures, bioelectrical impedance (BIA) values and laboratory data was performed between admission and discharge among those receiving a nutritional assessment at both times. A significant worsening at discharge was found in values of phase angle (PA) and fat-free mass (FFM), mid-upper arm circumference (MUAC), albumin and Prognostic Nutritional Index (PNI, calculated by combining albuminemia and lymphocyte count). In a multivariate regression model, PA reduction was significantly associated with PA value at admission and malnutrition diagnosis according to ESPEN criteria, but not with length of stay or the Charlson Comorbidity Index (CCI). No significant changes in weight, BMI and FM were found.

The study by Rinninella makes us consider the worsening of the nutritional status of our patients during hospitalization and suggests the need for greater attention and awareness by health professionals on this topic. Some methodological limitations should be recognized. The selection of the studied cohort was based on the availability of paired data; the sample size was small; the analyzed subgroup derived from a single Healthcare Unit. Furthermore, it is difficult to disentangle the actual role of hospitalization on the impact of the acute or chronic comorbidities on malnutrition. Nevertheless, regardless of the possible causes, the concerns raised by many conditions frequently occurring during in-hospital stay were well discussed, such as the poor food

quality, the inadequate food quantities, the inappropriate meal timing, the insufficient number of nurses to take care of the patient's nutrition and the relevant amount of food wasted in hospitals. Studies performed in other countries confirmed a worsening of malnutrition during the in-hospital stay in patients with malignancies [15] or stroke [16] and found that malnutrition prevalence increases along with length of stay [17]. These data outline the importance of continuously monitoring the nutritional status and performing a more careful assessment before discharge in order to give adequate nutritional support at home too.

Nutritional status can be assessed through many different feasible and validated tools, though none of them is universally accepted as a *gold standard* [18]. ESPEN recognizes the Malnutrition Universal Screening Tool (MUST), the Nutritional Risk Assessment (NRS-2002) and the Mini Nutritional Assessment (MNA) [18], and recommends the Subjective Global Assessment (SGA), which is extensively used worldwide [19, 20]. These tools are based on various combinations of registered or measured BMI, weight loss, food intake, disease severity and age. Blood biomarkers concentrations used for the assessment of nutritional risk should be properly interpreted in consideration of the fact that their variations often reflect the degree of catabolism/inflammation rather than the nutritional status [21]. The main anthropometric measures include body weight and BMI, triceps skinfold and MUAC. BIA allows the estimate of total body water, FFM, FM and PA. This latter has recently been validated as a reliable, easy and quick tool to monitor hospital malnutrition, even if some technical and physiological limitations of this tool should be considered [22, 23].

Indeed, standardized criteria rather than single nutritional status parameters should be employed in malnutrition evaluation both at admission and discharge. In addition to appropriate screening and nutritional assessment, attitude and mindset of health professionals should change. There are many aspects worth considering other than acute or chronic pathological conditions *per se*, such as drug-related anorexia, inappropriate *nil per os* medical prescription, dysphagia, unpalatable food, dysgeusia [1]. Furthermore, there are many issues easy to be improved that would have a favorable impact on the inpatients nutritional status, such as prolonged food avoiding for tests, staff interrupting or disturbing the patients at mealtime, patients' inability to reach meals, difficulties in opening packages or using cutlery, tray missing [24].

The possibility of having nutritional assessments and dietary advice for all inpatients, with specific personalized dietary regimens (for example, frequently administered energy dense small size

portion meals) together with a better assistance during meals would be the first step towards the solution to hospital malnutrition.

Therefore, the key word is probably *training* of health personnel and of all the people who revolve around the patient, such as relatives, volunteers or those who keep company or provide meal assistance, who are still now too little aware of the importance of ensuring a continuum of care to avoid the progressive deterioration of the nutritional status.

REFERENCES

1. Norman K, Pichard C, Lochs H, Pirlich M (2008) Prognostic impact of disease-related malnutrition. *Clin Nutr* 27:5–15. <https://doi.org/10.1016/j.clnu.2007.10.007>
2. Agarwal E, Ferguson M, Banks M, et al (2013) Malnutrition and poor food intake are associated with prolonged hospital stay, frequent readmissions, and greater in-hospital mortality: Results from the Nutrition Care Day Survey 2010. *Clin Nutr* 32:737–745. <https://doi.org/10.1016/j.clnu.2012.11.021>
3. Tobert CM, Mott SL, Nepple KG (2018) Malnutrition Diagnosis during Adult Inpatient Hospitalizations: Analysis of a Multi-Institutional Collaborative Database of Academic Medical Centers. *J Acad Nutr Diet* 118:125–131. <https://doi.org/10.1016/j.jand.2016.12.019>
4. Cereda E, Pedrolli C, Klersy C, et al (2016) Nutritional status in older persons according to healthcare setting: A systematic review and meta-analysis of prevalence data using MNA[®]. *Clin Nutr* 35:1282–1290. <https://doi.org/10.1016/j.clnu.2016.03.008>
5. Barker LA, Gout BS, Crowe TC (2011) Hospital malnutrition: Prevalence, identification and impact on patients and the healthcare system. *Int J Environ Res Public Health* 8:514–527. <https://doi.org/10.3390/ijerph8020514>
6. Lim SL, Ong KCB, Chan YH, et al (2012) Malnutrition and its impact on cost of hospitalization, length of stay, readmission and 3-year mortality. *Clin Nutr* 31:345–350. <https://doi.org/10.1016/j.clnu.2011.11.001>
7. Khalatbari-Soltani S, Marques-Vidal P (2015) The economic cost of hospital malnutrition in Europe; a narrative review. *Clin Nutr ESPEN* 10:e89–e94. <https://doi.org/10.1016/j.clnesp.2015.04.003>
8. Sherry CL, Sauer AC, Thrush KE (2017) Assessment of the Nutrition Care Process in US Hospitals Using a Web-Based Tool Demonstrates the Need for Quality Improvement in

Malnutrition Diagnosis and Discharge Care. *Curr Dev Nutr* 1:e001297.
<https://doi.org/10.3945/cdn.117.001297>

9. Lennard-Jones JE, Arrowsmith H, Davison C, et al (1995) Screening by nurses and junior doctors to detect malnutrition when patients are first assessed in hospital. *Clin Nutr* 14:336–340. [https://doi.org/10.1016/S0261-5614\(95\)80050-6](https://doi.org/10.1016/S0261-5614(95)80050-6)
10. Singh H, Watt K, Veitch R, et al (2006) Malnutrition is prevalent in hospitalized medical patients: Are housestaff identifying the malnourished patient? *Nutrition* 22:350–354. <https://doi.org/10.1016/j.nut.2005.08.009>
11. Feinberg J, Nielsen EE, Gluud C, et al (2015) Nutrition support in hospitalised adults at nutritional risk. *Cochrane Database Syst Rev* 2015:CD011598. <https://doi.org/10.1002/14651858.CD011598>
12. Muscaritoli M, Krznarić Z, Singer P, et al (2017) Effectiveness and efficacy of nutritional therapy: A systematic review following Cochrane methodology. *Clin Nutr* 36:939–957. <https://doi.org/10.1016/j.clnu.2016.06.022>
13. Cederholm T, Barazzoni R, Austin P, et al (2017) ESPEN guidelines on definitions and terminology of clinical nutrition. *Clin Nutr* 36:49–64. <https://doi.org/10.1016/j.clnu.2016.09.004>
14. Rinninella E, Cintoni M, Lorenzo A De, et al (2018) May nutritional status worsen during hospital stay? A sub - group analysis from a cross - sectional study. *Intern Emerg Med*. <https://doi.org/10.1007/s11739-018-1944-5>
15. Zhu MW, Wei JM, Chen W, et al (2018) Dynamic investigation of nutritional risk in patients with malignant tumor during hospitalization. *Zhonghua Yi Xue Za Zhi* 98:1093–1098. <https://doi.org/10.3760/cma.j.issn.0376-2491.2018.14.009>
16. Mosselman MJ, Kruitwagen CLJJ, Schuurmans MJ, Hafsteinsdóttir TB (2013) Malnutrition and risk of malnutrition in patients with stroke: Prevalence during hospital stay. *J Neurosci Nurs* 45:194–204. <https://doi.org/10.1097/JNN.0b013e31829863cb>
17. Correia MITD, Perman MI, Waitzberg DL (2017) Hospital malnutrition in Latin America: A systematic review. *Clin Nutr* 36:958–967. <https://doi.org/10.1016/j.clnu.2016.06.025>
18. Cederholm T, Bosaeus I, Barazzoni R, et al (2015) Diagnostic criteria for malnutrition - An ESPEN Consensus Statement. *Clin Nutr* 34:335–340. <https://doi.org/10.1016/j.clnu.2015.03.001>

19. van Bokhorst-de van der Schueren MAE, Guaitoli PR, Jansma EP, de Vet HCW (2014) Nutrition screening tools: Does one size fit all? A systematic review of screening tools for the hospital setting. *Clin Nutr* 33:39–58. <https://doi.org/10.1016/j.clnu.2013.04.008>
20. Parsons K, Fraser A, Bauer J, Collins C (2009) Evidence Based Practice Guidelines for the Nutritional Management of Malnutrition in Adult Patients Across the Continuum of Care. *J Dietitians Assoc Aust* 66:S1–S34. <https://doi.org/10.1111/j.1747-0080.2009.01383.x>
21. Zhang Z, Pereira SL, Luo M, Matheson EM (2017) Evaluation of blood biomarkers associated with risk of malnutrition in older adults: A systematic review and meta-analysis. *Nutrients* 9:829
22. Player EL, Morris P, Thomas T, et al (2018) Bioelectrical impedance analysis (BIA)-derived phase angle (PA) is a practical aid to nutritional assessment in hospital in-patients. *Clin Nutr* <https://doi.org/10.1016/j.clnu.2018.08.003>
23. Lukaski HC, Kyle UG, Kondrup J (2017) Assessment of adult malnutrition and prognosis with bioelectrical impedance analysis: Phase angle and impedance ratio. *Curr Opin Clin Nutr Metab Care* 20:330–339. <https://doi.org/10.1097/MCO.0000000000000387>
24. Keller H, Allard J, Vesnaver E, et al (2015) Barriers to food intake in acute care hospitals: A report of the Canadian Malnutrition Task Force. *J Hum Nutr Diet* 28:546–557. <https://doi.org/10.1111/jhn.12314>