

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Effect of different feeding sources on lifespan and fecundity in the biocontrol agent *Torymus sinensis***

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1705010> since 2019-06-27T12:00:04Z

*Published version:*

DOI:10.1016/j.biocontrol.2019.04.002

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**This is the author's final version of the contribution published as:**

[Picciau, L., Alma, A., Ferracini, C., 2019. Effect of different feeding sources on lifespan and fecundity in the biocontrol agent *Torymus sinensis*. *Biological Control* 134: 45-52, DOI: 10.1016/j.biocontrol.2019.04.002]

**The publisher's version is available at:**

[<https://www.sciencedirect.com/science/article/pii/S1049964419300921>]

**When citing, please refer to the published version.**

This full text was downloaded from iris-Aperto: <https://iris.unito.it/>

1 Effect of different feeding sources on lifespan and fecundity in the biocontrol agent *Torymus*  
2 *sinensis*

3

4 Picciau L., Alma A., Ferracini, C.\*

5

6 Affiliation:

7 University of Torino, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Largo P. Braccini 2,  
8 10095 Grugliasco (TO), Italy.

9

10 \*Corresponding author: Chiara Ferracini, e-mail: chiara.ferracini@unito.it

11

12 Compliance with ethical standards

13 All the insect rearing and experiments were conducted in accordance with the legislation and  
14 guidelines of the European Union for the protection of animals used for scientific purposes  
15 ([http://ec.europa.eu/environment/chemicals/lab\\_animals/legislation\\_en.htm](http://ec.europa.eu/environment/chemicals/lab_animals/legislation_en.htm)). All experimental  
16 protocols using insects were approved by the *ad-hoc* Committee of DISAFA of the University of  
17 Torino.

18

19 Conflict of interest

20 The authors declare that they have no conflict of interest.

21

22 Author contribution statement

23 CF, LP, and AA conceived and designed research. CF and LP carried out field and laboratory assays.  
24 All authors contributed to the writing of the manuscript and approved the final manuscript.

25

26 **Abstract**

27 *Torymus sinensis* Kamijo (Hymenoptera, Torymidae) is a biocontrol agent released to control  
28 outbreaks of the Asian chestnut gall wasp, *Dryocosmus kuriphilus* Yasumatsu (Hymenoptera,  
29 Cynipidae). This wasp is mass reared in controlled conditions in order to be released on a large scale  
30 in chestnut orchards and coppices, thus factors such as food quality may deeply influence the  
31 effectiveness of parasitoids. To identify different diets that could be used to increase mass rearing,  
32 performance, longevity and fecundity of *T. sinensis* were assessed using honey, honey plus pollen,  
33 pollen, and water as diet, compared to unfed wasps in laboratory experiments.

34 In the present study diet quality greatly influenced survival and reproductive output in *T. sinensis*.  
35 Adults using honey plus pollen as food had the longest life span, and significantly increased fecundity  
36 compared to pollen alone, water, and unfed wasps. Median lifespan (day at which 50% of the initial  
37 number of wasps are still alive) was 31, 31, 2.5, 4, 1.5 days, in the honey, honey plus pollen, pollen,  
38 water, control, respectively.

39 The egg production in wasps fed with honey (diet 1) and honey plus pollen (diet 2) observed over  
40 lifetime showed the newly emerged specimens with an average of  $2.40 \pm 0.21$  and  $0.14 \pm 0.12$   
41 respectively at day 0. The number of mature eggs increased rapidly reaching in diet 1 the highest  
42 average number,  $26.60 \pm 2.48$ , at day 6, and  $30.2 \pm 1.59$  at day 7 in diet 2. No significant difference  
43 in the mean number of mature eggs between diets 1 and 2 was observed during all 13 weeks except  
44 for week 11. The mean number of reabsorbed eggs was 0 both in diet 1 and 2 at day 0. The mean  
45 number increased till  $2.80 \pm 0.66$  at day 10 and  $1.40 \pm 0.24$  at day 14 in diet 1 and 2 respectively.

46 The cumulative lifetime mean egg load was 1012.67 and 1095.82 when feeding honey and honey  
47 plus pollen, respectively, which was significantly higher than 32.40 for pollen, 58.40 for water and  
48 43.57 for control.

49 Based on our studies, feeding the female wasps with honey and pollen increased lifespan by 33-fold,  
50 and fecundity by 2.3-fold at day 4 over unfed wasps.

51 This work contributes to a better understanding of the influence of different diets on lifespan and  
52 fecundity in the parasitoid wasp *T. sinensis*, to the refinement of mass rearing in controlled  
53 conditions, and to the optimization of classical biocontrol programs. Prior to the field release, it  
54 seems advisable to supply insects with sugar-rich diets to improve their performances in fields and  
55 to enhance parasitoid egg expenditure.

56 Key words: parasitoid feeding; honey-based diets; oosorption; mass rearing; Asian chestnut gall  
57 wasp.

58 **Introduction**

59 Life history traits of insects in nature, namely survival, development, and fecundity, are deeply  
60 influenced by food quality. Food sources of adult parasitoids include host food, host hemolymph and  
61 tissues. However, host-feeding female parasitoids can also use non-host foods to prolong their  
62 lifespan and increase their reproductive potential, as they usually contain carbohydrates, proteins, and  
63 other trace elements (Liu et al., 2015).

64 Proteins are often regarded as important nutritional elements for insect reproduction, enhancing  
65 offspring production (Bong et al., 2014). Conversely, the energy obtained from a carbohydrate-rich  
66 diet is mainly focused on insect life maintenance, thus prolonging life span. Potential sugar sources  
67 for parasitoids include nectar and homopteran honeydew. These sugar-meals carried over from  
68 juvenile stage can be used immediately to generate energy for metabolic purposes and even stored  
69 (Olson et al., 2000). Furthermore, pollens contain a number of nutrients and often high levels of free  
70 amino acids, instead of most nectars, there included all 10 essential amino acids. Compared to nectars,  
71 pollens have higher level of protein, lipid and polysaccharides (Thompson et al., 1999; Zhang et al.,  
72 2004).

73 Insects, like many animals, have the ability to respond plastically to environmental stress, resorbing  
74 oocytes that are not oviposited (Moore and Attisano, 2011). Sugar-feeding may help synovigenic  
75 species to mature additional eggs, and can prevent parasitoids from resorbing eggs (Lee et al., 2004).  
76 Honey has long been used as an alternate food source to sucrose solution for insects such as  
77 parasitoids mass reared in controlled conditions, but little is known about its actual properties in  
78 influencing reproduction and lifespan (Harvey et al., 2012). Honey is presumably the most studied  
79 artificial diet for parasitoids since it is palatable and cheap. Provision of honey solutions boosts  
80 mymarid and trichogrammatid parasitoid lifespan similarly to floral and extrafloral nectars, and even  
81 more than honeydew (Benelli et al., 2017). A lack of sugar availability may temporarily or  
82 permanently limit the reproductive success of parasitoids in agricultural system (Heimpel et al.,  
83 1997).

84 Pollen is a food for numerous insects, including many species of natural enemies, and parasitic  
85 Hymenoptera feed on pollen to varying degrees (Lundgren and Wiedenmann, 2004). It is one of the  
86 most nutritious non-prey food sources for parasitoids based on its protein levels, but pollen-feeding  
87 by parasitoids has been studied much less frequently than sugar-feeding (Lundgren, 2009).

88 *Torymus sinensis* Kamijo (Hymenoptera: Torymidae) is the biocontrol agent of the Asian chestnut  
89 gall wasp, *Dryocosmus kuriphilus* Yasumatsu (Hymenoptera: Cynipidae) (Moriya et al., 2003), and  
90 is considered as one of the most successful examples of recent European classical biocontrol  
91 programs (Avtzis et al., 2018; Ferracini et al., 2018a; Matošević et al., 2017). Although research on  
92 the reproductive biology and fecundity of this synovigenic parasitoid was already carried out (Picciau  
93 et al., 2017), the impact of diet on its life history has never been studied. In the present work, we  
94 investigated if lifespan and fecundity of *T. sinensis* are affected by different sugar diets. Since factors  
95 such as food quality are also likely to determine the effectiveness of parasitoids, wasps were allowed  
96 to feed on honey, honey plus pollen, pollen, and water. Depending on the different diet, lifespan, egg  
97 load and egg resorption of *T. sinensis* individuals were compared with unfed wasps in laboratory  
98 experiments. The results were expected to deepen knowledge about the factors affecting the  
99 biocontrol efficacy of this species in controlled conditions, thus improving mass rearing by insectaries  
100 and private companies.

## 101 **Material and Methods**

### 102 *Insect culture*

103 Galls were randomly collected during winter 2016-2017 according to the method described by  
104 Ferracini et al. (2015; 2018b) in chestnut-growing areas of NW Italy with a stable population of *T.*  
105 *sinensis*. Galls were individually isolated in plastic vials (120mm in length by 25mm in diameter),  
106 and kept outdoor until *T. sinensis* emergence.

### 107 *Food sources*

108 Insects were provided *ad libitum* one of the following different diets: 1) undiluted honey, 2) undiluted  
109 honey plus organic grinded pollen grains, 3) organic grinded pollen grains, 4) distilled water. The

110 diets have been compared to a control, with no food source provided. For the first diet we used a total  
111 of 0.02 g of honeydew honey, disposing few droplets on a cardboard, using a pipette. For the second  
112 diet, a stock mixture was prepared adding 1 g of organic pollen to 30 g of honeydew honey; few  
113 droplets were placed on a cardboard, for a total of 0.02 g. For the third diet, 0.02 g of organic pollen  
114 grain was weighted, grinded with a pestle and directly inserted in the plastic tube. For the fourth diet,  
115 the cotton plug of the tube was imbedded with 3 ml of distilled water. In the control, no source of  
116 food and water was provided. Each sugar source was replaced once a week, while the moistened  
117 cotton plug was changed twice a week.

#### 118 *Parasitoid lifespan*

119 Three newly emerged wasps for each sex, for a total of six individuals, were placed into plastic vials  
120 (120 mm in length by 25 mm in diameter), and provided with one of four different sugar diets  
121 throughout their adult life. Twenty replications were performed for each food source. The vials were  
122 maintained in climatic chamber under controlled conditions [ $15 \pm 1$  °C,  $60 \pm 5$  % RH, and a  
123 photoperiod of 16:8 (L:D) h], and checked twice a day for dead wasps every 10 a.m. and 16 p.m. each  
124 day until all individuals had died.

#### 125 *Parasitoid fecundity*

126 Emerging females only, used for egg load survey, were collected every day in number of 1,800  
127 specimens originating 30 age-cohorts for each feed-test for a total of 180 cohorts.

128 The insects were reared in plastic vials (120 mm in length by 25 mm in diameter) arranged in 5 groups  
129 each fed with different food combination: 1) honey, 2) honey plus pollen, 3) pollen, 4) water and 5)  
130 no food source (control). All of them were kept in a climatic chamber under controlled conditions ( $15$   
131  $\pm 1$  °C,  $60 \pm 5$ % RH, and a photoperiod of 16:8 L:D) according to Picciau et al. (2017) and maintained  
132 till dissection.

#### 133 *Egg load and oosorption quantifying*

134 Five females for each age cohort were collected daily and killed with ethyl acetate. The specimens  
135 were placed on a microscope slide ( $25 \times 75 \times 1$  mm) into a phosphate buffered saline (PBS) solution



136 for ovary dissection. Egg load was assessed by dissecting each female under a stereomicroscope at  
137  $25 \times$  magnification. The abdomen was separated from the rest of the body using a couple of needles.  
138 Ovaries were kindly removed from the abdomen by means of micro-pin needles and spread on a slide  
139 for egg load counts. Mature eggs were defined as described in Picciau et al. (2017). Egg resorption  
140 (oosorption) was also investigated by counting those eggs close to the lateral oviducts which  
141 displayed a collapsed pattern indicating a reabsorption process (Figure 1). The number of mature and  
142 reabsorbed eggs was counted under microscope at  $80 \times$  magnification.

#### 143 *Statistical analysis*

144 The data on the female longevity of the different diets were  $\log_{10}(x+1)$  transformed and transformed  
145 using analysis of variance (ANOVA). We used the Kaplan-Meier estimator (Kaplan and Meier, 1958)  
146 to visualize survivorship curves in each diet, and a log-rank test was used to determine differences in  
147 survival rates. The significance of each term in the model was analyzed using Wald  $\chi^2$  statistic.

148 After testing for homogeneity of variance (Levene test), data were analyzed by T test independent  
149 samples ( $P < 0.05$ ) to compare the egg load and the oosorption records comparing wasps fed with  
150 honey and with honey and pollen through 13 weeks. Furthermore, all data were analyzed within the  
151 first 8 days by a Generalized Linear Model (GLM) with Poisson distribution and Log link function.  
152 Averages among the diets were separated by a Bonferroni *post hoc* when significant difference were  
153 present. All analyses were performed using the software SPSS version 20.0 (SPSS, Chicago, IL,  
154 USA). The cumulative mean egg load was analyzed with a Kruskal Wallis test, and averages were  
155 separated by a Bonferroni *post hoc* when significant difference were present.

#### 156 **Results**

157 The results concerning the effects of different food sources on lifespan of the parasitoid *T. sinensis*  
158 are shown in Figure 2. Diet significantly affected lifespan, in particular wasps fed with honey (diet  
159 1) and honey plus pollen (diet 2), increased lifespan to the greatest extent. This lifespan, in fact, was  
160 more than 20 times as long as that of individuals starved (control) (Figure 2). Furthermore, lifespan  
161 of the wasps was significantly reduced when they were fed with pollen (diet 3), and water (diet 4) as

162 well. Even if the average lifespan was higher when adults were provided with honey plus pollen, no  
163 significant difference was found between this source and honey alone.

164 Median lifespan (day at which 50% of the initial number of wasps are still alive) was 31, 31, 2.5, 4,  
165 1.5 days, in the honey, honey plus pollen, pollen, water, and control, respectively. Furthermore,  
166 according to the diet all wasps were dead after 88, 108.5, 6, 7, 4 days in honey, honey plus pollen,  
167 pollen, water and control, respectively. The median survival time differed between female and male  
168 parasitoids, with females living longer (log rank test,  $\chi^2 = 5.42$ ,  $df = 1$ ,  $P = 0.0199$ ) (Figure 3).

169 The comparison of egg load and oosorption was performed through 13 weeks, considering only honey  
170 (diet 1) and honey plus pollen (diet 2), basing this choice on the diets with the longest lifespan. The  
171 egg production observed over lifetime showed that the newly emerged specimens had an average of  
172  $2.40 \pm 0.21$  mature eggs in specimens fed with diet 1 and  $0.14 \pm 0.12$  mature eggs in specimens fed  
173 with diet 2 at day 0. In the following days the number of mature eggs increased rapidly reaching in  
174 diet 1 the highest average number,  $26.60 \pm 2.48$ , at day 6 with a maximum value per individual of 34  
175 eggs. For this diet the highest number per individual was 40 and reached at day 17 when the average  
176 was  $20.8 \pm 5.21$ . The highest average number reached in diet 2 was  $30.2 \pm 1.59$  at day 7 with a  
177 maximum value per individual of 34 eggs. For this diet the highest number per individual was 43 and  
178 reached at day 9 when the average was  $26.5 \pm 3.12$ . No significant difference in the mean number of  
179 mature eggs between diets 1 and 2 was observed during all 13 weeks except for week 11 ( $t = -0.452$ ,  
180  $df = 7.842$ ,  $P = 0.663$ ) (Figure 4).

181 Concerning the oosorption, the number of reabsorbed eggs observed over lifetime showed that the  
182 newly emerged specimens had an average of 0 collapsed eggs both in diet 1 and 2. Subsequently the  
183 number increased, reaching in diet 1 the highest average number,  $2.80 \pm 0.66$ , at day 10 with a  
184 maximum value per individual of 5. For this diet the highest number per individual was 6 and reached  
185 only at day 14 when the average was  $2.60 \pm 0.87$ . The highest average number reached in diet 2 was  
186  $1.40 \pm 0.24$  at day 14 with a maximum value per individual of 2 reabsorbed eggs. For this diet the  
187 highest number per individual was 2 and reached for the first time at day 11 when the average was

188 0.40 ± 0.40. Significant differences were found only in weeks 1, 3, 12 and 13 (week 1:  $t = -1.364$ ,  $df$   
189 = 8.053,  $P = 0.209$ ; week 3:  $t = -4.418$ ,  $df = 6.075$ ,  $P = 0.004$ ; week 12:  $t = -0.926$ ,  $df = 1.010$ ,  $P =$   
190 0.523; week 13:  $t = -3.000$ ,  $df = 3.000$ ,  $P = 0.058$ ) (Figure 5).

191 The comparison among all the diets was performed only through the first 8 days when only the  
192 specimens provided with diet 1 and 2 remained alive (Figure 6). The unfed wasp, with the shortest  
193 lifespan, showed a peak at the third day with an egg load mean of 17.70. The specimens fed with  
194 pollen had the higher egg load during the last three days before dying reaching an average egg load  
195 of 8.53. In case of water, the egg load followed a normal distribution with the highest mean value of  
196 18.20 eggs at day 2. The mean egg load for wasps provided with honey and honey plus pollen showed  
197 a positive trend with lower values until day 3 with a subsequent increase in the following days  
198 overcoming the other feeding sources (Figure 6). The cumulative lifetime mean egg load was 1012.67  
199 and 1095.82 when feeding honey and honey plus pollen, respectively, which was significantly higher  
200 than 32.40 for pollen, 58.40 for water and 43.57 for control.

## 201 **Discussion**

202 Egg maturation in parasitoids depends on the diet provided to adult females (Onagbola et al., 2007).  
203 It is considered a fitness-related parameter, which may influence the reproductive potential of  
204 biocontrol agents and thus the population dynamics of host. However, different diets may have an  
205 effect on the egg maturation patterns through lifetime of synovigenic parasitoids species (Jervis et al.,  
206 2001). In fact, lipids and proteins, from either the diet and teneral reserves, are important for  
207 production of the eggs (Ellers and Van Alpen, 1997), and the balanced acquisition of nutrients has  
208 great impact on the size and number of eggs produced (Harvey et al., 2012; Wheeler 1996). However,  
209 egg production is also improved by various artificial diets based on carbohydrate solutions, and  
210 exploitation of sugar sources or pollen is widespread in a broad range of orders, including  
211 Hymenoptera (Wackers et al., 2008). Food quality greatly influences life history traits such as  
212 survival, development, and fecundity of insects. Its availability over time can affect fitness-related  
213 traits and may alter egg maturation (Benelli et al., 2017) in synovigenic species, there included *T.*

214 *sinensis* (Picciau et al., 2017). Carbohydrate-rich diet access, even as intermittent feeding and without  
215 additional proteins, has a strong effect on life span being able to increase longevity of about 5 times  
216 compared to water alone as demonstrated by Ellers et al. (2011). Among the different sugar source,  
217 honey is considered the most effective food to promote egg maturation (Hogervorst et al., 2007), and  
218 improve fecundity by prolonging parasitoid lifespan (Mutitu et al., 2013). Moreover, pollen can also  
219 be exploited as food source to improve fecundity when used in addition to honey (Zhang et al., 2004).  
220 In the present study diet quality equally influenced the survival and the reproductive output in *T.*  
221 *sinensis*. Adults fed with honey plus pollen had the longest life span, and this diet significantly  
222 increased fecundity compared to control. In particular, *T. sinensis* responded to a honey supplement,  
223 increasing the mean survival up to 49 days, considering female wasps provided with honey plus  
224 pollen, confirming previous data by Picciau et al. (2017). Conversely, when only pollen or water were  
225 provided similar effects to control were recorded. In fact, wasps reached a mean survival of 2.5, 4,  
226 and 1.5 days, respectively. Significant reduction in longevity, with no significant difference between  
227 sexes, was recorded highlighting how parasitoids wasps are especially sensitive to sugar deprivation  
228 as adults. As already pointed out by Olson et al. (2000), laboratory lifespan of many parasitoids  
229 species is typically less than 5 days in the absence of sugar but exceeds 2 to 3 weeks when sugar  
230 meals are provided, confirming our results.

231 Honey has long been used in controlled conditions as an alternate food source to sucrose solution for  
232 insects such as parasitoids, but its actual properties in influencing parasitoid longevity and  
233 reproduction have been little explored. The content of honey is dominated by sugars, although it also  
234 contains trace amounts of vitamins or minerals, with tiny amounts of several compounds thought to  
235 function as antioxidants (Harvey et al., 2012). Based on our studies, feeding the female wasps with  
236 honey plus pollen increased lifespan by 33-fold, and fecundity by 2.3-fold at day 4 over unfed wasps.  
237 Moreover, pollen of plants is known to be a source of food for a large number of insect species in  
238 nature, but pollen feeding by parasitoids has been studied less frequently than sugar feeding. Some  
239 groups of insects have been reported feeding on pollen (e.g., Bombyliidae, Eulophidae, Mutillidae,

240 Scoliidæ, Trichogrammatidæ) (Jervis et al., 2008; Patt et al., 1997; Zhang et al., 2004). In our  
241 experiment, pollen alone did not increase lifespan but a higher longevity was recorded when it was  
242 provided in addition to honey.

243 In case of scarcity of food, insects, like many animals, have the ability to adapt to environmental  
244 stress, resorbing oocytes that are not oviposited (Moore and Attisano, 2011). In our experiments the  
245 presence of egg deterioration related to oosorption was verified in *T. sinensis*, as previously  
246 hypothesized in Picciau et al. (2017). The comparison of egg load and egg maturation trend recorded  
247 in the specimens fed with honey and honey plus pollen through their lifetime suggest that the addition  
248 of pollen in the diet does not heavily affect the number of egg production. On the contrary it seems  
249 to have a strong impact on egg resorption probably due to the increased nutritional value of the food  
250 provided. Compared to males, females were in fact able to reallocate resources to somatic  
251 maintenance as evidenced by no reduction in life span, especially in their first month of life.  
252 Comparing the mean cumulative egg load recorded with all different diets to the unfed wasps, a  
253 significant increase in fecundity was detected. In particular feeding with honey increased the total  
254 egg load by 23.2-fold, and with honey plus pollen by 25.1-fold, whilst with only water only 1.4-fold  
255 and pollen alone showed no significant difference. As already stated by Zhang et al. (2004) for  
256 *Trichogramma* spp., honey and pollen should be considered a complete diet for *T. sinensis* as well  
257 conversely to pollen and water.

258 This knowledge about feeding sources influence on parasitoid life-history traits is essential to improve  
259 mass-rearing techniques. In particular, *T. sinensis*, being a biocontrol agents employed in classical  
260 biocontrol programs in Europe, may be positively affected by diet in laboratory conditions. A  
261 balanced acquisition of nutrients has indeed great influence on the size and number of eggs produced  
262 (Harvey et al., 2012; Wheeler 1996).

263 Modern agriculture has simplified the agro-ecosystem landscape, increasing the pest population  
264 density and reducing the alternative non-crop sources for natural enemies (Benelli et al., 2017). In a  
265 frame of a biocontrol programs based on conservative and augmentative actions, nectar and

266 honeydew accessibility in the field is very important to improve parasitoid effectiveness, as already  
267 pointed out in case of scarce flower resources (Pinheiro et al., 2015; Wäckers et al., 2008). Moreover,  
268 providing parasitoids with sugar-rich diets before their release to the field is recommendable to  
269 improve their performance and optimize the egg laying (Hougardy and Mills, 2007).  
270 The current findings have offered some important insights into the life history traits of *T. sinensis*,  
271 providing new cues to enhance methods of mass rearing for biocontrol. To overcome occurring field  
272 limitations, the employment of artificial sugar sources, flowering cover crops and the management of  
273 natural food resources may be helpful to improve natural populations and to enhance the fitness of  
274 parasitic wasps in both conservative and augmentative biocontrol programs.

275

#### 276 **Acknowledgements**

277 The authors would like to thank Ester Ferrari and Marianna Pontini for managing *T. sinensis* mass  
278 rearing, and Silvia T. Moraglio and Fabio Mazzetto for assistance with statistical analysis.

279

280 Funding: This research did not receive any specific grant from funding agencies in the public,  
281 commercial, or not-for-profit sectors.

282

283 **References**

- 284 Avtzis, D.N., Melika, G., Matošević, D., Coyle, D.R., 2018. The Asian chestnut gall wasp  
285 *Dryocosmus kuriphilus*: a global invader and a successful case of classical biological control. J. Pest  
286 Sci., DOI 10.1007/s10340-018-1046-1.
- 287 Benelli, G., Giunti, G, Tena A., Desneux, N., Caselli A., Canale, A., 2017. The impact of adult diet  
288 on parasitoid reproductive performance. J. Pest Sci., 90, 807–823.
- 289 Bong, L.J., Neoh, K.B., Lee, C.Y., Jaal, Z., 2014. Effect of diet quality on survival and reproduction  
290 of adult *Paederus fuscipes* (Coleoptera: Staphylinidae). J. Medical Entomol., 51(4), 752-759.
- 291 Ellers, J., van Alphen, J.J.M., 1997. Life history evolution in *Asobara tabida*: plasticity in allocation  
292 of fat reserves to survival and reproduction. J. Evol. Biol., 10, 771-785.
- 293 Ellers J., Ruhe, B., Visser, B., 2011. Discriminating between energetic content and dietary  
294 composition as an explanation for dietary restriction effects. J. Insect Physiol., 57, 1670-1676.
- 295 Ferracini, C., Gonella, E., Ferrari, E., Saladini, M.A., Picciau, L., Tota, F., Pontini, M., Alma, A.,  
296 2015. Novel insight in the life cycle of *Torymus sinensis*, biocontrol agent of the chestnut gall wasp.  
297 BioControl. 60, 169-177.
- 298 Ferracini, C., Ferrari, E., Pontini, M., Saladini, M.A., Alma, A., 2018a. Effectiveness of *Torymus*  
299 *sinensis*: a successful long-term control of the Asian chestnut gall wasp in Italy. J. Pest Sci., DOI:  
300 10.1007/s10340-018-0989-6.
- 301 Ferracini, C., Bertolino, S., Bernardo, U., Bonsignore, C., Faccoli, M., Ferrari, E., Lupi, D., Maini,  
302 S., Mazzon, L., Nugnes, F., Rocco, A., Santi, F., Tavella, L., 2018b. Do *Torymus sinensis*  
303 (Hymenoptera: Torymidae) and agroforestry system affect native parasitoids associated with the  
304 Asian chestnut gall wasp? Biol. Control., 121, 36–43.
- 305 Harvey, J.A., Cloutier, J., Visser, B., Ellers, J., Wäckers, F.L., Gols, R., 2012. The effect of different  
306 dietary sugars and honey on longevity and fecundity in two hyperparasitoids wasps. J. Insect Physiol.,  
307 58, 816-823.

308 Heimpel, G.E., Rosenheim, J.A, Kattari, D., 1997. Adult feeding and lifetime reproductive success  
309 in the parasitoid *Aphytis melinus*. Entomol. Exp. Appl., 83, 305–315.

310 Hogervorst, P.A.M., Wäckers, F.L., Romeis, J., 2007. Effects of honeydew sugar composition on the  
311 longevity of *Aphidius ervi*. Entomol. Exp. Appl., 122, 223-232.

312 Hougardy, E., Bezemer, T.M., Mills, N.J., 2005. Effects of host deprivation and egg expenditure on  
313 the reproductive capacity of *Mastrus ridibundus*, an introduced parasitoid for the biological control  
314 of codling moth in California. Biol. Control., 33, 96–106.

315 Hougardy, E., Mills, N.J., 2006. The influence of host deprivation and egg expenditure on the rate of  
316 dispersal of a parasitoid following field release. Biol. Control., 37, 206–213.

317 Jervis, M.A., Heimpel, G.E., Ferns, P.N., Harvey, J.A., Kidd, N.A.C., 2001. Life-history strategies  
318 in parasitoid wasps: a comparative analysis of “ovigeny”. J. Animal Ecol., 70, 442-458.

319 Kaplan, E.L., Meier, P., 1958. Nonparametric estimation from incomplete observations. J. Am. Stat.  
320 Assoc., 53, 457–481.

321 Lee, J.C., Heimpel, G.E., Leibe, G.L., 2004. Comparing floral nectar and aphid honeydew diets on  
322 the longevity and nutrient levels of a parasitoid wasp. Entomol. Exp. Appl., 111, 189–199.

323 Lundgren, J.G., 2009. The pollen feeders. In: Relationships of natural enemies and non-prey foods.  
324 Springer, Brookings, USA, pp. 454.

325 Lundgren, J.C., Wiedenmann, R.N., 2004. Nutritional suitability of corn pollen for the predator  
326 *Coleomegilla maculata* (Coleoptera: Coccinellidae), J. Insect Physiol., 50 (6), 567-575.

327 Matošević, D., Lacković, N., Kos, K., Kriston, E., Melika, G., Rot, M., Pernek, M., 2017. Success  
328 of classical biocontrol agent *Torymus sinensis* within its expanding range in Europe. J. Appl.  
329 Entomol., 141, 758–767.

330 Moore, P.J., Attisano, A., 2011. Oosorption in response to poor food: complexity in the trade-off  
331 between reproduction and survival. Ecol. Evol., 1(1), 37-45.



332 Moriya, S., Shiga, M., Adachi, I., 2003. Classical biological control of the chestnut gall wasp in  
333 Japan. In: van Driesche RG (ed). Proceedings of the 1st international symposium on biological  
334 control of arthropods. USDA Forest Service, Washington, DC, USA, pp 407–415.

335 Mutitu, E.K., Garnas, J.R., Hurley, B.P., Wingfield, M.J., Harney, M., Bush, S.J., Slippers, B., 2013.  
336 Biology and rearing of *Cleruchoides noackae* (Hymenoptera: Mymaridae), an egg parasitoid for the  
337 biological control of *Thaumastocoris peregrinus* (Hemiptera: Thaumastocoridae). J. Econ. Entomol.,  
338 106, 1979-1985.

339 Onagbola, E.O., Fadamiro, H.Y., Mbata, G.N., 2007. Longevity, fecundity and progeny sex ratio of  
340 *Pteromalus cerealellae* in relation to diet, host provision, and mating. Biol. Control., 40, 222-229.

341 Olson, D.M, Fadamiro, H., Lundgren, J.G., Heimpel, G.E., 2000. Effects of sugar feeding on  
342 carbohydrate and lipid metabolism in a parasitoid wasp. Physiol. Entomol., 25, 17-26.

343 Patt, J.M., Hamilton, G.C., Lashomb, J.H., 1997. Impact of strip-insectary intercropping with flowers  
344 on conservation biological control of the Colorado potato beetle. Adv. Hort. Sci. 11(4), 175-181.

345 Picciau, L., Ferracini, C., Alma, A., 2017. Reproductive traits in *Torymus sinensis*, biocontrol agent  
346 of the Asian chestnut gall wasp: implications for biological control success. Bull. Insectol., 70, 49-  
347 56.

348 Pinheiro, L.A., Torres, L.M., Raimundo, J., Santos, S.A.P., 2015. Effects of pollen, sugars and  
349 honeydew on lifespan and nutrient levels of *Episyrphus balteatus*. BioControl. 60, 47–57.

350 Thompson, S.N., Hagen, K.S., 1999. Nutrition of entomophagous insects and other arthropods. In  
351 Handbook of Biological Control (Bellows, T.S., Fisher, T.W., Eds), Academic Press, CA, pp. 594-  
352 652.

353 Zhang, G., Zimmermann, O., Hassan, S.H., 2004. Pollen as a source of food for egg parasitoids of  
354 the genus *Trichogramma* (Hymenoptera: Trichogrammatidae), Biocontrol Sci. Technol., 14(2), 201-  
355 209.

356 Wäckers, F.L., van Rijn, P.C.J., Heimpel, G.E., 2008. Honeydew as a food source for natural enemies:  
357 Making the best of a bad meal? Biol. Control. 45, 176–184.

- 358 Liu, W.X., Wang, W.X., Zhang, Y.B., Wang, W., Lu, S.L., Wan, F.H., 2015. Adult diet affects the  
359 life history and host-killing behavior of a host-feeding parasitoid. *Biol. Control.* 81, 58–64.
- 360 Wheeler, D., 1996. The role of nourishment in oogenesis. *Ann. Rev. Entomol.*, 41, 407-431.