

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

The endobacterium of an arbuscular mycorrhizal fungus modulates the expression of its toxin-antitoxin systems during the life cycle of its host

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1645358> since 2017-07-20T15:20:42Z

Published version:

DOI:10.1038/ismej.2017.84

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Salvioli di Fossalunga, Alessandra; Lipuma, Justine; Venice, Francesco; Dupont, Laurence; Bonfante, Paola. The endobacterium of an arbuscular mycorrhizal fungus modulates the expression of its toxin-antitoxin systems during the life cycle of its host. THE ISME JOURNAL. None pp: 1-5.
DOI: 10.1038/ismej.2017.84

The publisher's version is available at:

<http://www.nature.com/doifinder/10.1038/ismej.2017.84>

When citing, please refer to the published version.

Link to this full text:

<http://hdl.handle.net/>

The Endobacterium of an Arbuscular Mycorrhizal Fungus Modulates the Expression of its Toxin-Antitoxin Systems During the Life Cycle of its Host

Alessandra Salvioli di Fossalunga^{1*}, Justine Lipuma^{2*}, Francesco Venice¹, Laurence Dupont², Paola Bonfante¹

¹ Department of Life Sciences and Systems Biology, Torino, ITALY

² Institut Sophia Agrobiotech (ISA), INRA UMR 1355, CNRS UMR 7254, Université de Nice Sophia Antipolis, Sophia Antipolis, FRANCE

* These authors contributed equally to this work

Running title: TA systems in the genome of a fungal endobacterium

Correspondence: Alessandra Salvioli, viale Pier Andrea Mattioli 25, 10125 Torino, Italy.

Phone: +39 0116705775

Email: alessandra.salvioli@unito.it

Conflict of interest statement: The authors declare no conflict of interest.

Subject Categories: Microbe-microbe and microbe-host interactions

Abstract

25

Arbuscular mycorrhizal fungi (AMF) are widespread root symbionts that perform important 26
ecological services, such as improving plant nutrient and water acquisition. Some AMF from the 27
Gigasporaceae family host a population of endobacteria, *Candidatus Glomeribacter gigasporarum* 28
(*Cagg*). The analysis of the *Cagg* genome identified six putative toxin-antitoxin modules (TAs), 29
consisting of pairs of stable toxins and unstable antitoxins that affect diverse physiological 30
functions. Sequence analysis suggested that these TA modules were acquired by horizontal transfer. 31
Gene expression patterns of two TAs (*yoeB/yefM* and *chpB/chpS*) changed during the fungal life 32
cycle, with the expression during the presymbiotic phase higher than during the symbiosis with the 33
plant host. The heterologous expression in *Escherichia coli* demonstrated the functionality only for 34
the *YoeB-YefM* pair. Based on these observations, we speculate that TA modules might help *Cagg* 35
adapt to its intracellular habitat, coordinating its proliferation with the physiological state of the 36
AMF host. 37

Arbuscular mycorrhizal fungi (AMF) perform key ecological services, improving nutrient acquisition and water uptake by their plant hosts, while receiving fixed carbon from the host (Smith and Read, 2010). Many fungi, particularly from basal clades, harbor bacterial endosymbionts (Bonfante and Desiro, 2017) and AMF from the *Gigasporaceae* family host a population of *Burkholderia*-related microbes (Bianciotto *et al.*, 1996) named *Candidatus Glomeribacter gigasporarum* (*Cagg*). *Cagg* is vertically transmitted and currently uncultivable; although not essential for *Gigaspora* survival, *Cagg* can enhance fungal fitness (Bianciotto *et al.*, 2003; Lumini *et al.*, 2007; Salvioli *et al.*, 2016). Analysis of the 1.72-Mb *Cagg* genome revealed strong nutritional dependence on the fungal host (Ghignone *et al.*, 2012). We wondered whether the *Cagg* genome possesses genetic determinants involved in its endocellular lifestyle and environmental sensing. Our previous study identified potentially secreted proteins, which could act as effectors (Ghignone *et al.*, 2012); the present study identified genes encoding toxin-antitoxin (TA) systems.

Bacterial TA systems (TAs) are typically encoded in operons located on plasmids or on the bacterial chromosome. They can be classified into five types (I to V) with the antitoxin acting as a protein (types II, IV and V) or as a noncoding RNA (types I and III) (Schuster and Bertram, 2013). The well-studied type II TAs include a stable toxin protein and an unstable antitoxin that counteracts the effect of the toxin (Leplae *et al.*, 2011). Depending on the TA superfamily, the type II toxins either interfere with DNA replication (*e.g.* CcdB) or with the translation of mRNA (*e.g.* MazE and RelE) (Mruk and Kobayashi, 2014). It was originally proposed that obligate host-associated bacteria lost their TAs as a consequence of the reduction in genome size experienced during evolution (Pandey and Gerdes, 2005). For example, *Mycobacterium tuberculosis* possesses 88 TAs, but the genome of the obligate intracellular pathogen *Mycobacterium leprae* has no TAs, supporting the idea that TAs help free-living bacteria cope with the changing environment. However, data from high-throughput sequencing have challenged this view by revealing TAs in endocellular bacteria, including pathogens and beneficial symbionts (Sevin and Barloy-Hubler, 2007). By contrast, the genomes of insect endosymbionts do not seem to carry TA operons, except *Wolbachia*, the secondary symbiont of *Drosophila melanogaster*, which encodes seven relBE-like TAs (Sevin and Barloy-Hubler, 2007).

TAs have multiple roles, including in programmed cell death (Bayles, 2014), stress adaptation (Ramage *et al.*, 2009), and survival in host cells (Helaine *et al.*, 2014). For example, persistence of *Salmonella* within macrophages requires the presence of TAs (Helaine *et al.*, 2014). In the symbiotic nitrogen-fixing *Sinorhizobium meliloti*, a mutation of the toxin from the *ntrPR* system

improved symbiotic efficiency and mutations of the *vapBC-5* pair influenced nitrogen fixation 71
capacity and bacteroid senescence (Olah *et al.*, 2001; Lipuma *et al.*, 2014). These data suggest that 72
intracellular endobacteria may exploit TAs to modulate their interactions with host cells. 73

To date, characterization of TAs from fungal endobacteria has been limited to descriptions of their 74
presence in the genome. However, due to their involvement in modulation of growth under stress 75
conditions and survival in host cells, TAs have been hypothesized to play important roles in 76
regulating the life of fungal endobacteria (Lackner *et al.*, 2011). Here, we characterize TA systems 77
present in the *Cagg* genome and test the functionality of two of them. 78

Analysis of the *Cagg* genome revealed six complete TAs and three orphan coding sequences 79
(Supplementary Table 1). We also conducted a similar analysis of the genomes of *Cagg* relatives 80
with different lifestyles, considering two endofungal bacteria, namely *Burkholderia rhizoxinica*, 81
living inside *Rhizopus microsporus* (Partida Martinez *et al.*, 2005), and *Mycoavidus cysteinexigens* 82
living inside *Mortierella* (Ohshima *et al.*, 2016; Uehling *et al.*, 2017), as well as the obligate 83
endosymbiont of the citrus mealybug, *Candidatus Tremblaya princeps* (López-Madrugal *et al.*, 84
2011). The genomes of *B. rhizoxinica*, *M. cysteinexigens*, and *Cagg* encode TA systems, albeit 85
fewer than in their free-living relatives (Figure 1). By contrast, no TA could be confidently 86
predicted for *Ca. Tremblaya princeps*, probably due to its extremely reduced genome (only 139 kb). 87
Comparison of *Cagg* TAs with those from its close relative *B. rhizoxinica* revealed a low sequence 88
identity and a lack of collinearity between the two species (Supplementary Table 2). In most cases, 89
the sequences showed more similarity to phylogenetically distant bacteria than to each other 90
(Supplementary Table 1), as also supported by phylogenetic analyses (Supplementary Figure 1). 91
These results are consistent with the acquisition of TAs through lateral gene transfer, as extensively 92
demonstrated for other bacteria (Makarova *et al.*, 2009), including endocellular ones as *Rickettsia* 93
(Audoly *et al.*, 2011). 94

To test whether the *Cagg* TAs play a functional role, we selected two of them, a *yoeB-yefM*-like TA 95
pair (CAGGBEG34_v5_20154- CAGGBEG34_v5_20012) and a *chpB-chpS*-like pair 96
(CAGGBEG34_v5_60038- CAGGBEG34_v5_60039) from the complete TA systems. These 97
modules are likely to have a chromosomal and plasmidic localization, respectively (see 98
Supplementary Information) and they both belong to the type II TAs (RelE and MazF 99
superfamilies, respectively). Measurement of the expression of these TAs at different stages in the 100
life cycle of *G. margarita*, from the pre-symbiotic stages (non-germinating, germinating, 101
strigolactone-treated spores) to the formation of the symbiotic mycelium (see Supplementary Figure 102
2 for a depiction of the fungal life cycle), showed that TA gene expression changed throughout the 103

fungal life cycle (Figure 2a and b), with more expression during the fungal pre-symbiotic stages. *G. margarita* is a biotroph, requiring the plant environment to complete its life cycle. The pre-symbiotic stage, when fungal growth mostly depends on its endogenous nutrient supplies, represents a critical step for the fungus as it explores the surrounding soil to find and associate with a plant root. Under these conditions, excessive proliferation of the endobacterium, which acts as an energy sink, might be deleterious for the survival of the fungal/bacterial system. Such stressful conditions might trigger the activation of TA transcription, in contrast to the symbiotic stage, when there is a balanced plant- fungal nutrient exchange. Thus, it is possible that *Cagg* TAs are involved in the bacterial response to the stress experienced during the fungal pre-symbiotic stages and/or in the survival inside spores, by the induction of a state comparable to the TA-induced persistence state described in *Salmonella* (Helaine *et al.*, 2014). Interestingly, the expression of the *Cagg* cell division gene *ftsZ* inversely mirrors the expression of the tested TAs, suggesting that TA activity negatively correlates with bacterial growth (Anca *et al.*, 2009).

To test whether the *Cagg* TAs encode functional toxins and antitoxins, we heterologously expressed the proteins from an inducible promoter, as *Cagg* currently cannot be cultivated *in vitro*. The expression of the YoeB protein strongly affected the growth of *E. coli* cells, producing a decrease in numbers of living cells as early as 3 h after toxin induction. By contrast, the expression of YoeB toxin together with its cognate antitoxin YefM did not affect *E. coli* growth (Figure 2c and d, Supplementary Figure 3). These results demonstrate that YoeB affects *E. coli* cell viability, while YefM prevents its toxic effect, and that this system acts as a TA module. The activity of the ChpB toxin was also analyzed but ChpB showed no effect on *E. coli* growth and viability in our experimental conditions (data not shown).

TAs from the RelE and MazF superfamilies help bacteria cope with nutritional stresses (Gerdes *et al.*, 2005). Recent evidence indicates that *Mycobacterium tuberculosis Rel* loci also react to other stresses, such as oxidative and nitrogen-limiting conditions (Korch *et al.*, 2015). For this reason, we next tested whether an oxidative stress can induce the YoeB-YefM TA gene expression. *Cagg* has been shown to promote antioxidative responses in its fungal host (Vannini *et al.*, 2016); therefore, we challenged germinating spores with H₂O₂. This stress induced the up-regulation of TA gene expression, with statistically significant upregulation of the *yefM* antitoxin gene (Supplementary Figure 4).

The observations described above demonstrate the functionality of a TA module from an endobacterium that lives inside a fungus that lives inside a plant. TA gene expression is regulated throughout the fungal life cycle, and we speculate that it can respond to external stimuli by modulating bacterial cell division. In conclusion, our findings suggest that TAs might represent one

of the genetic determinants that coordinate the <i>Cagg</i> population dynamics with the life cycle of its	138
fungal	139
host.	140
	141
Acknowledgements	141
Research was funded by the local project, 60% to PB. Exchanges between Torino and Nice University were	142
supported by the Italian-French PRES Agreement. The authors wish to express their thanks to Jennifer Mach	143
for the language revision. They also thank Stefano Ghignone, Mara Novero, and Veronica Volpe, as well as	144
the RASTA software developers.	145
	146
Supplementary information is available at The ISME Journal's website (http://	147
www.nature.com/ismej).	148
	149
References	150
Anca IA, Lumini E, Ghignone S, Salvioli A, Bianciotto V, Bonfante P. (2009). The <i>ftsZ</i> gene of the	151
endocellular bacterium ' <i>Candidatus Glomeribacter gigasporarum</i> ' is preferentially expressed during the	152
symbiotic phases of its host mycorrhizal fungus. <i>Mol Plant Microbe Interact</i> 22:302-310.	153
Audoly G, Vincentelli R, Edouard S, Georgiades K, Mediannikov O, Gimenez G <i>et al.</i> (2011). Effect of	154
rickettsial toxin VapC on its eukaryotic host. <i>PLoS One</i> doi: 10.1371/journal.pone.0026528.	155
Bayles KW. (2014). Bacterial programmed cell death: making sense of a paradox. <i>Nat Rev Microbiol</i> 12:63-	156
69. doi: 10.1371/journal.pone.0026528.	157
Bianciotto V, Bandi C, Minerdi D, Sironi M, Tichy HV, Bonfante P. (1996). An obligately endosymbiotic	158
mycorrhizal fungus itself harbors obligately intracellular bacteria. <i>Appl Environ Microbiol</i> 62:3005–3010.	159
Bianciotto V, Lumini E, Bonfante P, Vandamme P. (2003). ' <i>Candidatus Glomeribacter gigasporarum</i> ' gen.	160
nov., sp. nov., an endosymbiont of arbuscular mycorrhizal fungi. <i>Int Syst Evol Micr</i> 53:121–124.	161
Bonfante P, Desirò A. (2017). Who lives in a fungus? The diversity, origins and functions of fungal	162
endobacteria living in the Mucoromycota. <i>ISME J</i> (in press).	163
Gerdes K, Christensen SK, Løbner-Olesen A. (2005). Prokaryotic toxin–antitoxin stress response loci. <i>Nat</i>	164
<i>Rev Microbiol</i> 3:371-382.	165
Ghignone S, Salvioli A, Anca I, Lumini E, Ortu G, Petit L <i>et al.</i> (2012). The genome of the obligate	166
endobacterium of an AM fungus reveals an interphylum network of nutritional interactions. <i>ISME J</i> 6:136–	167
145.	168
Helaine S, Cheverton AM, Watson KG, Faure LM, Matthews SA, Holden DW. (2014). Internalization of	169
<i>Salmonella</i> by macrophages induces formation of nonreplicating persisters. <i>Science</i> 343:204-208.	170
Korch SB, Malhotra V, Contreras H, Clark-Curtiss JE. (2015). The <i>Mycobacterium tuberculosis</i> <i>relBE</i>	171
toxin:antitoxin genes are stress-responsive modules that regulate growth through translation inhibition. <i>J</i>	172
<i>Microbiol</i> 53:783-795.	173

Lackner G, Moebius N, Partida-Martinez LP, Boland S, Hertweck C. (2011). Evolution of an endofungal lifestyle: deductions from the <i>Burkholderia rhizoxinica</i> genome. BMC Genomics doi: 10.1186/1471-2164-12-210.	174 175 176
Leplae R, Geeraerts D, Hallez R, Guglielmini J, Drèze P, Van Melderden L. (2011). Diversity of bacterial type II toxin-antitoxin systems: a comprehensive search and functional analysis of novel families. Nucleic Acids Res 39:5513-5525.	177 178 179
Lipuma J, Cinege G, Bodogai M, Olah B, Kiers A, Endre G <i>et al.</i> (2014). A vapBC-type toxin-antitoxin module of <i>Sinorhizobium meliloti</i> influences symbiotic efficiency and nodule senescence of <i>Medicago sativa</i> . Environ Microbiol 16:3714-3729.	180 181 182
López-Madrigal S, Latorre A, Porcar M, Moya A, Gil R. (2011). Complete genome sequence of " <i>Candidatus Tremblaya princeps</i> " strain PCVAL, an intriguing translational machine below the living-cell status. J Bacteriol 193:5587-5588.	183 184 185
Lumini E, Bianciotto V, Jargeat P, Novero M, Salvioli A, Faccio A <i>et al.</i> (2007). Presymbiotic growth and sporal morphology are affected in the arbuscular mycorrhizal fungus <i>Gigaspora margarita</i> cured of its endobacteria. Cell Microbiol 9:1716–1729.	186 187 188
Makarova KS, Wolf YI, Koonin EV. (2009). Comprehensive comparative-genomic analysis of type 2 toxin-antitoxin systems and related mobile stress response systems in prokaryotes. Biol Direct 4:19.	189 190
Mruk I, Kobayashi I. (2014). To be or not to be: regulation of restriction-modification systems and other toxin-antitoxin systems. Nucleic Acids Res 42:70-86.	191 192
Ohshima S, Sato Y, Fujimura R, Takashima Y, Hamada M, Nishizawa T <i>et al.</i> (2016). <i>Mycoavidus cysteinexigens</i> gen. nov., sp. nov., an endohyphal bacterium isolated from a soil isolate of the fungus <i>Mortierella elongata</i> . Int J Syst Evol Microbiol 66:2052-2057.	193 194 195
Olah B, Kiss E, Gyorgypal Z, Borzi J, Cinege G, Csanadi G <i>et al.</i> (2001). Mutation in the ntrR gene, a member of the vap gene family, increases the symbiotic efficiency of <i>Sinorhizobium meliloti</i> . Mol Plant Microbe Interact 14:887-894.	196 197 198
Pandey DP and Gerdes K. (2005). Toxin-antitoxin loci are highly abundant in free-living but lost from host-associated prokaryotes. Nucleic Acids Res 33:966-976.	199 200
Partida-Martinez LP, Hertweck C. (2005). Pathogenic fungus harbours endosymbiotic bacteria for toxin production. Nature 437:884-888.	201 202
Ramage HR, Connolly LE, Cox JS. (2009). Comprehensive functional analysis of <i>Mycobacterium tuberculosis</i> toxin-antitoxin systems: implications for pathogenesis, stress responses, and evolution. PLoS Genet 5: e1000767.	203 204 205
Salvioli A, Ghignone S, Novero M, Navazio L, Venice F, Bagnaresi P <i>et al.</i> (2016). Symbiosis with an endobacterium increases the fitness of a mycorrhizal fungus, raising its bioenergetic potential. ISME J 10:130-144.	206 207 208
Schuster CF, Bertram R. (2013). Toxin-antitoxin systems are ubiquitous and versatile modulators of prokaryotic cell fate. FEMS Microbiol Lett 340:73-85.	209 210

Sevin EW, Barloy-Hubler F. (2007). RASTA-Bacteria: a web-based tool for identifying toxin-antitoxin loci in prokaryotes. <i>Genome Biol</i> 8:R155.	211 212
Smith SE, Read DJ. (2010). <i>Mycorrhizal symbiosis</i> . Academic Press: London.	213
Uehling J, Gryganskyi A, Hameed K, Tschaplinski T, Misztal PK, Wu S <i>et al.</i> Comparative genomics of <i>Mortierella elongata</i> and its bacterial endosymbiont <i>Mycoavidus cysteinexigens</i> . <i>Environ Microbiol</i> 2017; e-pub ahead of print Jan 11, doi: 10.1111/1462-2920.13669.	214 215 216
Vannini C, Carpentieri A, Salvioli A, Novero M, Marsoni M, Testa L <i>et al.</i> (2016). An interdomain network: the endobacterium of a mycorrhizal fungus promotes antioxidative responses in both fungal and plant hosts. <i>New Phytol</i> 211:265-75.	217 218 219
Yamaguchi Y, Park JH, Masayori Inouye M. (2011). Toxin-antitoxin systems in bacteria and archaea. <i>Annu Rev Genet</i> 45:61-79.	220 221
	222
	223

Figure legends 224

Figure 1 Numbers of TA systems found in the genome of the *Cagg* endosymbiont and its closest relatives. The approximate genome size is given in circles and the number of TAs is given below the circles. Genomes were scanned for TAs using the RASTA and TA finder online tools. 225
226
227

**B. phymatum* is a free-living microbe that can nodulate legume roots (Vandamme *et al.*, 2002). 228

Figure 2 Relative expression of the bacterial gene pairs *yoeB/yefM* (a) and *chpB/chpS* (b) obtained by RT-qPCR according to the $2^{-\Delta\Delta ct}$ method. Means with different superscripts differ significantly (Kruskal-Wallis, $p < 0.05$). The biological material includes the fungal host *G. margarita*, which contains the endobacterium *Cagg*. The tested conditions were the following: non-germinating spores (control); germinating spores; strigolactone-treated spores; symbiotic intra-radical mycelium colonizing *Lotus japonicus* roots; extra-radical mycelium, developing at the root surface. Strigolactones are phytohormones that stimulate AMF hyphal growth and branching (Akyiama *et al.*, 2005). Compared with the pre-symbiotic stages (spores), the expression of the TA operons decreased when the fungus formed the symbiotic mycelium (extra- and intra-radical). 229
230
231
232
233
234
235
236
237

Growth (c) and viability (d) of *E. coli* expressing the YoeB toxin from plasmid pBAD24-YoeB and the YoeB and YefM proteins from plasmid pBAD24-YoeBYefM. The expression was induced by the addition of 1% arabinose to the bacterial cultures. Control cells carried the empty vector pBAD24. Samples taken at 0, 3, and 4 hours after induction were plated and living cell numbers were calculated from three independent experiments. Bars represent standard deviations. 238
239
240
241
242

243

244