

Original Article



Characteristics and patterns of care of endometrial cancer before and during COVID-19 pandemic

OPEN ACCESS

Received: Aug 31, 2021
Revised: Oct 3, 2021
Accepted: Oct 28, 2021
Published online: Nov 12, 2021

Correspondence to

Chiara Cimmino

Department of Obstetrics and Gynaecology,
University of Insubria, Piazza Birelli 1, Varese
21100, Italy.

E-mail: chia.cimmino@gmail.com

Giorgio Bogani

Gynecologic Oncology Unit, Fondazione IRCCS
Istituto Nazionale dei Tumori di Milano, Via
Giacomo Venezian, 1, Milan 20133, Italy.
E-mail: giorgiobogani@yahoo.it

Copyright © 2022. Asian Society of
Gynecologic Oncology, Korean Society of
Gynecologic Oncology, and Japan Society of
Gynecologic Oncology

This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>)
which permits unrestricted non-commercial
use, distribution, and reproduction in any
medium, provided the original work is properly
cited.

ORCID iDs

Giorgio Bogani
<https://orcid.org/0000-0001-8373-8569>
Giovanni Scambia
<https://orcid.org/0000-0002-9503-9041>
Chiara Cimmino
<https://orcid.org/0000-0003-4214-6326>
Francesco Fanfani
<https://orcid.org/0000-0002-4354-5735>
Barbara Costantini
<https://orcid.org/0000-0002-1414-0106>

Giorgio Bogani ¹, Giovanni Scambia ², Chiara Cimmino ³, Francesco Fanfani ²,
Barbara Costantini ², Matteo Loverro ², Gabriella Ferrandina ²,
Fabio Landoni ⁴, Luca Bazzurini ⁴, Tommaso Grassi ⁴, Domenico Vitobello ⁵,
Gabriele Siesto ⁵, Anna Myriam Perrone ⁶, Vanna Zanagnolo ⁷,
Pierandrea De Iaco ⁶, Francesco Multinu ⁷, Fabio Ghezzi ³, Jvan Casarin ³,
Roberto Berretta ⁸, Vito A Capozzi ⁸, Errico Zupi ⁹, Gabriele Centini ⁹,
Antonio Pellegrino ¹⁰, Silvia Corso ¹⁰, Guido Stevenazzi ¹¹, Serena Montoli ¹¹,
Anna Chiara Boschi ¹², Giuseppe Comerci ¹², Pantaleo Greco ¹³,
Ruby Martinello ¹³, Francesco Sopracordevole ¹⁴, Giorgio Giorda ¹⁴,
Tommaso Simoncini ¹⁵, Marta Caretto ¹⁵, Enrico Sartori ¹⁶, Federico Ferrari ¹⁶,
Antonio Cianci ¹⁷, Giuseppe Sarpietro ¹⁷, Maria Grazia Matarazzo ¹⁷,
Fulvio Zullo ¹⁸, Giuseppe Bifulco ¹⁸, Michele Morelli ¹⁹, Annamaria Ferrero ²⁰,
Nicoletta Biglia ²⁰, Fabio Barra ²¹, Simone Ferrero ²¹,
Umberto Leone Roberti Maggiore ¹, Stefano Cianci ²², Vito Chiantera ²³,
Alfredo Ercoli ²², Giulio Sozzi ²³, Angela Martocchia ²⁴, Sergio Schettini ²⁴,
Teresa Orlando ²⁴, Francesco G Cannone ²⁵, Giuseppe Ettore ²⁵,
Andrea Puppo ²⁶, Martina Borghese ²⁶, Canio Martinelli ²², Ludovico Muzii ²⁷,
Violante Di Donato ²⁷, Lorenza Driul ²⁸, Stefano Restaino ²⁸, Alice Bergamini ²⁹,
Giorgio Candotti ²⁹, Luca Bocciolone ²⁹, Francesco Plotti ³⁰, Roberto Angioli ³⁰,
Giulia Mantovani ³¹, Marcello Ceccaroni ³¹, Chiara Cassani ³²,
Mattia Dominoni ³², Laura Giambanco ³³, Silvia Amodeo ³³, Livio Leo ³⁴,
Raphael Thomasset ³⁴, Diego Raimondo ³⁵, Renato Seracchioli ³⁵,
Mario Malzoni ³⁶, Franco Gorlero ³⁷, Martina Di Luca ³⁷, Enrico Busato ³⁸,
Sami Kilzie ³⁸, Andrea Dell'Acqua ³⁹, Giovanna Scarfone ³⁹, Paolo Vercellini ³⁹,
Marco Petrillo ⁴⁰, Salvatore Dessole ⁴⁰, Giampiero Capobianco ⁴⁰,
Andrea Ciavattini ⁴¹, Giovanni Delli Carpini ⁴¹, Luca Giannella ⁴¹,
Liliana Mereu ⁴², Saverio Tateo ⁴², Flavia Sorbi ⁴³, Massimiliano Fambrini ⁴³,
Stefania Cicogna ⁴⁴, Federico Romano ⁴⁴, Giuseppe Ricci ^{44,45},
Giuseppe Trojano ⁴⁶, Roberto Consonni ⁴⁷, Simona Cantaluppi ⁴⁷,
Antonio Lippolis ⁴⁸, Raffaele Tinelli ⁴⁸, Giovanni D'Ippolito ⁴⁹,
Lorenzo Aguzzoli ⁴⁹, Vincenzo D Mandato ⁴⁹, Stefano Palomba ⁵⁰,
Davide Calandra ⁵¹, Maurizio Rosati ^{51,52}, Cinzia Gallo ⁵³, Daniela Surico ⁵⁴,
Valentino Remorgida ⁵⁴, Francesco Ruscitto ⁵⁵, Paolo Beretta ⁵⁵,
Pierluigi Benedetti Panici ²⁷, Francesco Raspagliesi ¹

¹Gynecologic Oncology Unit, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Milan, Italy

²Gynecologic Oncology Unit, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy

³Department of Obstetrics and Gynaecology, University of Insubria, F. Del Ponte Hospital, Varese, Italy

⁴Department of Obstetrics and Gynaecology, San Gerardo Hospital, Monza, Italy

⁵Unit of Gynecology, Humanitas Cancer Center, Humanitas Clinical and Research Center - IRCCS, Milan, Italy

⁶Unit of Gynecology, AOU S. Orsola-Malpighi, Bologna, Italy

⁷Department of Gynecologic Oncology, IEO, European Institute of Oncology IRCCS, Milan, Italy

⁸Department of Obstetrics and Gynaecology, University of Parma, Parma, Italy

⁹Department of Obstetrics and Gynaecology, University of Siena, Siena, Italy

¹⁰Department of Obstetrics and Gynaecology, ASST Lecco - Ospedale Alessandro Manzoni, Lecco, Italy

¹¹Department of Obstetrics and Gynaecology, ASST OVEST MI, Legnano (Milan) Hospital, Legnano, Italy

Matteo Loverro 
<https://orcid.org/0000-0002-9740-3169>
 Gabriella Ferrandina 
<https://orcid.org/0000-0003-4672-4197>
 Fabio Landoni 
<https://orcid.org/0000-0001-6734-739X>
 Luca Bazzurini 
<https://orcid.org/0000-0001-7644-6971>
 Tommaso Grassi 
<https://orcid.org/0000-0002-3019-1077>
 Domenico Vitobello 
<https://orcid.org/0000-0001-7455-1831>
 Gabriele Siesto 
<https://orcid.org/0000-0003-1316-2589>
 Anna Myriam Perrone 
<https://orcid.org/0000-0003-3140-4772>
 Vanna Zanagnolo 
<https://orcid.org/0000-0002-7123-191X>
 Pierandrea De Iaco 
<https://orcid.org/0000-0002-8841-6531>
 Francesco Multinu 
<https://orcid.org/0000-0001-8535-4059>
 Fabio Ghezzi 
<https://orcid.org/0000-0003-3949-5410>
 Jvan Casarin 
<https://orcid.org/0000-0001-9519-1097>
 Roberto Berretta 
<https://orcid.org/0000-0003-2324-5409>
 Vito A Capozzi 
<https://orcid.org/0000-0003-4720-5663>
 Errico Zupi 
<https://orcid.org/0000-0003-0735-6301>
 Gabriele Centini 
<https://orcid.org/0000-0002-6113-7401>
 Antonio Pellegrino 
<https://orcid.org/0000-0001-7918-783X>
 Silvia Corso 
<https://orcid.org/0000-0001-7258-6600>
 Guido Stevenazzi 
<https://orcid.org/0000-0003-2855-9829>
 Serena Montoli 
<https://orcid.org/0000-0002-4553-460X>
 Anna Chiara Boschi 
<https://orcid.org/0000-0001-7837-2697>
 Giuseppe Comerchi 
<https://orcid.org/0000-0003-3723-5391>
 Pantaleo Greco 
<https://orcid.org/0000-0003-2461-6777>
 Ruby Martinello 
<https://orcid.org/0000-0003-1979-8721>
 Francesco Sopracordevole 
<https://orcid.org/0000-0001-5562-4353>
 Giorgio Giorda 
<https://orcid.org/0000-0001-6386-3565>
 Tommaso Simoncini 
<https://orcid.org/0000-0002-2971-0079>
 Marta Caretto 
<https://orcid.org/0000-0002-1157-9930>

¹²Department of Obstetrics and Gynaecology, AUSL Romagna, Ospedale "Santa Maria delle Croci", Ravenna, Italy
¹³Clinica Ostetrica e Ginecologica - Dipartimento Scienze Mediche - Università di Ferrara, Ferrara, Italy
¹⁴Gynecological Oncology Unit, Centro di Riferimento Oncologico - National Cancer Institute, Aviano, Italy
¹⁵Division of Obstetrics and Gynecology, Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Italy
¹⁶Department of Clinical and Experimental Sciences, University of Brescia, Brescia, Italy
¹⁷Department of General Surgery and Medical Surgical Specialties, Gynecological Clinic, University of Catania, Catania, Italy
¹⁸Department of Obstetrics and Gynaecology, Azienda Ospedaliera Universitaria - Federico II, Naples, Italy
¹⁹Department of Obstetrics and Gynaecology, AO "S.S. Annunziata", Cosenza, Italy
²⁰Academic Department of Obstetrics and Gynecology, Mauriziano Hospital, Torino, Italy
²¹Academic Unit of Obstetrics and Gynecology, IRCCS Ospedale Policlinico San Martino, Genova, Italy
²²Unit of Gynecology and Obstetrics, Department of Human Pathology in Adulthood and Childhood "G. Barresi", University Hospital G. Martino, University of Messina, Messina, Italy
²³Department of Gynecologic Oncology, University of Palermo, Palermo, Italy
²⁴Department of Obstetrics and Gynaecology, AOR San Carlo, Potenza, Italy
²⁵Department of Obstetrics and Gynaecology, ARNAS Garibaldi Catania, Catania, Italy
²⁶Department of Obstetrics and Gynaecology, ASO Santa Croce e Carle, Cuneo, Italy
²⁷Department of Maternal and Child Health and Urological Sciences, Sapienza University of Rome, Policlinico Umberto I, Rome, Italy
²⁸Department of Maternal and Child Health, University-Hospital of Udine, Udine, Italy
²⁹Department of Obstetrics and Gynaecology, IRCCS San Raffaele Hospital, Milan, Italy
³⁰Department of Obstetrics and Gynecology, Campus Bio-Medico University of Rome, Rome, Italy
³¹Department of Obstetrics and Gynecology, Gynecology Oncology and Minimally-Invasive Pelvic Surgery, International School of Surgical Anatomy, Sacred Heart Hospital Negrar, Verona, Italy
³²Department of Obstetrics and Gynecology, IRCCS Foundation Policlinico San Matteo and University of Pavia, Pavia, Italy
³³Department of Obstetrics and Gynecology, S. Antonio Abate Hospital, Trapani, Italy and Department of Health Promotion, Mother and Child Care, Internal Medicine and Medical Specialties (PROMISE), University of Palermo, Palermo, Italy
³⁴Departments of Gynecology & Obstetrics, Hopital Beauregard, AUSL Valleè d'Aoste, Aosta, Italy
³⁵Division of Gynaecology and Human Reproduction Physiopathology, Department of Medical and Surgical Sciences (DIMEC), IRCCS Azienda Ospedaliero-Universitaria di Bologna. S. Orsola Hospital, University of Bologna, Bologna, Italy
³⁶Endoscopica Malzoni, Center for Advanced Endoscopic Gynecologic Surgery, Avellino, Italy
³⁷Department of Obstetrics and Gynaecology, Ente Ospedaliero Ospedali Galliera, Genova, Italy
³⁸Department of Obstetrics and Gynaecology, Ospedale di Treviso, Treviso, Italy
³⁹Gynaecology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy
⁴⁰Gynecologic and Obstetric Unit, Department of Medical, Surgical and Experimental Sciences, University of Sassari, Sassari, Italy
⁴¹Gynecologic Section, Department of Odontostomatologic and Specialized Clinical Sciences, Università Politecnica delle Marche, Ancona, Italy
⁴²Gynecological Oncology Unit, Santa Chiara Hospital, Trento, Italy
⁴³Gynecology Unit, Careggi University Hospital, Department of Biomedical, Experimental and Clinical Sciences "Mario Serio", University of Florence, Florence, Italy
⁴⁴Department of Obstetrics and Gynaecology, Institute for Maternal and Child Health, IRCCS 'Burlo Garofolo', Trieste, Italy
⁴⁵Department of Medicine, Surgery and Health Sciences, University of Trieste, Trieste, Italy
⁴⁶Department of Obstetrics and Gynaecology, Madonna delle Grazie Hospital ASM, Matera, Italy
⁴⁷Gynecology Unit, Ospedale Valduce, Como, Italy
⁴⁸Unit of Obstetrics and Gynaecology, Valle D'Itra Hospital, Martina Franca, Taranto, Italy
⁴⁹Unit of Obstetrics and Gynecology, Azienda Unità Sanitaria Locale - IRCCS, Reggio Emilia, Italy
⁵⁰Unit of Obstetrics and Gynecology, GOM of Reggio Calabria & Magna Graecia University of Catanzaro, Catanzaro, Italy
⁵¹Unit of Obstetrics and Gynecology, University G. D'Annunzio of Chieti, Pescara, Italy
⁵²Unit of Obstetrics and Gynecology, Santo Spirito Hospital, Pescara, Italy
⁵³Unit of Obstetrics and Gynecology, Università "Magna Graecia" di Catanzaro - AO "Pugliese - Ciaccio" Catanzaro, Italy
⁵⁴Unit of Obstetrics and Gynecology, University of Eastern Piedmont, Novara, Italy
⁵⁵Gynecology Unit, Ospedale Valduce, Como - ASST Lariana, S. Anna, Como, Italy

Enrico Sartori 
<https://orcid.org/0000-0003-4076-303X>
 Federico Ferrari 
<https://orcid.org/0000-0001-7065-2432>
 Antonio Cianci 
<https://orcid.org/0000-0003-2758-3413>
 Giuseppe Sarpietro 
<https://orcid.org/0000-0003-4891-9000>
 Maria Grazia Matarazzo 
<https://orcid.org/0000-0002-5183-4422>
 Fulvio Zullo 
<https://orcid.org/0000-0003-3726-7561>
 Giuseppe Bifulco 
<https://orcid.org/0000-0002-1788-5170>
 Michele Morelli 
<https://orcid.org/0000-0003-1759-571X>
 Annamaria Ferrero 
<https://orcid.org/0000-0003-1544-6016>
 Nicoletta Biglia 
<https://orcid.org/0000-0003-1009-5309>
 Fabio Barra 
<https://orcid.org/0000-0003-4117-6603>
 Simone Ferrero 
<https://orcid.org/0000-0003-2225-5568>
 Umberto Leone Roberti Maggiore 
<https://orcid.org/0000-0002-3744-2668>
 Stefano Cianci 
<https://orcid.org/0000-0002-0548-9891>
 Vito Chiantera 
<https://orcid.org/0000-0002-6294-3720>
 Alfredo Ercoli 
<https://orcid.org/0000-0002-3377-2803>
 Giulio Sozzi 
<https://orcid.org/0000-0001-5847-8822>
 Angela Martocchia 
<https://orcid.org/0000-0001-5876-1089>
 Sergio Schettini 
<https://orcid.org/0000-0003-3886-478X>
 Teresa Orlando 
<https://orcid.org/0000-0002-3689-4611>
 Francesco G Cannone 
<https://orcid.org/0000-0001-7681-1801>
 Giuseppe Ettore 
<https://orcid.org/0000-0002-5765-7682>
 Andrea Puppo 
<https://orcid.org/0000-0003-1714-7765>
 Martina Borghese 
<https://orcid.org/0000-0002-6504-9978>
 Canio Martinelli 
<https://orcid.org/0000-0002-0587-8467>
 Ludovico Muzii 
<https://orcid.org/0000-0001-7195-9583>
 Violante Di Donato 
<https://orcid.org/0000-0002-9254-5790>
 Lorenza Driul 
<https://orcid.org/0000-0003-1896-9410>
 Stefano Restaino 
<https://orcid.org/0000-0002-7848-0329>

ABSTRACT

Objective: Coronavirus disease 2019 (COVID-19) outbreak has correlated with the disruption of screening activities and diagnostic assessments. Endometrial cancer (EC) is one of the most common gynecological malignancies and it is often detected at an early stage, because it frequently produces symptoms. Here, we aim to investigate the impact of COVID-19 outbreak on patterns of presentation and treatment of EC patients.

Methods: This is a retrospective study involving 54 centers in Italy. We evaluated patterns of presentation and treatment of EC patients before (period 1: March 1, 2019 to February 29, 2020) and during (period 2: April 1, 2020 to March 31, 2021) the COVID-19 outbreak.

Results: Medical records of 5,164 EC patients have been retrieved: 2,718 and 2,446 women treated in period 1 and period 2, respectively. Surgery was the mainstay of treatment in both periods ($p=0.356$). Nodal assessment was omitted in 689 (27.3%) and 484 (21.2%) patients treated in period 1 and 2, respectively ($p<0.001$). While, the prevalence of patients undergoing sentinel node mapping (with or without backup lymphadenectomy) has increased during the COVID-19 pandemic (46.7% in period 1 vs. 52.8% in period 2; $p<0.001$). Overall, 1,280 (50.4%) and 1,021 (44.7%) patients had no adjuvant therapy in period 1 and 2, respectively ($p<0.001$). Adjuvant therapy use has increased during COVID-19 pandemic ($p<0.001$).

Conclusion: Our data suggest that the COVID-19 pandemic had a significant impact on the characteristics and patterns of care of EC patients. These findings highlight the need to implement healthcare services during the pandemic.

Keywords: Endometrial Cancer; COVID-19; Uterine cancer; SARS-CoV-2


























Synopsis

The prevalence of patients with early-stage endometrial cancer (EC) has been lower during coronavirus disease 2019 (COVID-19) pandemic than before its onset. Further evidence is needed to assess the impact of COVID-19 pandemic on survival outcomes of EC patients.

INTRODUCTION

Endometrial cancer (EC) is one of the most common gynecological cancers in developed countries [1]. It is estimated that more than 55,000 new EC cases are diagnosed every year in the United States [1]. Over the last decade, the incidence of EC has increased by more than 20,000 cases/year [2]. Similarly, the incidence of EC in Europe is increasing due to the aging of populations and increased prevalence of obesity [3]. Generally, EC is considered a disease with good prognosis, since the majority of patients are diagnosed at early stage of disease. The main reason for this is that EC frequently produces symptoms, namely abnormal vaginal bleeding. Although no screening activities are approved for early detection of EC, regular visits and prompt assessments in patients with new-onset symptoms have been useful in improving early detection of uterine malignancies [4].

Over the last year, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) respiratory disease (coronavirus disease 2019, COVID-19) has been spreading worldwide, dramatically changing everyday life. On February 3, 2020, the World Health Organization (WHO) proposed the '2019 COVID-19 Strategic Preparedness and Response Plan,' which includes accelerating research and development processes as one of the main strategies against the

Alice Bergamini 
<https://orcid.org/0000-0002-1915-8313>
 Giorgio Candotti 
<https://orcid.org/0000-0002-6475-7278>
 Luca Boccione 
<https://orcid.org/0000-0001-5098-0500>
 Francesco Plotti 
<https://orcid.org/0000-0002-6218-5616>
 Roberto Angioli 
<https://orcid.org/0000-0002-8528-4318>
 Giulia Mantovani 
<https://orcid.org/0000-0001-6372-0046>
 Marcello Ceccaroni 
<https://orcid.org/0000-0002-6797-4876>
 Chiara Cassani 
<https://orcid.org/0000-0002-8055-5086>
 Mattia Dominoni 
<https://orcid.org/0000-0002-6322-607X>
 Laura Giambanco 
<https://orcid.org/0000-0002-7755-6037>
 Silvia Amodeo 
<https://orcid.org/0000-0002-1836-4211>
 Livio Leo 
<https://orcid.org/0000-0002-3452-0608>
 Raphael Thomasset 
<https://orcid.org/0000-0002-4655-8005>
 Diego Raimondo 
<https://orcid.org/0000-0002-3235-4378>
 Renato Seracchioli 
<https://orcid.org/0000-0001-7487-1333>
 Mario Malzoni 
<https://orcid.org/0000-0003-2514-7159>
 Franco Gorlero 
<https://orcid.org/0000-0003-1166-6951>
 Martina Di Luca 
<https://orcid.org/0000-0003-2184-749X>
 Enrico Busato 
<https://orcid.org/0000-0001-6739-2326>
 Sami Kilzie 
<https://orcid.org/0000-0002-2511-7040>
 Andrea Dell'Acqua 
<https://orcid.org/0000-0002-7589-5019>
 Giovanna Scarfone 
<https://orcid.org/0000-0001-9838-9998>
 Paolo Vercellini 
<https://orcid.org/0000-0003-4195-0996>
 Marco Petrillo 
<https://orcid.org/0000-0003-0306-4328>
 Salvatore Dessole 
<https://orcid.org/0000-0001-8287-9079>
 Giampiero Capobianco 
<https://orcid.org/0000-0002-5523-8943>
 Andrea Ciavattini 
<https://orcid.org/0000-0003-0074-5996>
 Giovanni Delli Carpini 
<https://orcid.org/0000-0003-2849-4690>
 Luca Giannella 
<https://orcid.org/0000-0002-6405-8522>




COVID-19 outbreak. On March 11, 2020, the WHO declared COVID-19 a pandemic [5]. Italy was the first European country suffering the spread of COVID-19. In order to flatten the growth curve and to face the growing need for assistance of COVID-19 patients, physicians have started delaying non-urgent procedures. This has partly translated in the risk of diagnosis and treatment procrastination, with significant negative impacts on the outcomes of patients with cancer. Indeed, COVID-19 pandemic has negatively influenced the timing of screening activities and regular periodic visits [6]. Disruption of screening activities, periodic visits, and diagnostic assessments have delayed regular medical investigations [6-9]. This is of paramount importance in subjects at risk, and especially to those patients who developed new symptoms and need to be investigated in the suspect of cancer diagnosis. To date, several opinions and surveys have been published on this topic [5,10,11]. Emerging data suggest that the COVID-19 outbreak might influence patterns of disease presentation, compromising the possibility of early access to care for patients with malignancies [6-10]. However, so far, no studies specifically evaluated the real impact of COVID-19 on the attitudes, practice, and the workflow in the setting of oncology. We evaluated patterns of presentation and treatment of EC patients before and during the COVID-19 pandemic. EC represents an ideal target to assess the impact of COVID-19 for three main reasons: i) it is a common gynecological occurrence; ii) it is generally detected at an early stage of disease; and iii) it is usually related to favorable oncologic outcomes. Hence, we performed a large multi-institutional study evaluating the COVID-19 pandemic on patients diagnosed with EC, with the aim to verify whether diagnostic and treatment's attitudes have changed across these 2 periods.

MATERIALS AND METHODS

1. Methods

This is a multi-institutional retrospective study coordinated by the Fondazione IRCCS Istituto Nazionale dei Tumori. As coordinator center the Institutional Review Board of the Fondazione IRCCS Istituto Nazionale dei Tumori approved this investigation (#62/20). For the present study, we collected medical records of consecutive patients with newly diagnosed EC treated in Italy before (period 1: from March 1, 2019 to February 29, 2020) and during (period 2: from April 1, 2020 to March 31, 2021) the COVID-19 outbreak. This study involved 54 high-volume centers in Italy. We collected data from any regional referral centers or cancer centers in Italy. **Table S1** displays the centers participating in the study.

The primary endpoint measures were: i) the prevalence of patients affected by International Federation of Obstetrics and Gynecologists (FIGO) stage >I disease at presentation; ii) the prevalence of adjuvant therapy indication in the 2 periods. As secondary endpoints, we sought to report changes in EC management during the COVID-19 outbreak. We included consecutive patients receiving treatment (i.e., surgery, radiotherapy, and/or anticancer systemic treatment) in period 1 and 2. To minimize possible biases we decided to exclude cases treated in March 2020, focusing only on cases treated before and during the COVID-19 outbreak. In March 2020 only few regions in the North of Italy were affected by the pandemic. Since April 2020, the whole Italian healthcare care system was impacted by COVID-19 [12]. We included all patients aged ≥ 18 years old, with a confirmed histological diagnosis of EC, regardless of the type of treatment. In all included centers, data concerning surgical procedures, peri-operative details, adjuvant therapy, as well as follow-up evaluations were recorded in computerized databases, updated by trained residents and nurses on a regular basis. The taxonomy proposed by the WHO was used to designate EC histological

Liliana Mereu 
<https://orcid.org/0000-0002-0610-5132>
 Saverio Tateo 
<https://orcid.org/0000-0002-5824-8020>
 Flavia Sorbi 
<https://orcid.org/0000-0001-7277-6420>
 Massimiliano Fambrini 
<https://orcid.org/0000-0003-0461-6390>
 Stefania Cicogna 
<https://orcid.org/0000-0003-4432-0826>
 Federico Romano 
<https://orcid.org/0000-0003-2157-8330>
 Giuseppe Ricci 
<https://orcid.org/0000-0002-8031-1102>
 Giuseppe Trojano 
<https://orcid.org/0000-0002-8737-9943>
 Roberto Consonni 
<https://orcid.org/0000-0003-0184-2384>
 Simona Cantaluppi 
<https://orcid.org/0000-0002-0476-189X>
 Antonio Lippolis 
<https://orcid.org/0000-0002-3700-5513>
 Raffaele Tinelli 
<https://orcid.org/0000-0001-5597-3035>
 Giovanni D'Ippolito 
<https://orcid.org/0000-0003-0050-4498>
 Lorenzo Aguzzoli 
<https://orcid.org/0000-0002-0723-151X>
 Vincenzo D Mandato 
<https://orcid.org/0000-0001-6537-1046>
 Stefano Palomba 
<https://orcid.org/0000-0003-2767-8295>
 Davide Calandra 
<https://orcid.org/0000-0001-9984-5350>
 Maurizio Rosati 
<https://orcid.org/0000-0002-2887-5576>
 Cinzia Gallo 
<https://orcid.org/0000-0001-8064-7012>
 Daniela Surico 
<https://orcid.org/0000-0001-6801-0043>
 Valentino Remorgida 
<https://orcid.org/0000-0002-8660-0044>
 Francesco Ruscitto 
<https://orcid.org/0000-0002-4928-824X>
 Paolo Beretta 
<https://orcid.org/0000-0001-9111-174X>
 Pierluigi Benedetti Panici 
<https://orcid.org/0000-0001-6752-2039>
 Francesco Raspagliesi 
<https://orcid.org/0000-0001-8953-1657>

Conflict of Interest

The first author is a principal editor of the Journal of Gynecologic Oncology. No other potential conflict of interest relevant to this article was reported

subtypes [13,14]. The degree of glandular differentiation and cytologic atypia to determine architectural grade and stage were reported according to the FIGO criteria [13,14]. Details about surgical techniques, adjuvant therapies, and follow-up schedules are reported elsewhere [15-18]. During the two study periods, there were no significant differences in the facilities available for patients care and in the referral patterns of our service. Other features of patient management remained consistent in the two periods.

2. Statistical methods

Basic descriptive statistics were used to describe the study populations. Differences in categorical variables were analyzed using the Fisher exact and χ^2 test when comparing 2 and 3 (or more) groups, respectively. When indicated odds ratio (OR) and 95% confidence intervals (95% CI) were calculated. Student's t-test and Mann-Whitney test were used to compare continuous variables as appropriate. The p-values <0.05 were considered statistically significant. Statistical analysis was performed with GraphPad Prism version 6.0 (GraphPad Software, San Diego CA, USA) and IBM-Microsoft SPSS version 20.0 (IBM Corp., Armonk, NY, USA) for Mac.

RESULTS

Charts of 5,164 EC patients were retrieved from 54 Italian centers over the whole study period. Overall, 2,718 and 2,446 women with EC received treatment in period 1 and 2, respectively.

Table 1 shows the main characteristics of the study population in the 2 time periods. The prevalence of patients aged > 65 years was similar between the 2 study periods (1,400 [51.5%] in period 1 vs. 1,248 [51.0%]; p=0.726). Similarly, the prevalence of elderly patients (i.e., aged >85 years) was comparable between groups (189 [6.9%] vs. 180 [7.4%]; p=0.572).

Considering data on the histological characterization, the prevalence of endometrioid FIGO grade 1, 2, and 3 was consistent over the study period (p=0.855). However, the prevalence

Table 1. Characteristics of EC patients included in the study

Characteristics	Period*		p-value
	Period 1	Period 2	
No. of cases	2,718	2,446	-
Age of patients (yr)			0.742
<50	306 (11.3)	262 (10.7)	-
50-64	1,012 (37.2)	936 (38.2)	-
65-84	1,211 (44.6)	1,068 (43.7)	-
>84	189 (6.9)	180 (7.4)	-
Histology			0.178
Endometrioid FIGO G1	808 (29.7)	719 (29.3)	-
Endometrioid FIGO G2	1,019 (37.5)	878 (35.9)	-
Endometrioid FIGO G3	447 (16.4)	400 (16.4)	-
Non-endometrioid	425 (15.6)	438 (17.9)	-
Unknown	19 (0.7)	11 (0.5)	-
FIGO stage			0.003
Stage I	2,021 (74.3)	1,754 (71.7)	-
Stage II	179 (6.6)	176 (7.2)	-
Stage III	348 (12.8)	349 (14.3)	-
Stage IV	129 (4.7)	167 (6.8)	-

Data are reported in number (%).

EC, endometrial cancer; COVID-19, coronavirus disease 2019; FIGO, International Federation of Obstetrics and Gynecologists.

*This study has evaluated patterns of presentation and treatment of EC patients before (period 1: March 1, 2019 to February 29, 2020) and during (period 2: April 1, 2020 to March 31, 2021) the COVID-19 outbreak.

Author Contributions

Conceptualization: B.G.¹; Data curation: C.B., L.M., B.L.¹, G.T., V.D., S.G.², P.A.M., Z.V., D.I.P., M.F., C.J., B.R., C.V.A., Z.E., C.G.¹, P.A.¹, C.S.¹, M.S., B.A.C., C.G.², G.P., M.R., S.F.¹, G.G., S.T., C.M.¹, S.E., F.F.², C.A.¹, S.G.⁴, M.M.G., Z.F., B.G.², M.M.¹, F.A., B.N., B.F., F.S., C.S.², S.G.⁵, M.A., S.S., O.T., C.F.G., E.G., P.A.², B.M., M.C., D.L., R.S., B.A., C.G.³, B.L.², P.F., M.G., C.M.², C.C.², D.M., G.L.¹, A.S., L.L., T.R.¹, R.D., M.M.², G.F.², D.L.M., B.E., K.S., D.A.A., S.G.⁶, P.M., D.S., C.G.⁴, C.A.², D.C.G., G.L.², M.L.², T.S., S.F.², F.M., C.S.³, R.F.¹, R.G., T.G., C.R., C.S.⁴, L.A., T.R.², D.J.G., A.L., M.V.D., P.S., C.D., R.M., G.C., S.D., R.V., R.F.²; Investigation: C.C., L.R.M.U.; Methodology: B.G.¹; Project administration: B.G.¹; Supervision: B.G.¹, S.G.¹, F.F.¹, F.G., L.F., G.F.¹, S.G.³, C.V., E.A., M.L.¹, D.D.V., A.R., S.R., B.P., R.F.³; Validation: F.F.¹, S.G.², V.P., B.P.P., R.F.³; Writing - original draft: B.G.¹; Writing - review & editing: S.G.¹, C.C.¹, C.B., C.J., R.F.³

¹B.G., Giorgio Bogani; ²B.G., Giuseppe Bifulco

¹B.L., Luca Bazzurini; ²B.L., Luca Boccione

¹C.A., Antonio Cianci; ²C.A., Andrea Ciavattini

¹C.C., Chiara Cimmino; ²C.C., Chiara Cassani

¹C.G., Gabriele Centini; ²C.G., Giuseppe

Comerci; ³C.G., Giorgio Candotti; ⁴C.G.,

Giampiero Capobianco

¹C.M., Marta Caretto; ²C.M., Marcello

Ceccaroni

¹C.S., Silvia Corso; ²C.S., Stefano Cianci; ³C.S.,

Stefania Cicogna; ⁴C.S., Simona Cantaluppi

¹F.F., Francesco Fanfani; ²F.F., Federico Ferrari

¹G.F., Fabio Ghezzi; ²G.F., Franco Gorlero

¹G.L., Laura Giambanco; ²G.L., Luca Giannella

¹M.L., Ludovico Muzii; ²M.L., Liliana Mereu

¹M.M., Michele Morelli; ²M.M., Mario Malzoni

¹P.A., Antonio Pellegrino; ²P.A., Andrea Puppo

¹R.F., Federico Romano; ²R.F., Francesco

Ruscitto; ³R.F., Francesco Raspagliesi

¹S.F., Francesco Sopracordevole; ²S.F., Flavia

Sorbi

¹S.G., Giovanni Scambia; ²S.G., Gabriele Siesto;

³S.G., Guido Stevenazzi; ⁴S.G., Giuseppe

Sarpietro; ⁵S.G., Giulio Sozzi; ⁶S.G., Giovanna

Scarfone

¹T.R., Raphael Thomasset; ²T.R., Raffaele Tinelli

Table 2. Details of treatment modalities adopted for managing EC before and during COVID-19 pandemic

Variables	Period*		p-value
	Period 1	Period 2	
No. of cases	2,718	2,446	-
Primary treatment			0.361
Surgery	2,539 (93.4)	2,286 (93.5)	-
Other therapies	169 (6.2)	156 (6.4)	-
No treatment/palliation	10 (0.4)	4 (0.2)	-
Other non-surgical treatments			0.048
Radiotherapy	24 (0.9)	21 (0.9)	-
Chemotherapy	29 (1.1)	49 (2.0)	-
Radio + chemotherapy	25 (0.9)	16 (0.7)	-
Hormonal therapy	19 (0.7)	14 (0.6)	-
IUD/hysteroscopic resection	72 (2.6)	56 (2.3)	-
Type of surgery†			0.096
Laparoscopy	1,400 (55.1)	1,273 (55.7)	-
Robotic assisted	448 (17.6)	390 (17.1)	-
Open surgery	666 (26.2)	582 (25.5)	-
Vaginal	25 (0.9)	41 (1.7)	-
Waiting time between diagnosis and surgery (days)†	25 (7–41)	23 (6–53)	0.654
Type of nodal assessment at surgery†			<0.001
Sentinel node mapping	961 (37.8)	973 (42.5)	-
Sentinel node mapping + lymphadenectomy	224 (8.9)	234 (10.2)	-
Pelvic Lymphadenectomy	456 (17.9)	422 (18.5)	-
Pelvic + para-aortic lymphadenectomy	206 (8.1)	173 (7.6)	-
None	692 (27.3)	484 (21.2)	-

Data are reported in number (%), or median (range).

EC, endometrial cancer; COVID-19, coronavirus disease 2019; IUD, intra-uterine devices

*This study has evaluated patterns of presentation and treatment of EC patients before (period 1: March 1, 2019 to February 29, 2020) and during (period 2: April 1, 2020 to March 31, 2021) the COVID-19 outbreak; †This analysis is restricted only to patients having surgery.

of non-endometrioid EC was lower in period 1 than in period 2 (15.6% vs. 17.9%; $p=0.032$).

Table 2 reports details on the treatment of patients in the 2 study periods. Surgery was the mainstay of treatment before and during the COVID-19 pandemic. Overall, 2,539 and 2,286 women received surgery in period 1 and 2, respectively (93.4% vs. 93.5%; $p=0.948$). Primary conservative attempts (i.e., progesterone-based therapy) was performed in 72 (2.7%) and 56 (2.3%) patients in period 1 and 2, respectively ($p=0.406$).

The adoption of minimally invasive surgery was consistent in the two study periods ($p=0.976$). Before COVID-19 pandemic, 1,848 (72.8%), 666 (26.3%), and 25 (0.9%) patients had minimally invasive, open and vaginal surgery, respectively. During the COVID-19 pandemic, 1,663 (72.8%), 582 (25.5%), and 41 (1.7%) patients had minimally invasive, open, and vaginal surgery, respectively. Restricting the analysis to patients treated with surgery, we observed that sentinel node mapping was the most adopted method before and during the COVID-19 pandemic. Before the COVID-19 pandemic, sentinel node mapping, sentinel node mapping plus backup lymphadenectomy, and lymphadenectomy (pelvic and/or para-aortic) were performed in 961 (37.0%), 224 (8.9%), and 662 (26.0%) patients, respectively. During the COVID-19 pandemic, sentinel node mapping, sentinel node mapping plus backup lymphadenectomy, and lymphadenectomy (pelvic and/or para-aortic) were performed in 973 (42.5%), 234 (10.2%), and 595 (26.1%) patients, respectively. Nodal disease assessment was omitted in 692 (27.3%) and 484 (21.2%) patients treated in periods 1 and 2, respectively ($p<0.001$). Conversely, the prevalence of patients undergoing sentinel node mapping (with or without backup lymphadenectomy) has increased during the COVID-19 pandemic (46.7% in period 1 vs. 52.8% in period 2; $p<0.001$).

Table 3. Details of treatment modalities adopted for managing EC before and during COVID-19 pandemic

Variables	Period*		p-value
	Period 1	Period 2	
No. of patients having surgery	2,539	2,286	-
Adjuvant therapy [†]			<0.001
Yes	1,259 (49.6)	1,265 (55.3)	-
No	1,280 (50.4)	1,021 (44.7)	-
Type of adjuvant therapy [‡]			0.064
Vaginal brachytherapy (VB)	304 (11.9)	253 (11.1)	-
External radiotherapy (+/-VB)	317 (12.5)	323 (14.1)	-
Chemotherapy (+/-VB)	274 (10.8)	310 (13.5)	-
Chemo-radiotherapy	364 (14.3)	379 (16.6)	-

Data are reported in number (%).

EC, endometrial cancer; COVID-19, coronavirus disease 2019; VB, vaginal brachytherapy.

*This study has evaluated patterns of presentation and treatment of EC patients before (period 1: March 1, 2019 to February 29, 2020) and during (period 2: April 1, 2020 to March 31, 2021) the COVID-19 outbreak; [†]This analysis is restricted only to patients having surgery; [‡]This analysis is restricted only to patients having surgery plus adjuvant therapy.

Table 3 shows the details of adjuvant treatment used in the study population, before and during COVID-19 pandemic. Overall, 1,280 (50.4%) and 1,021 (44.7%) patients had no adjuvant therapy in period 1 and 2, respectively ($p < 0.001$). The adoption of vaginal brachytherapy as adjuvant treatment remained stable in the study periods (11.9% vs. 11.1%; $p = 0.325$). Adjuvant therapies indication has increased during the COVID-19 pandemic ($p < 0.001$). In particular, the use of adjuvant radiotherapy (26.8% vs. 30.7%; $p = 0.001$) and chemotherapy (25.1% vs. 30.1%; $p < 0.001$) alone or in combination increased from period 1 to 2.

DISCUSSION

The present study investigated the characteristics and patterns of care of patients diagnosed with EC, before and during the COVID-19 pandemic. This large retrospective analysis reported several noteworthy findings. First, we observed that during the COVID-19 pandemic patients were more likely to be treated for advanced-stage disease (FIGO stage $>I$, with a high rate of patients with FIGO stage III–IV disease). Second, there was a higher proportion of patients treated with adjuvant therapy among those treated with surgery during the COVID-19 pandemic, as compared with those undergoing surgery before the pandemic. Third, the number of EC patients treated per year has decreased during the COVID-19 pandemic.

Generally, EC is an indolent entity. The short-term time interval, starting from the onset of the pandemic, is not fully explaining the high prevalence of advanced disease observed during COVID-19 pandemic than before. Possibly, we can suppose that is not the high incidence of advanced disease, but the low prevalence of early-stage disease during the pandemic that is driving these findings. Hence, the “real” high prevalence of advanced disease could be expected in the next years. COVID-19 has posed a significant challenge to worldwide health care systems. One of the main indirect consequences of COVID-19 pandemic lies in the limited access to health care services. Reluctance to report symptoms, potentially owing to fear of COVID-19, might become one of the main drivers of lower detection rates of early-stage EC. As aforementioned, the COVID-19 outbreak has correlated with the disruption of screening activities, regular follow up visits, and diagnostic assessments [6-10]. Many patients are delaying or missing their visits, even in presence of symptoms. However, to date the impact of COVID-19 on patients with newly diagnosed cancer is unclear. In the present paper, we decided to focus on the impact of COVID-19 in

EC patients. EC is often diagnosed in the early phase of the disease (FIGO stage I) and it is often characterized by a good prognosis. In our study, we observed that during the COVID-19 pandemic patients were more likely to be diagnosed with more advanced disease, as compared with the period before the pandemic. Similarly, the need for adjuvant treatments was higher during COVID-19 pandemic than before, while the prevalence of patients receiving non-surgical treatments remained steadily stable over time. Delayed diagnosis might potentially explain our results. Unexpectedly, the crude number of EC cases receiving medical and/or surgical treatment has decreased during the pandemic. Of note, we are missing several patients with early-stage disease. Few features might explain these findings: i) the number of patients treated in the two periods reflects a physiological variation of EC incidence over time ii) patients with more favorable disease are treated in more peripheral centers (e.g., low volume hospitals), thus meaning there might have been an allocation bias; iii) some patients with clinical stage I disease might be treated with intra-uterine devices (IUD) by general practitioners and not referred to the hospitals; and iv) patients are not diagnosed with EC since they are missing their visits. The reason is likely related with a multifactorial process. Further evidence will be necessary to assess the characteristics and patterns of presentation of EC patients during the next years. We are expecting that our results would be more evident in the next future, however with a potential opposite trend due to the implementation of COVID-19 vaccination among patients with cancer.

Interestingly, an Israeli Gynecologic Oncology Group retrospective study evaluated the role of EC diagnosis in asymptomatic patients [19]. The Authors compared data of 1,374 patients presenting with postmenopausal bleeding with 233 asymptomatic patients (diagnosed with EC after instrumental finding of thickened endometrium or polyps). Although the authors observed that EC diagnosed in asymptomatic postmenopausal patients is not associated with a survival advantage, the prevalence of patients diagnosed with more advanced disease stages and adjuvant therapy administration rate is lower in asymptomatic patients [19]. These findings seem to corroborate and partially explain our results.

Six points of the present investigation deserve to be addressed: i) Due to the absence of follow-up, we are not able to evaluate the impact of the COVID-19 pandemic on the oncologic outcomes of EC patients involved in this study. ii) We arbitrarily decided to omit data of March 2020 from the analysis, due to the limited impact of COVID-19 in that period (i.e., only few regions in the North of Italy were affected by the pandemic by March 2020) [5,12]. iii) We collected a huge amount of data (more than 5,000 patients) from the whole Italian territory, with a potential missing of EC cases diagnosed and treated in low volume centers. iv) Data about the prevalence of COVID-19 infections in EC patients (before or after treatments) is lacking. However, the main outcome measure of this research was not to assess the impact of COVID-19 disease on patients, but to assess how the COVID-19 outbreak impacted on patients' access to care. v) Data about the time between symptom presentation and date of a check-up at hospital, and the time between first histological diagnosis and date of surgery are important variables impacting outcomes. Further studies have to assess how waiting time impacted on survival outcomes, according to various histological features. vi) In Italy, the vaccination campaign against COVID-19 was managed by the Ministry of Health and (for patients) started on March 1, 2021. Hence, our data are reflecting the pre-vaccination era. We can speculate that the implementation of vaccination might improve the patients' access to care.

The inherent biases related to the retrospective nature of the study design are the main weaknesses of the present paper. Additionally, selection biases might impact the

interpretation of our results. Possibly, advanced and more challenging cases are more likely to be referred to high-volume centers, while peripheral centers are more likely to treat more simple cases. However, if this is true this evidence should have been detected even in the pre-pandemic period. The main strengths of this paper consist in: i) the large series of patients enrolled in this study and ii) the strong collaboration among a high number gynecological centers in Italy, suggesting the proactive and cooperative approach to the actual COVID-19 pandemic. Interestingly, about 8,000 new EC cases are diagnosed every year in Italy [20]. Hence, our study collected data about one third of EC patients having diagnosis and treatment in Italy in the last 2 years.

In conclusion, our study shows that the characteristics and patterns of care are changing during the COVID-19 pandemic. Compared with the pre-pandemic period, patients are more likely diagnosed with advanced stage disease (FIGO stage >I) during COVID-19 pandemic, with a consequently higher indication for adjuvant therapies. Due to the absence of mature post-treatment data, the impact of COVID-19 on survival outcomes of EC cannot be assessed yet. However, we have to highlight that possible physiological variations in pattern of EC presentation and allocation biases are influencing these results. Additionally, we have to point out that the modifications in EC presentation pattern presentation are minimal and not clinically meaningful. Owing to the indolent nature of EC we are not expected that delaying primary treatment of few months correlated with a high proportion of advanced stage of disease. Possibly, the number of patients with advanced stage is less diluted due to the decrease of number of patients with early-stage disease. Further analysis of our collaborative dataset will clarify these features in the next future. Similarly, further prospective evidence is necessary to corroborate our preliminary results. Attempts are warranted to improve risk-based strategies to recover, preserve, and implement healthcare services during the COVID-19 pandemic.

SUPPLEMENTARY MATERIAL

Table S1

Centers participating in this study

[Click here to view](#)

REFERENCES

1. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer Statistics, 2021. *CA Cancer J Clin* 2021;71:7-33.
[PUBMED](#) | [CROSSREF](#)
2. Siegel R, Ward E, Brawley O, Jemal A. Cancer statistics, 2011: the impact of eliminating socioeconomic and racial disparities on premature cancer deaths. *CA Cancer J Clin* 2011;61:212-36.
[PUBMED](#) | [CROSSREF](#)
3. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2021;71:209-49.
[PUBMED](#) | [CROSSREF](#)
4. Lu KH, Broaddus RR. Endometrial cancer. *N Engl J Med* 2020;383:2053-64.
[PUBMED](#) | [CROSSREF](#)
5. Bogani G, Apolone G, Ditto A, Scambia G, Panici PB, Angioli R, et al. Impact of COVID-19 in gynecologic oncology: a Nationwide Italian Survey of the SIGO and MITO groups. *J Gynecol Oncol* 2020;31:e92.
[PUBMED](#) | [CROSSREF](#)

6. Castanon A, Rebolj M, Burger EA, de Kok IM, Smith MA, Hanley SJ, et al. Cervical screening during the COVID-19 pandemic: optimising recovery strategies. *Lancet Public Health* 2021;6:e522-7.
[PUBMED](#) | [CROSSREF](#)
7. Piedimonte S, Li S, Laframboise S, Ferguson SE, Bernardini MQ, Bouchard-Fortier G, et al. Gynecologic oncology treatment modifications or delays in response to the COVID-19 pandemic in a publicly funded versus privately funded North American tertiary cancer center. *Gynecol Oncol* 2021;162:12-7.
[PUBMED](#) | [CROSSREF](#)
8. COVIDSurg Collaborative; GlobalSurg Collaborative. Timing of surgery following SARS-CoV-2 infection: an international prospective cohort study. *Anaesthesia* 2021;76:748-58.
[PUBMED](#) | [CROSSREF](#)
9. Glasbey JC, Nepogodiev D, Simoes JFF, Omar O, Li E, Venn ML, et al. Elective cancer surgery in COVID-19-free surgical pathways during the SARS-CoV-2 pandemic: an international, multicenter, comparative cohort study. *J Clin Oncol* 2021;39:66-78.
[PUBMED](#) | [CROSSREF](#)
10. Ciavattini A, Delli Carpini G, Giannella L, De Vincenzo R, Frega A, Cattani P, et al. Expert consensus from the Italian Society for Colposcopy and Cervico-Vaginal Pathology (SICPCV) for colposcopy and outpatient surgery of the lower genital tract during the COVID-19 pandemic. *Int J Gynaecol Obstet* 2020;149:269-72.
[PUBMED](#) | [CROSSREF](#)
11. Ramirez PT, Chiva L, Eriksson AG, Frumovitz M, Fagotti A, Gonzalez Martin A, et al. COVID-19 global pandemic: options for management of gynecologic cancers. *Int J Gynecol Cancer* 2020;30:561-3.
[PUBMED](#) | [CROSSREF](#)
12. Indini A, Cattaneo M, Ghidini M, Rijavec E, Bareggi C, Galassi B, et al. Triage process for the assessment of coronavirus disease 2019-positive patients with cancer: the ONCOVID prospective study. *Cancer* 2021;127:1091-101. Epub 2020 Dec 3.
[PUBMED](#) | [CROSSREF](#)
13. Concin N, Matias-Guiu X, Vergote I, Cibula D, Mirza MR, Marnitz S, et al. ESGO/ESTRO/ESP guidelines for the management of patients with endometrial carcinoma. *Int J Gynecol Cancer* 2021;31:12-39.
[PUBMED](#) | [CROSSREF](#)
14. Bogani G, Ray-Coquard I, Concin N, Ngoi NY, Morice P, Enomoto T, et al. Uterine serous carcinoma. *Gynecol Oncol* 2021;162:226-34.
[PUBMED](#) | [CROSSREF](#)
15. Malzoni M, Tinelli R, Cosentino F, Perone C, Rasile M, Iuzzolino D, et al. Total laparoscopic hysterectomy versus abdominal hysterectomy with lymphadenectomy for early-stage endometrial cancer: a prospective randomized study. *Gynecol Oncol* 2009;112:126-33.
[PUBMED](#) | [CROSSREF](#)
16. Casarin J, Bogani G, Piovano E, Falcone F, Ferrari F, Odicino F, et al. Survival implication of lymphadenectomy in patients surgically treated for apparent early-stage uterine serous carcinoma. *J Gynecol Oncol* 2020;31:e64.
[PUBMED](#) | [CROSSREF](#)
17. Bogani G, Papadia A, Buda A, Casarin J, Di Donato V, Gasparri ML, et al. Sentinel node mapping vs. sentinel node mapping plus back-up lymphadenectomy in high-risk endometrial cancer patients: results from a multi-institutional study. *Gynecol Oncol* 2021;161:122-9.
[PUBMED](#) | [CROSSREF](#)
18. Perrone E, Capasso I, Pasciuto T, Gioè A, Gueli Alletti S, Restaino S, et al. Laparoscopic vs. robotic-assisted laparoscopy in endometrial cancer staging: large retrospective single-institution study. *J Gynecol Oncol* 2021;32:e45.
[PUBMED](#) | [CROSSREF](#)
19. Gemer O, Segev Y, Helpman L, Hag-Yahia N, Eitan R, Raban O, et al. Is there a survival advantage in diagnosing endometrial cancer in asymptomatic postmenopausal patients? An Israeli Gynecology Oncology Group study. *Am J Obstet Gynecol* 2018;219:181.e1-6.
[PUBMED](#) | [CROSSREF](#)
20. Associazione Italiana di Oncologia Medica (AIOM). Neoplasie dell'utero: endometrio e cervice. Milan: AIOM; 2019 [cited 2021 Aug 31]. Available from: https://www.aiom.it/wp-content/uploads/2019/10/2019_LG_AIOM_Utero.pdf.