



LJMU Research Online

Fernandes, L, Mason, AC, Horta, D, Schiavon, RP, Hayes, C, Hasselquist, S, Feuillet, D, Beaton, RL, Jönsson, H, Kisku, S, Lacerna, I, Lian, J, Minniti, D and Villanova, S

A comparative analysis of the chemical compositions of Gaia-Enceladus/Sausage and Milky Way satellites using APOGEE

<http://researchonline.ljmu.ac.uk/id/eprint/18534/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Fernandes, L, Mason, AC, Horta, D, Schiavon, RP, Hayes, C, Hasselquist, S, Feuillet, D, Beaton, RL, Jönsson, H, Kisku, S, Lacerna, I, Lian, J, Minniti, D and Villanova, S A comparative analysis of the chemical compositions of Gaia-Enceladus/Sausage and Milky Way satellites using APOGEE. Monthly

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

A Comparative Analysis of the Chemical Compositions of Gaia-Enceladus/Sausage and Milky Way Satellites using APOGEE

Laura Fernandes¹, * Andrew C. Mason¹, Danny Horta¹, Ricardo P. Schiavon¹, Christian Hayes², Sten Hasselquist³, Diane Feuillet⁴, Rachael L. Beaton^{5,6}, Henrik Jönsson⁷, Shobhit Kisku¹, Ivan Lacerna^{8,9}, Jianhui Lian¹⁰, Dante Minniti^{11,12}, Sandro Villanova¹³

¹ Astrophysics Research Institute, Liverpool John Moores University, 146 Brownlow Hill, Liverpool L3 5RF, UK

² NRC Herzberg Astronomy and Astrophysics, 5071 West Saanich Road, Victoria, B.C., Canada, V9E 2E7

³ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁴ Lund Observatory, Department of Astronomy and Theoretical Physics, Box 43, SE-221 00 Lund, Sweden

⁵ Department of Astrophysical Sciences, 4 Ivy Lane, Princeton University, Princeton, NJ 08544

⁶ The Observatories of the Carnegie Institution for Science, 813 Santa Barbara St., Pasadena, CA 91101

⁷ Materials Science and Applied Mathematics, Malmö University, SE-205 06 Malmö, Sweden

⁸ Instituto de Astronomía y Ciencias Planetarias, Universidad de Atacama, Copayapu 485, Copiapó, Chile

⁹ Millennium Institute of Astrophysics, Nuncio Monsenor Sotero Sanz 100, Of. 104, Providencia, Santiago, Chile

¹⁰ Department of Physics and Astronomy, University of Utah, 115 S. 1400 E., Salt Lake City, UT 84112, USA

¹¹ Departamento de Ciencias Físicas, Facultad de Ciencias Exactas, Universidad Andres Bello, Fernández Concha 700, Las Condes, Santiago, Chile

¹² Vatican Observatory, Vatican City State, V-00120, Italy

¹³ Departamento de Astronomía, Universidad de Concepción, Casilla 160-C, Concepción, Chile

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

We use data from the 17th data release of the Apache Point Observatory Galactic Evolution Experiment (APOGEE 2) to contrast the chemical composition of the recently discovered Gaia Enceladus/Sausage system (GE/S) to those of ten Milky Way (MW) dwarf satellite galaxies: LMC, SMC, Boötes I, Carina, Draco, Fornax, Sagittarius, Sculptor, Sextans and Ursa Minor. Our main focus is on the distributions of the stellar populations of those systems in the [Mg/Fe]-[Fe/H] and [Mg/Mn]-[Al/Fe] planes, which are commonly employed in the literature for chemical diagnosis and where dwarf galaxies can be distinguished from *in situ* populations. We show that, unlike MW satellites, a GE/S sample defined purely on the basis of orbital parameters falls almost entirely within the locus of “accreted” stellar populations in chemical space, which is likely caused by an early quenching of star formation in GE/S. Due to a more protracted history of star formation, stars in the metal-rich end of the MW satellite populations are characterized by lower [Mg/Mn] than those of their GE/S counterparts. The chemical compositions of GE/S stars are consistent with a higher early star formation rate than MW satellites of comparable and even higher mass, suggesting that star formation in the early universe was strongly influenced by other parameters in addition to mass. We find that the direction of the metallicity gradient in the [Mg/Mn]-[Al/Fe] plane of dwarf galaxies is an indicator of the early star formation rate of the system.

Key words: Galaxies: dwarf spheroidal – stars; abundances

1 INTRODUCTION

Our current understanding of galaxy formation is immersed in the framework of the Lambda Cold Dark Matter (ΛCDM) model. In this model, galaxies are an ensemble of stars, gas, and dust bound by massive dark matter haloes, built largely by the merging of smaller systems (e.g., White & Rees 1978; Blumenthal et al. 1984; Frenk & White 2012). In this context, the Milky Way (MW) is of great importance, as the galaxy for which we can obtain the most detailed information. Naturally, one should expect that our knowledge of the MW should be used to constrain galaxy formation theory. While

most of the stellar mass in the MW is contained in its bar and both thin and thick disks (Morris & Serabyn 1996; Barbuy et al. 2018; Nataf 2017; Rich 2013; Beraldo e Silva et al. 2021), these structures are enveloped by a large stellar halo which contains about $1.3 \times 10^9 M_{\odot}$ in the form of stars (Mackereth & Bovy 2020a).

Galaxy mergers were very common in the early universe. Local evidence of the merging activity of the MW has been accumulating over the years, with the identification of the remnants of multiple accretion events, starting with the discovery of the Sagittarius dwarf spheroidal (Sgr dSph) by Ibata et al. (1994). In addition, various stellar streams have been identified as remnants of past accretion events (e.g., Belokurov et al. 2006). Due to the stellar halo’s long dynamical timescale, such relatively recent accretion events can be

* E-mail: Lfernandes2@msn.com (LF); R.P.Schiavon@ljmu.ac.uk (RPS)

identified in the form of spatial substructure. However, discerning the remnants of earlier accretion requires additional information, in the form of, e.g., kinematics, chemical compositions, and ages (e.g., Nissen & Schuster 2010, 2011; Schuster et al. 2012; Hayes et al. 2018)

With the advent of large spectroscopic surveys, we entered the golden age of Galactic archaeology. The information acquired by past, ongoing and future surveys such as RAVE, (Steinmetz et al. 2006), SEGUE (Yanny et al. 2009), LAMOST (Cui et al. 2012), GALAH (De Silva et al. 2015), *Gaia* mission (Gaia Collaboration et al. 2016), WEAVE (Dalton 2016), APOGEE (Majewski et al. 2017), H3 (Conroy et al. 2019), MOONS (Cirasuolo et al. 2011; Gonzalez et al. 2020), 4MOST (de Jong et al. 2019) is very quickly advancing our understanding of the history of the MW. Chemical compositions and orbital information is becoming available for millions of stars in the MW and its Local Group companions. In addition, with the launching of the *Gaia* satellite (Gaia Collaboration et al. 2016), detailed phase space information has become available, enabling the identification of further substructures in the stellar halo of the Galaxy, associated with various accretion events, including Gaia-Enceladus/Sausage, a massive dwarf galaxy accreted to the MW about ~ 10 Gyr ago (Belokurov et al. 2018; Haywood et al. 2018; Helmi et al. 2018; Mackereth et al. 2019), as well as Heracles (Horta et al. 2021a), and various other substructures (e.g., Koppelman et al. 2019a; Belokurov et al. 2018; Kruijssen et al. 2020; Naidu et al. 2020; Horta et al. 2022a).

Hawkins et al. (2015) and Das et al. (2020) have recently proposed that a combination of the abundances of Mn, Mg, Al, and Fe can be used to chemically discriminate accreted populations from their counterparts formed *in situ*. However, Horta et al. (2021a) used chemical evolution models to show that the region occupied by accreted populations in the $[\text{Mg}/\text{Mn}]$ vs $[\text{Al}/\text{Fe}]$ plane has a non-negligible presence of stars formed *in situ*.

It is difficult to disentangle metal-poor *in situ* populations from their accreted counterparts based on purely kinematic or orbital criteria. To minimise inter-sample contamination, one has to often resort to additional criteria based on chemical compositions, introducing sample biases that prevent a clean analysis of the chemical compositions of accreted and *in situ* populations. That difficulty can be overcome by resorting to data from external systems, on the assumption that the chemistry of their stellar populations is dominated by intrinsic evolutionary effects.

In this paper we contrast the distribution of stars from GE/S system in key chemical composition planes with those of dwarf satellites of the MW. The latter include the Sagittarius dSph, which is currently in the process of being engulfed by the Milky Way, as well as gas rich, irregular dwarf galaxies, such as the Large and Small Magellanic clouds (LMC & SMC); the gas-deficient, classical dwarf spheroidal galaxies (dSph), Carina, Draco, Fornax, Sculptor, Sextans, and Ursa-Minor; and the ultra-faint dwarf spheroidal galaxy, Boötes I. Our goal is to make inferences regarding the history of star formation and chemical enrichment of GE/S by contrasting it with MW satellites. We note that Hasselquist et al. (2021) have performed a similar comparative analysis of GE/S and massive satellites of the MW. In this paper we extend the analysis to lower mass satellites, with a focus on the distribution of dwarf galaxy stellar populations on the $[\text{Mg}/\text{Mn}]$ - $[\text{Al}/\text{Fe}]$ plane and how it can be used to constrain the history of star formation of those systems. In addition, we examine the behaviour of chemical evolution models calculated for a range of relevant input parameters on that same chemical plane.

This paper is organised as follows. Section 2 presents the data and sample adopted. In Section 3 we present an examination of the




Figure 1. Selecting GE/S candidates in IoM in the plane of APOGEE DR17 stars, the number density is represented by the grey-scale colour bar and does not include the following stellar populations: thin-thick disk selection ($L_z > 0$, eccentricity < 0.3), globular clusters [VAC by Schiavón et al. (2021, in prep)], the large and small Magellanic clouds, Sagittarius dSph and low-mass dSph galaxies. The GE/S stars in this selection (pink thistle circles) are within $-1.6 < E/10^5 < -1.3 \text{ km}^2 \text{ s}^{-2}$ and $-0.4 < L_z/10^3 < 0.3 \text{ kpc km s}^{-1}$. The selection criteria of disk stars (light steel blue circles) in this analysis have been placed on top of the plane for reference.

distribution of the stellar populations of the various systems in key chemical planes. In Section 4 we contrast the chemical composition characteristics of GE/S with those of MW satellites of various masses, using chemical evolution models to guide the discussion. Our conclusions are presented in Section 5.

2 DATA AND SAMPLE

2.1 APOGEE DR17

The second generation Apache Point Observatory Galactic Evolution Experiment (APOGEE-2, Majewski et al. 2017) is part of the Sloan Digital Sky Survey IV (Blanton et al. 2017). APOGEE-2 surveys the stellar populations of the MW with high resolution ($R \sim 22,500$), high S/N, spectroscopy in the H-band spectral region ($\lambda \sim 1.51\text{--}1.70 \mu\text{m}$), using two twin multi-fibre spectrographs (Wilson et al. 2019). Less hampered by interstellar dust than optical surveys its observations cover both Northern and Southern hemispheres, based primarily on the 2.5 m Sloan Foundation telescope (Gunn et al. 2006) at Apache Point Observatory and the 2.5 m Irénée du Pont telescope (Bowen & Vaughan 1973) at Las Campanas Observatory.

The data employed in this paper consist of chemical compositions, stellar parameters and integrals of motion obtained from the 17th data release (DR17, Abdurro'uf et al. 2021, Holtzman et al. in prep.) of SDSS-IV/APOGEE-2. Chemical compositions and stellar parameters were generated by the APOGEE Stellar Parameter and Chemical Abundance Pipeline (ASPCAP, García Pérez et al. 2016), whereas the integrals of motion result from application of the `galpy` package (Bovy 2015; Mackereth & Bovy 2018) to 6D phase space information resulting from combination of *Gaia* eDR3 proper motions (Gaia Collaboration et al. 2021), APOGEE-2 DR17 radial velocities (Nidever et al. 2015, Holtzman et al. in prep.), and `astroNN` machine learning-based distances (Leung & Bovy 2019). Calculations were performed adopting a McMillan (2017) potential for the Milky Way.

Table 1. Total sample selection.

Stellar Structures	No. of Stars in Sample
Disks	193,220
LMC	4,610
SMC	1,660
Sgr	291
Boötes I	3
Carina	35
Draco	31
Fornax	140
Sculptor	85
Sextans	18
Ursa Minor	25
GE/S	1,952

2.2 SELECTION CRITERIA

Our goal is to examine the chemical compositions of stars belonging to 10 dwarf satellites of the Milky Way, namely LMC, SMC, Boötes I, Carina, Draco, Fornax, Sculptor, Sagittarius, Sextans and Ursa Minor, contrasting their distributions in chemical diagnostic planes with those of the accreted system GE/S and the MW high- and low- α disks. The APOGEE-2 DR17 catalogue contains data for 733,901 stars, selected according to criteria extensively discussed by Zasowski et al. (2013), Zasowski et al. (2017), and Santana et al. (2021). Before selecting stars belonging to the above systems, we must apply a number of criteria to certify the quality of the data for our analysis. We first cleaned the sample from stars with unreliable parameters by removing all stars with STARFLAG or ASPCAPFLAG=BAD (see definitions in Holtzman et al. 2015). We then limit the data to red giant stars with $S/N > 50$, stellar effective temperatures (T_{eff}) between 3750 – 5500 K and surface gravity ($\log(g)$) < 3.0.

Next, we removed from the sample a total of 7,562 stars that are deemed to be members of globular clusters, as listed in the Value Added Catalogue by Schiavon et al. (2022, in prep.). Application of the above filters left us with a sample of 300,389 stars. The data for the objects of interest were extracted from this surviving catalogue according to the criteria described in the following sub-sections. The final sample sizes for each system are given in Table 1.

2.2.1 Magellanic Clouds

Our sample selection for the LMC and SMC members mimics that of Nidever et al. (2020) and is summarised in Table 2. We focus on the bright and faint red giant branch (RGB) stellar populations in the MCs (see their figures 3 and 5). In this way we expect to restrict our sample to stars in approximately the same evolutionary stage as those in the MW and other satellites.

2.2.2 Sgr dSph

Our sample for the Sgr dSph stars comes from the study by Hasselquist et al. (2017) and was selected by the methods described in Majewski et al. (2013). Further sampling of Sgr core and stream

Table 2. MCs Sample Selection. Table summaries the sky positions (α, δ), projected distance on the sky (d_{proj}), Gaia proper motions (PM), radial velocities (RV), and magnitudes (H) for the LMC and SMC.

LMC $\alpha \delta : (80.893860, -69.756126)$	SMC $\alpha \delta : (13.18667, -72.8286)$
$d_{\text{proj}} \leq 12$	$d_{\text{proj}} \leq 8$
$RV \geq 125$	$RV \geq 100$
$2.7 \leq \alpha\text{PM} \leq 1$	$2.0 \leq \alpha\text{PM} \leq 0$
$2 \leq \delta\text{PM} \leq -1.2$	$-0.5 \leq \delta\text{PM} \leq -2.0$
$J-K \geq 0.5$	$J-K \geq 0.5$
$12.35 < H < 14.6$	$12.9 < H < 14.7$

members are covered extensively in Hasselquist et al. (2019a) and Hayes et al. (2020)¹. Table 3 summaries the selection criteria.

2.2.3 Dwarf Spheroidal Galaxies

APOGEE-2 has targeted a number of dwarf spheroidal galaxies. The field placement and target selection criteria adopted are described by Zasowski et al. (2017) and Santana et al. (2021). To identify dwarf spheroidal members we first selected all stars observed within the fields of each dwarf galaxy and filtered out foreground contaminants on the basis of radial velocity and surface gravity. Stars considered members are giants ($\log g < 3.0$) whose heliocentric radial velocities differ from the central values for each galaxy by less than twice its velocity dispersion. By proceeding in this way we prioritise sample purity over completeness. Radial velocities and velocity dispersions of the sample dSph galaxies are taken from Table 1 from McConnachie & Venn (2020) and Table 4 from McConnachie (2012)² with the exception of Boötes I, for which values were taken from Martin et al. (2007).

2.2.4 Gaia-Enceladus/Sausage

Stars from the accreted system GE/S are distributed throughout the MW and can be discriminated through a range of chemical, kinematical and/or orbital selection criteria. In order to obtain a GE/S sample devoid of chemical composition biases, we base our selection purely on integral of motion (IoM) measurements.

After removal from the sample of all stars associated to the MW satellites, the stellar populations of the accreted system GE/S are selected on the basis of their position in the energy vs. angular momentum ($E-L_z$) plane.

A star is considered a member of the Gaia-Enceladus system if its energy and angular momentum fall within the following intervals:

- $-1.6 < E/10^5 < -1.3 \text{ km}^2 \text{ s}^{-2}$
- $-0.4 < L_z/10^3 < 0.3 \text{ kpc km s}^{-1}$

This region of $E-L_z$ is shown in Figure 1. The above selection criteria mimic those adopted in previous work (e.g., Koppelman et al. 2019b; Massari et al. 2019; Feuillet et al. 2021; Horta et al. 2021a). They are designed to take advantage of the overdensity

¹ <http://vizier.u-strasbg.fr/viz-bin/VizieR?-source=J/ApJ/889/63>

² http://www.astro.uvic.ca/~alan/Nearby_Dwarf_Database.html

Table 3. Sgr Sample Selection. Table summaries the sky positions (α, δ), radial velocities (RV), surface gravity ($\log g$), effective temperature (T_{eff}), and signal-to-noise (S/N) selection criteria for the Sagittarius dSph galaxy core and tail.

Sagittarius dSph Galaxy $\alpha \delta : (284, -30)$
$90 \text{ km s}^{-1} < \text{RV} < 220 \text{ km s}^{-1}$
$d > 5 \text{ kpc}$
$J - K_0 > 0.8$
$S/N > 70$
$3550 \text{ K} < T_{\text{eff}} < 4200 \text{ K}$
$\log g < 4$

in the $E-L_Z$ plane around $L_Z \sim 0$ and at relatively high energy, that is easily identifiable in Figure 1. We deliberately adopted a relatively high lower energy limit for our GE/S selection with an eye towards minimising disk contamination. Yet because we imposed no chemical composition cuts, we expect a small contamination of our GE/S sample by disk stars. See discussion in Section 3.1. This contamination by disk stars is further enhanced by the existence of “Splash” stars, which are early disk stars whose orbits were perturbed by the collision with GE/S (Belokurov et al. 2020). For a discussion of the impact of selection criteria on the chemical properties of GE/S, see Horta et al. (2022b).

2.2.5 Thin & Thick Disk

The stellar populations of the the MW disk are selected using orbital parameters in IoM, focusing on stars with circular, prograde orbits. We thus retain disk stars with the following criteria:

- $L_z > 0$
- eccentricity < 0.3
- $S/N > 70$

The eccentricity cut employed is used to select stellar populations with disc orbits. The adopted eccentricity threshold is arbitrary. Placing the cut at, e.g., eccentricity < 0.2 or < 0.4 would cause no impact on our analysis. The detailed choice is not critical because the disc population is used solely as a reference for comparison with the dwarf satellite data.

3 CHEMICAL PROPERTIES

In the next two sections we analyse the chemical properties of the dwarf galaxies and GE/S members alongside the MW disks in the chemical planes of Mg, Mn, Al and [Fe/H].

3.1 Magnesium

Magnesium is an α -element synthesised during carbon burning in massive stars, and injected into the interstellar medium during supernovae type II (SNe II) explosions (Kobayashi et al. 2006; Woosley & Weaver 1995). The distribution of the stellar populations on the α -Fe plane provides important clues on the star formation history and IMF of the system (e.g., Matteucci & Greggio 1986; Wheeler et al. 1989; McWilliam 1997).

In Figure 2, the distributions of the stellar populations in the dwarf galaxies and GE/S stars are shown in the chemical plane of [Fe/H]

Table 4. Properties of dSph Galaxies in the sample, including ID, stellar mass, velocity dispersion, radial velocity, and original references. All data from compilation by McConnachie (2012).

Galaxy	M_\star ($10^6 M_\odot$)	R.V. (km s^{-1})	σ (km s^{-1})	References
Boötes I	0.029	99.9 ± 2.4	$6.5^{+2.1}_{-1.3}$	(1)
Carina	0.38	222.9 ± 0.1	6.6 ± 1.2	(2)(6)(7)
Draco	0.29	-291 ± 0.1	9.1 ± 1.2	(2)(3)(4)
Fornax	20	55.3 ± 0.1	11.7 ± 0.9	(2)(6)(7)(9)
Sculptor	2.3	111.4 ± 0.12	9.2 ± 1.4	(2)(6)(7)(8)
Sextans	0.44	224.2 ± 0.1	7.9 ± 1.3	(2)(6)(7)
Ursa Minor	0.29	-246.9 ± 0.1	9.5 ± 1.2	(2)(4)(5)

References: (1) Martin et al. (2007), (2) Grcevich & Putman (2009), (3) Walker et al. (2007), (4) Wilkinson et al. (2004), (5) Walker et al. (2009b), (6) Walker et al. (2009a), (7) Walker et al. (2008), (8) Carignan et al. (1998), (9) Bouchard et al. (2006)

- [Mg/Fe] across eleven panels, each displayed alongside the stellar populations of the low- and high- α disks of the MW, whose chemical compositions range roughly between $-1.2 < [\text{Fe}/\text{H}] < +0.65$ and $-0.2 < [\text{Mg}/\text{Fe}] < +0.4$. These numbers are in good agreement with independent studies based on abundance analysis of high resolution optical spectra, such as those by Aguado et al. (2021), Matsuno et al. (2021), and Carrillo et al. (2022).

We start by pointing out that, for the reasons discussed in Section 2.2, our GE/S sample contains a small, yet non-negligible, contamination by *in situ* stars, which can be easily spotted as they fall on the loci defined by the low- and high- α disks. It is fair to assume that the metal-rich stars in our GE/S sample that overlap with the high- and low- α disc sequences are contaminants because, on one hand, their chemical properties associate them strongly with the disc, and moreover it has been argued by Mackereth et al. (2018) and Mason et al. (2022, in prep.) that low mass galaxies do not host a bimodal α distribution. It is also likely that our sample may be contaminated on the metal-poor end, as *in situ* and accreted populations overlap on the [Mg/Fe]-[Fe/H] plane at low metallicity, (e.g., Horta et al. 2020). However, the disc sequence becomes very thinly populated at $[\text{Fe}/\text{H}] \lesssim -1.2$, and moreover our selection criteria prioritizing high energy stars also helps minimising disk contaminations. Thus, unless otherwise stated, our discussion henceforth ignores metal-rich contaminants overlapping with the high- and low- α disk sequences on both chemical planes under study.

The latter are characterised by predominantly low [Fe/H] and lower [Mg/Fe] than high- α disk stars at $[\text{Fe}/\text{H}] \gtrsim -1$, and overall decreasing with increasing metallicity. On the metal-poor end ($[\text{Fe}/\text{H}] \lesssim -1.8$), GE/S stars reach [Mg/Fe] values comparable or even slightly higher than those of the high- α disk, whereas on the metal-rich end ($[\text{Fe}/\text{H}] \sim -0.6$) GE/S stars have slightly lower [Mg/Fe] than that of low- α disk stars of same metallicity. The slope of the [Mg/Fe]-[Fe/H] relation undergoes a slight change, forming the so-called “ α -knee” at $[\text{Fe}/\text{H}] \sim -1.2$ (see also Mackereth et al. 2019; Horta et al. 2021a), which indicates the increased contribution of SN Ia to the chemical enrichment of the interstellar medium.

By and large, the locus occupied by GE/S stars in the Mg-Fe plane is somewhat similar to that where stars from the massive MW satellites LMC, SMC, and Sgr dSph are located (see also Hasselquist et al. 2021). The distributions however differ in important details. All three of the massive satellites show, at a given [Fe/H], a positive




Figure 2. Stellar populations of the dwarf galaxies and GE/S in the plane of MW disk (marginal density 2D hexagonal binning - the grey-scale of each hexbin denotes the number of points) in the chemical plane of $[\text{Mg}/\text{Fe}]$ versus $[\text{Fe}/\text{H}]$. From top to bottom; the selected GE/S stellar population (thistle) as shown in figure 1, LMC (light blue), SMC (coral), Sgr dSph (blue), Sculptor (green), Fornax (teal), Carina (crimson), Draco (cyan), Sextans (fuchsia), Ursa Minor (brown) and Boötes I (purple).

change in the slope of the $[\text{Mg}/\text{Fe}]-[\text{Fe}/\text{H}]$ relation, whereby $[\text{Mg}/\text{Fe}]$ starts increasing with increasing metallicity. This is likely associated with the occurrence of a burst of star formation in those systems, which causes an increase in the contribution of SNI_{II} to the chemical enrichment of the interstellar medium, resulting in a jump in $[\text{Mg}/\text{Fe}]$ for increasing $[\text{Fe}/\text{H}]$ (for a discussion, see Mason et al. 2022, in prep.). Because star formation in GE/S was quenched at the time of accretion, no similar change in slope can be seen in its stellar populations.

In contrast, for most low-mass dwarf spheroidal galaxies (except for Fornax), $[\text{Mg}/\text{Fe}]$ decreases monotonically with increasing $[\text{Fe}/\text{H}]$. The slope of the relation is steeper than that of GE/S and the more massive satellites, and the mean $[\text{Mg}/\text{Fe}]$ is substantially lower in low-mass satellites. At $[\text{Fe}/\text{H}]\sim-1.2$ the stellar populations of low mass satellites are lower in $[\text{Mg}/\text{Fe}]$ by $\sim 0.2-0.4$ dex than those from their massive counterparts.

The absence of a clear “knee” in the α -Fe plane of low mass satellites is likely due to it being located in those systems at metallicities that are lower than the values spanned by our sample. Indeed, previous studies of stars in Draco, Sextans and Ursa Minor (Shetrone et al. 2003), Sculptor (Hill et al. 2019), and Sextans (Theiler

et al. 2020) suggest the presence of a “knee” at $[\text{Fe}/\text{H}]\sim-2$. Based on a compilation of chemical compositions from various works (Cohen & Huang 2009, 2010; Starkenburg et al. 2013), Hendricks et al. (2014) quote $[\text{Fe}/\text{H}]_{\text{knee}} \sim -1.9$ for Sculptor, Ursa Minor, and Fornax and $[\text{Fe}/\text{H}]_{\text{knee}} \lesssim -2.5$ for Draco and Carina (see also de Boer et al. 2014). The low metallicity limit of our sample prohibits any statement on the existence of a change of slope for the latter two galaxies. As for the others, while our data do not rule out the presence of a change of slope in the Mg-Fe relation for Sculptor, Ursa Minor, and Fornax at $[\text{Fe}/\text{H}]\sim-2$, they cannot confirm it either, due to increased uncertainties and relatively small sample sizes in the low metallicity end.

The above result, taken together with the overall lower $[\text{Mg}/\text{Fe}]$ of lower mass satellites, suggests that the contribution by SN Ia to the chemical enrichment of the interstellar medium of those satellites was more dominant than in their massive counterparts, possibly indicating a lower star formation rate (SFR) throughout their histories (e.g., Mason et al. 2022, in prep.). The one exception is Fornax, for which a sharp positive change of slope can be seen at $[\text{Fe}/\text{H}]\sim-1.2$, similar to the case of more massive satellites, suggesting also in the case of Fornax the occurrence of a burst of star formation in the recent past.

Most importantly for the goals of this study, when we contrast the position of GE/S in the Mg-Fe plane with those of dwarf satellites of the MW, we conclude that it has undergone an intense early history of star formation leading up to the build up of a relatively metal-rich and α -enhanced stellar population, akin to those of the massive satellites of the MW. This is not surprising, considering current estimates for the original mass of the GE/S progenitor ($\sim 10^8 - 10^9 M_{\odot}$, e.g., [Helmi et al. 2018](#); [Belokurov et al. 2018](#); [Mackereth et al. 2019](#); [Deason et al. 2019](#); [Mackereth & Bovy 2020b](#); [Feuillet et al. 2020](#)).

3.2 Aluminium & Manganese [Al, Mn]

It is well known that stellar population diagnosis is substantially improved by the consideration of the abundances of elements associated with distinct nucleosynthetic pathways. In that vein, it has been proposed by [Hawkins et al. \(2015\)](#) and [Das et al. \(2020\)](#) that the combination of the abundances of Fe, Mg, Mn, and Al can aid in the discrimination of accreted stellar populations in the Galactic halo. Manganese is an Fe-peak element generated in Type Ia supernovae (SNIa) ([Iwamoto et al. 1999](#); [Hillebrandt & Niemeyer 2000](#); [Weinberg et al. 2019](#)). According to [Hawkins et al. \(2015\)](#), manganese is a more pure indicator of enrichment by SNe Ia than iron, which makes that element particularly useful for chemical diagnosis. Unlike the case of [Mg/Fe], [Mn/Fe] correlates positively with metallicity ([Kobayashi et al. 2020](#)). This sharp distinction between the dependence of Mg and Mn on metallicity makes the ratio between these two elements a powerful discriminator between stellar populations with different chemical evolution histories.

Aluminium, in turn, is referred to as an odd-Z element. Although similarly to magnesium, Al is produced predominantly by SNe II ([Buchmann et al. 1984](#); [Prantzos & Diehl 1996](#)), it can be contributed relevantly by a number of other nucleosynthetic sources. Aluminium primarily forms during H burning phases in the CNO, NeNa and MgAl cycles ([Samland 1998](#); [Guelin et al. 1995](#)). A small amount of Al is also created in white dwarf binary collisions ([Nofar et al. 1991](#); [Weiss & Truran 1990](#)). Other sources of Al are observed from the winds of Wolf-Rayet ([Limongi & Chieffi 2006](#)) and AGB stars ([Nomoto et al. 1984](#)). Furthermore, viable sources of ^{26}Al in the ISM are thought to originate from the accretion of hydrogen-rich gas in white dwarf binaries following novae explosions. ([Gehrz et al. 1998](#); [Kamiński et al. 2018](#); [Clayton 1984](#)).

[Das et al. \(2020\)](#) used the SDSS-III/APOGEE DR14 sample ([Abolfathi et al. 2018](#); [Holtzman et al. 2018](#)) to show that stars belonging to the GE/S system occupy a distinct locus in the [Al/Fe] vs. [Mg/Mn] plane, characterised by low [Al/Fe] and high [Mg/Mn]. More recently, [Horta et al. \(2021a\)](#) used chemical evolution models to show that this particular locus of chemical space is actually the home of *chemically unevolved* stellar populations. In other words, any early stellar populations inhabit that region of chemical space, regardless of where they are formed. As chemical evolution proceeds, the elemental abundances of *in situ* populations move away from that locus of chemical space, whereas the star formation of early accreted systems is quenched, so that chemical compositions are frozen in their early, pre-accretion state.

4 STELLAR POPULATIONS: CHEMICALLY EVOLVED OR UN-EVOLVED

In this section we examine the distribution of GE/S stars and MW dwarf satellites in the [Al/Fe] vs. [Mg/Mn] plane, in order to check whether the above scenario is supported by an entirely empirical

examination of the data. For that purpose we compare the distribution of GE/S stars in that plane with those of the low- and high-mass satellites of the MW.

4.1 The detailed chemistry of a kinematically selected GE/S

Our first goal is to check whether the locus of a GE/S sample selected purely on the basis of kinematics would be concentrated in the “chemically unevolved” region of the [Mg/Mn]-[Al/Fe] space. We recall that, for this analysis to be meaningful, it is critical that the selection of the stars from all systems is entirely free of any chemical composition criterion (for details, see Section 2.2.4).

Inspection of Figure 3 shows that the majority of GE/S stars selected purely on the basis of orbital parameters fall within the “chemically unevolved” locus of the [Mg/Mn]-[Al/Fe] space where 82% of GE/S stars are located. It is important, however, to keep in mind that this line is arbitrary so that it is possible that some stars in the “evolved” region actually belong to GE/S. There is a small amount of contamination by high- α disk stars, spreading towards the high [Al/Fe] and high [Mg/Mn] (upper right) locus of the chemical space. In contrast to the case of [Mg/Fe] abundance ratios, all the dwarf galaxies and GE/S are characterised by similarly low [Al/Fe]. In fact, at $[\text{Fe}/\text{H}] \lesssim -1.0$, the bulk [Al/Fe] abundances in the dwarf galaxies are lower than those in the low- and high- α disks.

Our GE/S sample stars fall squarely within the “chemically unevolved” region of the plot, which we interpret as indicating an early quenching of star formation taking place during the merger of that galaxy with the MW (see discussion in [Feuillet et al. 2021](#)). This result confirms our interpretation of the distribution of the stellar populations in the [Mg/Mn]-[Al/Fe] plane, as well as the chemical evolution calculations presented in [Horta et al. \(2021b\)](#).

4.2 GE/S vs Dwarf Satellites

We next compare the distribution of GE/S stars in the [Al/Fe] vs. [Mg/Mn] plane with those of the massive MW satellites (MCs and Sgr dSph). In contrast to the case of GE/S, a substantial fraction of the stars in the MCs and the Sgr dSph lie outside that locus. This is not surprising, as these massive satellites continued forming stars long after star formation in GE/S had ceased. The chemically evolved, more metal-rich, stars in those massive satellites spread towards the low [Mg/Mn] region of the plane, at approximately constant [Al/Fe]. It is also worth noticing that those among the stars belonging to the massive satellite that do inhabit the “chemically unevolved” locus of the plot are located towards a substantially lower mean [Mg/Mn] than the stars from GE/S. While the GE/S stars within the “chemically unevolved” region have mean [Mg/Mn] ~ 0.50 , those in the MCs and Sgr dSph have mean values between ~ 0.35 and 0.25 dex lower. We speculate whether this difference is due to a selection effect caused by the fact that the massive satellites samples may be biased towards the high metallicity end of the MDF.

Differences between the loci occupied by GE/S and lower mass satellites are even more pronounced, particularly for Draco, Carina, Sextans and Ursa Minor, whose sample stars are predominantly located outside the “chemically unevolved” region. While this difference may partly reflect selection biases, it is qualitatively consistent with these lower mass satellites having undergone an evolutionary history characterised by a low, yet more prolonged, SFR than those of GE/S and the massive MW satellites, which would naturally lead to a stronger contribution to enrichment by SN Ia and




Figure 3. Diagnostic plot $[Al/Fe] - [Mg/Mn]$. Stellar populations of the dwarf galaxies and GE/S in the plane of MW disk (marginal density 2D hexagonal binning - the grey-scale of each hexbin denotes the number of points). From top to bottom; the selected GE/S stellar population (thistle) as shown in figure 1, LMC (light blue), SMC (coral), Sgr dSph (blue), Sculptor (green), Fornax (teal), Carina (crimson), Draco (cyan), Sextans (fuchsia), Ursa Minor (brown) and the disk selection (steel blue) as shown in figure 1. Black lines separate *in-situ* high- α , *in-situ* low- α stars and the unevolved region.

a consequently lower mean $[Mg/Mn]$. This is further discussed in Section 4.5.

4.3 Comparison with Chemical Evolution Models

In Figure 4 we build on the diagnostic $[Al/Fe]-[Mg/Mn]$ plane with the addition of two chemical evolution models calculated using the flexCE code from Andrews et al. (2017). The orange line shows the evolution of a model made to match the properties of the stellar populations of the solar neighbourhood, representing an *in situ* population, and the yellow model line shows a chemical evolution model built to match the chemical properties of Gaia-Enceladus/Sausage, characterising the chemical evolution of a relatively massive satellite galaxy. The parameters adopted for the chemical evolution models are shown in Table 5. The *in situ* MW model is outlined in Horta et al. (2020). The model for GE/S was built to match the distribution of the data on the Si-Fe plane. For details on the model parameters, see Tables 3 and 4 of Hasselquist et al. (2021). The models evolve for approximately 13 Gyr: the filled circles on the models mark the evolutionary times at 0.3, 1

and 5 Gyr. The black cross marks the position at which the iron abundance reaches $[Fe/H]=-1$. The star formation efficiencies in the two models differ by an order of magnitude, at 1.5 Gyr^{-1} in the *in situ* case (Horta et al. 2021b) and 0.14 Gyr^{-1} for the best fitting GE/S model (Hasselquist et al. 2021). As a result, the solar neighbourhood model reaches $[Fe/H]=-1$ a mere 0.12 Gyr after the beginning of the evolution, whereas the GE/S model takes ~ 1.18 Gyr to reach the same metallicity. The models provide a qualitatively good description of the data for the *in situ*, accreted, and satellite stellar populations on the $[Mg/Mn]$ vs. $[Al/Fe]$ plane. These model calculations are an important tool for the interpretation of the data. In both cases, the early chemical enrichment drives the evolution towards the right due to the contribution by Type II/core collapse supernovae, because $[Mg/Mn]$ remains approximately constant while $[Al/Fe]$ grows due to the metallicity dependence of Al yields. By the same token, downward/left evolution on this plane reflects the increased contribution by SN Ia.

The data for the various accreted/satellite stellar populations in Figure 4 are colour-coded by metallicity. One can immediately notice a difference between GE/S and all the MW satellites, which




Figure 4. Diagnostic plot $[Al/Fe] - [Mg/Mn]$, colour-coded by metallicity. The MW Chemical Evolution Model (orange) and Dwarf Chemical Evolution Model (yellow) in the $[Mg/Mn]$ vs $[Al/Fe]$ abundance plane. Coloured circles mark the evolutionary times at 0.3, 1.0 and 5.0 Gyrs. Black cross is the position at which the models reach $[Fe/H] \sim -1$: $t = 0.1$ Gyrs for the MW chemical evolution model and $t = 1.2$ Gyrs for the dwarf chemical evolution model. From top to bottom; GE/S, LMC, SMC, Sgr dSph, Sculptor, Fornax, Carina, Draco, Sextans, Ursa Minor and the disk. Black lines separate *in-situ* high- α , *in-situ* low- α stars and the unevolved region.

Table 5. Summary of parameters used in the chemical evolution models.

Model Parameters	MW	GE/S	LMC-like	Sgr dSph-like	Sculptor-like
Initial Gas Mass	$2 \times 10^{10} M_{\odot}$	$3 \times 10^9 M_{\odot}$	$3 \times 10^9 M_{\odot}$	$3 \times 10^9 M_{\odot}$	$3 \times 10^9 M_{\odot}$
Inflow Mass Scale	$3.5 \times 10^{11} M_{\odot}$	$6 \times 10^{10} M_{\odot}$	$6 \times 10^{10} M_{\odot}$	$6 \times 10^{10} M_{\odot}$	$6 \times 10^{10} M_{\odot}$
Outflow Mass Loading Factor	2.5	6	5.4	20	40
Star Formation Efficiency	$1 \times 10^{-9} \text{yr}^{-1}$	$1.5 \times 10^{-10} \text{yr}^{-1}$	$1.25 \times 10^{-11} \text{yr}^{-1}$	$2.5 \times 10^{-11} \text{yr}^{-1}$	$1 \times 10^{-11} \text{yr}^{-1}$
Exponential Inflow Timescale	6 Gyr	2.5 Gyr	2.5 Gyr	2.5 Gyr	2.5 Gyr

have lower $[Al/Fe]$ on average than GE/S. Indeed, the model is a good match to GE/S, by construction, while failing to reproduce the main locus of satellites, which is particularly noteworthy in $[Al/Fe]$. Given the clear dependence of the value of $[Al/Fe]$ on star formation efficiency indicated by the models, this difference suggests that GE/S formed stars more vigorously early in its history than the

MW satellites. In Section 4.5 we discuss the behaviour of chemical evolution models on this plane in more detail.




Figure 5. Diagnostic plot $[Al/Fe]$ - $[Mg/Mn]$, colour-coded by metallicity. From top to bottom: GE/S stellar population (excluding the contaminating thick-disk/high- α stars), LMC, SMC, Sgr dSph, Sculptor, Fornax and the disk selection as shown in figure 1. Black lines separate *in-situ* high- α , *in-situ* low- α stars and the unevolved region. In each panel, the direction of the arrow represents the metallicity vector of the stellar population.

4.4 Metallicity gradients on the $[Mg/Mn]$ - $[Al/Fe]$ plane

We call attention to an important feature in the distribution of the data for different stellar populations in Figure 4. As the points are colour-coded by metallicity, one can see that the colour gradient on the $[Mg/Mn]$ vs $[Al/Fe]$ plane, varies widely from galaxy to galaxy, with more metal-rich stars being distributed towards the lower right in more massive systems and towards the lower left in low mass satellites. This suggests that GE/S and the MCs seem to have evolved more quickly in $[Al/Fe]$ than all the other surviving satellites of the MW (this is true even after correction for contamination by *in situ* stars, see Section 4.4).

We quantified this effect by measuring the metallicity gradients of our sample galaxies in the $[Mg/Mn]$ - $[Al/Fe]$ plane as follows. We first reduced the sample in two ways. For GE/S, we minimised contamination by *in situ* stars by selecting them in the Mg-Fe plane in the same way as (Horta et al. 2021a, see their appendix). In addition, we reduced the overall sample to stars with the most reliable elemental abundances. The APOGEE/ASPCAP abundance measurements for Fe, Mg, and Al are quite reliable in the range of metallicities spanned by our data. However, the abundance of Mn relies on a few relatively weak lines, becoming more uncertain in the low metallicity end. We examined the spectra of sample stars in a range of metallicities and S/N ratios and decided to restrict the sample to stars with $[Fe/H] > -1.6$ and, for those with $[Fe/H] < -1.5$ we only kept spectra with $S/N > 70$. As a result the samples for Carina, Draco, Sextans, and Ursa Minor become too small and they are not included in this analysis. On the basis of this reduced sample we fit linear relations to the data for each galaxy in the $[Mg/Mn]$ - $[Fe/H]$

and $[Al/Fe]$ - $[Fe/H]$ planes and derive the coefficients of the relations in the $[Mg/Mn]$ - $[Al/Fe]$ plane.

The result is displayed in Figure 5, where arrows indicating the direction (but not the modulus) of the metallicity gradients are overlaid on the reduced samples described above. The arrows confirm the visual impression about the direction of chemical evolution on the diagnostic plane of Figures 3-5. In massive Milky Way satellites the arrow points towards the lower right, whereas in lower mass satellites and the Milky Way disk it points towards the lower left. Along the same lines, the direction of the metallicity gradient in the disk population points strongly towards the lower left of the chemical plane. That is also the case of the disk of the Milky Way.

As indicated by the models in Figure 4, the overall direction of the metallicity gradient vector in the $[Mg/Mn]$ - $[Al/Fe]$ plane seems to be dependent on the star formation history of the system, which in turn is associated with its mass. It is therefore instructive to examine the behaviour of stellar populations on this chemical plane on the basis of chemical evolution models sampling a wider range of input parameters. That is the topic of the next Sub-section.

4.5 Chemical Evolution on the $[Mg/Mn]$ - $[Al/Fe]$ plane

In this Section we further examine the hypothesis that the metallicity gradient in the $[Mg/Mn]$ - $[Al/Fe]$ plane is sensitive to the SFR. For that purpose we calculated further chemical evolution models building on those presented in Figure 4, based on flexCE (Andrews et al. 2017). The same yields and SNe Ia delay time distribution are adopted in all calculations. In order to produce a spectrum of models that replicate the behaviour of dwarfs of different masses in this chemical plane, we adopt a range of values for the star formation

efficiency (SFE) and the wind mass-loading factor (η), which are respectively positively and negatively correlated with galaxy mass. For GE/S and MW we adopt the same models as discussed in Section 4.3. Table 5 summarises how these parameters differ between the models. They are chosen to cover, in a qualitative fashion, the range of properties characteristic of the dSph galaxies included in our sample. Specific parameters adopted are tuned to approximately match the properties of the MCs (MC-like), Sgr dSph (Sgr-like) and the lower mass galaxies (Sculptor-like).

The results are displayed in Figure 6. The top panel shows evolutionary tracks on the $[\text{Mg}/\text{Mn}]$ - $[\text{Al}/\text{Fe}]$ plane, whereas the star formation histories are plotted in the bottom panel. As in Figure 4, evolutionary times of 0.3, 1.0, and 5.0 Gyr are indicated for each model.

An examination of the behaviour of these various models is quite informative. We start by considering the fiducial MW model (Table 5), represented by the solid blue line. It is the one for which the SFR is by far the largest, exceeding those of the other models by at least an order of magnitude. On account of such a very high SFR, enrichment is initially entirely dominated by massive stars. Thus the $[\text{Mg}/\text{Mn}]$ ratio remains very high in the first few 100 Myr of evolution, whereas $[\text{Al}/\text{Fe}]$ builds up very quickly. With the increasing contribution by SN Ia at $t \gtrsim 300$ Myr, $[\text{Mg}/\text{Mn}]$ starts declining steadily. As the star formation rate declines further, so does the $[\text{Mg}/\text{Mn}]$ ratio, and eventually the contribution by SN Ia becomes important enough that the $[\text{Al}/\text{Fe}]$ ratio starts declining, after about 1 Gyr of evolution.

The remainder of this discussion contrasts the behaviour of various models within the first Gyr of cosmic evolution, where enrichment of the elements involved is dominated by CCSNe and SN Ia. In all of the dwarf-like models, the initial SFR is substantially lower than for the MW model. As a result, within the first few 100 Myr the evolution in $[\text{Al}/\text{Fe}]$ is slower while $[\text{Mg}/\text{Mn}]$ is more strongly influenced by the SN Ia enrichment. Hence, with decreasing SFR the early evolutionary tracks switch from pointing straight to the right to instead pointing at increasing degrees towards the lower right, and finally towards the lower left. Indeed, as suggested by the discussion in the previous section, within the first Gyr of evolution, there is a clear correlation between the orientation of the track on the $[\text{Mg}/\text{Mn}]$ - $[\text{Al}/\text{Fe}]$ plane and the SFR.

This qualitative analysis can inform an interpretation of broad trends that one can promptly grasp from even a perfunctory evaluation of the data. For instance, it is easy to see from Figures 3 and 4 that the data for GE/S are shifted towards higher $[\text{Al}/\text{Fe}]$ and $[\text{Mg}/\text{Mn}]$ than those of the MCs, which in turn have higher values than the lower mass MW satellites. The numbers are summarised in Table 6. The mean values for $[\text{Mg}/\text{Mn}]$ and $[\text{Al}/\text{Fe}]$ are substantially larger in GE/S than in all the MW satellites which, according to our interpretation of the models, suggests a stronger SFR in the early stages of evolution, in agreement with the results by Hasselquist et al. (2019b). The same conclusion can be drawn when comparing the massive MW satellites with their less massive counterparts, whose data suggest a slower rate of chemical evolution, associated with a weaker star formation rate.

Subsequent evolution, beyond ~ 1 Gyr seems to be dictated by the slope of the SFR. Models with strongly decreasing SFR (e.g., MW, Sausage-, and Sgr-like) tend to evolve more strongly towards lower $[\text{Mg}/\text{Mn}]$ with more or less constant $[\text{Al}/\text{Fe}]$, whereas those with more approximately constant SFR (e.g., Sculptor- and LMC-like) display a slight turn over of $[\text{Al}/\text{Fe}]$ and slower decline in $[\text{Mg}/\text{Mn}]$. These trends ultimately reflect the balance between the contribution




Figure 6. Chemical evolution and star formation histories for a series of one-zone open box models ran using the FLEXCE code. *Top panel:* tracks of $[\text{Mg}/\text{Mn}](t)$ plotted as a function of $[\text{Al}/\text{Fe}](t)$. *Bottom panel:* star formation rate as a function of time, $\text{SFR}(t)$. The solid blue tracks illustrate the fiducial model of Andrews et al. (2017) as shown in Fig. 4. For the other models, for a fixed history of gas inflow we vary the star formation efficiency and the wind mass loading factor (η) so as to qualitatively capture the behaviour of a selection of the Local Group dwarfs in our sample with varying stellar masses. Parameters adopted are listed in Table 5.

Table 6. Mean abundances and their dispersions of MW satellites in our sample.

System	$\langle[\text{Al}/\text{Fe}]\rangle$	$\sigma[\text{Al}/\text{Fe}]$	$\langle[\text{Mg}/\text{Mn}]\rangle$	$\sigma[\text{Mg}/\text{Mn}]$
GE/S	-0.21	0.17	0.50	0.20
GE/S (clean)	-0.23	0.14	0.51	0.20
SMC	-0.42	0.13	0.27	0.14
LMC	-0.31	0.12	0.23	0.13
Sgr	-0.46	0.08	0.14	0.10
Sculptor	-0.64	0.20	0.28	0.31
Fornax	-0.59	0.17	0.17	0.18

by CCSNe and SN Ia (and AGB stars in the case of AI) to the chemical enrichment of the interstellar medium (see Mason et al. 2022, in prep., for a detailed discussion).

The contrast between the histories of star formation of GE/S and massive MW satellites, and in particular the LMC and SMC is interesting in light of the fact that the mass of GE/S, according to various studies, is of the order of a few to several times $10^8 M_{\odot}$ (e.g., Lane et al. 2021; Deason et al. 2019; Mackereth et al. 2019), which is comparable to that of the SMC ($\sim 5 \times 10^8 M_{\odot}$) and smaller than that of the LMC ($\sim 1.5 \times 10^9 M_{\odot}$, McConnachie 2012). That galaxies with similar masses have undergone such vastly different histories of star formation indicates a physical variable other than mass is at play at regulating the star formation histories of dwarf galaxies. Hasselquist et al. (2021) suggest it is the environment in which the dwarf galaxies formed and evolved. We tackle this problem from the point of view of numerical cosmological simulations in a forthcoming paper (Mason et al. 2022, in prep.).

5 CONCLUSIONS

We present a comparative study of the distribution in chemical diagnostic planes of the stellar populations of the Gaia-Enceladus/Sausage (GE/S) system and those of satellites of the Milky Way. Our main conclusions are the following:

- We investigate the location on the [Mg/Mn] vs. [Al/Fe] plane of a GE/S sample defined purely on the basis of orbital properties. When selected in this way, GE/S stars lie almost entirely within the locus of that chemical plane deemed to contain “accreted” populations by (Das et al. 2020; Hawkins et al. 2015). While this result validates previous use of that method for the identification of stellar populations formed *ex situ*, caution is recommended, since old populations formed *in situ* share the same locus, as shown by Horta et al. (2021a). We therefore propose adopting a “chemically unevolved” nomenclature when referring to that particular locus of chemical space.

- The stellar populations of the satellites of the Milky Way are mostly divided between the chemically evolved and unevolved loci on this plane. The chemically evolved, more metal-rich stars are located towards the region of lower [Mg/Mn] and approximately constant, or slightly different, [Al/Fe].

- The distribution of GE/S stars on the [Mg/Mn]–[Al/Fe] plane differs from those of MW satellites in an important respect. The chemical evolution of its stellar populations in this plane suggest a higher *early* star formation rate than MW satellites with comparable or even higher masses, as suggested by Hasselquist et al. (2019b).

- The direction of the metallicity vector on the [Mg/Mn]–[Al/Fe] plane is an indicator of the early star formation rate of a system. Higher mass galaxies and/or those undergoing high star formation rates evolve more quickly in [Al/Fe] than in [Mg/Mn]. The existence of this trend is suggested by the APOGEE data on the stellar populations of the systems under study, and is boldly confirmed by the predictions of analytical chemical evolution models. The ensuing interpretation of our data on MW satellites in the light of such models leads to the conclusion that the early star formation rates of these systems was strongly affected by parameters other than galaxy mass.

ACKNOWLEDGEMENTS

Funding for the Sloan Digital Sky Survey IV has been provided by the Alfred P. Sloan Foundation, the U.S. Department of Energy Office of Science, and the Participating Institutions.

SDSS-IV acknowledges support and resources from the Center for High Performance Computing at the University of Utah. The SDSS website is www.sdss.org.

SDSS-IV is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS Collaboration including the Brazilian Participation Group, the Carnegie Institution for Science, Carnegie Mellon University, Center for Astrophysics | Harvard & Smithsonian, the Chilean Participation Group, the French Participation Group, Instituto de Astrofísica de Canarias, The Johns Hopkins University, Kavli Institute for the Physics and Mathematics of the Universe (IPMU) / University of Tokyo, the Korean Participation Group, Lawrence Berkeley National Laboratory, Leibniz Institut für Astrophysik Potsdam (AIP), Max-Planck-Institut für Astronomie (MPIA Heidelberg), Max-Planck-Institut für Astrophysik (MPA Garching), Max-Planck-Institut für Extraterrestrische Physik (MPE), National Astronomical Observatories of China, New Mexico State University, New York University, University of Notre Dame, Observatório Nacional / MCTI, The Ohio State University, Pennsylvania State University, Shanghai Astronomical Observatory, United Kingdom Participation Group, Universidad Nacional Autónoma de México, University of Arizona, University of Colorado Boulder, University of Oxford, University of Portsmouth, University of Utah, University of Virginia, University of Washington, University of Wisconsin, Vanderbilt University, and Yale University.

This work has made use of data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

Software used in this research: Astropy (Astropy Collaboration et al. 2018; Astropy Collaboration et al. 2013), SciPy (Virtanen et al. 2020), NumPy (Oliphant 2006; Harris et al. 2020), Matplotlib (Hunter 2007), Galpy (Bovy 2015), TOPCAT (Taylor 2005), flexCE (Andrews et al. 2017).

DATA AVAILABILITY

This research was made possible with data from the SDSS-IV/APOGEE-2, 17th data release and Gaia eDR3, publicly available at https://www.sdss.org/dr17/irspec/spectro_data/ and <https://gea.esac.esa.int/archive/>, respectively.

REFERENCES

- Abdurro'uf et al., 2021, arXiv e-prints, p. [arXiv:2112.02026](https://arxiv.org/abs/2112.02026)
- Abolfathi B., et al., 2018, *ApJS*, **235**, 42
- Aguado D. S., et al., 2021, *ApJ*, **908**, L8
- Andrews B. H., Weinberg D. H., Schönrich R., Johnson J. A., 2017, *ApJ*, **835**, 224
- Astropy Collaboration et al., 2013, *A&A*, **558**, A33
- Astropy Collaboration et al., 2018, *AJ*, **156**, 123
- Barbuy B., Chiappini C., Gerhard O., 2018, *Annual Review of Astronomy and Astrophysics*, **56**, 223
- Belokurov V., et al., 2006, *ApJ*, **642**, L137
- Belokurov V., Erkal D., Evans N. W., Koposov S. E., Deason A. J., 2018, *MNRAS*, **478**, 611
- Belokurov V., Sanders J. L., Fattahi A., Smith M. C., Deason A. J., Evans N. W., Grand R. J., 2020, *Monthly Notices of the Royal Astronomical Society*, **494**, 3880
- Beraldo e Silva L., Debattista V. P., Nidever D., Amarante J. A. S., Garver B., 2021, *MNRAS*, **502**, 260
- Blanton M. R., et al., 2017, *AJ*, **154**, 28
- Blumenthal G. R., Faber S. M., Primack J. R., Rees M. J., 1984, *Nature*, **311**, 517
- Bouchard A., Carignan C., Staveley-Smith L., 2006, *AJ*, **131**, 2913
- Bovy J., 2015, *ApJS*, **216**, 29
- Bowen I. S., Vaughan A. H. J., 1973, *Appl. Opt.*, **12**, 1430
- Buchmann L., Hilgemeier M., Krauss A., Redder A., Rolfs C., Trautvetter H. P., Donoghue T. R., 1984, *Nuclear Phys. A*, **415**, 93
- Carignan C., Beaulieu S., Côté S., Demers S., Mateo M., 1998, *AJ*, **116**, 1690
- Carrillo A., Hawkins K., Jofré P., de Brito Silva D., Das P., Lucey M., 2022, *MNRAS*, **513**, 1557
- Cirasuolo M., Afonso J., Bender R., Bonifacio P., Evans C., Kaper L., Oliva E., Vanzi L., 2011, *The Messenger*, **145**, 11
- Clayton D. D., 1984, *ApJ*, **280**, 144
- Cohen J. G., Huang W., 2009, *ApJ*, **701**, 1053
- Cohen J. G., Huang W., 2010, *ApJ*, **719**, 931
- Conroy C., et al., 2019, *ApJ*, **883**, 107
- Cui X.-Q., et al., 2012, *Research in Astronomy and Astrophysics*, **12**, 1197
- Dalton G., 2016, in Skillen I., Balcells M., Trager S., eds, *Astronomical Society of the Pacific Conference Series Vol. 507, Multi-Object Spectroscopy in the Next Decade: Big Questions, Large Surveys, and Wide Fields*. p. 97
- Das P., Hawkins K., Jofré P., 2020, *MNRAS*, **493**, 5195
- De Silva G. M., et al., 2015, *MNRAS*, **449**, 2604
- Deason A. J., Belokurov V., Sanders J. L., 2019, *MNRAS*, **490**, 3426
- Feuillet D. K., Feltzing S., Sahlholdt C. L., Casagrande L., 2020, *MNRAS*, **497**, 109
- Feuillet D. K., Sahlholdt C. L., Feltzing S., Casagrande L., 2021, *MNRAS*, **508**, 1489
- Frenk C. S., White S. D. M., 2012, *Annalen der Physik*, **524**, 507
- Gaia Collaboration et al., 2016, *A&A*, **595**, A1
- Gaia Collaboration et al., 2021, *A&A*, **649**, A1
- García Pérez A. E., et al., 2016, *AJ*, **151**, 144
- Gehrz R. D., Truran J. W., Williams R. E., Starrfield S., 1998, *PASP*, **110**, 3
- Gonzalez O. A., et al., 2020, *The Messenger*, **180**, 18
- Greive J., Putman M. E., 2009, *ApJ*, **696**, 385
- Guelin M., Forestini M., Valiron P., Zdziarski L. M., Anderson M. A., Cernicharo J., Kahane C., 1995, *A&A*, **297**, 183
- Gunn J. E., et al., 2006, *AJ*, **131**, 2332
- Harris C. R., et al., 2020, *Nature*, **585**, 357
- Hasselquist S., et al., 2017, *ApJ*, **845**, 162
- Hasselquist S., et al., 2019a, *ApJ*, **872**, 58
- Hasselquist S., et al., 2019b, *The Astrophysical Journal*, **872**, 58
- Hasselquist S., et al., 2021, *ApJ*, **923**, 172
- Hawkins K., Jofré P., Masseron T., Gilmore G., 2015, *Monthly Notices of the Royal Astronomical Society*, **453**, 758–774
- Hayes C. R., et al., 2018, *ApJ*, **852**, 49
- Hayes C. R., et al., 2020, *ApJ*, **889**, 63
- Haywood M., Di Matteo P., Lehnert M. D., Snaith O., Khoperskov S., Gómez A., 2018, *ApJ*, **863**, 113
- Helmi A., Babusiaux C., Koppelman H. H., Massari D., Veljanoski J., Brown A. G. A., 2018, *Nature*, **563**, 85
- Hendricks B., Koch A., Lanfranchi G. A., Boeche C., Walker M., Johnson C. I., Peñarrubia J., Gilmore G., 2014, *ApJ*, **785**, 102
- Hill V., et al., 2019, *A&A*, **626**, A15
- Hillebrandt W., Niemeyer J. C., 2000, *Annual Review of Astronomy and Astrophysics*, **38**, 191
- Holtzman J. A., et al., 2015, *AJ*, **150**, 148
- Holtzman J. A., et al., 2018, *AJ*, **156**, 125
- Horta D., et al., 2020, *MNRAS*, **493**, 3363
- Horta D., et al., 2021a, *MNRAS*, **500**, 1385
- Horta D., et al., 2021b, *MNRAS*, **500**, 1385
- Horta D., et al., 2022a, *MNRAS*,
- Horta D., et al., 2022b, arXiv e-prints, p. [arXiv:2204.04233](https://arxiv.org/abs/2204.04233)
- Hunter J. D., 2007, *Computing in Science and Engineering*, **9**, 90
- Ibata R. A., Gilmore G., Irwin M. J., 1994, *Nature*, **370**, 194
- Iwamoto K., Brachwitz F., Nomoto K., Kishimoto N., Umeda H., Hix W. R., Thielemann F.-K., 1999, *ApJS*, **125**, 439
- Kamiński T., et al., 2018, *Nature Astronomy*, **2**, 778
- Kobayashi C., Umeda H., Nomoto K., Tominaga N., Ohkubo T., 2006, *ApJ*, **653**, 1145
- Kobayashi C., Leung S.-C., Nomoto K., 2020, *ApJ*, **895**, 138
- Koppelman H. H., Helmi A., Massari D., Price-Whelan A. M., Starkenburg T. K., 2019a, *Astronomy & Astrophysics*, **631**, L9
- Koppelman H. H., Helmi A., Massari D., Price-Whelan A. M., Starkenburg T. K., 2019b, *A&A*, **631**, L9
- Kruijssen J. M. D., et al., 2020, *MNRAS*, **498**, 2472
- Lane J. M. M., Bovy J., Mackereth J. T., 2021, arXiv e-prints, p. [arXiv:2106.09699](https://arxiv.org/abs/2106.09699)
- Leung H. W., Bovy J., 2019, *MNRAS*, **489**, 2079
- Limongi M., Chieffi A., 2006, *ApJ*, **647**, 483
- Mackereth J. T., Bovy J., 2018, *PASP*, **130**, 114501
- Mackereth J. T., Bovy J., 2020a, *MNRAS*, **492**, 3631
- Mackereth J. T., Bovy J., 2020b, *MNRAS*, **492**, 3631
- Mackereth J. T., Crain R. A., Schiavon R. P., Schaye J., Theuns T., Schaller M., 2018, *MNRAS*, **477**, 5072
- Mackereth J. T., et al., 2019, *MNRAS*, **482**, 3426
- Majewski S. R., et al., 2013, *ApJ*, **777**, L13
- Majewski S. R., et al., 2017, *AJ*, **154**, 94
- Martin N. F., Ibata R. A., Chapman S. C., Irwin M., Lewis G. F., 2007, *MNRAS*, **380**, 281
- Massari D., Koppelman H. H., Helmi A., 2019, *A&A*, **630**, L4
- Matsuno T., et al., 2021, *ApJ*, **912**, 72
- Matteucci F., Greggio L., 1986, *A&A*, **154**, 279
- McConnachie A. W., 2012, *AJ*, **144**, 4
- McConnachie A. W., Venn K. A., 2020, *Research Notes of the American Astronomical Society*, **4**, 229
- McMillan P. J., 2017, *MNRAS*, **465**, 76
- McWilliam A., 1997, *ARA&A*, **35**, 503
- Morris M., Serabyn E., 1996, *Annual Review of Astronomy and Astrophysics*, **34**, 645
- Naidu R. P., Conroy C., Bonaca A., Johnson B. D., Ting Y.-S., Caldwell N., Zaritsky D., Cargile P. A., 2020, *ApJ*, **901**, 48
- Nataf D. M., 2017, *Publications of the Astronomical Society of Australia*, **34**
- Nidever D. L., et al., 2015, *AJ*, **150**, 173
- Nidever D. L., et al., 2020, *ApJ*, **895**, 88
- Nissen P. E., Schuster W. J., 2010, *A&A*, **511**, L10
- Nissen P. E., Schuster W. J., 2011, *A&A*, **530**, A15
- Nofar I., Shaviv G., Starrfield S., 1991, *ApJ*, **369**, 440
- Nomoto K., Thielemann F. K., Yokoi K., 1984, *ApJ*, **286**, 644
- Oliphant T., 2006, *A Guide to NumPy*. Trelgol Publishing, <https://books.google.co.uk/books?id=fKulSgAACAAJ>
- Prantzos N., Diehl R., 1996, *Phys. Rep.*, **267**, 1
- Rich R. M., 2013, *The Galactic Bulge*. p. 271, doi:10.1007/978-94-007-5612-0_6
- Samland M., 1998, *ApJ*, **496**, 155
- Santana F. A., et al., 2021, arXiv e-prints, p. [arXiv:2108.11908](https://arxiv.org/abs/2108.11908)

- Schuster W. J., Moreno E., Nissen P. E., Pichardo B., 2012, *A&A*, **538**, A21
- Shetrone M., Venn K. A., Tolstoy E., Primas F., Hill V., Kaufer A., 2003, *AJ*, **125**, 684
- Starkenburger E., et al., 2013, *A&A*, **549**, A88
- Steinmetz M., et al., 2006, *AJ*, **132**, 1645
- Taylor M. B., 2005, **347**, 29
- Theler R., et al., 2020, *A&A*, **642**, A176
- Virtanen P., et al., 2020, *Nature Methods*, **17**, 261
- Walker M. G., Mateo M., Olszewski E. W., Gnedin O. Y., Wang X., Sen B., Woodroffe M., 2007, *ApJ*, **667**, L53
- Walker M. G., Mateo M., Olszewski E. W., 2008, *ApJ*, **688**, L75
- Walker M. G., Mateo M., Olszewski E. W., 2009a, *AJ*, **137**, 3100
- Walker M. G., Mateo M., Olszewski E. W., Peñarrubia J., Evans N. W., Gilmore G., 2009b, *ApJ*, **704**, 1274
- Weinberg D. H., et al., 2019, *ApJ*, **874**, 102
- Weiss A., Truran J. W., 1990, *A&A*, **238**, 178
- Wheeler J. C., Sneden C., Truran James W. J., 1989, *ARA&A*, **27**, 279
- White S. D. M., Rees M. J., 1978, *MNRAS*, **183**, 341
- Wilkinson M. I., Kleyna J. T., Evans N. W., Gilmore G. F., Irwin M. J., Grebel E. K., 2004, *ApJ*, **611**, L21
- Wilson J. C., et al., 2019, *PASP*, **131**, 055001
- Woosley S. E., Weaver T. A., 1995, *ApJS*, **101**, 181
- Yanny B., et al., 2009, *AJ*, **137**, 4377
- Zasowski G., et al., 2013, *The Astronomical Journal*, **146**, 81
- Zasowski G., et al., 2017, *AJ*, **154**, 198
- de Boer T. J. L., Tolstoy E., Lemasle B., Saha A., Olszewski E. W., Mateo M., Irwin M. J., Battaglia G., 2014, *A&A*, **572**, A10
- de Jong R. S., et al., 2019, *The Messenger*, **175**, 3

ORIGINAL UNEDITED MANUSCRIPT