

Article

The Effect of Extraction Methods on the Total Phenols and Total Flavonoids Content of Jackfruit (*Artocarpus heterophyllus* Lamk) Peels Extract

Article Info

Article history :

Received November 24, 2022
Revised December 05, 2022
Accepted December 10, 2022
Published March 30, 2023
(In-Press)

Keywords :

Extraction methods, total phenolic content, total flavonoid content, jackfruit peel.

Dyan Wigati¹, Lindawati Setyaningrum^{1*}, Dwi Koko Pratoko²

¹Bachelor in Pharmacy Program, Faculty of Health Science, Universitas dr Soebandi, Jember, Indonesia

²Drug Utilisation and Discovery Research Group, Faculty of Pharmacy, Universitas Jember, Jember, Indonesia

Abstract. Jackfruit (*Artocarpus heterophyllus* Lamk), especially the flesh usually consumed in a ripe state. The jackfruit peel is a non-edible waste. The research aims to determine the effect of different extraction methods on the total phenolic content and total flavonoid content of jackfruit peel extract. Extraction methods are percolation, maceration, and digestion. The total phenolic content (TPC) and total flavonoid content (TFC) using a spectrophotometer with $AlCl_3$ and Folin-cialcalteu reagents. Rutin and gallic acid as a standarts. The data from various extraction methods were analyzed statistically by One-Way Anova. The results showed TPC values by percolation, maceration, and digestion respectively 5.94 ± 0.92 ; 3.03 ± 0.16 and 2.72 ± 0.22 (GAE mg/g). The TFC respectively 13.04 ± 0.38 ; 5.49 ± 0.33 and 4.16 ± 0.28 (RTE mg/g). Statistical analysis showed that there were significant differences ($p < 0.05$) in the levels of flavonoids and total phenolics between various extraction methods from the ethanol extract of jackfruit peel. The percolation method showed the highest TPC, TFC, and yield rather than other methods.

This is an open access article under the [CC-BY](https://creativecommons.org/licenses/by/4.0/) license.



This is an open access article distributed under the Creative Commons 4.0 Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. ©2023 by author.

Corresponding Author :

Lindawati Setyaningrum

Bachelor in Pharmacy Program, Faculty of Health Science, Universitas dr Soebandi, Jember, Indonesia

Email : linda.w.setyaningrum@uds.ac.id

1. Introduction

Indonesia is a country with a diversity of medicinal plants and has the potential to be developed as a source of new drugs. Secondary metabolic compounds extracted from plants have been reported effective in preventing or treating chronic diseases, such as natural antioxidants that can prevent

oxidative stress or cell damage [1-2]. The leaves are commonly used as traditional medicine, but the fruit, flowers, seeds, and even the peel also contain lots of bioactive compounds. Sundai lime peel ethyl acetate fraction has antioxidant activity against DPPH caused by flavonoids and phenolic compounds.

Jackfruit (*Artocarpus heterophyllus* Lamk) is a fruit that is usually consumed in a ripe state, especially the flesh. The unripe (young) jackfruit can also be cooked as a vegetable. The part of the jackfruit such as the peel and central axis is a non-edible waste [4-6]. The right way to reduce disposal costs of non-edible waste is the utilization of by-products, which will increase economic value and environmental protection [7-9]. The ethanol extract of jackfruit peel reported contains a group of alkaloids, flavonoids, phenols and flavonoids terpenoids. One of them is stigmasterol which is a flavonoid group. Jackfruit peel reportedly contains pectin, cellulose, ferulic acid, 4-hydroxybenzoic acid, tryptophan N-glucoside and caffeic acid [10-12].

Jackfruit peel extract has a higher concentration of total phenolics and flavonoids than pulp, flakes, or seed extract. It also has the strongest glucosidase inhibitor and DPPH and ABTS scavenging abilities [13]. The ethanolic extract of jackfruit peel had an antioxidant activity with an IC₅₀ of 87.09 g/ml [14]. The total value of polyphenols and flavonoids is correlated with their antioxidant activity, mainly due to the redox properties that allow phenolic compounds to act as reducing agents or hydrogen donors [15-16]. The jackfruit peel was suggested as a new source of natural antioxidants.

The main step in obtaining bioactive compounds from natural materials is extraction. Extraction is the process of separating active secondary metabolites from plants using suitable solvents, commonly used to quantify or isolate compounds. Several factors affect the level of bioactive metabolites in the extract including the total phenolic and flavonoid content. The selection of extraction methods and solvents is a critical point in producing extracts with high levels of bioactive compounds [17-18]. To standardize extracts and increase yield, it is important to select an appropriate extraction method because the different extraction methods will produce different yields and bioactivity. Conventional methods like maceration, percolation, infusion, decoction, hot continuous extraction, etc., are often used for extraction [19-20].

Maceration is one of the simple methods commonly used to extract the metabolite bioactive and this method is commonly used for the extraction of thermolabile compounds. A modification of the maceration method is digestion where extraction involves churning and low heat (Kinetic maceration). The other method is percolation which is extraction using a percolator with constant solvent changes during the extraction process. Research on the effect of different extraction methods on the total phenol and total flavonoid content of jackfruit peel has never been reported. Therefore, to find the effect of different extraction methods to the yield, the total phenol and the total flavonoid content of jackfruit peel ethanolic extract, the present investigation was carried out.

2. Experimental Section

2.1. Materials

Jackfruit collected from a local market in Jember, East Java. Ethanol 96 %, rutin (Sigma), Gallic acid (Sigma), Follin ciaocalteu (Merck), AlCl₃, CH₃COONa anhydrous, sodium carbonate, methanol, distilled water.

2.2. Instrumentation

A macerator container, percolator, separating funnel, glassware, volumetric flask, filler, UV-Vis spectrophotometer, and rotary evaporator.

2.3. Methods

2.3.1. Sample Preparation

The jackfruit is peeled and separated between the skin, flesh, and seeds, then cut into small pieces and washed with clean water. The next process is to dry the peel in an oven at a temperature of 50 °C until dry then mashed and sieved through a 30 mesh. Dried jackfruit peel powder is ready to be extracted.

2.3.2. Extraction

2.3.2.1 Maceration

200 g jackfruit peel dried powder mixed with ethanol 96 % (1:10) in a macerator container, soaked for 3 days with continuous stirring. The liquid extract is separated from the residue by filtration and then concentrated using a rotary evaporator. The yield was calculated.

2.3.2.2 Percolation

Soaking 60 g of jackfruit peel dried powder with 96% ethanol for 24 hours then putting it into the percolator and percolating with 96% ethanol (1:10). The supernatant was concentrated with a rotary evaporator then the yield was calculated.

2.3.2.3 Digestion

Mixed 100 g of jackfruit peel dry powder with 1 L ethanol 96 % (1:10). Extraction was carried out at 50 °C with a rotation speed of 120 rpm (Kinetic maceration- digestion) [21]. The supernatants separated from the residue by filtration and then concentrated using a rotary evaporator, and calculated the yield.

2.3.3. Analysis of Total Phenolic Content

Total phenolic test using Folin-Ciocalteu reagent. The TPC value is expressed as milligrams of gallic acid equivalent (mg GAE/g extract) [22-23]. 0.1 ml from 1.000 ppm extract solution was mixed with 7.9 ml of distilled water and 0.5 ml of Folin-ciaocalteu reagent for 15 minutes before adding 1.5 ml of Na₂CO₃ solution and incubating for 30 minutes. The absorbance was measured at 760 nm with a UV Vis spectrophotometer and compared to the gallic acid calibration curve. The test was three replicated

2.3.4. Analysis of Total Flavonoid Content

Total flavonoid levels were measured using a colorimetric method using AlCl₃ [24-25]. The sample was made at a concentration of 10,000 ppm and then 0.5 mL mixed with 1.5 mL methanol, 0.1 mL AlCl₃ solution, 0.1 mL 20% Na acetate anhydrous, and 2.8 mL of distilled water, homogenized. The mixture was incubated for 30 minutes and the absorbance was measured at 421 nm. The total flavonoid content was calculated as rutin equivalent (mg rutin/g extract). This test was three replicated.

2.3.5. Statistic Analysis

Results are expressed as mean ± standard deviation (SD) of three replicates. Regression analysis to calculate the TPC and TFC. Data analysis between extracts with various methods was carried out using one-way ANOVA.




Figure 1. Design experimental

3. Results and Discussion

Biologically active substances are typically found in small amounts in plants. An effective extraction technique can produce high yields of extracts with minimal changes in the extracts' properties. Several studies reported that different extraction techniques influence the biological activity also the levels of compounds in the extract, including total phenolic and total flavonoid levels. Selecting the appropriate extraction method is necessary.

Total phenolics and total flavonoids are secondary metabolites that have the potential as antioxidants. Jackfruit peel has high total phenolic and total flavonoid content [13]. There was a high correlation between the extraction method and total phenolic content on antioxidant activity [26]. The higher the total phenolic content in extracts will increase an antioxidant activity.

The jackfruit peel extraction in this study used maceration, percolation, and digestion methods (kinetic maceration) with ethanol 96 % as solvent. Ethanol is a universal solvent and can dissolve a large number of secondary metabolites. It is also less toxic than other organic solvents and more volatile [27]. Other solvents such as water, methanol, ethanol, and acetonitrile even cyclohexane could be used in the extraction of phenolic compounds [28-30] or a combination of several solvents.

The highest yield was obtained from extraction using the percolation method, followed by maceration and digestion. The different extraction methods impacted the yield describes in figure 1.




Figure 1. The yield of the jackfruit peel extract in various methods of extraction

In the percolation method, the sample is moistened with solvent several times and then put into the percolator and the solvent is allowed to flow from top to bottom until the active ingredients are extracted. Solvents can be used up to their saturation point [31-32]. The benefit of this percolation method is the extraction of secondary metabolites is more optimal because the solvent flows constantly and is always new, increasing the degree of concentration difference and the yield higher than other methods. Similar to Safitri et al [33] and Anwar et al, (2022) [34] who reported that the yield of extract by the percolation method produced the highest yield compared to other extraction methods.

Digestion is maceration with kinetic and heat presence at a low temperature. Digestion is an option when extraction at a high enough temperature will damage the compounds in the sample [35-36] but extraction time is faster than maceration so secondary metabolites may not be extracted completely. In this study, the digestion methods showed the lowest yield. A high yield value indicates that a large amount of extract is produced, but quantity does not always correlate with extract quality [37-38].

To determine the total phenolic content (TPC), the first step is to calculate the standard curve equation using gallic acid as a standard. The regression equation obtained is $y = 0.00402x + 0.0095$ with the correlation coefficient ($r = 0.9993$) shown in Figure 2. Gallic acid is a phenolic compound that is stable, pure, cheap, and widely available [39-41]. Suitable for use as an assay standard. The Folin-Ciocalteu reagent was used to determine TPC levels. The principle is the reducing power of phenolic hydroxyl groups. A phenolic compound with an aromatic core can reduce phosphotungstic phosphorus to molybdenum blue [42]. The total phenolic content expressed in GAE is the number of milligram equivalents of gallic acid per gram of sample.




Figure 2. Standard curve equation of gallic acid

The total phenolic and total flavonoid content of ethanol extract of jackfruit (*Artocarpus heterophyllus*) peels with various extraction methods are described in Table 1.

Table 1. Total phenolic and total flavonoid content of ethanol extract of jackfruit (*Artocarpus heterophyllus*) peels with various extraction methods

Methods	Organoleptic	Total Phenolic (GAE mg/g)	Total Flavonoid (RTE mg/g)
Maceration		3.03 ± 0.16 ^a	5.49 ± 0.33 ^d
Percolation	thick extract, brownish green, and specific aroma	5.94 ± 0.92 ^b	13.04 ± 0.38 ^e
Digestion		2.72 ± 0.22 ^c	4.16 ± 0.28 ^f

GAE = gallic acid equivalent; RTE = rutin equivalent;

Values with different superscripts in one column show significant differences ($p < 0.05$).

Based on table 1, the highest TPC was obtained in the percolation method, followed by maceration and digestion. The results are in agreement with the TPC of bay leaf extract [39] and beluntas leaf methanol extract [33] that showed the percolation methods have the highest TPC rather than other methods. The lowest TPC value in the digestion method indicates that the phenolic compounds are thermolabile, and heating during the extraction process affects the degradation and destruction of these phenolic compounds. In the maceration method (cold extraction), the solvent does not change continuously, it is affected by solvent saturation and the phenolic in a sample may not be completely extracted [31].

Total flavonoid levels are measured by colorimetry, based on the formation of color complexes. $AlCl_3$ forms complexes with the hydroxyl group on flavonoids. The total flavonoid content (TFC) expressed in RTE, which is the number of milligram equivalents of rutin per gram of sample. The total flavonoid content is calculated using the regression equation $y = 0.0031x + (-0.0151)$ with correlation coefficient = 0.9996 using rutin as a reference shown on Figure 3 below.




Figure 3. Standard curve equation of rutin

The result of the determination of TFC (table 1), showed relatively similar to TPC that the percolation method showed the highest TFC compared to other methods. The different extraction methods resulted in different total phenol and total flavonoid values. Data on TPC and TFC were then analyzed statistically using One-Way Anova. The results showed that the levels of flavonoids and total phenolic ethanol extract of jackfruit peel with various extraction methods have differences significant with a significance value <0.05 .

Maceration and percolation are cold extraction methods, whereas digestion is a heating and stirring modification of maceration. Settharaksa et al [43] reported that temperature and heating during the extraction process can affect the levels of flavonoids and phenolic compounds. The heating process either directly or indirectly can affect the levels of these two compounds. This research showed a correlation between the levels of total phenol, total flavonoids, and the yield of each extract. Extracts by percolation method had the highest levels of TPC, TFC, and yield compared to the maceration and digestion methods. However, several studies have reported that the amount of yield with the levels of secondary metabolites extracted using different methods was unrelated [39-40]. Extraction method, extraction temperature, extraction time, and solvent selection are factors that affect the quality of the extract. Optimization is needed to get the expected active compounds.

Flavonoids may be phenolic compounds that contribute to various biological activities such as antioxidants [44] but there are many other phenolic compounds including saponins, terpenoids, etc. Phenolic compounds can prevent cell damage because of their ability to scavenge free radicals with reduction mechanisms and electron donors [45-46]. Selection of the appropriate extraction method is the first step for large-scale process implementation to produce high-quality active compounds.

4. Conclusion

The research shows that different extraction methods were applied on the same plant with the same solvent affect significantly in terms of the parameters evaluated. This might be useful in deciding on the best extraction technique for a later assessment of biological activity. The percolation method showed the highest TPC, TFC, and yield rather than maceration and digestion. This method can be developed for the extraction of Jackfruit peel.

5. Acknowledgement

This study was supported by Hibah Penelitian Internal 2021 Universitas dr Soebandi, Jember

References

- [1] Yatma., S. Falah., L. Ambarsari., S. I. Aisyah., and W. Nurcholis. (2023). Optimization of Extraction of Phenolic and Antioxidant Activities from *Celosia cristata* Seeds Using Response Surface Methodology. *Biointerface Res. Appl. Chem.*, vol. 13, no. 2, p. 148
- [2] Sayeed, R., Thakur, M., & Gani, A. (2020). *Celosia cristata* Linn. flowers as a new source of nutraceuticals-A study on nutritional composition, chemical characterization and in-vitro antioxidant capacity. *Heliyon*, 6(12), e05792.
- [3] M. M. Melindra., M. efdi., A. Afrizal., and N. L. Pernadi. (2022). Antioxidant Activities Extracts N-Hexane, Ethyl Acetate and Methanol of Limau Sundai Peels (*Citrus Nobilis* Lour). *EKSAKTA Berk. Ilm. Bid. MIPA*, vol. 23, no. 04, pp. 291–297
- [4] F. Akter and M. Haque. (2020). Jackfruit Waste: A Promising Source of Food and Feed. *Ann. Bangladesh Agric.*, vol. 23, no. 1, pp. 91–102
- [5] Alves, J. L. F., da Silva, J. C. G., Mumbach, G. D., Di Domenico, M., da Silva Filho, V. F., de Sena, R. F., ... & Marangoni, C. (2020). Insights into the bioenergy potential of jackfruit wastes considering their physicochemical properties, bioenergy indicators, combustion behaviors, and emission characteristics. *Renewable Energy*, 155, 1328-1338.
- [6] Kalse, S. B., & Swami, S. B. (2022). Recent application of jackfruit waste in food and material engineering: A review. *Food Bioscience*, 101740.
- [7] A. A. Sundarraj and T. V. Ranganathan. (2017). Physicochemical characterization of Jackfruit (*Artocarpus integer* (Thumb.)) Peel. *Res. J. Pharm. Biol. Chem. Sci.*, vol. 8, no. 3, pp. 2285–2295
- [8] Calderón-Chiu, C., Calderón-Santoyo, M., Herman-Lara, E., & Ragazzo-Sánchez, J. A. (2021). Jackfruit (*Artocarpus heterophyllus* Lam) leaf as a new source to obtain protein hydrolysates: Physicochemical characterization, techno-functional properties and antioxidant capacity. *Food Hydrocolloids*, 112, 106319.
- [9] Morelos-Flores, D. A., Montalvo-González, E., Chacón-López, M. A., Santacruz-Varela, A., Zamora-Gasga, V. M., Torres-García, G., & de Lourdes García-Magaña, M. (2022). Comparative Study of Four Jackfruit Genotypes: Morphology, Physiology and Physicochemical Characterization. *Horticulturae*, 8(11), 1010.
- [10] A. M. N. Lal *et al.* (2021). Pulsed electric field combined with microwave-assisted extraction of pectin polysaccharide from jackfruit waste. *Innov. Food Sci. Emerg. Technol.*, vol. 74, p. 102844
- [11] D. A. R. Lokhande., D. K. S. Wani., and M. S. Siddiqui. (2016). 'Study of Pectin from Peels of *Magnifera Indica* and *Artocarpus Hetrophyllus*. *Interna. Conf. Glob. Trends Engg., Tech. Manag*, pp. 375–381
- [12] S. min Pu *et al.* (2022). Comparative Investigation on the Phytochemicals and Biological Activities of Jackfruit (*Artocarpus heterophyllus* Lam.) pulp from Five Cultivars. *Plant Foods Hum. Nutr.*
- [13] L. Zhang *et al.* (2017). Jackfruit (*Artocarpus heterophyllus* Lam.) peel: A better source of antioxidants and a-glucosidase inhibitors than pulp, flake and seed, and phytochemical profile by HPLC-QTOF-MS/MS. *Food Chem.*, vol. 234, pp. 303–313
- [14] M. Raihan., N. Taqwa., A. R. Hanifah., S. Lallo., I. Ismail., and M. N. Amir. (2020). Skrining Fitokimia Ekstrak Kulit Buah Nangka (*Artocarpus heterophyllus*) Dan Aktifitas Antioksidannya Terhadap [2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonate)] (ABTS). *Maj. Farm. dan Farmakol.*, vol. 23, no. 3, pp. 101–105
- [15] S. A. Baba and S. A. Malik. (2015). Determination of total phenolic and flavonoid content, antimicrobial and antioxidant activity of a root extract of *Arisaema jacquemontii* Blume. *J. Taibah Univ. Sci.*, vol. 9, no. 4, pp. 449–454

-
- [16] Shehzadi, S., Khan, S. M., Mustafa, G., Abdullah, A., Khan, I., Ahmad, Z., ... & Raposo, A. (2022). Antiviral COVID-19 protein and molecular docking: In silico characterization of various antiviral compounds extracted from *Arisaema jacquemontii* Blume. *Frontiers in Public Health*, 10.
- [17] T. Dhanani., S. Shah., N. A. Gajbhiye., and S. Kumar. (2017). Effect of extraction methods on yield, phytochemical constituents and antioxidant activity of *Withania somnifera*. *Arab. J. Chem.*, vol. 10, pp. S1193–S1199
- [18] B. P. Pratama., Y. Pranoto., Supriyadi., and R. T. Swasono. (2023). The Properties of Salam Leaf Extract (*Syzygium Polyanthum*) with Different Solvent Ratio and Processing Time Using Ultrasonication-Assisted Extraction Method. *J. Appl. Sci. Eng.*, vol. 26, no. 4, pp. 581–587,
- [19] O. A. O. Adam., R. S. M. Abadi., and S. M. H. Ayoub. (2019). The Effect of Extraction method and Solvents on yield and Antioxidant Activity of Certain Sudanese Medicinal Plant Extracts. *J. Phytopharm.*, vol. 8, no. 5, pp. 248–252
- [20] E. El Maaiden *et al.* (2022). A Comparative Study between Conventional and Advanced Extraction Techniques: Pharmaceutical and Cosmetic Properties of Plant Extracts. *Molecules*, vol. 27, no. 7
- [21] N. L. Rahmah., B. S. D. Dewanti., and F. Azizah. (2018). Combination of kinetic maceration - digestion in the extraction of areca seeds (*Areca catechu* L.). *Adv. Food Sci. Sustain. Agric. Agroindustrial Eng.*, vol. 1, no. 2, pp. 27–33
- [22] A. M. Abdel-Aty., R. I. Bassuiny., A. Z. Barakat., and S. A. Mohamed. (2019). Upgrading the phenolic content, antioxidant and antimicrobial activities of garden cress seeds using solid-state fermentation by *Trichoderma reesei*. *J. Appl. Microbiol.*, vol. 127, no. 5, pp. 1454–1467,
- [23] Alyileili, S. R., Hussein, A. S., Ibrahim, W., & El-Tarabily, K. A. (2020). Phytochemical composition and antioxidant activity of *Trichoderma reesei* degraded date (*Phoenix dactylifera* L.) pits. *Current Bioactive Compounds*, 16(4), 528-536.
- [24] A. M. Shraim., T. A. Ahmed., M. M. Rahman., and Y. M. Hijji. (2021). Determination of total flavonoid content by aluminum chloride assay: A critical evaluation. *LWT*, vol. 150, p. 111932
- [25] Tristantini, D., & Amalia, R. (2019, December). Quercetin concentration and total flavonoid content of anti-atherosclerotic herbs using aluminum chloride colorimetric assay. In *AIP Conference Proceedings* (Vol. 2193, No. 1, p. 030012). AIP Publishing LLC.
- [26] L. Hartanti., S. M. K. Yonas., J. J. Mustamu., S. Wijaya., H. K. Setiawan., and L. Soegianto. (2019). Influence of extraction methods of bay leaves (*Syzygium polyanthum*) on antioxidant and HMG-CoA Reductase inhibitory activity. *Heliyon*, vol. 5, no. 4
- [27] L. Arum Utami., and D. Hilda Putri. (2020). The Effect of Ethanol Solvent Concentration on Antimicrobial Activities The Extract of Andalas Endophytic Bacteria (*Morus Macroura* Miq.) Fermentation Product. *EKSAKTA Berk. Ilm. Bid. MIPA*, vol. 21, no. 1, pp. 1–6
- [28] S. Luliana., H. Riza., and E. N. Indriyani. (2019). The Effect of Extraction Method on Total Phenolic Content and Antioxidant Activity of Salam Leaves (*Syzygium polyanthum*) using DPPH (1,1-Diphenyl-2-Picrylhidrazil). *Maj. Obat Tradis.*, vol. 24, no. 2, p. 72
- [29] W. Cai., H. Wang., Q. G. Yang., T. Liu., and Y. Wang. (2022). Extracting phenolic compounds from aqueous solutions by cyclohexanone, a highly efficient extractant. *J. Ind. Eng. Chem.*, vol. 116, pp. 393–399
- [30] N. L. Rahmah., S. M. M. Kamal., A. Sulaiman., F. S. Taip., and S. I. Siajam. (2023). Optimization Of Phenolic Compounds And Antioxidant Extraction From Piper Betle Linn. Leaves Using Pressurized Hot Water. *J. Appl. Sci. Eng.*, vol. 26, no. 2, pp. 175–184
- [31] G. Oshadie., D. Silva., A. T. Abeyundara., M. Minoli., and W. Aponso. (2017). Extraction methods, qualitative and quantitative techniques for screening of phytochemicals from plants. *Am. J. Essent. Oils Nat. Prod.*, vol. 5, no. 2, pp. 29–32
-

- [32] Panchal Mital, D., & Jha, C. V. (2021). Qualitative and quantitative phytochemical screening of three plants stem bark and leaves from sapotaceae family. *Int. J. Multidiscip. Educ. Res*, 6(8), 1-6.
- [33] I. Safitri., M. C. Nuria., and A. D. Puspitasari. (2018). Perbandingan Kadar Flavonoid Dan Fenolik Total Ekstrak Metanol Daun Beluntas (*Pluchea Indica L.*) Pada Berbagai Metode Ekstraksi. *Inov. Tek. Kim.*, vol. 3, no. 1, pp. 31–36
- [34] K. Anwar., A. Lutpi., A. Melinda., and S. Hadi. (2022). Extraction methods effect on antioxidant activity of ethanol extract of 'pasak bumi' (*Eurycoma longifolia Jack.*) root. *IOP Conf. Ser. Earth Environ. Sci.*, vol. 976, no. 1
- [35] M. Hait. (2019). Extraction techniques of herbal drugs. *Res. Trends Med. Plant Sci.*, no. pp. 17–33
- [36] Bagade, S. B., & Patil, M. (2021). Recent advances in microwave assisted extraction of bioactive compounds from complex herbal samples: A review. *Critical reviews in analytical chemistry*, 51(2), 138-149.
- [37] Y. Noviyanty., Hepiyansori., and T. . Insani. (2021). Uji Aktivitas Senyawa Flavonoid Dari Ekstrak Etanol Kulit Buah Mangga (*Mangifera Indica L.*) Terhadap Bakteri *Staphylococcus aureus*. *Ocean. Biomed. J.*, vol. 4, no. 1, pp. 38–52
- [38] Badshah, S. L., Faisal, S., Muhammad, A., Poulson, B. G., Emwas, A. H., & Jaremko, M. (2021). Antiviral activities of flavonoids. *Biomedicine & Pharmacotherapy*, 140, 111596.
- [39] V. Verawati., D. Nofiandi., and P. Petmawati. (2017). Pengaruh Metode Ekstraksi Terhadap Kadar Fenolat Total Dan Aktivitas Antioksidan Daun Salam (*Syzygium polyanthum (Wight) Walp.*). *J. Katalisator*, vol. 2, no. 2, pp. 53–60
- [40] Dirar, A. I., Alsaadi, D. H. M., Wada, M., Mohamed, M. A., Watanabe, T., & Devkota, H. P. (2019). Effects of extraction solvents on total phenolic and flavonoid contents and biological activities of extracts from Sudanese medicinal plants. *South African Journal of Botany*, 120, 261-267.
- [41] Oreopoulou, A., Tsimogiannis, D., & Oreopoulou, V. (2019). Extraction of polyphenols from aromatic and medicinal plants: an overview of the methods and the effect of extraction parameters. *Polyphenols in plants*, 243-259.
- [42] M. R. R. Rahardian., B. T. Murti., D. Wigati., R. Suharsanti., and C. N. Putri. (2019). Solvent concentration effect on total flavonoid and total phenolic contents of *Averrhoa bilimbi* leaf extract. *Pharmaciana*, vol. 9, no. 1, pp. 137–144
- [43] S. Settharaksa., A. Jongjareonrak., P. Hmadhlu., and W. Chansuwan. (2012). Flavonoid, phenolic contents and antioxidant properties of Thai hot curry paste extract and its ingredients as affected of pH, solvent types and high temperature. *nternational Food Res. J. 1*, vol. 19, no. 4, pp. 1581–1587
- [44] D. Wigati and D. K. Pratoko. (2019). Total Flavonoid dan Aktivitas Penangkapan Radikal Bebas dari Ekstrak Etanolik Daun Dan Buah Mengkudu. *J. Farm. (Journal Pharmacy)*, vol. 5, no. 1
- [45] D. Wigati., K. Anwar., Sudarsono., and A. E. Nugroho. (2017). Hypotensive Activity of Ethanollic Extracts of *Morinda citrifolia L.* Leaves and Fruit in Dexamethasone-Induced Hypertensive Rat. *J. Evidence-Based Complement. Altern. Med.*, vol. 22, no. 1, pp. 107–113,
- [46] A. L. Dela Cruz., M. A. M. Feliciano., D. S. Paragas., J. A. Detablan., and P. W. Tsai. (2023). Isolation, Characterization, and Antioxidant Activity of *Selliguea taeniata* Secondary Metabolites. *Biointerface Res. Appl. Chem.*, vol. 13, no. 4