

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



ISSN: 0015-5659

e-ISSN: 1644-3284

Branching pattern of the internal iliac artery accompanied by a venous anastomosis: rare vascular variations

Authors: Maciej Kula, Łukasz Olewnik, Kacper Ruzik, Richard Shane Tubbs, Adrian Balcerzak, Nicol Zielinska

DOI: 10.5603/FM.a2022.0111

Article type: Case report

Submitted: 2022-07-10

Accepted: 2022-08-21

Published online: 2022-12-23

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Articles in "Folia Morphologica" are listed in PubMed.

Branching pattern of the internal iliac artery accompanied by a venous anastomosis: rare vascular variations

Maciej Kula et al., Internal iliac artery

Maciej Kula¹, Łukasz Olewnik¹, Kacper Ruzik¹, Rihard Shane Tubbs²⁻⁷, Adrian Balcerzak¹, Nicol Zielinska¹

¹Department of Anatomical Dissection and Donation, Medical University of Lodz, Poland

²Department of Anatomical Sciences, St. George's University, Grenada

³Department of Neurosurgery, Tulane University School of Medicine, New Orleans, Louisiana, United States

⁴Department of Neurology, Tulane University School of Medicine, New Orleans, Louisiana, United States

⁵Department of Structural and Cellular Biology, Tulane University School of Medicine, New Orleans, Louisiana, United States

⁶Department of Surgery, Tulane University School of Medicine, New Orleans, Louisiana, United States

⁷Department of Neurosurgery, Ochsner Medical Center, New Orleans, Louisiana, United States

Address for correspondence: Łukasz Olewnik, MD, PhD, Department of Normal and Clinical Anatomy, Interfaculty Chair of Anatomy and Histology, Medical University of Lodz, Poland, e-mail: lukasz.olewnik@umed.lodz.pl

ABSTRACT

The ability to navigate the complex and often deceptive branching patterns of the internal iliac artery can be decisive in planning and performing surgeries within the lesser pelvis. The following case report presents a peculiar quadruple division of the internal iliac artery, accompanied by a venous anastomotic structure. Apart from the posterior and anterior trunks, the superior vesicle and iliolumbar arteries arose independently from the internal iliac artery. The division was surrounded by a venous oval, compressing certain branches and potentially complicating surgical access. Due to the uncommon course of the internal iliac artery and the

presence of the anastomosis, a possible nerve root compression has been identified. Both clinical significance and classification method of the case are discussed. Knowledge of this anatomical variation is valuable for both diagnosis and surgery, especially within the specialties of urology, gynecology and general surgery.

Key words: anatomical variations, internal iliac artery, case report

INTRODUCTION

The internal iliac artery (IIA), also known as the hypogastric artery, is the main blood vessel supplying the walls and organs of the pelvis, the reproductive organs, buttocks, muscles of the lumbar region and the medial section of the thigh. About 3-4 cm long, it is significantly smaller than the external iliac artery[1]. The IIA arises at the bifurcation of the common iliac artery, anterior to the pelvic brim, at the level of the L5-S1 intervertebral disc[2]. It descends posteriorly to the upper margin of the sciatic foramen, where it divides into two branches, the posterior and anterior trunks.

The posterior trunk bends posteriorly and passes towards the greater sciatic foramen, exiting the pelvis above the piriformis muscle. It most commonly gives off three parietal branches, the iliolumbar, lateral sacral and superior gluteal arteries, the last-named being its termination. The anterior trunk passes proximally to the piriformis muscle and the sacral plexus until it reaches the lower section of the sciatic foramen. It gives rise to six visceral branches: the umbilical, superior vesical, inferior vesical and middle rectal arteries and, in females, the vaginal and uterine arteries. It also gives off two parietal branches, the obturator and inferior gluteal arteries, the latter being its termination.

The following case study describes an unusual branching pattern of the IIA. Instead of giving rise to two main trunks that further divided into its terminations, the IIA divided into four lower caliber branches, the iliolumbar and superior vesical arteries arised directly from its main trunk. The division was encircled by a venous anastomotic structure derived from the internal iliac vein. The anastomosis passed between the posterior and anterior trunks of the IIA, surrounding the artery's division and coming back beneath the iliolumbar artery before ending its course, reaching the external iliac artery.

Understanding the elusive branching pattern of the IIA is invaluable for clinicians, especially gynecologists[3]. This knowledge is likely to prove essential during surgical

procedures in which a certain part of the artery needs to be ligated in order to control pelvic hemorrhage.

CASE REPORT

A female cadaver 75 years old at death was subjected to routine anatomical dissection for research and educational purposes at the Department of Anatomical Dissection and Donation, Medical University of Lodz. During a routine assessment of the lumbosacral plexus, an anomaly of IIA branching was identified (Fig. 1).

The lengths and widths of the branches were measured using a digital caliper. A detailed analysis of the branching pattern revealed an unusual division, which has not previously been described to our knowledge. Additionally, a venous anastomotic structure surrounding the division was exposed (Fig. 2).

The IIA begins its course at the bifurcation of the common iliac artery. It is 38.6 mm long and its diameter is 7.21 mm. After branching out of the common iliac artery, it descends medially alongside the posterior pelvic wall towards the external iliac vein (EIV). Upon passing the EIV it descends towards the pelvic floor for approximately 16 mm before giving rise to its five branches: the iliolumbar artery (ILA), the superior lateral sacral artery (SLSA), the posterior trunk (PT), the anterior trunk (AT), and the superior vesical artery (SVA) from which the medial umbilical ligament (MUL) arises. Immediately after branching from the IIA, the ILA (7.94 mm) bends upwards and runs on top of the anastomosis of the internal iliac vein, then passes beneath the EIV and divides into its terminations, the iliac and lumbar branches. The uterine artery (UtA) begins at the anterior trunk 3.12 mm below the origin of the umbilico-vesical trunk and descends towards the uterus. The internal pudendal artery (IPA) and the middle rectal artery (MRA) share a short (6.09 mm) common trunk, which begins at the anterior trunk 20 mm below the beginning of the UtA and almost immediately bifurcates, creating points of origin for the aforementioned vessels. After giving off the IPA and the MRA, the anterior trunk passes anteriorly for 18 mm and then bends posteriorly, exiting the lesser pelvic cavity between the L5 and S1 spinal nerves and below the piriformis muscle to become the inferior gluteal artery (IGA). Further dissection and assessment of the IGA ruled out the presence of a sciatic artery. The posterior trunk arises from the IIA below the anterior trunk and descends anteriorly. Before exiting the pelvic cavity above the piriformis muscle and becoming the superior gluteal artery, it gives off the inferior lateral sacral artery (ILSA) 7 mm below the SLSA, and the obturator artery (OA), which arises from the PT directly opposite the ILSA. Further dissection and analysis of the OA and the inferior

epigastric artery ruled out the presence of a “corona mortis” anastomosis[4]. The posterior trunk is 4.43 mm in diameter.

The venous anastomotic structure originates from the internal iliac vein and is oval. The anastomosis is 9.48 mm wide and 19.76 mm long and encircles the division of the IIA. The posterior trunk of the IIA passes through the venous circle and descends towards the pelvic floor, whereas the anterior trunk runs anteriorly, resting on top of the anastomosis, pressing it down against the posterior trunk. Applied pressure forces the PT down, causing it to rest on the L5 root of the sciatic nerve.

Table 1. Measurements of the structures of interest

Structure	Length (mm)	Width (mm)
Internal iliac artery	38.6	7.21
Anterior trunk	41.97	3.97
Posterior trunk	22	4.43
Iliolumbar artery	7.94	3.12
Uterine artery	84.41	2.22
Umbilico-vesical trunk	27	2
Umbilical ligament	90	1.34
Vesical artery	79.4	1.47
Obturator artery	47.3	1.67
Middle rectal artery	29	1.22
Internal pudendal artery	19.4	1.25
Internal iliac vein anastomosis	19.76	9.48

DISCUSSION

Since the pelvic vascularization is intricate, it is essential for a wide range of specialists such as gynecologists, urologists, proctologists and general surgeons to understand the possible anatomical variations of the IIA to maximize the efficiency of certain procedures[5-8]. Especially in laparoscopic surgery, there is a huge demand for precise descriptions of pelvic vascularization to help preclude iatrogenic injuries[9-11].

The branching pattern described here resembles type III in the Balcerzak classification[12] because three main branches (AT, PT, ILA) exit the common trunk. However, certain nuances such as a superior vesical artery arising directly from the IIA instead of the anterior trunk, and two sacral lateral arteries, distinguish it significantly since none of them are considered in Balcerzak’s system. The artery described in the present article has a further uncommon anatomical variation: the obturator artery branches out of the posterior division of the IIA. Pai et al. estimated the prevalence of a similar variant at around

8-10%[13]. This has some clinical value: during procedures such as embolization or ligation of the OA, it is vital for the operator to be aware of the possible points of origins of the vessel. Such knowledge can reduce the duration of the procedure and thereby minimize potential complications[14].

Since the IIA provides the main blood supply to the uterus, it is also crucial in the development of a range of uterine pathologies such as adenomyosis[15] and uterine fibroids[16]. Both those conditions are exacerbated by an excessive blood supply from the uterine artery, which is why uterine artery embolization (UAE) provides effective treatment. During this procedure, a catheter is inserted through the femoral or radial artery into the IIA and then into the UtA. It is therefore important for the surgeon to have a good understanding of the highly variable branching pattern of IIA in order to avoid dire complications. For instance, in the case presented here, the procedure could prove difficult because of the close proximity (3.12 mm) of the UtA to the superior vesical artery and the difference in vessel diameters: UtA: 1.72 mm, SVA: 2.2 mm. This could lead to accidental insertion of the catheter into the SVA; or puncture of the anterior trunk, inevitably causing profuse hemorrhage.

Because the IIA follows a sinuous course and is close to major nerves, anatomical variations often correlate directly with compression syndromes. Conditions such as aneurysms involving a drastic increase of the artery's diameter are the most common causes of such pathologies. In a recent case study, de Bruijn et al. described a type IA (Balcerzak, 2021) IIA containing an aneurysm located between the L5 and S1 nerve roots[17]. The location of the aneurysm combined with the artery's branching pattern elicited buttock ischemia and neurological symptoms such as impairment of active knee and plantar foot flexion, absent dorsal flexion, and sensitivity disorders below the knee, all caused by nerve root compression[17]. In the case presented here, owing to the venous anastomosis, the posterior trunk of the IIA applied considerable pressure on the L5 nerve, causing it to deform. It is unclear whether this caused any discomfort during life, but it certainly aggravated or increased the risk of closely-related conditions such as wallet neuritis, lotus neuropathy or piriformis syndrome.

Rupture of the IIA and its branches is a rare but potentially lethal complication that can occur during pregnancy[18, 19]. In a case report describing three instances of utero-ovarian ruptures, Ginsburg et al. estimate the initial overall mortality at up to 49%. However, owing to the rapid development of intensive intraoperative and postoperative treatments, a decline to 3.6% has been noted[20].

Table 2. The 2021 Balcerzak classification. “Main vessels” include the superior gluteal, inferior gluteal, internal pudendal and sciatic arteries (in group V).

TYPE	DESCRIPTION
Group I	Main vessels have two points of origin: the first is a single branch, the second a common trunk
Subgroup A: Common trunk divides inside the pelvic cavity	
Subgroup B: Common trunk divides outside the pelvic cavity	
Group II	Main vessels have two points of origin: the first is a common trunk, the second a single branch
Subgroup A: Common trunk divides inside the pelvic cavity	
Subgroup B: Common trunk divides outside the pelvic cavity	
Group III	Main vessels have three points of origin; order of origin is irrelevant
Group IV	Main vessels have one point of origin; order of origin is irrelevant
Group V	Main vessels have four points of origin; additionally includes the sciatic artery

CONCLUSIONS

In conclusion, variations of the IIA are common and it is nearly impossible to classify them comprehensively. This underlines the importance of supplementing existing classifications with case studies like the present one. Preoperative knowledge of anatomical variants and their relationships to neighboring structures is crucial in planning and performing surgical procedures on the IIA and in its vicinity.

Declarations

Ethical approval and consent to participate: The cadavers belonged to the Department of Anatomical Dissection and Donation, Medical University of Lodz.

Consent to publish: Not applicable

Competing interests: The authors declare that they have no competing interests.

Funding: The authors have no financial or personal relationship with any third party whose interests could be positively or negatively influenced by the article’s content. This research

did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

REFERENCES

1. Fătu C, Puișoru M, Fătu IC (2006) Morphometry of the internal iliac artery in different ethnic groups. *Annals of Anatomy = Anatomischer Anzeiger: Official Organ of the Anatomische Gesellschaft* 188:541–546. <https://doi.org/10.1016/J.AANAT.2006.05.016>
2. Sakthivelavan S, Aristotle S, Sivanandan A, et al (2014) Variability in the Branching Pattern of the Internal Iliac Artery in Indian Population and Its Clinical Importance. *Anatomy Research International* 2014:1–6. <https://doi.org/10.1155/2014/597103>
3. Giordano G, Meo D, Magnano San Lio V. Internal iliac artery aneurysm embolization with direct percutaneous puncture and thrombin injection. *Radiology Case Reports*. 2020;15(3):210-213. doi:10.1016/j.radcr.2019.11.018
4. Boutefnouchet T, Bassett J, Patil S. Anatomy and Clinical Relevance of the ‘Corona Mortis’: A Review of the Literature and Current Aspects of Management. *J Orthopedics Rheumatol*. 2016; 3(2):5.
5. Kemmochi R, Matsumoto M, Kubo Y, Yamasawa T (2013) Hypogastric and lumbar artery bypasses to restore gluteal perfusion after open aortoiliac aneurysm repair. *Journal of Vascular Surgery* 58:1073–1075. <https://doi.org/10.1016/j.jvs.2012.12.079>
6. O’Brien MC, Schell BA, Lands H, et al (2018) Massive Gluteal Muscle Necrosis after Iliac Arterial Embolization in Pelvic Trauma: A Literature Review and Illustrative Case Report. *Journal of orthopaedic case reports* 8:23–27. <https://doi.org/10.13107/jocr.2250-0685.1092>
7. Chayen D, Copeliovitch L, Itzhakov Z, Zaretsky M, Rabin I. An original external iliac artery reconstruction with internal iliac artery translocation in a blunt injury of the pelvic vessels in a 4-year-old child: A 12-year follow-up study. *Journal of Vascular Surgery Cases and Innovative Techniques*. 2019;5(4):492-496. doi:10.1016/j.jvscit.2019.09.003
8. Das BN, Biswas AK. Ligation of internal iliac arteries in pelvic haemorrhage. *J Obstet Gynaecol Res*. 1998 Aug;24(4):251-4. doi: 10.1111/j.1447-0756.1998.tb00085.x. PMID: 9798353.
9. Lu Gao J, Lortie K, Singh SS (2010) Laparoscopic Internal Iliac Artery Ligation for Postpartum Spontaneous Hemoperitoneum. *Journal of Obstetrics and Gynaecology Canada* 32:1172–1175. [https://doi.org/10.1016/S1701-2163\(16\)34742-9](https://doi.org/10.1016/S1701-2163(16)34742-9)
10. Jeon SH, Lee SH, Choi WC (2007) Iliac artery perforation following lumbar discectomy with microsurgical carbon dioxide laser: A report of a rare case and discussion on the treatment. *Spine* 32:. <https://doi.org/10.1097/01.BRS.0000254078.88358.33>
11. Siedhoff M, Gubernick L, Ronen I, Wright K. Internal Iliac Artery Ligation in Laparoscopic Myomectomy. *Journal of Minimally Invasive Gynecology*. 2019;26(7):S1. doi:10.1016/J.JMIG.2019.09.007
12. Balcerzak A, Hajdys J, Shane Tubbs R, Karauda P, Georgiev GP, Olewnik Ł. Clinical importance of variability in the branching pattern of the internal iliac artery - An updated and comprehensive review with a new classification proposal. *Ann Anat*. 2022;239:151837. doi:10.1016/j.aanat.2021.151837
13. Pai MM, Krishnamurthy A, Prabhu L v., et al (2009) Variability in the origin of the obturator artery. *Clinics* 64:897–901. <https://doi.org/10.1590/S1807-59322009000900011>
14. Wojciechowski J, Znaniński L, Brzeziński M (2005) Obturator Artery Aneurysm. *EJVES Extra* 9:129–130. <https://doi.org/10.1016/j.ejvsextra.2005.04.002>
15. Sharara FI, Kheil MH, Feki A, et al (2021) Current and Prospective Treatment of Adenomyosis. *Journal of Clinical Medicine* 10:10. <https://doi.org/10.3390/JCM10153410>
16. Levy BS (2008) Modern management of uterine fibroids. *Acta Obstetrica et Gynecologica Scandinavica* 87:812–823. <https://doi.org/10.1080/00016340802146912>

17. de Bruijn MT, Verbelen T, Kralt CP, van Baal JG (2015) An internal iliac artery aneurysm causing sudden buttock ischemia and nerve root compression. *Journal of Vascular Surgery Cases* 1:151–153. <https://doi.org/10.1016/j.jvsc.2015.04.015>
18. Cole M, Elton C, Bosio P, Waugh JJS (2005) Spontaneous rupture of utero-ovarian vessels in labour - A rare case of obstetric haemoperitoneum. *Journal of Obstetrics and Gynaecology* 25:301–303. <https://doi.org/10.1111/j.1440-1711.2005.01324.x>
19. Vellekoop J, de Leeuw JP, Neijenhuis PA (2001) Spontaneous rupture of a utero-ovarian vein during pregnancy. *American journal of obstetrics and gynecology* 184:241–2. <https://doi.org/10.1067/mob.2001.110310>
20. Ginsburg KA, Valdes C, Schnider G (1987) Spontaneous utero-ovarian vessel rupture during pregnancy: Three case reports and a review of the literature. *Obstetrics and Gynecology* 69:474–47618.

Figure 1. Branching point of a left internal iliac artery seen from the medial perspective.

Section of a left internal iliac vein has been resected for better exposure. CIA – common iliac artery EIA – external iliac artery, IIA – internal iliac artery, CIV – common iliac vein, EIV – external iliac vein, IIV – internal iliac vein, AT – anterior trunk of internal iliac artery, PT – posterior trunk of internal iliac artery, ILA – iliolumbar artery, OA – obturator artery, UtA – uterine artery, SVA – superior vesical artery, MRA – medial rectal artery, IPA – interior pudendal artery UVT – umbilico-vesical trunk SLSA – superior lateral sacral artery, ILSA – inferior lateral sacral artery, ON – obturator nerve MUL – median umbilical ligament. A – point of anastomosis of the EIV (beginning of the venous circle).

Figure 2. Branching of the internal iliac artery surrounded by a venous anastomosis originating from the internal iliac vein. Top-down perspective. CIA – common iliac artery EIA – external iliac artery, IIA – internal iliac artery, CIV – common iliac vein, EIV – external iliac vein, IIV – internal iliac vein, AT – anterior trunk of internal iliac artery, ILA – iliolumbar artery, OA – obturator artery, UtA – uterine artery, MRA – medial rectal artery, IPA – interior pudendal artery UVT – umbilico-vesical trunk SLSA – superior lateral sacral artery, ILSA – inferior lateral sacral artery, ON – obturator nerve.



