



2022

Deconstructing Stereotypes: Stature, Match-playing Time, and Performance in Elite Women's World Cup Soccer

Ciara N. Manning
University of Connecticut

Yasuki Sekiguchi
Texas Tech University

Courteney L. Benjamin
Samford University

McKenna R. Spaulding
Dickinson College

Erin E. Dierickx
University of Connecticut

See next page for additional authors

Follow this and additional works at: https://digitalcommons.sacredheart.edu/pthms_honors



Part of the [Rehabilitation and Therapy Commons](#)

Recommended Citation

Manning, C. N., Sekiguchi, Y., Benjamin, C. L., Spaulding, M. R., Dierickx, E. E., Spaulding, J. M., Davenport, D. M., Picard-Busky, J. R., Chiampas, G. T., Nassis, G. P., & Casa, D. J. (2022). Deconstructing stereotypes: Stature, match-playing time, and performance in elite women's world cup soccer. *Frontiers in Sports and Active Living*, 4, 1067190. Doi.org/10.3389/fspor.2022.1067190

This Thesis is brought to you for free and open access by the Physical Therapy & Human Movement Science at DigitalCommons@SHU. It has been accepted for inclusion in PTHMS Undergraduate Publications by an authorized administrator of DigitalCommons@SHU. For more information, please contact lysobeyb@sacredheart.edu.

Author

Ciara N. Manning, Yasuki Sekiguchi, Courteney L. Benjamin, McKenna R. Spaulding, Erin E. Dierickx, Jayson M. Spaulding, Dayshia M. Davenport, Jillian R. Picard-Busky, George T. Chiampas, George P. Nassis, and Douglas J. Casa



OPEN ACCESS

EDITED BY

Jason Vescovi,
University of Toronto, Canada

REVIEWED BY

Antonio Tessitore,
Foro Italico University of Rome, Italy
Serdar Bayrakdaroglu,
Gumushane University, Turkey

*CORRESPONDENCE

C. N. Manning
ciara.manning@uconn.edu

SPECIALTY SECTION

This article was submitted to Elite Sports and Performance Enhancement, a section of the journal Frontiers in Sports and Active Living

RECEIVED 11 October 2022

ACCEPTED 23 November 2022

PUBLISHED 14 December 2022

CITATION

Manning CN, Sekiguchi Y, Benjamin CL, Spaulding MR, Dierickx EE, Spaulding JM, Davenport DM, Picard-Busky JR, Chiampas GT, Nassis GP and Casa DJ (2022) Deconstructing stereotypes: Stature, match-playing time, and performance in elite Women's World Cup soccer.

Front. Sports Act. Living 4:1067190.
doi: 10.3389/fspor.2022.1067190

COPYRIGHT

© 2022 Manning, Sekiguchi, Benjamin, Spaulding, Dierickx, Spaulding, Davenport, Picard-Busky, Chiampas, Nassis and Casa. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Deconstructing stereotypes: Stature, match-playing time, and performance in elite Women's World Cup soccer

Ciara N. Manning^{1*}, Yasuki Sekiguchi^{1,2},
Courtney L. Benjamin^{1,3}, McKenna R. Spaulding⁴,
Erin E. Dierickx¹, Jayson M. Spaulding⁵, Dayshia M. Davenport⁶,
Jillian R. Picard-Busky⁷, George T. Chiampas^{8,9},
George P. Nassis^{10,11} and Douglas J. Casa¹

¹Department of Kinesiology, Korey Stringer Institute, University of Connecticut, Storrs, CT, United States, ²Sports Performance Lab, Department of Kinesiology and Sport Management, Texas Tech University, Lubbock, TX, United States, ³Department of Kinesiology, Samford University, Birmingham, AL, United States, ⁴Department of Health and Physical Education and Recreation, Dickinson College, Carlisle, PA, United States, ⁵Department of Exercise Science and Athletic Training, Ithaca College, Ithaca, NY, United States, ⁶School of Kinesiology, Louisiana State University, Baton Rouge, LA, United States, ⁷Department of Exercise Science, Sacred Heart University, Fairfield, CT, United States, ⁸Department of Emergency Medicine, Northwestern University, Chicago, IL, United States, ⁹Chief Medical Officer, United States Soccer Federation, Chicago, IL, United States, ¹⁰Physical Education Department, College of Education, United Arab Emirates University, Al Ain, UAE, ¹¹Department of Sports Science and Clinical Biomechanics, SDU Sport and Health Sciences Cluster (SHSC), University of Southern Denmark, Odense, Denmark

Recruiting companies recommend elite female soccer players be ≥ 165 cm (5'5") in stature. This study investigated if stature limits match-playing time and performance in elite World Cup soccer among players, positions, and countries. We hypothesized stature would not affect match-playing time or performance. Descriptive data were collected on 552 players from 2019 FIFA Women's World Cup. Odds ratios determined likelihood of starting for players < 165 cm and ≥ 165 cm. ANOVAs compared playing time between stature groups, among positions, and between countries. Performance factors including assists, goals, attempts, corners, shots blocked, and defending blocks were reported. Independent t-tests compared differences between players (≥ 165 cm, < 165 cm). Data are reported, mean difference [95% confidence interval] [MD (95%CI)] and effect sizes (ES). On average, 32.3% of players were < 165 cm. Of total players, no differences existed in total minutes ($F = 0.98$, $p = 0.32$), matches ($F = 0.27$, $p = 0.59$), or average minutes per match ($F = 0.48$, $p = 0.49$) between stature groups, regardless of position. No differences existed in playing time between players < 165 cm and ≥ 165 cm among any positions ($p > 0.05$), or between countries ($p > 0.05$). Taller mid-fielders exhibited greater performance in goals, assists, attempts, shots blocked, and defending blocks (MD [95%CI] ES; assists, $-0.44[-0.76, -0.11]0.59$, $p = 0.009$; goals, $-0.35[-0.69, -0.01]0.44$, $p = 0.047$); attempts, $3.14[1.38, 4.90]0.80$, $p = 0.001$; corners, $2.04[0.12, 3.95]0.48$, $p = 0.037$; shots blocked, $0.96[0.40, 1.51]0.75$, $p = 0.001$; defending blocks, $0.43[0.32, 0.82]0.48$, $p = 0.035$), however, actual differences were minimal. Our findings indicate stature does not inhibit playing and performing elite women's soccer, as nearly one-third of players were < 165 cm.

KEYWORDS

women's soccer, height, performance, FIFA world cup, playing time

Introduction

The International Federation of Association Football (FIFA), the largest governing body in the sport, organizes the FIFA World Cup tournaments for qualifying teams all over the world to compete in every four years (1). Many youth athletes may aspire to play at this elite level of soccer. The number of high-school soccer players in the United States has increased 10.7% for females and 17.2% for males in the past decade (1). A small number of these high-school athletes make it to the collegiate and professional level (1, 2). Of the 390,482 female high-school soccer players reported in 2017–2018, only 27,811 (7.1%) continued playing at the National Collegiate Athletic Association (NCAA) level (2). The estimated probability of performing at the collegiate level after high-school in 2019 was 2.4% for Division I, 1.9% for Division II, and 2.8% for Division III (2). This low probability raises the question of what it takes to reach collegiate and elite level success, and what coaches and colleges look for in potential athletes.

College recruiting companies list minimal stature guidelines of ≥ 165 cm for female high-school athletes who aspire to play at the NCAA level (3). Provisions also include that Division I players be of the following statures: 170 cm–188 cm (5'7"–6'2") for goalkeepers, ≥ 168 cm (5'6"+) for outside defenders, 170 cm–188 cm (5'7"–6'2") for center defenders, and ≥ 165 cm (5'5"+) for outside mid/wing forwards (3). No stature is listed for center midfielders or forwards but can be implied based on requirements for other positions (3). These guidelines set the basis and understanding of what to strive for, for both athletes and coaches. Stating such guidelines and creating stereotypes for soccer players may discourage talented athletes from pursuing soccer at the elite level, strictly on the basis of their stature (4). However, it's not currently known if stature impacts match-playing time or performance in elite level soccer.

Generally, it is a common assumption that players of taller stature are more likely to succeed in elite-level soccer (5). However, in contradiction, some studies have found sub-elite soccer players to be of greater stature than their elite counterparts, suggesting that stature may not always characterize elite status or performance (6). Sports performance research has yet to fully define main attributes and actions that differentiate successful soccer players from unsuccessful soccer players (7–13). Limited research has been done assessing what constitutes match-playing time in soccer and why some players receive more playing time than others.

Many researchers have tried to determine what results in athletes being selected to play at an elite level, and what should constitute the selection process to ensure that selection is based on objective rather than subjective measures (14–17). In a recent systematic review, researchers operationalized multiple criterion variables that exist in soccer to determine their level of application in the actual playing soccer (18). Of

the criterion variables, stature was indicated as having little overlap in the actual playing of soccer (18). Regardless of its low applicability in relation to match-play, stature is commonly used by recruiting companies and coaches to identify talented soccer players (3, 18).

Anatomical and biomechanical differences have been considered partially influential in soccer performance in regards to the skill of kicking and certain running motion patterns (19). Longer limbs have the ability to produce more force, torque, and can produce a greater change in momentum.(19–24) This is believed to relay to soccer performance in kicking, jumping, and running (19). The anthropometric profile of soccer players has also been found to differ between positions suggesting that physical characteristics vary by playing position (17, 25–27). Certain statures and ranges of body mass have been deemed beneficial to players of different positions in elite-level soccer (25, 28). Anthropometric factors including stature, body mass, % body fat, and sum of skinfolds tend to decrease among positions ranging from goal keepers with the highest values, followed by defenders, forwards, and midfielders having the lowest values (25). Previous research suggests that heavier and taller players had greater vertical jump and 30-m sprinting abilities required by positions of goal keepers, defenders, and forwards (25, 28). This research also reported that players of lower BMI had greater aerobic endurance required of external defenders and central and external midfielders (25, 28).

No research has been done on the duration of match-playing time in relation to one's stature. While recruiting companies list minimal stature provisions of ≥ 165 cm (5'5") for aspiring female soccer players, it is not currently known whether or not stature impacts the duration of match-playing time in soccer at the elite World Cup level. Therefore, the primary aim of this study is to determine if stature is a limitation to the duration of match-playing time on an elite World Cup soccer team. We hypothesized that stature (≥ 165 cm or < 165 cm) would have no effect in the duration of match-playing time among players. Additionally, no research has been done assessing the impact of stature on performance in elite women's World Cup soccer. Therefore, the secondary aim of this study is to determine if differences in performance exist among elite female soccer players of different stature (≥ 165 cm and < 165 cm) among all positions in the 2019 FIFA Women's World Cup. We hypothesized that no performance differences would exist among players of different stature within all positions.

Materials and methods

Participants

This study utilized an observational, cross-sectional design. Data were obtained from the official website of FIFA World Cup

and the records of the 2019 Women's World Cup (29). Statistics from 552 female soccer players from 24 teams who played in the 2019 FIFA World Cup games were reported on the website and manually recorded in an Excel sheet for this analysis. Variables obtained from the website included player's name, age, stature, position, minutes played, matches played, and performance factors (29). Stature was self-reported by each team. On average players were 27.14 ± 3.96 years old. Players were categorized as those ≥ 165 cm (5'5") and those < 165 cm (5'5"). Since the collection of data for this manuscript, the website has been updated and no longer reports individual demographics of individual players, but rather general team statistics (29). Physical analysis of player demographics can be found in the Physical Analysis of the FIFA Women's World Cup 2019 report and on the U.S. soccer website for the U.S. team (30, 31).

Likelihood of playing

All 552 female soccer players were included to assess the amount of time spent in match-play in relation to stature. Descriptive data on matches played and minutes played for each player were analysed. The data were categorized into those below and those at or above the 165-cm stature criterion. Odds ratios were calculated to determine if differences existed in the likelihood of being a starter for those players on the team < 165 cm tall and those on the team ≥ 165 cm tall. Categorizing players as starters or non-starters allowed for identification and comparison of players who typically receive the longest duration of match-play time with players who receive less match-play time. Researchers have used various criteria when classifying players as starters and non-starters with some researchers classifying starters as those who play for 83% of total game time and non-starters as

those who play for 17% of total game time (32). Other researchers have classified starters as those who regularly participate in practices (33), or those who play $> 60\%$ of games (34). Additionally, researchers have considered starters as those playing for 45 min of total game time and non-starters as those who play less (35, 36). In our analysis, players were considered starters if they played ≥ 60 min when entered into a match. We believe this is an appropriate classification as this would equate to players who played a majority (roughly 67%) of the duration of a match. This was defined as a significant amount of playing time. A one-way ANOVA was used to examine if stature impacted the number of minutes played, number of matches played, average minutes played when entered into a match, average minutes played in relation to the total number of matches the team played during the entire 2019 FIFA Women's World Cup, and percent playing time between all players < 165 cm and ≥ 165 cm (Table 1). Normality was not necessary for these analyses given the large sample size (37).

Match-playing time

For each player, average minutes played per match, average minutes played in relation to the total number of matches their team played in during the World Cup, and percentage of playing time were manually computed (29, 38). Average minutes played per match was calculated by dividing the total number of minutes an individual played by the number of matches that the individual played (9, 38, 39). This was calculated to assess how much playing time, if given the opportunity by being entered into a match, individuals received. Average minutes played in relation to the total number of matches the team played in during World Cup was calculated by dividing the total

TABLE 1 Playing time and performance measures for all players ≥ 165 cm and < 165 cm.

Playing Time Measures	≥ 165 cm	< 165 cm	Statistics
Total minutes played	195.10 ± 185.66	179.46 ± 146.99	$F = 0.98, p = 0.32, ES = 0.09$
Number of matches played	2.65 ± 2.08	2.56 ± 1.66	$F = 0.27, p = 0.59, ES = 0.05$
Average minutes played per match	52.73 ± 37.64	55.04 ± 34.45	$F = 0.48, p = 0.49, ES = 0.05$
Average minutes played in relation to total minutes team played in	42.26 ± 36.50	45.12 ± 34.14	$F = 0.78, p = 0.38, ES = 0.08$
Percentage of playing time	$46.95\% \pm 40.56$	$50.14\% \pm 37.93$	$F = 0.78, p = 0.38, ES = 0.06$
Performance Measures			
Goals (number)	$0.57 \pm 1.17^*$	0.25 ± 0.62	$F = 16.98, p = 0.01, MD = -0.32, 95\% CI [-0.58, -0.07]$
Assists (number)	0.32 ± 0.69	0.22 ± 0.55	$F = 4.71, p = 0.25, MD = -0.10, 95\% CI [-0.26, 0.07]$
Attempts (number)	$4.55 \pm 5.27^*$	3.22 ± 3.56	$F = 9.57, p = 0.03, MD = -1.33, 95\% CI [-2.52, -0.13]$
Corners (number)	1.79 ± 4.53	1.18 ± 3.07	$F = 6.00, p = 0.25, MD = -0.61, 95\% CI [-1.64, 0.42]$
Distance covered per match (km)	9.13 ± 1.27	8.97 ± 1.84	$F = 5.32, p = 0.41, MD = -0.16, 95\% CI [-0.54, 0.22]$

Data are presented as averages \pm standard deviation, F -value, p -value, mean difference, and 95% confidence interval.

*Indicates statistically significant differences ($p < 0.05$) between stature groups.

number of minutes an individual played by the number of matches that the team played in during the World Cup (9, 29, 38, 39). This was calculated to analyse playing time on the basis of team opportunity rather than on an individual level. Percentage of playing time was calculated by dividing the number of minutes an individual played by the total number of minutes their team played during the World Cup and multiplying by 100. This was calculated as another means to compare playing time between height groups.

Positions and playing time

Players were separated by position to determine how many players in each position were <165 cm and ≥165 cm

tall. Four positions were included in this analysis: forwards, defenders, midfielders, and goal keepers. Ratios were calculated to determine percentages of players in each position and each stature group. A multivariate ANOVA was used to examine differences in stature, position, and stature among positions (Table 2). This determined if stature impacted the number of minutes played, number of matches played, average minutes individuals played when entered into a match, average minutes individuals played in relation to the total number of matches their team played in during the entire 2019 FIFA Women's World Cup, and percentage of playing time between players <165 cm and ≥165 cm tall, and within each position. LSD *post hoc* comparisons were used when appropriate to determine where specific differences occurred.

TABLE 2 Playing time for defenders, forwards, and midfielders ≥165 cm and <165 cm.

Position	≥165 cm	<165 cm	Statistics
Defenders			
Minutes played	220.73 ± 195.08	179.78 ± 149.98	MD = -45.78, 95% CI = [-99.65, 8.08], ES = 0.25, <i>p</i> = 0.10, <i>t</i> = -1.68,
Matches played	2.73 ± 2.06	2.57 ± 1.56	MD = -0.53, 95% CI = [-1.16, 0.95], ES = 0.25, <i>p</i> = 0.10, <i>t</i> = -1.67
Average minutes per match	60.24 ± 37.95	56.79 ± 33.58	MD = -5.25, 95% CI = [-17.55, 7.05], ES = 0.14, <i>p</i> = 0.40, <i>t</i> = -0.84
Average minutes per total matches team played	47.49 ± 37.84	46.42 ± 34.60	MD = -2.93, 95% CI = [-14.98, 9.12], ES = 0.08, <i>p</i> = 0.63, <i>t</i> = -0.48
Forwards			
Minutes played	206.35 ± 156.57	188.24 ± 38.47	MD = -18.11, 95% CI = [-71.97, 35.75], ES = 0.12, <i>p</i> = 0.51, <i>t</i> = -0.67
Matches played	3.18 ± 1.86	3.04 ± 1.65	MD = -0.14, 95% CI = [-0.78, 0.50], ES = 0.08, <i>p</i> = 0.66, <i>t</i> = -0.44
Average minutes per match	55.53 ± 30.65	52.62 ± 28.21	MD = -2.90, 95% CI = [-13.57, 7.76], ES = 0.10, <i>p</i> = 0.59, <i>t</i> = -0.54
Average minutes per total matches team played	45.74 ± 31.39	44.63 ± 29.46	MD = -1.11, 95% CI = [-12.10, 9.88], ES = 0.04, <i>p</i> = 0.84, <i>t</i> = -0.20
Midfielders			
Minutes played	194.69 ± 186.69	179.78 ± 149.98	MD = -14.92, 95% CI = [-67.12, 37.27], ES = 0.20, <i>p</i> = 0.57, <i>t</i> = -0.57
Matches played	2.93 ± 2.02	2.57 ± 1.56	MD = -0.36, 95% CI = [-0.92, 0.19], ES = 0.09, <i>p</i> = 0.20, <i>t</i> = -1.29
Average minutes per match	51.11 ± 34.06	56.79 ± 33.58	MD = 5.68, 95% CI = [-4.87, 16.23], ES = 0.17, <i>p</i> = 0.29, <i>t</i> = 1.06
Average minutes per total matches team played	40.27 ± 34.72	46.42 ± 34.60	MD = 6.14, 95% CI = [-4.66, 16.95], ES = 0.18, <i>p</i> = 0.26, <i>t</i> = 1.12

Data are presented as averages ± standard deviation, mean difference (MD), and 95% confidence interval (95%CI), effect size (ES), *p*-value, and *t*-value.

Performance factors

Data for players was further separated from stature, by position. Of the total 552 female soccer players, 288 players (93 players <165 cm) were included when assessing performance. Players were only included if they played an average of ≥ 60 min when entered into a match as this was representative of athletes who play a majority (roughly 67%) of the duration of a match (32–36). Positions were classified as forwards, midfielders, and defenders. No goal keepers were <165 cm, therefore, they were not included in the analysis. Performance was determined in terms of number of assists, number of goals, number of attempts, attempts on target, attempts off target, attempts inside the area, attempts outside the area, attempts on target inside area, attempts on target outside the area, number of corners, shots blocked, defensive blocks, and average distance covered per match played. Average distance covered per match played was a manually computed variable calculated for each player by dividing the total distance covered in kilometres during the World Cup games by the number of matches that individual played in during the World Cup. The averages of each performance factor per player were calculated to compare differences between the two stature groups. Independent t-tests were used to examine differences in each of these factors in players ≥ 165 cm and <165 cm of all positions (Table 1).


Countries

Each of the 24 teams that competed in the 2019 FIFA Women's World Cup were included in this analysis to determine the percentage of players from each team that were <165 cm and ≥ 165 cm tall. Each team consisted of 23 players. Ratios were calculated to determine the percentage of starters on each team that were <165 cm. A one-way ANOVA was used to determine differences in playing time between countries.

All statistical analyses were performed using SPSS Statistics Version 26 (IBM Corp., Armonk, NY). Statistical significance was set at $p < 0.05$, *a priori*. Hedge's *g* effect size was determined. Effect size was identified as small (0.05–0.49), moderate (0.50–0.79), or large (≥ 0.80) (40).

Results

Data are reported as mean difference [95% confidence interval] [MD (95%CI)] and effect sizes (ES), or in some cases (*F*-value, *p*-value) and effect sizes (ES). In total, there were 178 players <165 cm and 374 players ≥ 165 cm. Of the total 552 female soccer players, 288 players were starters. Of the starters, 93 players (32.29%) were <165 cm (Figure 1).




Playing time

For players on the team rosters, the odds of starting if they were <165 cm (52.20% of all players <165 cm) were equivalent to the odds of starting if they were ≥ 165 cm (52.10% of all players ≥ 165 cm); odds ratio [95% confidence interval] (OR [95%CI], 1.0 [0.70, 1.42]). No statistically significant differences were found in playing time between those <165 cm and those ≥ 165 cm (Table 1).

Position

Figure 2 gives a visual representation of the number of players in each position. Midfielders had significantly lower stature than defenders [MD = -2.83 (-4.07, -1.58) cm $p = 0.00$], forwards [MD = -2.02 (-3.35, -0.68) cm, $p = 0.003$], and goal keepers [MD = -8.35 (4.66, 8.01) cm, $p < 0.001$],



respectively. Goal keepers had significantly greater stature than all other positions; defenders, [MD = 5.53 (3.92, 7.14) cm, $p < 0.001$]; forwards, [MD = 6.34 (4.66, 8.01) cm, $p < 0.001$], midfielders, (refer to previous sentence). No statistically significant differences were found between the two stature groups and playing time among defenders, forwards, midfielders, or goal keepers ($p > 0.05$). ES was small (**Table 2**).

Performance

Of the 93 athletes who played ≥ 60 min when entered into a match and were < 165 cm tall, 31 (33.33%) were defenders, 18 (19.35%) were forwards, and 44 (47.31%) were midfielders. 20.34% of all goals (24 out of 118), and 28.77% of all assists (21 out of 73) were made by players < 165 cm. In comparison of the two stature groups, 93 players < 165 cm scored 24 goals and made 21 assists, whereas 195 players ≥ 165 cm scored 94 goals and made 52 assists. There were no significant differences between players ≥ 165 cm and players < 165 cm in regard to the number of assists among both forwards and defenders (MD [95%CI] ES; forwards, 0.17[-0.21,0.54] 0.29, $p = 0.375$; defenders, 0.03[-0.19,0.24] 0.04, $p = 0.804$). No statistically significant difference was found in the number of goals scored among forwards of the two stature groups (MD [95%CI] ES; 0.41[-1.02,0.20]0.38, $p = 0.183$). Among midfielders in the two stature groups, a significant difference was found in the number of assists and the number of goals made (MD [95%CI] ES; number of assists, -0.44[-0.76,-0.11]0.59, $p = 0.009$; number of goals, -0.35[-0.69,-0.01]0.44, $p = 0.047$), with players ≥ 165 cm showing a higher number of assists and goals. (**Supplementary File 1** shows the performance differences reported above among midfielders in each stature group).

Forwards

No statistically significant difference was found between stature and any factor of performance, $p > 0.05$.

Midfielders

Midfielders of greater stature showed significantly higher performance in all variables except distance covered per match played. A statistically significant difference was found with moderate and large effects in number of attempts, attempts on target, attempts off target, attempts on target inside area, attempts on target outside of area, number of corners, shots blocked, and defending blocks (MD [95%CI] ES; number of attempts, 3.14 [1.38, 4.90] 0.80, $p = 0.001$; attempts on target, 1.18 [0.41, 1.94] 0.69, $p = 0.003$; attempts off target, 1.01 [0.19, 1.83] 0.55, $p = 0.016$; attempts on target inside area, 0.83 [0.26, 1.41] 0.65, $p = 0.005$; attempts on target outside area, 0.36 [0.05,

0.67] 0.51, $p = 0.001$; number of corners, 2.04 [0.12, 3.95] 0.48, $p = 0.037$; shots blocked, 0.96 [0.40, 1.51] 0.75, $p = 0.001$; defending blocks, 0.43 [0.32,0.82] 0.48, $p = 0.035$). No statistically significant difference was found in average distance covered per match played (MD [95%CI] ES; average distance covered per match played, 0.29 [-0.52, 1.10] 0.15, $p = 0.484$).

Defenders

No statistical significance was found in any performance variables for defenders between stature groups, $p > 0.05$.

Countries

On average, 32.25% of players on the 2019 World Cup team rosters were < 165 cm, ranging from China with 4.35% of total players < 165 cm, to South Africa with 69.57% of total players < 165 cm. Six out of the 24 teams were composed of 50% or more of their starters being < 165 cm. Twelve out of the 24 teams had a 35% or greater number of starters that were < 165 cm. **Supplementary File 2** shows the number and percentage of players < 165 cm from each country. On average, athletes from all teams and countries played 190.03 ± 174.09 total minutes, 2.62 ± 1.95 matches, 53.48 ± 36.62 average minutes per match, 43.19 ± 35.75 average minutes in relation to total number of matches team played, and 47.98 ± 39.72 percent of matches. There were no statistically significant differences between countries in average minutes played per match, average minutes played in relation to the total number of matches the team played, or percent playing time, respectively; (F -value, p -value); average minutes played per match ($F = 0.34$, $p = 1.00$); average minutes played in relation to the total number of matches the team played ($F = 0.11$, $p = 1.00$); percent playing time ($F = 0.11$, $p = 1.00$). **Supplementary File 3** shows the breakdown of playing time for players from each country in the 2019 FIFA Women's World Cup.

Discussion

The primary aim of this study was to determine if a stature of 165 cm (5'5") is a limitation to the duration of match-playing time on an elite World Cup soccer team, to explore the idea that elite success is in some way related to match-playing time. The secondary aim of this study was to determine if differences in performance exist among elite female soccer players of different stature (≥ 165 cm and < 165 cm) and position in the 2019 FIFA Women's World Cup. Our findings indicate that stature is not a limitation to the duration of match-playing time on an elite World Cup team. Stature did not impact match-playing time at the elite World Cup level in any variables including the odds of starting, minutes played, matches played, average minutes played

when entered into a match, average minutes played in relation to the total number of matches the team played in during the World Cup, or percentage of playing time. One-third of players in the 2019 FIFA World Cup were <165 cm. Furthermore, the average height of females ages 20–29 in the world is 162.56 cm (5'4") (41). In fact, only females in the top 25th percentile are above this height criterion of 165 cm (5'5") (41). Our data do not justify the criterion value of 165 cm that recruiting companies are using.

Additionally, our findings suggest that stature does not have an impact on performance as minimal differences were observed between stature groups in midfielders, and no differences were observed between stature groups in forwards and defenders. Midfielders of greater stature only exhibited a 0.59 greater number of assists and a 0.44 greater number of goals on average compared to their shorter counterparts. Stature was not an imperative factor in any variable when assessing performance in both forwards and defenders. Although goals scored by players ≥ 165 cm were more than double that of players <165 cm, the actual difference was minimal when applied practically, with <1.0 goal difference on average per match between players <165 cm and players ≥ 165 cm (Table 2). Additionally, many players scored zero performance variables, which is reflected in the large confidence intervals found in statistical analyses. The minimal differences found between stature in midfielders may be a result of biomechanics, as longer limbs may attribute to different magnitudes of movement patterns in running, kicking, jumping, and changing direction (cutting ability) (19–21, 23, 42). However, many factors aside from biomechanics have been studied in relation to soccer performance, therefore the contribution of these biomechanical factors alone is not enough to conclude that these elicited the minimal increase found in number of assists and goals among midfielders of greater stature.

A statistically significant difference among midfielders was seen in number of assists, goals scored, attempts, attempts on target, attempts on target inside area, attempts on target outside of area, number of corners, shots blocked, and defending blocks. However, considering players in the 2019 FIFA Women's World Cup ≥ 165 cm ($n = 135$) were greater in numbers compared to players <165 cm ($n = 93$), this could also explain why midfielders of greater stature exhibited higher performance in all variables. This also addresses the performance of forwards <165 cm, as there were no significant differences in any performance variables regardless of there being more forwards ≥ 165 cm. In total, there were more attempts made in general by forwards ≥ 165 cm, however, when compared to the proportion of attempts that were on target and attempts that were off target, the trend was almost identical to that of forwards <165 cm, indicating no difference in performance among players of different stature.

Our findings support previous researchers Lago-Peñas et al. 2011 and Leão et al. 2019 indicating that forwards and midfielders tend to be of smaller, leaner stature and defenders and goal keepers tend to be of greater stature (25, 28). More

importantly, our findings support that shorter stature in forwards and midfielders, and being of smaller stature, does not inhibit performance in this position (5, 28, 43). Our study found controversial results from Rebelo et al. 2013 and Buchheit et al. 2010, that have reported taller and heavier anthropometry is critical for success among defenders, with our findings suggesting that stature is not imperative in performance among defenders (5, 43). The aforementioned researchers assessed stature and body mass in youth and adolescent elite male soccer players under the age of 19 and 15 years, whereas our analysis was among world classified elite female soccer players with an average age of 27.14 ± 3.96 years of age, therefore maturation may explain the difference in results (5, 28, 43–45). Biological maturity may have also been cause for the difference in our study compared to others, as mentioned by Malina et al., this accounts for 21% to 50% of variance seen in youth soccer performance (45). While our analysis suggests stature is of minimal influence in performance, it is only one variable. There are many other variables that need to be accounted for when determining if a player will be successful at the elite level including technical match performance, attack factors, and cognitive, perceptual, and motor skills (7, 25, 46, 47). Taking all of this into account, it is clear that there are many factors that influence performance in soccer.

The importance of stature for elite success in terms of match-playing time and performance in an international soccer competition is not clear in scientific research thus far. There is currently no consensus on which variables have the most impact in determining which potential student athletes are selected or receive the most match-playing time once selected (16). Researchers have recently determined the level of crossover that several criterion variables have in the actual playing of soccer, and physical variables such as stature were deemed to have little implication (18). Our findings support previous literature that emphasizes individual skill capabilities rather than physical variables such as stature when identifying talent in potential student athletes, as no difference in match-playing time exists once athletes are selected (15, 16, 48). Our study adds to current soccer literature as no studies have assessed the influence of stature on match-playing time.

The main limitation of this study is the lack of inclusion of other anthropometric variables to consider in addition to height. There are many other variables that need to be accounted for when determining if a player will be successful at the elite level. In addition, the height was captured from the database and it was not directly measured. No body mass was listed on the FIFA website for any players, therefore anthropometric measures of body mass were not able to be accounted for in our analysis. A second limitation of this study is that the analysis mainly included offensive performance variables typically outside the role of defenders. An analysis of more defensive variables may be more representative of performance in defenders. The amount of time each team spent defending was not considered which could

have impacted performance variables if teams spent a majority of time during matches defending rather than attacking. Finally, this analysis included players from all teams in the 2019 FIFA Women's World Cup, which included 24 different teams from 24 different countries. Training experience and background may have varied greatly between different teams as players may have learned many different approaches and techniques when performing different skills that could have impacted the performance outcome measures used in this study. This may have influenced the performance of players in their ability of making assists, goals, and other offensive performance variables, regardless of stature. Performance of players of different positions between different countries was not analysed, therefore, training experience and background was not accounted for in this analysis.

These findings make clear that stature does not impact the duration of match-playing time or performance on an elite Women's World Cup soccer team. Recruiting companies suggest that aspiring female soccer players be ≥ 165 cm, however, 32.3% of elite female soccer players in the 2019 World Cup are below this stature criterion. These guidelines could potentially discourage coaches from selecting players < 165 cm and discourage aspiring youth female soccer players from pursuing and advancing in the sport (4). Conveying this information and educating coaches on such scientific evidence will help them better understand that stature does not impact an athlete's performance when playing in a match. Furthermore, deconstructing obsolete stereotypes, such as stature, that are placed on female soccer players is essential in developing the future direction of the sport. Additionally, this will help evolve the sport of soccer ensuring that all female athletes are given a chance to acquire and prove their skill, before being excluded merely based on stature.

Data availability statement

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding author/s.

Ethics statement

Ethical review and approval was not required for the study on human participants in accordance with the local legislation and

institutional requirements. Written informed consent was not required for this study in accordance with the national legislation and the institutional requirements.

Author contributions

CM: data analysis, manuscript planning/development, manuscript writing, figure/table creation. YS: data analysis, manuscript planning/development, feedback on writing/figures/tables. CB: data analysis, manuscript planning/development, feedback on writing/figures/tables. MS: data collection, data analysis, manuscript planning/development, abstract writing. ED: feedback on writing/figures/tables. JS: data collection. DD: data collection. JPB: data collection. GC: manuscript perspective, feedback on writing/figures/tables. GN: manuscript perspective, feedback on writing/figures/tables. DC: manuscript oversight and guidance, manuscript planning, development, analyses feedback and perspective. All authors contributed to the article and approved the submitted version.

Conflict of interest

Author GC was employed by the United States Soccer Federation. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fspor.2022.1067190/full#supplementary-material>.

References

1. U.S. girls/boys high school soccer participation 2009–2019. Statista. Available at: <https://www.statista.com/statistics/267963/participation-in-us-high-school-soccer/> (Accessed February 13, 2020).
2. smeyers@ncaa.org. Estimated probability of competing in college athletics. NCAA.org—The Official Site of the NCAA. Published March 2, 2015. Available

at: <http://www.ncaa.org/about/resources/research/estimated-probability-competing-college-athletics> (Accessed February 14, 2020).

3. College recruiting guidelines for women's soccer | D1 requirements. Available at: <https://www.ncsasports.org/womens-soccer/recruiting-guidelines> (Accessed February 14, 2020).

4. Traill RD, And Others. *Girls Playing Soccer: Resistance or Submission? A Case Study of Women's Soccer in the ACT. A Report to the National Sports Research Centre, Australian Sports Commission* (1993). Available at: <https://eric.ed.gov/?id=ED366564> (Accessed July 23, 2020).
5. Rebelo A, Brito J, Maia J, Coelho-e-Silva MJ, Figueiredo AJ, Bangsbo J, et al. Anthropometric characteristics, physical fitness and technical performance of under-19 soccer players by competitive level and field position. *Int J Sports Med.* (2013) 34(4):312–7. doi: 10.1055/s-0032-1323729
6. Reilly T, Williams AM, Nevill A, Franks A. A multidisciplinary approach to talent identification in soccer. *J Sports Sci.* (2000) 18(9):695–702. doi: 10.1080/02640410050120078
7. Zambom-Ferraresi F, Rios V, Lera-López F. Determinants of sport performance in European football: what can we learn from the data? *Decis Support Syst.* (2018) 114:18–28. doi: 10.1016/j.dss.2018.08.006
8. Slimani M, Nikolaidis PT. Anthropometric and physiological characteristics of Male soccer players according to their competitive level, playing position and age group: a systematic review. *J Sports Med Phys Fitness.* (2019) 59(1):141–63. doi: 10.23736/S0022-4707.17.07950-6
9. Grazer JL. *Identifying determinants of match performance in division I Women's Collegiate soccer players.* Johnson City, TN: Proquest Dissertations Publishing (2016). 133 p.
10. Silva JR, Magalhães JF, Ascensão AA, Oliveira EM, Seabra AF, Rebelo AN. Individual match playing time during the season affects fitness-related parameters of Male professional soccer players. *J Strength Cond Res.* (2011) 25(10):2729–39. doi: 10.1519/JSC.0b013e31820da078
11. França C, Inhlle A, Marques A, Sarmento H, Martins F, Henriques R, et al. Physical development differences between professional soccer players from different competitive levels. *Appl Sci.* (2022) 12(14):7343. doi: 10.3390/app12147343
12. Malina RM, Martinho DV, Valente-dos-Santos J, Coelho-e-Silva MJ, Kozielec SM. Growth and maturity Status of female soccer players: a narrative review. *Int J Environ Res Public Health.* (2021) 18(4):1448. doi: 10.3390/ijerph18041448
13. Spehnyak M, Gušić M, Molnar S, Baić M, Andrašić S, Selimi M, et al. Body composition in elite soccer players from youth to senior squad. *Int J Environ Res Public Health.* (2021) 18(9):4982. doi: 10.3390/ijerph18094982
14. Sedano S, Vaeyens R, Redondo JC. The relative age effect in spanish female soccer players. Influence of the competitive level and a playing position. *J Hum Kinet.* (2015) 46(1):129–37. doi: 10.1515/hukin-2015-0041
15. Castillo D, Pérez-González B, Raya-González J, Fernández-Luna Á, Burillo P, Lago-Rodríguez Á. Selection and promotion processes are not associated by the relative age effect in an elite Spanish soccer academy. *PLoS One.* (2019) 14(7):e0219945. doi: 10.1371/journal.pone.0219945
16. Aquino R, Alves IS, Padilha MB, Casanova F, Puggina EF, Maia J. Multivariate profiles of selected versus non-selected elite youth Brazilian soccer players. *J Hum Kinet.* (2017) 60(1):113–21. doi: 10.1515/hukin-2017-0094
17. Carling C, Gall FL, Reilly T, Williams AM. Do anthropometric and fitness characteristics vary according to birth date distribution in elite youth academy soccer players? *Scand J Med Sci Sports.* (2009) 19(1):3–9. doi: 10.1111/j.1600-0838.2008.00867.x
18. Bergkamp TLG, Niessen ASM, den Hartigh RJR, Frencken WGP, Meijer RR. Methodological issues in soccer talent identification research. *Sports Med.* (2019) 49(9):1317–35. doi: 10.1007/s40279-019-01113-w
19. Putnam CA. A segment interaction analysis of proximal-to-distal sequential segment motion patterns. *Med Sci Sports Exerc.* (1991) 23(1):130–44. PMID: 1997807.
20. Chrisman SP, O'Kane JW, Polissar NL, Tencer AF, Mack CD, Levy MR, et al. Strength and jump biomechanics of elite and recreational female youth soccer players. *J Athl Train.* (2012) 47(6):609–15. doi: 10.4085/1062-6050-47.6.01
21. Dörge HC, Anderson TB, Sørensen H, Simonsen EB. Biomechanical differences in soccer kicking with the preferred and the non-preferred leg. *J Sports Sci.* (2002) 20(4):293–9. doi: 10.1080/026404102753576062
22. Orloff H, Sumida B, Chow J, Habibi L, Fujino A, Kramer B. Ground reaction forces and kinematics of plant leg position during instep kicking in Male and female collegiate soccer players. *Sports Biomech.* (2008) 7(2):238–47. doi: 10.1080/14763140701841704
23. Marshall BM, Franklyn-Miller AD, King EA, Moran KA, Strike SC, Falvey ÉC. Biomechanical factors associated with time to complete a change of direction cutting maneuver. *J Strength Cond Res.* (2014) 28(10):2845–51. doi: 10.1519/JSC.0000000000000463
24. Kellis E, Katis A. Biomechanical characteristics and determinants of instep soccer kick. *J Sports Sci Med.* (2007) 6(2):154–65. PMID: 24149324.
25. Leão C, Camões M, Clemente FM, Nikolaidis PT, Lima R, Bezerra P, et al. Anthropometric profile of soccer players as a determinant of position specificity and methodological issues of body composition estimation. *Int J Environ Res Public Health.* (2019) 16(13):1–10. doi: 10.3390/ijerph16132386
26. Deprez DN, Franssen J, Lenoir M, Philippaerts RM, Vaeyens R. A retrospective study on anthropometrical, physical fitness, and motor coordination characteristics that influence dropout, contract Status, and first-team playing time in high-level soccer players aged eight to eighteen years. *J Strength Cond Res.* (2015) 29(6):1692–704. doi: 10.1519/JSC.0000000000000806
27. Malina RM, Bouchard C, Bar-Or O. *Growth, maturation, and physical activity.* Austin, TX: Human Kinetics (2004).
28. Lago-Peñas C, Casais L, Dellal A, Rey E, Domínguez E. Anthropometric and physiological characteristics of young soccer players according to their playing positions: relevance for competition success. *J Strength Cond Res.* (2011) 25(12):3358–67. doi: 10.1519/JSC.0b013e318216305d
29. FIFA Women's World Cup France 2019™. Available at: <https://www.fifa.com/tournaments/womens/womensworldcup/france2019/origin1904-p.cxm.fifa.com/tournaments/womens/womensworldcup/france2019/teams> (Accessed November 21, 2022).
30. Players | USWNT & USMNT | U.S. Soccer Official Website. Available at: <https://www.ussoccer.com/players> (Accessed November 21, 2022).
31. Bradley P, Scott D. Physical Analysis of the FIFA Women's World Cup France (2019). Available at: <https://digitalhub.fifa.com/m/4f40a98140d305e2/original/zijqly40ednqa5gffgaz-pdf.pdf> (Accessed November 21, 2022).
32. Kraemer WJ, French DN, Paxton NJ, Häkkinen K, Volek JS, Sebastianelli WJ, et al. Changes in exercise performance and hormonal concentrations over a big ten soccer season in starters and nonstarters. *J Strength Cond Res.* (2004) 18(1):121–8.
33. Giménez JV, Leicht AS, Gomez MA. Physical performance differences between starter and non-starter players during professional soccer friendly matches. *J Hum Kinet.* (2019) 69(1):283–91. doi: 10.2478/hukin-2019-0018
34. Anderson L, Orme P, Michele RD, Close GL, Milsom J, Morgans R, et al. Quantification of seasonal-long physical load in soccer players with different starting Status from the English premier league: implications for maintaining squad physical fitness. *Int J Sports Physiol Perform.* (2016) 11(8):1038–46. doi: 10.1123/ijssp.2015-0672
35. Arcos A L, Mendez-Villanueva A, Martínez-Santos R. In-season training periodization of professional soccer players. *Biol Sport.* (2017) 34(2):149–55. doi: 10.5114/biolsport.2017.64588
36. Raya-González J, Castillo D, Yanci J, Los Arcos A. Assessing the perceived exertion in elite soccer players during official matches according to situational factors. *Int J Environ Res Public Health.* (2020) 17(2):410. doi: 10.3390/ijerph17020410
37. Ghasemi A, Zahediasl S. Normality tests for statistical analysis: a guide for non-statisticians. *Int J Endocrinol Metab.* (2012) 10(2):486–9. doi: 10.5812/ijem.3505
38. Gómez MÁ, Lago C, Gómez MT, Furley P. Analysis of elite soccer players' performance before and after signing a new contract. *PLoS One.* (2019) 14(1):e0211058. doi: 10.1371/journal.pone.0211058
39. Wells AJ, Hoffman JR, Beyer KS, Hoffman MW, Jajtner AR, Fukuda DH, et al. Regular- and postseason comparisons of playing time and measures of running performance in NCAA division I women soccer players. *Appl Physiol Nutr Metab.* (2015) 40(9):907–17. doi: 10.1139/apnm-2014-0560
40. McGough JJ, Faraone SV. Estimating the size of treatment effects. *Psychiatry.* (2009) 6(10):21–9. PMID: 20011465.
41. Vital and Health Statistics, Series 3, Number 46:44.
42. Strength and speed characteristics of elite, subelite, and recreational young soccer players.—PubMed—NCBI. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/?term=strength+and+speed+characteristics+of+elite%2C+subelite%2C+and+recreational+young+soccer+players> (Accessed February 13, 2020).
43. Buchheit M, Mendez-Villanueva A, Simpson BM, Bourdon PC. Match running performance and fitness in youth soccer. *Int J Sports Med.* (2010) 31(11):818–25. doi: 10.1055/s-0030-1262838
44. Mirkov DM, Kukolj M, Ugarkovic D, Koprivica VJ, Jaric S. Development of anthropometric and physical performance profiles of young elite Male soccer players: a longitudinal study. *J Strength Cond Res.* (2010) 24(10):2677–82. doi: 10.1519/JSC.0b013e3181e27245
45. Malina RM, Eisenmann JC, Cumming SP, Ribeiro B, Aroso J. Maturity-associated variation in the growth and functional capacities of youth football (soccer) players 13–15 years. *Eur J Appl Physiol.* (2004) 91(5):555–62. doi: 10.1007/s00421-003-0995-z
46. Rowat O, Fenner J, Unnithan V. Technical and physical determinants of soccer match-play performance in elite youth soccer players. *J Sports Med Phys Fitness.* (2017) 57(4):369–79. doi: 10.23736/S0022-4707.16.06093-X
47. Ali A. Measuring soccer skill performance: a review. *Scand J Med Sci Sports.* (2011) 21(2):170–83. doi: 10.1111/j.1600-0838.2010.01256.x
48. Deprez D, Coutts AJ, Franssen J, Deconinck F, Lenoir M, Vaeyens R, et al. Relative age, biological maturation and anaerobic characteristics in elite youth soccer players. *Int J Sports Med.* (2013) 34(10):897–903. doi: 10.1055/s-0032-1333262