

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

6,200

Open access books available

168,000

International authors and editors

185M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Chapter

Conservation Agriculture: Climate Proof and Nature Positive Approach

*Rachid Mrabet, Akashdeep Singh, Tarun Sharma,
Amir Kassam, Theodor Friedrich, Gottlieb Basch,
Rachid Moussadek and Emilio Gonzalez-Sanchez*

Abstract

The development pathways of countries and regions have impacted land-climate interactions and shaped challenges, opportunities and actions. Adverse impacts of climate change increasingly threaten livelihoods and resilience of people around the globe, food security and the stability of environmental resources. Globally, the current food systems are not fit for purpose. Land-based options such as Conservation Agriculture (CA) were found to mitigate climate change, regenerate soils and ensure durable food systems. Achieving sustained results using CA systems, under climate change and social pressures, while maximizing co-benefits related to food and nutrient security, social and biological diversity, ecosystem restoration and services and sustainable development, requires appropriate country-specific policies and significant investment. CA implementation is challenging and context specific and necessitates an integrated framework and road map to enable deeper ambitions for social equity and development and inclusive economic growth.

Keywords: no-till, soil mulch cover, climate change, sustainability, environment, carbon sequestration

1. Introduction

The interaction between land and climate is a complex system thoroughly influencing the agriculture production systems around the globe [1]. Agricultural production systems are the largest single source of environmental degradation, responsible between 21 and 37% of global greenhouse gas (GHG) emissions through deforestation, depletion of soil carbon, release of nitrous oxide and enteric fermentation. Without intervention, these are likely to increase by about 30–40% by 2050, due to increasing demand based on population and income growth and dietary change. Agriculture is also responsible for 70% of freshwater use, 30% of energy use and 80% of land conversion [2]. Conventional agricultural practices revolved around the

burning of crop residues to facilitate land preparation for the succeeding crop, regular plowing and tillage of the land for preparing seedbeds and controlling weeds. Reduced natural soil productivity and pest control were corrected with new high yielding breeds, fertilizers and pesticides. These practices initially had a positive effect on production and yield of crops but at the cost of continuous land degradation, erosion by wind or water, underground water pollution, oxidation of the soil organic matter due to tillage and emitting carbon dioxide (CO₂) in large amounts [3]. Like in other aspects of the economy a trickle-down of benefits to poor farmers is assumed, but rarely materialized.

By the year 2050, the global population is expected to increase to 9.1 billion which would mean that the existing production systems need to gear up and increase their food production by 70 per cent by the year 2050, assuming food waste and change of consumer preferences continue unchanged. Producing sufficient food with finite resources to feed the growing global population while having a smaller impact on the environment has always been a great challenge. In addition, the 2022 IPCC reports generated enormous attention as a demand for immediate actions across all sectors and regions. There is a need for rethinking the actual food systems and address all the connected challenges and threats and explore the root causes of unsustainability. Consequently, healthy growth and stable productivity of crops and livestock require innovative models of food production for resource-saving, environmentally friendly agriculture. Conservation Agriculture (CA) has proven to overcome the shortcomings of tillage-based agriculture in terms of sustainability as a promising system-based approach [4]. Here, we review the environmental impacts of CA that should lead to a paradigm shift in goals and models of food production for promoting sustainable and regenerative agriculture worldwide.

2. Conservation agriculture: adoption evolution and trends

The need for a transformation of conventional tillage-based agriculture became obvious in the early 1930s after the 'Dust Bowl' trembled the mid-west farming communities of the United States and obliged the scientific community to reorient its research agenda and focus more on erosion mitigation and soil conservation through no-tillage systems (later called Conservation Agriculture or CA systems) [5]. CA is a resource-conserving agricultural concept that is steadily gaining ground and covers an estimated area of 205 million hectares (14.7% of global cropland) (also see **Table 1** for regional distribution). This represents an increase of 93% in global CA cropland area since 2008/09 and represents an annual increase of about 10 Mha.

The major countries practicing CA in 2018/19 are the USA (44.0 Mha), Brazil (43.9 Mha), Argentina (32.9 Mha), Australia (22.9 Mha), Canada (21.7 Mha) and others (39.6 Mha) [5]. In other terms, the total CA area is approaching 70% and 75% of the total cropland area in South America and in Australia, respectively. However, since 2008/09, percentage change in CA adoption has been greater in Asia, Africa and Europe than in the other continents, and corresponds to 33.1 Mha or about 16% of the global CA cropland area [5]. CA as climate proof agriculture and its roles for soil sustainability and resilience are widely recognized and should favor increase in its adoption by mainstreaming the concept in agricultural and environmental policies.

Region	CA cropland area 2008/2009	CA cropland area 2013/2014	CA cropland area 2015/2016	CA cropland area 2018/2019	Percent change in CA area since 2015/2016	Percent change in CA area since 2013/2014	Percent change in CA area since 2008/2009	Percent CA cropland area in the region 2018/2019
S and C America	49,564.10	66,377.00	69,895.00	82,996.18	18.7	25.0	67.5	68.7
North America	40,003.80	53,967.00	63,181.00	65,937.22	4.4	22.2	64.8	33.6
Australia and New Zealand	12,162.00	17,857.00	22,665.00	23,293.00	2.8	30.4	91.5	74.0
Russia and Ukraine	100.00	5200.00	5700.00	6900.00	21.1	32.7	6800.0	4.5
Europe	1560.10	2075.97	3558.20	5601.53	57.4	169.8	259.0	5.2
Asia	2630.00	10,288.65	13,930.20	17,529.02	25.8	70.4	566.5	3.6
Africa	485.23	993.44	1509.24	3143.09	108.3	216.4	547.8	1.1
Total	106,505.23	156,759.06	180,438.64	205,400.04	13.8	31.0	92.9	14.7

Table 1. Global spread of CA cropland area ('000 ha) in different regions for 2008/2009, 2014/2015, and 2018/2019, and corresponding percent change (source: [5]).

3. CA as climate proof agriculture

Since food production is sensitive to weather conditions, the very existence of mankind is being threatened by an unseen force referred to as climate change. Climate change is expected to adversely affect the climatic/weather phenomena thereby impacting the global food supply system [6, 7]. The evidence is irrefutable that GHG are choking our planet and placing billions of people in danger. An increasing number of people are not able to realize their right to adequate food. In 2020, between 720 and 811 million people in the world faced hunger, up to 161 million more than in 2019 ([8]). In other words, climate change, food security and biodiversity are the “trilemma of land use”. Solutions to these challenges should be integrated to combine and tackle multiple goals. Hence, considering the world is at stake, various international organizations such as Food and Agriculture Organization (FAO), World Bank, and many more, have come together to tackle climate change and food insecurity and search for a reasonable, economical, and sustainable solution. In addition, these urgencies should also rely on the need for grassroots led structural change to stay within the ecological boundaries of the planet, several of which have already been exceeded.

Ingenuous and meritoriously employed land-based measures [9], including specific measures to protect and enhance soil organic carbon stocks, can directly support the global environmental and sustainability goals under the UNFCCC [1], the UNCCD [10], and the CBD [11]. In lieu, among these measures, CA works as a systemic approach with its key contributions to sustainability, climate change adaptation and mitigation as well as food security [12–15].

From its wide-ranging adaptation and adoption, CA systems are being practiced in rainfed and irrigated systems, annual, perennial and mixed cropping systems, orchards and plantation systems, agroforestry systems, pasture systems, organic and nonorganic systems, and rice-based systems [16, 17].

Abundant literature and multi-stakeholder innovation platforms across various farming systems showed that CA is a climate proofing agriculture [18–20]. In fact, several recent studies have found that fully-implemented CA can improve crop yield stability—a measure of climate resilience—in different soil types, climates and cropping systems. Worldwide, CA has helped bolstering productivity, augmenting resilience to weather shocks, and tumbling negative externalities (i.e. [21] in USA; [22] in Australia; [23, 24] [25] in India; [26, 27] in China; [28–31] in Africa; [32, 33] in West Asia and North Africa, [34] in Europe). As sometimes observed, decrease in crop yield following the adoption of CA largely depends on whether CA has been correctly implemented, with the use of appropriate seeders, seed rates, fertilizer applications and management practices followed to manage weeds and pests. Some yield reductions in initial years were also due to problems of drainage and stagnation of water in cool and humid regions as result of poorly structured soils from a tillage-based farming history.

Experiences in drought conditions have shown that CA yields can be twice as much as conventional agriculture, peaking up to 4-fold higher yields in wheat [19, 35]. Sun et al. [36] found that in arid regions, CA permitted both increased carbon sequestration and crop yields. Based upon a meta-analysis comprising 610 studies, 48 crops and 63 countries, Pittelkow et al. [37] found variable responses from CA compared to conventional tillage systems. The authors concluded that CA are better performing under a range of crop species in arid regions – particularly where water is limiting to crop growth. The authors also reported that yield gaps are due to partial use of CA

principles, which obviously will not produce all the CA benefits. When no-till is combined with residue retention and crop rotation, which is the full implementation of the CA principles, no significant yield reduction is noticed: indeed, this combination of techniques significantly increases crop yields in dry climates. A dataset containing 4403 paired (CA vs. CT) yield observations collected between 1980 and 2017 for eight major staple crops in 50 countries presented by Su et al. [38] also confirmed this trend. In addition, selecting high-efficiency crop varieties and optimizing agronomic (nitrogen) management practices to increase water/nitrogen use efficiency is an effective way to increase crop yield with less associated environmental costs under CA. In order to achieve increased yield stability across climate and soil gradients, it is of paramount importance to grow mixtures of crop species or mixtures of genotypes to exploit positive interaction effects and thus reduce the risk of crop failure [39].

A meta-analysis using data from 9686 paired site-year comparisons across South Asia in a variety of cropping systems found that, CA systems provided 5.8% higher mean yield than conventional agricultural practices [40]. In another study by Laik et al. [41], under CA systems, yields of wheat and rice increased by 46–54 and 10–24%, respectively, over conventional tillage, thereby obtaining ~53% higher total output from the CA system. In a review study by Das et al. [24], the CA systems increased yields of crops from 2% to 200% depending on crop rotations and years of implementation.

Through a meta-analysis of 933 observations from 16 different countries in sub-Saharan African studies, Corbeels et al. [42] showed that average yields under CA are only slightly higher than those of conventional tillage systems (3.7% for six major crop species and 4.0% for maize). Larger yield responses for maize result from mulching and crop rotations/intercropping. They also concluded that when CA principles are implemented concomitantly, maize yield increases by 8.4%, which proves the fact, that the lower yield benefits reported in the study resulted from mixing CA systems with systems that only adopted some of the CA principles.

One of the most entrenched benefits of CA systems is their ability to improve soil water storage. The maintenance of crop residues and mulches at the surface of soils under CA systems improves the water balance of the soil-cropping system. CA systems improve the uptake, conservation, and use of available water in the soil by the crops [43, 44]. All this increases the responsiveness of CA systems to changes in climate, meaning crops under these systems have a much better capacity for coping and adapting to drought. Under rainfed ecologies of eastern and south African countries, CA systems reduced the yield variability by 11% over CT [45].

When CA systems are implemented in warmer and drier regions, higher crop yields are often observed due to a lowering of soil temperatures in addition to increases in soil water storage. In irrigated regions, higher water storage and better water management under CA systems can reduce the amount of water required for crop production and help conserve water resources [46–48].

4. Environmental sustainability: soil re-carbonization, conservation, health and security

Soil's multi-functionality and health were generally neglected to address food and climate security challenges [3, 49]. However, after the Paris Agreement was signed, stakeholders committed in a voluntary action plan to implement farming systems and practices that maintain or enhance soil carbon stocks in agricultural soils and to

preserve carbon-rich soils [50]. Global technical potential of SOC sequestration is 1.45–3.44 Pg C/year (2.45 Pg C/year) but varies with type of soils, management and ecologies [51].

The push to the CA-based system is due to its environmental and productivity sustainability and especially its ability to (i) reduce soil degradation, erosion and runoff, (ii) mitigate greenhouse gas emissions and (iii) sequester atmospheric CO₂ in the form of soil organic carbon, tackle climate change, (iv) improve biodiversity below and above the soil surface, and (v) enhance production system resilience to abiotic and biotic stresses [52–57]. In fact, CA systems were initially adopted for soil conservation and erosion control benefits, but they are gaining more and more attention as a practice to maintain and/or increase SOC and harness ecosystem services in agroecosystems [58].

CA aims to implement soil-based strategies and long-term soil fertility dynamics that restore soil functions and health [3, 56, 59] and increase carbon storage reversing consequently the food insecurity spiral [30, 60–62]. In other terms, it is beneficial for crop production and soil health and functions and hence to global food security and adaptation of agriculture to climate change [16, 17, 38, 62–65]. Lal et al. [66] concluded that evidence-based strategy based on CA can allow re-carbonization of depleted soils. Studies by Blanco-Canqui and Ruis [67] confirmed that when CA systems are applied in an integrative way, synergic effects of the principles give rise to levels of soil organic matters.

It goes without saying that carbon stored in the soil is the most stable carbon (C) pool, an essential part of ecosystem services and a tool to tackle climate change [68]. In view of its role in soil aggregation and erosion control, in availability of plant nutrients and in ameliorating other forms of soil degradation than erosion, CA systems have proven to reduce soil degradation and rebuild soil quality. However, in areas with low fertility, integrated nutrient management is essential to ensure a build-up of SOC and the success of CA systems (i.e., in Africa). This cycle can be broken by judicious addition of nutrients to the soil/crop system via organic or synthetic fertilizers and/or the incorporation of legumes into cropping rotations [56, 69].

In lieu of climate change, sequestering CO₂ has become inevitable. CA systems in comparison to the conventional practices saw an increase in SOC in top-soil (0–15 cm) by 3.8 Mg ha⁻¹, in the deepest layer (70–100 cm) by 2.5 Mg ha⁻¹ and mean C sequestration rates of 0.09 and 0.27 Mg ha⁻¹ yr⁻¹ [70]. Soil carbon sequestration bids to improve soil fertility and reduce carbon dioxide levels in the atmosphere. Among continents, Africa is the smallest contributor to greenhouse gas emissions but is highly susceptible to climate change, which is mainly responsible for rising temperatures, fluctuating rainfall patterns, increased frequency of disastrous events such as droughts and floods leading to heavy losses in terms of resources. Gonzalez-Sanchez et al. [71] reported that an estimate of the potential annual carbon sequestration in African agricultural soils through CA amounts to 143 Tg of C per year, that is 524 Tg of CO₂ per year. This figure represents about 93 times the current sequestration figures. In addition, this potential is almost 3 times higher than the one found for Europe by Gonzalez-Sanchez et al. [72], which amounts to 189 Tg CO₂ per year.

In the rice-wheat cropping system, an improvement in carbon stocks by 20% and 40% at a depth of 0–15 and 15–30 cm was realized by following the CA principles [73]. A worldwide meta-analysis by Li et al. [64] found that, on average, the number of water stable aggregates in CA systems are 31% greater compared to conventionally tilled systems. Such soil quality improvements are based on greater SOM content which provides greater abundance of habitats to support microbial, micro- and

meso-fauna activity. By enhancing soil health and re-carbonizing the soils [3], CA systems establish dynamic ecological conditions in the soil/plant/landscape continuum which offers resilient performance with maximum productivity (water and nutrient use efficiency and water productivity) [4].

Several authors reported that CA systems minimize on-site and off-site effects with regards to soil degradation and that benefits to soil health and ecosystems follow a chain-like process. Under CA systems, erosion is lessened, infiltration is improved, and water losses either through evaporation or runoff are reduced, allowing the crop to have more water in dry periods or years [64]. In other terms, CA also contributes to the environment by mitigating pollution as it reduces off-site transport of residual agrochemicals through runoff and soil sediments. This reduces the surface transport of nitrate and phosphorus from agricultural fields and the eutrophication of water bodies. Also leaching of nutrients under CA is usually reduced, as the water is mainly transported through macro pores (bypass flow) and not washing the soil matrix as long as synthetic or organic fertilizer or slurry is not applied directly before a heavy rainstorm, which can potentially increase leaching of nitrate to groundwater through the macropores [58, 74].

According to Lal [74], in addition to carbon sequestration and erosion control, adoption of CA systems accentuates several other ecosystem services such as biodiversity, elemental cycling, and resilience to natural and anthropogenic perturbations, all of which can affect food security. It was also reported that CA systems do not lead to significant compaction and higher bulk densities than traditional systems based on soil disturbance [64].

In addition, when combined with frontier technologies (precision agriculture, plant breeding and biotechnology, microbial biotechnology, smart fertilizers, biochar additions etc.), CA systems can help to soak up even more carbon in the soil, create soil resilience to achieve food security and mitigate climate change and allow higher and stable yields [54, 75, 76].

5. Economics under CA systems: no regret options

Countries seek to and should improve the well-being of people and especially farmers. The conventional system of agricultural production is hugely dependent on intensive tillage operations with the support of much labor or heavy farm machinery. The latter results in higher CO₂ emissions and both in higher production costs [77]. Reducing the tillage operations has the potential of reducing emissions and fuel consumption. CA systems can save up to 80% of fossil fuel energy used by tillage [24].

The farmers, and mainly the resource-poor ones, need production systems that are regenerative, reliable, financially viable and profitable. However, many scientific studies agree that CA systems are cost-effective, energy efficient and allow farmers higher and more stable incomes [24, 34]. The major factor leading to lower costs in CA systems is attributed to bypassing soil manipulation and disturbance unlike conventional tillage systems, where 4–5 primary and secondary tillage operations are performed for seedbed preparation and weed control, which acquire higher costs [78, 79].

Even if CA systems in the beginning might have undesirable effects on crop yield levels, the cost of cultivating crops decreases with fewer use of machinery and compensates for eventual initial yield declines. Subsequently, continuous use of such practices improves soil properties, sustains crop productivity and ultimately

economic returns [80]. In fact, according to several authors, there is mounting evidence that when CA is inconsistently applied, it leads to lower yields and higher costs than expected [38, 62, 81–83]. Arenas-Calle et al. [84] showed that the lack of climate-smartness resulted in yield penalties in early stages of CA implementation. However, in eastern and southern Africa the highest financial returns (90–95%) from CA investments by small-holder farmers were realized under low-rainfall conditions (<700 mm), thereby providing clear evidence of the climate smartness of CA systems under soil moisture-stressed conditions [45].

With the reduced expenses in terms of labor, energy and monetary inputs, CA practices reduce the cost of cultivation. Reduced expenditure in such a pattern was observed in winter wheat for no-tillage practices (1300 Yuan ha⁻¹), reduced tillage (2250 Yuan ha⁻¹) as compared to conventional tillage practices (2500 Yuan ha⁻¹) [85]. Especially in the case of small and resource poor farmers, with reduced usage of machinery cost (<65.52%) under CA, farmers spend less (14.46%) on different cultivation practices, increasing their net returns as compared to conventional agriculture practices [86, 87]. In sub-Saharan Africa, scientific studies revealed that with systematic use of practices such as no-tillage, residue retention and crop rotation the costs of cultivating maize or soybean were reduced (20–29%) and the net returns, the benefit-cost ratio increased to a greater extent [88–98]. A similar impact of less soil disturbing practices such as permanent beds and zero tillage was obtained on net returns or profitability of maize-chickpea rotation in India (28.8% and 24% respectively) [99–103].

In a regional study in Ethiopia, CA was found to have reduced the labor usage by 32–41% whereas 50–60% labor was replaced at the critical periods of crop production due to reduced tillage operations in the maize-soybean intercropping system. Further, a maximum return of 15,545 ETH birr ha⁻¹ and 12,693 ETH birr ha⁻¹ was obtained when soybean and haricot were intercropped in maize [104]. The net returns in production of the rice CA systems were 581 USD ha⁻¹ in comparison to 412 USD ha⁻¹ under the conventional system. The gross returns in the rice-wheat system were highest (2456 USD ha⁻¹) under the CA system [79].

Choudhary et al. [90] found 22.3 and 24.5% higher grain yield of pearl millet [*Cenchrus americanus* (L.) Morrone] and Indian mustard [*Brassica juncea* (L.) Czernj.] under CA systems, respectively, compared to conventional systems, which ultimately led to higher net returns (US\$ 1270 ha⁻¹).

From a meta-analysis carried out by Ogle et al. [98], it was concluded that CA systems drastically reduce the number of field operating hours and associated fuel use by about 69%. From these studies, it is clear that broadening access to finance, including international and climate finance will catalyze adoption and accelerate the shift towards CA systems.

6. Conclusion

Research and development efforts in agriculture have been increasingly oriented towards improving modern, industrial or corporate agriculture—new chemicals, hybrid and genetically modified seeds, mechanization, factory farming, etc. Hence, the agriculture sector is replete with innovations but not all of them were found sustainable.

The CA systems backed by various institutions, research scholars, policymakers were found able to adapt to the fast-changing environment thus making the food

system healthy, flexible, productive and profitable. Further, CA helps to extenuate the greenhouse gas emissions and increase the carbon stocks making soils resilient, reliable and sustainable. In other words, the main benefits of CA systems cover numerous areas and contribute to a number of SDGs. In addition, CA feasibility or adoption was assessed in contrasted biophysical, social and economic environments.

CA systems are alternative pathways for agriculture to be more conducive to durable food systems and longer-term sustainability. Especially soil carbon sequestration and health improvement allowed by CA systems can support various ecosystem services related to climate change adaptation, food security and biodiversity due to enhanced soil fertility and nutrient pools, increased moisture retention, improved water availability to plants and reduced soil erosion and runoff [58]. The number of countries explicitly including SOC in agricultural land (including wetlands) in the Nationally Determined Contributions (NDCs) increased from 28 (15% of first-round NDCs assessed) to 35 (24% of latest NDCs assessed) [101] which is still insufficient. An international agenda for restoring soil health and inclusion of soil carbon sequestration in policies and actions should be advocated and supported [88, 102]. Policies promoting the target of land degradation neutrality can support food security, human wellbeing and climate change adaptation and mitigation [2].

Barriers to the adoption of CA system are more related to farmers' attributes (adopter's characteristics, limited availability of resources, level of perception, mind-set, cultural values, illiteracy, willingness for change, etc.) and their enabling environment (e.g., legal compliance, governance, lack of training and capacity building, stakeholder communication, lack of financial support, insufficient economic and social incentives) than to technical concerns (i.e. herbicide and machinery availability and costs, energy use and price, competitive uses of crop residues and livestock etc.). The science related to CA systems is currently advanced enough to inform the formulation of policy and incentive programs for CA adoption at a scale large enough to result in the radical transformation of mainstream agricultural production systems CA [5, 103].

Dis-adoption, accumulating challenges and difficulties of mainstreaming CA by additional farmers arise from two main issues: (i) CA is dynamic, meaning that it should respond to simultaneous changes in environment, social and/or economic contexts, (ii) CA is also a holistic concept based on a system-wide approach to solving farm management shifts and problems while considering the integrality of the food system. In addition, agriculture functions are changing over time and getting more complex with increasing socio-economic and environmental stresses and social and institutional shocks [33]. Approaches for upscaling CA range from sophisticated decision support systems to improved enabling environments (i.e., through land policies and subsidies focused on water, environment, and poverty) and promotion of social or sustainability-oriented learning processes [16].

The new Green Revolution (GR) of the twenty-first century must be: (i) soil-centric, based on soil health and resilience, (ii) ecosystem-centric, based on eco-efficiency of inputs, (iii) knowledge or innovation-centric, based on scientific principles, and (iv) nature-centric, based on nature positive solutions which restore and enhance nature [95]. The new GR should also recognize the "One Health" concept, which states that the "health of soil, plants, animals, people, ecosystems, and the planetary processes is one and indivisible [94].

The 8th World Congress on Conservation Agriculture (WCCA), which inspired from these paradigms of the new GR, set a goal to increase the global CA cropland

area to 50% of the total cropland by 2050, in particular to respond to the global challenge to mitigate the advancing climate change and land degradation and reduce gaps in food security and nutrition (as well as other sustainable development goals). This represents an area of 700 M ha [92, 103]. In achieving such goals, policy and economic incentives should be enforced and augmented in most countries. In addition, the integration of CA benefits in the farming system (e.g., value chain design, marketing, labeling), can lead to giving carbon both economic and environmental values and thus increasing farmer income and stewardship. Social norms as well as psychological and behavioral factors must be considered for widespread adoption of CA systems. Accordingly, a multi-stakeholder engagement and joint coordination (i.e., science-policy dialog and engaged civil society) are major issues in the development and implementation of a CA Road Map for wide mainstreaming and large-scale adoption by farmers in markedly diverse ecologies. Implementing CA Road maps enable governments, landowners and land managers, and the community to share responsibility for land-based challenges mitigation and hence in achieving or reaching SDGs. According to Lal [93], sustainable intensification of agroecosystems (which includes CSA systems) can produce enough food grains to feed one person for a year on 0.045 ha of arable land. Hence, another issue of prime importance concerns socializing CA for the small land size farmers while integrating livestock and trees mainly in Africa and Asia. There is great momentum in merging principles of CA with those regenerative types of farming and especially those related to tillage, synthetic fertilizers and pesticide use [97]. However, issues related to GMOs are still largely debated within the agroecological stream. Kassam and Kassam [4] proposed an inclusive ethical and responsible system to integrate CA systems with plant-based diets and organic farming practices in order to move from corporate agriculture.

IntechOpen

IntechOpen

Author details

Rachid Mrabet¹, Akashdeep Singh², Tarun Sharma², Amir Kassam^{3*},
Theodor Friedrich⁴, Gottlieb Basch⁵, Rachid Moussadek⁶
and Emilio Gonzalez-Sanchez^{7,8,9}

1 National Institute of Agricultural Research, INRA, Rabat, Morocco

2 Department of Agronomy, CSK Himachal Pradesh Agricultural University,
Palampur, India

3 University of Reading, UK

4 Former Residential Representative, FAO, Bolivia

5 University of Evora, Portugal

6 International Center for Agricultural Research in Dry Areas (ICARDA), Rabat,
Morocco


7 Universidad de Córdoba, Spain

8 European Conservation Agriculture Federation (ECAAF), Brussels, Belgium

9 Asociación Española Agricultura de Conservación Suelos Vivos (AEACSV), Spain

*Address all correspondence to: amirkassam786@googlemail.com

IntechOpen

© 2022 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Jia G, Shevliakova E, Artaxo P, De Noblet-Ducoudré N, Houghton R, House J, et al. Land–climate interactions. In: *Climate Change and Land: An IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*. Geneva, Switzerland: IPCC; 2019
- [2] IPCC. *Climate Change and Land: An IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*. Geneva, Switzerland: IPCC; 2019
- [3] Montgomery DR. Soil health and the revolutionary potential of Conservation Agriculture. In: Kassam A, Kassam L, editors. *Rethinking Food and Agriculture: New Ways Forward*. Elsevier; 2020. pp. 219-229
- [4] Kassam A, Kassam L. *Rethinking Food and Agriculture: New Ways Forward*. Switzerland: Elsevier; 2020
- [5] Kassam A, Friedrich T, Derpsh R. State of the global adoption and spread of conservation agriculture. In: Kassam A, editor. *Advances in Conservation Agriculture, Adoption and Spread*. Cambridge, UK: Burleigh Dodds; 2022
- [6] Aryal JP, Sapkota TB, Khurana R, Khatri-Chhetri A, Rahut DB, Jat ML. Climate change and agriculture in South Asia: Adaptation options in smallholder production systems. *Environment, Development and Sustainability*. 2019;2019:1-31
- [7] Myers SS, Zanolotti A, Kloog I. Increasing CO₂ threatens human nutrition. *Nature*. 2014;510:139-142
- [8] FAO. *The State of the World's Land and Water Resources for Food and Agriculture – Systems at Breaking Point*. Rome: FAO; 2021
- [9] Roe S, Streck C, Beach R, Busch J, Chapman M, et al. Land-based measures to mitigate climate change: Potential and feasibility by country. *Global Change Biology*. 2021;27:6025-6058. DOI: 10.1111/gcb.15873
- [10] Chotte J, Aynekulu E, Cowie A, Campbell E, Vlek P, Lal R, et al. *Realising the Carbon Benefits of Sustainable Land Management Practices: Guidelines for Estimation of Soil Organic Carbon in the Context of Land Degradation Neutrality Planning and Monitoring*. Bonn, Germany: United Nations Convention to Combat Desertification (UNCCD); 2019
- [11] IPBES. *Summary for Policymakers of the Assessment Report on Land Degradation and Restoration of the Intergovernmental Science Policy Platform on Biodiversity and Ecosystem Services*. Bonn, Germany: IPBES; 2018
- [12] Barasa PM, Botai CM, Botai JO, Mabhaudhi T. A review of climate-smart agriculture research and applications in Africa. *Agronomy*. 2021;11:1255. DOI: 10.3390/agronomy11061255
- [13] Corbeels M, Naudin K, Guibert H, Torquebiau E, Cardinael R. Is the 4 per thousand initiative for soil organic carbon storage achievable in sub-Saharan Africa? Insights from agroforestry and conservation agriculture. *Soil Till Research*. 2018;188:16-26. DOI: 10.1016/j.still.2018.02.015
- [14] Mbow C, Rosenzweig C, Barioni LG, et al. Food security. In: Shukla PR, Skea J, Calvo Buendia E,

- Masson-Delmotte V, Pörtner H-O, et al., editors. *Climate Change and Land: An IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*. Geneva, Switzerland: IPCC; 2019
- [15] Neufeldt H, Jahn M, Campbell BM, Beddington JR, DeClerck F, De Pinto A. Beyond climate-smart agriculture: Toward safe operating spaces for global food systems. *Agriculture & Food Security*. 2013;2(1):12
- [16] Kassam A, editor. *Advances in Conservation Agriculture: Systems and Science*. Cambridge, UK: Burleigh Dodds; 2020a
- [17] Kassam A, editor. *Advances in Conservation Agriculture: Practice and Benefits*. Cambridge, UK: Burleigh Dodds; 2020b
- [18] Kassam AH, Mkomwa S, Friedrich T. Weather-proofing agriculture with conservation agriculture. In: Kassam A, Mkomwa S, Friedrich T, editors. *Conservation Agriculture for Africa: Building Resilient Farming Systems in a Changing Climate*. Wallingford, UK: CAB International; 2017
- [19] Michler JD, Baylis K, Arends-Kuenning M, Mazvimavi K. Conservation agriculture and climate resilience. *Journal of Environmental Economics and Management*. 2019;93:148-169
- [20] Ngachan V, Das A. Climate proofing of agriculture for Food security. In: Das et al, editor. *Conservation Agriculture for Advancing Food Security in Changing Climate*. New Delhi, India: Today & Tomorrow's Printers and Publishers; 2018. pp. 37-57
- [21] Williams A, Jordan NR, Smith RG, Hunter MC, Kammerer M. A regionally-adapted implementation of conservation agriculture delivers rapid improvements to soil properties associated with crop yield stability. *Scientific Reports*. 2018;8:8467
- [22] Bellotti B, Rochecouste JF. The development of conservation agriculture in Australia—Farmers as innovators. *International Soil Water Conservation Research*. 2014;2:21-34. DOI: 10.1016/S2095-6339(15)30011-3
- [23] DasTK, SaharawatYS, BhattacharyyaR, Sudhishri S, Bandyopadhyay K, Sharma AR, et al. Conservation agriculture effects on crop and water productivity, profitability and soil organic carbon accumulation under a maize-wheat cropping system in the North-western Indo-Gangetic Plains. *Field Crops Research*. 2018;215:222-231
- [24] Das TK, Ghosh S, Das A, Sen S, Datta D, Ghosh S, et al. Conservation agriculture impacts on productivity, resource-use efficiency and environmental sustainability: A holistic review. *Indian Journal of Agronomy*. 2021;66:111-127
- [25] Pradhan A, Chan C, Roul PK, Halbrendt J, Sipes B. Potential of conservation agriculture (CA) for climate change adaptation and food security under rainfed uplands of India: A transdisciplinary approach. *Agricultural Systems*. 2018;163:27-35. DOI: 10.1016/j.agry.2017.01.002
- [26] Du Z, Angers DA, Ren T, Zhang Q, Li G. The effect of no-till on organic C storage in Chinese soils should not be overemphasized: A meta-analysis. *Agriculture, Ecosystems and Environment*. 2017;236:1-11
- [27] Lal R. Sustainable intensification of China's agroecosystems by conservation

agriculture. *International Soil Water Conservation Research*. 2018b;**6**:1-12. DOI: 10.1016/j.iswcr.2017.11.001

[28] Corbeels M, de Graaff J, Ndah TH, Penot E, Baudron F, Naudin K, et al. Understanding the impact and adoption of conservation agriculture in Africa: A multi-scale analysis. *Agricultural Ecosystem Environment*. 2014;**187**:155-170. DOI: 10.1016/j.agee.2013.10.011

[29] Kuyah S, Sileshi GW, Nkurunziza L, Chirinda N, Ndayisaba PC, Dimobe K, et al. Innovative agronomic practices for sustainable intensification in sub-Saharan Africa: A review. *Agronomy for Sustainable Development*. 2021;**41**:16. DOI: 10.1007/s13593-021-00673-4

[30] Mkomwa S, Kassam A, editors. *Conservation Agriculture in Africa: Climate Smart Agricultural Development*. Wallingford, UK: CAB International; 2021

[31] Mupangwa W, Mutenje M, Thierfelder C, Nyagumbo I. Are conservation agriculture (CA) systems productive and profitable options for smallholder farmers in different agro-ecoregions of Zimbabwe? *Renewable Agriculture and Food Systems*. 2016;**32**(1):87-103

[32] Devkota M, Singh Y, Yigezu YA, Bashour I, Mussadek R, Mrabet R. Conservation agriculture in the drylands of the Middle East and North Africa (MENA) region: Past trend, current opportunities, challenges, and future outlook. *Advances in Agronomy*. 2021;**172**. DOI: 10.1016/bs.agron.2021.11.001

[33] Mrabet R, Moussadek R, Devkota M, Lal R. No-tillage farming in Maghreb region: Enhancing agricultural productivity and sequestering carbon in soils. In: Lal R, editor. *Advances in*

Soil Science: Soil Organic Matter and Feeding the Future: Environmental and Agronomic Impacts. United States: CRC Press Taylor and Francis Group; 2021

[34] Gonzalez-Sanchez EJ, Kassam A, Basch G, Streit B, Holgado-Cabrera A, Trivino-Tarradas P. Conservation agriculture and its contribution to the achievement of Agri-environmental and economic challenges in Europe. *Aims Agricultural Food*. 2016;**1**:387-408. DOI: 10.3934/agrfood.2016.4.387

[35] Mrabet R. No-tillage agriculture in West Asia & North Africa. In: Tow PG, Cooper IM, Partridge I, Birch CJ, editors. *Rainfed Farming Systems*. Dordrecht Netherlands: Springer; 2011. pp. 1015-1042

[36] Sun WJ, Canadell JG, Yu LJ, Yu LF, Zhang W, Smith P, et al. Climate drives global soil carbon sequestration and crop yield changes under conservation agriculture. *Global Change Biology*. 2020;**26**(6):3325-3335

[37] Pittelkow CM, Linquist BA, Lundy ME, Liang X, van Groenigen KJ, Lee J, et al. When does no-till yield more? A global meta-analysis. *Field Crops Research*. 2015a;**183**:156-168

[38] Su Y, Gabrielle B, Makowski D. The impact of climate change on the productivity of conservation agriculture. *Nature Climate Change*. 2021a;**11**:628-633. DOI: 10.1038/s41558-021-01075-w

[39] Knapp S, van der Heijden MGA. A global meta-analysis of yield stability in organic and conservation agriculture. *Nature Communications*. 2018;**9**:3632

[40] Jat HS, Choudhary KM, Nandal DP, Yadav AK, Poonia T, Singh Y, et al. Conservation agriculture-based sustainable intensification of cereal systems leads to energy conservation, higher productivity

and farm profitability. *Environmental Management*. 2020;**65**(6):774-786

[41] Laik R, Sharma S, Idris M, Singh AK, Singh SS, Bhatt BP, et al. Integration of conservation agriculture with best management practices for improving system performance of the rice–wheat rotation in the Eastern Indo–Gangetic Plains of India. *Agriculture, Ecosystems and Environment*. 2014;**195**:68-82

[42] Corbeels M, Naudin K, Whitbread AM, Kuhne R, Letourmy P. Limits of conservation agriculture to overcome low crop yields in sub-Saharan Africa. *Nature Food*. 2020;**1**:447-454

[43] Basch G, Kassam A, Friedrich T, Santos FL, Gubiani PI, Calegari A, et al. Sustainable soil water management systems. In: Lal R, Stewart BA, editors. *Soil Water and Agronomic Productivity, Advances in Soil Science*. Boca Raton, FL: CRC Press; 2012. pp. 229-289

[44] Gonzalez-Sanchez EJ, Veroz-Gonzalez O, Moreno-Garcia M, Gomez-ArizacMR, Ordoñez-FernandezR, Trivino-Tarradas P, et al. Climate change adaptability and mitigation with conservation agriculture. In: Kassam A, Kassam L, editors. *Rethinking Food and Agriculture: New Ways Forward*. Switzerland: Elsevier; 2020. pp. 231-246

[45] Nyagumbo I, Mupangwa W, Chipindu L, Rusinamhodzi L, Craufurd P. A regional synthesis of seven–year maize yield responses to conservation agriculture technologies in Eastern and Southern Africa. *Agriculture, Ecosystems and Environment*. 2020;**295**:106898

[46] Basch G, Saavedra N. Poupança de água na cultura do milho através da cobertura do solo. *Revista Agrotec*. 2022;**42**:16-19

[47] Mrabet R, Moussadek R, Fadlaou A, van Ranst E. Conservation agriculture

in dry areas of Morocco. *Field Crops Research*. 2012;**132**:84-94

[48] Page KL, Dang YP, Dalal RC, Reeves S, Thomas G, Wang W. Changes in soil water storage with no-tillage and crop residue retention on a vertisol: Impact on productivity and profitability over a 50-year period. *Soil Tillage Research*. 2019;**194**:104319

[49] Shahane AA, Shivay YS. Soil health and its improvement through novel agronomic and innovative approaches. *Frontier in Agronomy*. 2021;**3**:680456. DOI: 10.3389/fagro.2021.680456

[50] Lal R. Beyond COP 21: Potential and challenges of the “4 per thousand” initiative. *Journal of Soil Water Conservation*. 2016a;**71**:20A-25A

[51] Lal R. Digging deeper: A holistic perspective of factors affecting SOC sequestration. *Global Change Biology*. 2018a;**24**:3285-3301. DOI: 10.1111/gcb.14054

[52] Chatterjee S, Chakraborty R, Chakraborty H. Economic impact assessment of conservation agriculture on small and marginal farm households in eastern India. *Agricultural Economics Research Review*. 2020a;**33**:127-138

[53] Day S, Santos A, Cremonesi M, Maia L, Bartz MLC. Biodiversity management practices and benefits in conservation agriculture systems. In: Kassam A, editor. *Advances in Conservation Agriculture*. Vol. 2. Cambridge, UK: Burleigh Dodds; 2020. pp. 267-302

[54] Kassam A. Conservation agriculture for regenerative and resilient production systems. In: Uphoff N, Thies JE, editors. *Biological Approaches to Regenerative Soil Systems*. United States: CRC Press; 2022

- [55] Page KL, Dang YP, Menzies NW, Dalal RC. No-till Systems to sequester soil carbon: Potential and reality. In: Dang YP, Dalal RC, Menzies NW, editors. *No-till Farming Systems for Sustainable Agriculture: Challenges and Opportunities*. Cham, Switzerland: Springer; 2020a. pp. 301-317
- [56] Page KL, Dang YP, Dalal RC. The ability of conservation agriculture to conserve soil organic carbon and the subsequent impact on soil physical, chemical, and biological properties and yield. *Frontiers in Sustainable Food System*. 2020b;4:31
- [57] Valkama E, Kunyapiyaeva Zhapayev R, Karabayev M, Zhusupbekov E, Perego A, Schillaci C, et al. Can conservation agriculture increase soil carbon sequestration? A modelling approach. *Geoderma*. 2020;369:114298
- [58] Jayaraman S, Dang YP, Naorem A, Page KL, Dalal RC. Conservation agriculture as a system to enhance ecosystem services. *Agriculture*. 2021;11:718. DOI: 10.3390/agriculture11080718
- [59] Somasundaram J, Reeves S, Wang W, Heenan M, Dalal RC. Impact of 47 years of no tillage and stubble retention on soil aggregation and carbon distribution in a vertisol. *Land Degradation and Development*. 2017;28:1589-1602
- [60] Jat HS, Datta A, Choudhary M, Sharma PC, Yadav AK, Choudhary V, et al. Climate smart agriculture practices improve soil organic carbon pools, biological properties and crop productivity in cereal-based systems of North-West India. *Catena*. 2019;181:104059. DOI: 10.1016/j.catena.2019.05.005
- [61] Richards M, Sapkota T, Stirling C, Thierfelder C, Verhulst N, Friedrich T, et al. Conservation Agriculture: Implementation Guidance for Policymakers and Investors. Rome: FAO; 2014
- [62] Su Y, Gabrielle B, Beillouin D, Makowski D. High probability of yield gain through conservation agriculture in dry regions for major staple crops. *Scientific Reports*. 2021b;11:3344. DOI: 10.1038/s41598-021-82375-1
- [63] Demenois J, Torquebiau E, Arnoult MH, Eglin T, Masse D, Assouma MH, et al. Barriers and strategies to boost soil carbon sequestration in agriculture. *Frontiers in Sustainable Food System*. 2020;4:37
- [64] Li Y, Li Z, Cui S, Jagadamma S, Zhang QP. Residue retention and minimum tillage improve physical environment of the soil in croplands: A global meta-analysis. *Soil and Tillage Research*. 2019;194:104292
- [65] Lipper L, Thornton P, Campbell BM, Baedeker T, Braimoh A, Bwalya M, et al. Climate-smart agriculture for food security. *Nature Climate Change*. 2014;4:1068-1072
- [66] Lal R, Monger C, Nave L, Smith P. The role of soil in regulation of climate. *Philosophical Transactions on Royal Society London B Biological Science*. 2021;376(1838):20210420
- [67] Blanco-Canqui H, Ruis SJ. No-tillage and soil physical environment. *Geoderma*. 2018;326:164-200. DOI: 10.1016/j.geoderma.2018.03.011
- [68] Navarro-Pedreño J, Almendro-Candel MB, Zorpas AA. The increase of soil organic matter reduces global warming, myth or reality? *Science*. 2021;3(18):18. DOI: 10.3390/sci3010018
- [69] Veloso MG, Angers DA, Tiecher T, Giacomini S, Dieckow J, Bayer C. High

carbon storage in a previously degraded subtropical soil under no-tillage with legume cover crops. *Agriculture, Ecosystems and Environment*. 2018;**268**:15-23. DOI: 10.1016/j.agee.2018.08.024

[70] Krauss M, Wiesmeier M, Don A, Cuperus F, Gattinger A, Gruber S, et al. Reduced tillage in organic farming affects soil organic carbon stocks in temperate Europe. *Soil and Tillage Research*. 2022;**216**:105262. DOI: 10.1016/j.still.2021.105262

[71] Gonzalez-Sanchez EJ, Veroz-Gonzalez O, Conway G, Moreno-Garcia M, Kassam A, Mkomwa S, et al. Meta-analysis on carbon sequestration through conservation agriculture in Africa. *Soil & Tillage Research*. 2019;**190**:22-30

[72] González-Sánchez EJ, Moreno-García M, Kassam A, Holgado-Cabrera A, Triviño-Tarradas P, Carbonell Bojollo R, et al. Conservation agriculture: Making climate change mitigation and adaptation real in Europe. In: *European Conservation Agriculture Federation (ECAAF)*. Brussels, Belgium; 2017

[73] Dey A, Dwivedi BS, Bhattacharyya R, Datta SP, Meena MC, Jat RK, et al. Effect of conservation agriculture on soil organic and inorganic carbon sequestration and lability: A study from a rice-wheat cropping system on a calcareous soil of the eastern Indo-Gangetic Plains. *Soil Use and Management*. 2020;**36**:429-438

[74] Lal R. Enhancing ecosystem services with no-till. *Renewable Agricultural Food System*. 2013;**28**:102-114

[75] Paustian K, Lehmann J, Ogle S, Reay D, Robertson GP, Smith P. Climate-smart soils. *Nature*. 2016;**532**:49-57

[76] Paustian K, Larson E, Kent J, Marx E, Swan A. Soil C sequestration as a biological negative emission strategy. *Frontiers in Climate*. 2019;**1**:8

[77] Voltr V, Wollnerová J, Fuksa P, Hruška M. Influence of tillage on the production inputs, outputs, soil compaction and GHG emissions. *Agriculture*. 2021;**11**:456

[78] El Gharras O, El Mourid M, Boulal H. Conservation agriculture in North Africa: Experiences, achievements and challenges. In: Kassam A, Mkomwa S, Friedrich T, editors. *Conservation Agriculture for Africa: Building Resilient Farming Systems in a Changing Climate*. Wallingford, UK: CAB International; 2017. pp. 127-138

[79] Jat RK, Sapkota TB, Singh RG, Jat ML, Kumar M, Gupta RK. Seven-year-old conservation agriculture in rice-wheat rotation of eastern Gangetic Plains of South Asia: Yield trends and economic profitability. *Field Crops Research*. 2014;**164**:199-210. DOI: 10.1016/j.fcr.2014.04.015

[80] Fowler R, Rockstrom J. Conservation tillage for sustainable agriculture an agrarian revolution gathers momentum in Africa. *Soil & Tillage Research*. 2001;**61**:93-107

[81] Brown B, Nuberg I, Llewellyn R. Stepwise frameworks for understanding the utilisation of conservation agriculture in Africa. *Agricultural Systems*. 2017;**153**:11-22

[82] Rosenstock TS, Lamanna C, Chesterman S, Bell P, Arslan A, Richards M, et al. *The Scientific Basis of Climate-smart Agriculture: A Systematic Review Protocol*. Copenhagen, Denmark: CGIAR; 2016

[83] Yigezu YA, El-Shater T, Boughlala M, Devkota M, Mrabet R, Moussadek R. Can

- an incremental approach be a better option in the dissemination of conservation agriculture? Some socioeconomic justifications from the drylands of Morocco. *Soil & Tillage Research*. 2021;**212**:105067
- [84] Arenas-Calle LN, Whitfield S, Challinor AJ. A climate smartness index (CSI) based on greenhouse gas intensity and water productivity: Application to irrigated Rice. *Frontier in Sustainable Food System*. 2019;**3**:1-13
- [85] Su Z, Zhnag J, Wu W, Cai D, Lv J, Jiang G, et al. Effects of conservation tillage practices on winter wheat water-use efficiency and crop yield on the Loess Plateau, China. *Agricultural Water Management*. 2007;**87**:307-314
- [86] Chatterjee S, Ghosh S, Pal P. *Soil Carbon Restoration through Conservation Agriculture*. London, UK: IntechOpen; 2020b
- [87] Khan R, Biswas S, Kundu CK, Jana K. Effect of conservation tillage on yield and economics of fodder crops. *International Journal of Environment and Climate Change*. 2020;**10**(12):529-539
- [88] Amelung W, Bossio D, de Vries W, Kögel-Knabner I, Lehmann J, Amundson R, et al. Towards a global-scale soil climate mitigation strategy. *Nature Communications*. 2020;**11**:5427. DOI: 10.1038/s41467-020-18887-7
- [89] Bhan S, Behera UK. Conservation agriculture in India-problems, prospects and policy issues. *International Soil and Water Conservation Research*. 2014;**2**(4):1-12
- [90] Choudhary M, Rana KS, Bana RS, Ghasal PC, Choudhary GL, Jakhar P, et al. Energy budgeting and carbon footprint of pearl millet–mustard cropping system under conventional and conservation agriculture in rainfed semi–arid agro–ecosystem. *Energy*. 2017;**141**:1052-1058
- [91] FAO, IFAD, UNICEF, WFP, WHO. *The State of Food Security and Nutrition in the World 2020*. Italy: Rome, FAO; 2020
- [92] Kassam A, Friedrich T, Derpsch R. *Successful Experiences and Learnings from Conservation Agriculture Worldwide*. Online 8th World Congress on Conservation Agriculture (8WCCA). *The Future of Farming: Profitable and Sustainable Farming with Conservation Agriculture*. Bern (Switzerland); 2021
- [93] Lal R. Feeding 11 billion on 0.5 billion hectare of area under cereal crops. *Food and Energy Security*. 2016b;**5**(4):239-251. DOI: 10.1002/fes3.99
- [94] Lal R, Mohtar RH, Assi AT, Ray R, Baybil H, Jahn M. Soil as a basic Nexus tool: Soils at the Center of the Food–Energy–Water Nexus. *Current Sustainable/Renewable Energy Reports*. 2017;**4**(3):117-129
- [95] Lal R. Feeding the world and returning half of the agricultural land back to nature. *Journal of Soil and Water Conservation*. 2021;**76**(4):75. DOI: 10.2489/jswc.2021.0607A
- [96] Naab JB, Mahama GY, Yahaya I, Prasad PVV. Conservation agriculture improves soil quality, crop yield, and incomes of smallholder farmers in North-Western Ghana. *Frontiers in Plant Science*. 2017. DOI: doi.org/10.3389/fpls.2017.00996
- [97] Newton P, Civita N, Frankel-Goldwater L, Bartel K, Johns C. What is regenerative agriculture? A review of scholar and practitioner definitions

based on processes and outcomes. *Frontiers in Sustainable Food Systems*. 2020;4. DOI: 10.3389/fsufs.2020.577723

[98] Ogle S, Alsaker C, Baldock J, Bernoux M, Breidt F, McConkey BG, et al. Climate and soil characteristics determine where No-till management can store carbon in soils and mitigate greenhouse gas emissions. *Scientific Reports*. 2019;9:11665. DOI: 10.1038/s41598-019-47861-7

[99] Pooniya V, Zhiipao RR, Biswakarma N, Jat SL, Kumar D, Parihar CM, et al. Long-term conservation agriculture and best nutrient management improves productivity and profitability coupled with soil properties of a maize–chickpea rotation. *Scientific Reports*. 2021. DOI: doi.org/10.1038/s41598-021-89737-9

[100] Pittelkow CM, Liang X, Linquist BA, Van Groenigen KJ, Lee J, Lundy ME, et al. Productivity limits and potentials of the principles of conservation agriculture. *Nature*. 2015b;517:365-368

[101] Rose S, Khatri-Chhetri A, Stier M, Wiese-Rozanova L, Shelton S, Wollenberg E. *Ambition for Soil Organic Carbon Sequestration in New and Updated Nationally Determined Contributions. Analysis of Agricultural Sub-sectors in National Climate Change Strategies*. Wageningen, The Netherlands: CGIAR; 2021

[102] Vermeulen S, Bossio D, Lehmann J, Luu P, Paustian K, Webb C, et al. A global agenda for collective action on soil carbon. *Nature Sustainability*. 2019;2(1):2-4. DOI: 10.1038/s41893-018-0212-z

[103] WCCA. Declaration. The 8th World Congress on Conservation Agriculture 21-23 June. Bern, Switzerland; 2021

[104] Zerihun A, Tadesse B, Shiferaw T, Kifle D. Conservation agriculture: Maize-legume intensification for yield, profitability and soil fertility improvement in maize belt areas of Western Ethiopia. *International Journal of Plant and Soil Science*. 2014;3(8):969-985