

Research Article

Correlation of liver enzymes with serum ferritin levels in β -thalassemia major

Rameshwar L. Suman*, Anuradha Sanadhya, Pradeep Meena, Suresh Goyal

Department of Pediatrics, RNT Medical College, Udaipur, Rajasthan, India

Received: 01 June 2016

Accepted: 01 July 2016

***Correspondence:**

Dr. Rameshwar L Suman,

E-mail: sumanrl@yahoo.co.in

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Liver is the earliest site of iron deposition in transfusion dependent β -thalassemia major and iron induced liver injury is the common cause of morbidity. Liver enzymes are raised and indicative of liver injury in transfusion dependant β -thalassemia major patients. Objective of the study was to find out the correlation of serum ferritin with liver enzymes serum glutamic oxalocetic transaminase (SGOT) and serum glutamic pyruvic transaminase (SGPT) in thalassemia major children.

Methods: Hospital based study of fifty five (55) children of β -Thalassemia major in the age group of 4-20 years who were regularly transfused and were on oral iron chelators since at least one year were enrolled. Serum ferritin levels and serum SGOT and SGPT levels were estimated and results were correlated.

Results: Out of total fifty five (55) children, most were in the age group of 4 to 8 years. Mean rate of blood transfusion in subjects was 157.01ml/kg/year and mean duration of chelation therapy was 2.34 years. Serum ferritin levels were increased in β -thalassemic children with average of 2130.33 \pm 859.85ng/ml. The SGOT and SGPT were also raised significantly (p value <0.05) with mean of 71.37 \pm 24.16 IU/L and 62.3 5 \pm 25.75 IU/L respectively. The values of SGOT & SGPT becomes highly significant (p value <0.001) when serum ferritin becomes more than 2000 ng/ml. Onset of liver enzyme derangement starts at serum ferritin level of more than 1000 ng/ml We found positive correlation between serum ferritin and deranged liver enzymes (Pearson's bivariate correlation coefficient r = 0.84 \pm 84).

Conclusions: As soon as the serum ferritin level crosses the value of 1000 ng/ml and number of transfusions are more than 30, derangement in liver enzymes starts occurring in β -thalassemia major.

Keywords: Hemolytic anemia, Liver enzymes, Serum ferritin, β -Thalassemia

INTRODUCTION

β -Thalassemia major is an inherited haemoglobin disorder resulting in chronic haemolytic anemia. There is defect in globin chain synthesis leading to inability of individual to make normal haemoglobin thus leading to transfusion dependency throughout life. Frequent blood transfusion causes progressive iron overload which is major complication of treatment.¹

Thus thalassemia is secondary iron overload condition and this iron overload is both due to increased absorption of iron from gut and from frequent blood transfusions. The state of iron overload affects almost all systems of body directly or indirectly such as endocrines, liver and heart. Liver is the earliest site of iron overload in regularly transfused children and common cause of morbidity. Iron overload occurs both in hepatocytes and reticuloendothelial cells. The iron induced liver injury is characterized by development of fibrosis and eventually cirrhosis.²

Liver enzymes such as SGOT, SGPT are raised in transfusion dependent β -thalassemia major patients. Though liver biopsy is gold standard test to know iron overload state in liver but it is invasive method and T2 MRI is best non-invasive method of determining liver iron.³ Relatively simpler way of knowing the liver damage is by estimation of liver enzymes which are raised due to oxidative injury and direct toxic effect of iron on liver cells.² This study was planned to study the correlation of liver enzymes (SGOT and SGPT) with serum ferritin levels in children with transfusion dependent β -thalassemia major.

METHODS

The study was single centre tertiary hospital based prospective study carried out in Thalassemia Unit of Medical College Hospital. Fifty five (55) β -thalassemia major children of 4-20 years who were on regular chelation therapy with Deferasirox since at least 1 year were taken for study. All children were transfusion dependent at the rate of once or twice a month. These children were seronegative for HIV, HCV and HbsAg and were not having any other significant medical or surgical condition. Other than routine Complete Blood Count (CBC), venous blood was taken for serum ferritin and serum SGOT, SGPT estimation after blood transfusion.

Serum ferritin level was measured with Elecys 210. Ferritin specific antibody sandwich principle is used by this test which takes 18 minutes. The sample passes through two faces in which two different ferritin specific antibody coats the ferritin. Serum SGOT and SGPT estimation was done by semi auto analyser. Serum level of SGOT >50 IU/L and SGPT >40 IU/L were considered abnormal.

Institutional ethical committee permission was sought and written consent was taken from parents of children before entering into the study. Results were correlated and statistical analysis was done using Pearson's bivariate coefficient test.

Table 2: Liver enzymes at different serum ferritin levels.

S. Ferritin(ng/ml) (No.)	SGOT	SD	p Value	SGPT	SD	p Value
<1000 (3)	45.34	25.37	>0.05	35.15	24.2	>0.05
1000 - <2000 (23)	53.45	24.17	>0.05	45.92	23.8	>0.05
2000 - <3000 (20)	79.94	24.57	<0.001	71.78	23.7	<0.001
>3000 (9)	106.75	24.71	<0.001	92.47	23.7	<0.001
Mean 2130±859.85	71.37	24.16	<0.05	62.35	23.7	<0.05

When liver enzymes were analysed at different serum ferritin levels, they were continuously rising as the serum ferritin was increasing and after the level of 2000 ng/ml, there was steep rise in liver enzyme levels as shown in Figure 1. When serum ferritin levels were correlated with

RESULTS

Out of fifty five (55) children, majority of children 33 (60%) were in the age group of 4-8 years. Only 11 (20%) children were more than 12 years old. Female preponderance in subjects 31 (56.4%) was seen. Most of the patients belonged to Hindu religion 48 (87.3%) and rest were Muslims. Mean Hemoglobin Concentration after Blood transfusion was 9.7gm/dl. Various basic variables of β -thalassemia patients were shown in Table 1.

Table1: Basic general variables in β -thalassemic children.

Variables	Mean	Standard deviation
Age (year)	8.80	3.88
Age at Diagnosis(in Months)	29.91	31.04
No. of blood transfusions	84.65	85.71
Rate of Blood transfusion(in ml/Kg/yr)	157.02	21.33
Dose of Deferasirox (in mg/kg/day)	34.40	26.86
Duration of chelation therapy (in years)	2.35	1.87

Mean serum ferritin was 2130±859.85 ng/ml. 23 (41.81%) patients revealed serum ferritin levels between 1000 and 2000 ng/ml while 20 (36.36%) children had even higher serum ferritin levels between 2000 and 3000 ng/ml inspite of chelation. Only 3 (5%) patients maintained serum ferritin levels <1000 ng/ml. Mean SGOT was 71.37±24.16 IU/L and mean SGPT was 62.35±23.75IU/L which were higher than normal value (p <0.05) Serum liver enzymes at various levels of serum ferritin levels were as shown in Table 2. The statistically highly significant difference in SGOT and SGPT was observed once the serum ferritin crossed level of 2000 ng/ml (p Value <0.001).

no. of blood transfusions, we found that after 30 blood transfusions the serum ferritin increases to large extent and it continues to rise as no. of transfusion further increases. The pearson bivariate coefficient correlation was positive (r = +0.33) as depicted in Figure 2.

When we correlated serum ferritin levels with liver enzymes we found a positive correlation. As the serum ferritin level increases more than 1000 ng/ml, the levels of SGOT and SGPT rise significantly as shown in Figure 3.




Figure 1: Trend of liver enzymes with rising serum ferritin.




Figure 2: Correlation of serum ferritin with no. of blood transfusions in beta-thalassemic children.




Figure 3: Correlation of liver enzymes (SGOT, SGPT) with serum ferritin in beta-thalassemic children.

DISCUSSION

As liver is the earliest organ affected by iron overload in thalassemia children and serum SGOT and SGPT are raised due to peroxidative injury and direct toxic effect of iron on liver cells. So this study was conducted to know

the effect of iron toxicity on liver enzymes and their correlation with serum ferritin levels.

In present study the serum ferritin concentration was very high in beta-thalassemic children inspite of chelation therapy. A positive correlation was noted between number of transfusions and serum ferritin level with correlation coefficient ($r=+.33$). As Iron deposition in liver takes place, its functions are affected which are predicted by raised SGOT and SGPT. SGOT and SGPT were raised significantly (p Value <0.05) and continue to rise as ferritin crosses 1000 ng/ml. There was a positive correlation between serum ferritin and liver enzymes (Pearson's bivariate correlation coefficient $r=+0.87\pm.84$).

Patients with iron overload have increased levels of thiobarbituric acid reactant and increased hepatic level of aldehyde protein adduct indicating lipid peroxidation. Collagen formation and portal fibrosis starts as early as 2 years of onset of transfusion. In absence of chelation, cirrhosis may develop in first decade of life.⁴

Chekir KA et al conducted a study on 56 thalassemic children and determined various metabolic parameters. The study suggested that in beta thalassemia first organ impaired was liver. Plasma thiobarbituric acid reactive substances (TBARS) were significantly raised leading to deranged liver functions.⁵ Similar finding were seen by Ameli M et al by study conducted in Iran (2006). In their study they found that mean serum ALT (Alanine aminotransferase) was significantly high in thalassemic children with high serum ferritin and high transfusion index.⁶

Sedigheh shams et al noted the similar findings from study in Iran with significantly raised liver enzymes (ALT, AST) in homozygous thalassemia major patients than in controls.⁷ Asharaf soliman et al observed during a study that some disturbances occur in liver functions in hepatitis negative thalassemia patients with iron overload. Use of Desferrioxamine is associated with decrease in liver enzymes.⁸ Barton JC et al noted similar results as serum ferritin increases liver enzymes also increases.⁹ Limitation of our study is that we did not included controls which could have better given us comparative results as compared to normal subjects.

CONCLUSION

From present study it is concluded that poorly chelated beta-thalassemics had hepatic dysfunctions in the form of abnormal liver enzymes and this starts occurring as soon as serum ferritin level crosses 1000ng/ml and becomes highly significant after the serum ferritin levels of >2000 ng/ml due to potential toxic effect of iron overload on hepatocytes. Hence we recommend that liver functions should be carefully monitored in patients with transfusion dependent beta-thalassemia major. Every beta-thalassemic child should be screened for liver dysfunction as soon as

serum ferritin level exceeds 1000ng/ml or number of transfusion >30.

What is already known: Raised serum ferritin leads to liver dysfunction but it is not clear at what levels it starts.

What this study adds: Liver dysfunction starts when serum ferritin increases beyond 1000 ng/ml and number of blood transfusions >30.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

1. Michel RD, Melissa JF, Elliott. Thalassemia syndrome. In kleigman RM, Stanton BF, St Geme JW, Behrman RE (editors) Nelson's textbook of paediatrics. 20th edition. Elsevier publication; 2015:2349-53.
2. Richa J, Sachdeva A. Iron overload and its manifestations. In: sachdeva A, Jain R, Aggarwal RK, Yadav SP, Broker A, eds. Manual of Thalassemia IAP. Pediatric hemato oncology chapter of IAP. 2009:76-115.
3. Hillenbra CM, Loeffler RB, Carville M. Mc. Evaluation of hepatic iron concentrations by T2 magnetic resonance imaging in patients with iron overload: comparison of T2 measurements at 1.5T and 3T and validation with liver biopsies. Proc Intl Soc Mag Reson Med. 2005;105:855-61.
4. Livrea MA, Tesoriere L, Piantaudi AM. Oxidative stress and antioxidant status in beta-thalassemia major: iron overload and depletion of lipid-soluble antioxidants. Blood .1996; 88(9):3608-14.
5. Kassab CA, Laradi S. Oxidant antioxidant status and metabolic data in patients with beta-thalassemia. Pediatr Blood Cancer. 2003;338.
6. Ameli M, Besharati S, Nemati K, Zamani F. Relationship between elevated liver enzymes with iron overload and viral hepatitis in thalassemia major patients in northern Iran. Saudi Med J. 2008;29(11):1611-15.
7. Sedigheh S, Mohammed TA, Maryam MM, Leili K, Heshmat I, Fahimah J et al. Evaluation of serum insulin, glucose, lipid profile and liver functions in beta-thalassemia major patients and their correlation with iron overload. Labmedicine. 2010;41:486-9.
8. Asharaf S, Mohammad Y, Fauzia al Y, Lolwa al N, Noora al M, Aml S, Vincenzo de S. Longitudinal study on liver functions in patients with thalassemia major before and after desferrioxamine therapy. Mediterranean journal of hematology and infectious diseases. 2014;6.
9. Barton JC. Chelation therapy for iron overload Curr Gastroenterology. 2007;23:74-82.

Cite this article as: Suman RL, Sanadhya A, Meena P, Goyal S. Correlation of liver enzymes with serum ferritin levels in β -thalassemia major. Int J Res Med Sci 2016;4:3271-4.