20500904, 2022, 12, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cct3.6694 by Universität Bern, Wiley Online Library on [08/01/2023]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms

CASE VIDEO

Endoscopic resection of Intralabyrinthine Schwannoma followed by Cochlear implantation

Vera Lohnherr^{1,2} | Lukas Anschuetz^{1,3} | Georgios Mantokoudis^{1,3} | Marco Caversaccio^{1,3} | Stefan Weder^{1,3}

²Department of Otorhinolaryngology, Head and Neck Surgery, University Hospital Heidelberg, Ruprecht-Karls-University, Heidelberg, Germany

³Hearing Research Laboratory, ARTORG Center for Biomedical Engineering Research, University of Bern, Bern, Switzerland

Correspondence

Vera Lohnherr, Department of Otorhinolaryngology, Head and Neck Surgery, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland.

Email: vera.lohnherr@med.uniheidelberg.de

Abstract

We present the case of a 65 years old patient who developed a complete, sensorineural hearing loss on the right side due to an intravestibular schwannoma. Our video shows a transcanal, endoscopic approach with complete schwannoma removal on the right side, subsequent e-BERA recordings, and cochlear implantation.

KEYWORDS

cochlear implantation, endoscopic ear surgery, Intralabyrinthine Schwannoma

We present the case of a 65 years old patient who developed a complete, sensorineural hearing loss on the right side due to an intravestibular schwannoma. Our video shows a transcanal, endoscopic approach with complete schwannoma removal on the right side, subsequent e-BERA recordings, and cochlear implantation (case Video S1). The use of an endoscope allows improved visualization of anatomical structures. If cochlear and neural structures remain intact, simultaneous cochlear implantation with hearing restoration is feasible.¹

AUTHOR CONTRIBUTIONS

Lohnherr V: video editing, abstract. Anschuetz L: surgery, review and editing. Mantokoudis G: review and editing. Caversaccio M: resources, supervision. Weder S: surgery, review and editing.

ACKNOWLEDGMENT

None.

CONFLICT OF INTEREST

There are no conflicts of interest, financially, or otherwise.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. Clinical Case Reports published by John Wiley & Sons Ltd.

-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

¹Department of Otorhinolaryngology, Head and Neck Surgery, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

20500904, 2022, 12, Downloaded from https://onlinelibrary.wiley.com/doi/10.1002/cct3.6694 by Universität Bern, Wiley Online Library on [08/01/2023]. See the Terms

and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

CONSENT

Written informed consent was obtained from the patient to publish this report in accordance with the journal's patient consent policy.

ORCID

Vera Lohnherr https://orcid.org/0000-0002-5655-7671

REFERENCE

 Choudhury B, Carlson ML, Jethanamest D. Intralabyrinthine Schwannomas: disease presentation, tumor management, and hearing rehabilitation. J Neurol Surg B Skull Base. 2019;80(2):196-202.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Lohnherr V, Anschuetz L, Mantokoudis G, Caversaccio M, Weder S. Endoscopic resection of Intralabyrinthine Schwannoma followed by Cochlear implantation. *Clin Case Rep.* 2022;10:e06694. doi:10.1002/ccr3.6694