

The Heart Is at Risk: Understanding Stroke–Heart–Brain Interactions with Focus on Neurogenic Stress Cardiomyopathy—A Review

Mairi Ziaka,^aAristomenis Exadaktylos^b

^aDepartment of Internal Medicine, Thun General Hospital, Thun, Switzerland

^bDepartment of Emergency Medicine, Inselspital, University Hospital, University of Bern, Bern, Switzerland

In recent years, it has been convincingly demonstrated that acute brain injury may cause severe cardiac complications—such as neurogenic stress cardiomyopathy (NSC), a specific form of takotsubo cardiomyopathy. The pathophysiology of these brain–heart interactions is complex and involves sympathetic hyperactivity, activation of the hypothalamic–pituitary–adrenal axis, as well as immune and inflammatory pathways. There have been great strides in our understanding of the axis from the brain to the heart in patients with isolated acute brain injury and more specifically in patients with stroke. On the other hand, in patients with NSC, research has mainly focused on hemodynamic dysfunction due to arrhythmias, regional wall motion abnormality, or left ventricular hypokinesia that leads to impaired cerebral perfusion pressure. Comparatively little is known about the underlying secondary and delayed cerebral complications. The aim of the present review is to describe the stroke–heart–brain axis and highlight the main pathophysiological mechanisms leading to secondary and delayed cerebral injury in patients with concurrent hemorrhagic or ischemic stroke and NSC as well as to identify further areas of research that could potentially improve outcomes in this specific patient population.

Correspondence: Mairi Ziaka
Department of Internal Medicine, Thun
General Hospital, Krankenhausstrasse 12,
3600, Thun, Switzerland
Tel: +0041636582967
E-mail: mairi.ziaka@gmail.com
<https://orcid.org/0000-0001-5298-7925>

Received: July 4, 2022
Revised: November 4, 2022
Accepted: November 14, 2022

Keywords Stroke; Blood–brain barrier; Stroke–heart–brain interactions; Neuroinflammation; Neurogenic stress cardiomyopathy; Takotsubo cardiomyopathy.

Introduction

In recent decades, experimental and clinical studies have demonstrated that there is an “invisible” cross-talk between the heart and the brain.¹ Even in the absence of cardiac disease,² acute brain injury (ABI) can induce cardiac dysfunction, which in turn can increase mortality and lead to potentially long-lasting cardiac complications, such as heart failure.³ Indeed, ABI such as ischemic stroke, seizures,⁴ aneurysmal subarachnoid hemorrhage (SAH),⁵ and traumatic brain injury (TBI)⁶ may result in challenging heart pathologies, such as neurogenic stress cardiomyopathy (NSC).^{4,7,8}

In addition to the axis from the brain to the heart in patients with acute neurological injury, “heart to brain” connections must be seriously considered,⁹ especially in patients with cardiac dysfunction secondary to acute brain damage. Unfortunately, the pathophysiology of these heart–brain interactions is still poorly understood¹ and various factors have been suggested to explain the observed “heart to brain” cross-talk, including hemodynamic disturbances and impaired cerebral perfusion pressure (CPP) with subsequent cerebral hypoperfusion, development of secondary cerebral cardioembolic complications, oxygenation disturbances, neurohormonal mechanisms, systemic and cerebral inflammation,¹⁰ disruption of the blood–brain barrier (BBB), and activation

of glial cells.^{2,11-15}

The present review contains a detailed discussion of heart-brain connection in patients with NSC due to isolated ABI with focus on hemorrhagic and ischemic stroke. In addition, the review highlights underlying pathophysiological mechanisms regarding the stroke-heart-brain axis leading to secondary and delayed cerebral injury and identifies further areas of research that could potentially improve outcomes in this specific patient population.

NSC: a complex definition

Cardiac dysfunction from neurological injury has been classified as neurogenic stunned myocardium (NSM),⁷ which refers to a neurocardiogenic interaction in patients with ABI and results in reversible myocardial dysfunction^{7,16} or NSC (i.e., takotsubo cardiomyopathy [TTC], secondary to neurological disorders).¹⁷⁻¹⁹ However, it is still unclear whether NSM and NSC are distinct pathological entities or divergent manifestations of the same pathophysiological process.¹ In addition, the terminology of myocardial dysfunction in patients with acute neurological injury is confusing and is still the subject of scientific debate.¹⁹ Furthermore, it has been suggested that, since the terminology "stunned myocardium" generally refers to reversible myocardial dysfunction

in the context of coronary artery occlusion with subsequent reperfusion, it would be preferable to use the term "stressed myocardium" in order to emphasize the pathophysiological causal association of the catecholamine storm phenomenon observed in these patients.⁵

Moreover, the term "neurogenic stunned myocardium" comes from a time when clinicians were unfamiliar with the different types of TTC. As the definition of NSM is in accordance with the diagnostic criteria of TTC, a recent review¹⁹ has recommended the use of the term "takotsubo cardiomyopathy secondary to neurological disorders," together with a reference to its exact type, that is apical, basal, midventricular, or focal.¹⁹

The pathophysiology of neurogenic cardiac injury in patients with and without pre-existing heart diseases is complex^{18,19} and includes the principle pathophysiological mechanisms, such as autonomic dysregulation, excessive catecholamine release, myocyte injury, mitochondrial dysfunction, and perpetuated inflammatory cascades.⁹ Given the etiological association of the development of TTC in patients with acute neurological or psychiatric disease (Figure 1)²⁰ and in order to emphasize the underlying pathophysiological mechanisms, the authors of the present study propose the use of the term "neurogenic stress cardiomyopathy" as a specific variant of TTC in acutely affected neurological patients.




Figure 1. Pathophysiology of neurogenic stress cardiomyopathy and secondary brain damage in patients with acute brain injury. BBB, blood-brain barrier; HPA, hypothalamic-pituitary-adrenal.

NSC in patients with stroke

Pathophysiology of cardiac dysfunction in stroke

As mentioned above, the pathophysiology of neurogenic cardiac injury is complex¹⁹ and, in addition to the above-mentioned pathophysiological mechanisms, includes stroke-induced disturbances of hypothalamic-pituitary-adrenal (HPA) axis and perpetuated local and systemic inflammatory cascades.^{9,18,19,21} The consequences are increased myocardial demand, coronary microvascular spasm, acute and chronic structural and functional abnormalities, and molecular signaling myocardial alterations including microvascular endothelial disturbances, altered calcium metabolism, myocyte injury, contraction band necrosis, myocardial inflammation and fibrosis, oxidative stress, altered mitochondrial function, apoptosis and autophagy, and impaired gene expression.^{19,21-23}

Regarding NSC, despite extensive research, many questions remain poorly understood. However, the most prevailing theory trying to explain its pathophysiology is a sympathetic hyperstimulation due to physical and emotional triggers with an excessive release of catecholamines leading to direct catecholamine-toxicity, epicardial and microvascular coronary vasoconstriction, and adrenoceptor-mediated damage.^{22,24} Indeed, experimental models of SAH suggest an abrupt elevation of catecholamines within minutes following stroke, which positively correlates with the peak values of troponin T and creatine kinase-MB,²⁵ phenomenon that may explain the increased frequency and severity of cardiovascular complications in the first three days after a stroke onset.²⁶⁻²⁸ The typical distribution of left ventricular (LV) β -adrenergic receptors with apical-basal gradients and sympathetic innervation might explain the LV wall motion abnormalities seen in NSC as the apex characterized by increased density of receptors.²⁹ Indeed, the specific topographic distribution of β -receptors seems to be largely causally related to the increased apical responsiveness observed in states of catecholamine excess and especially epinephrine.² At the cellular level, the local action of catecholamines is facilitated by the close proximity of cardiomyocytes and nerve endings leading to impaired calcium homeostasis and altered β -adrenergic signal transduction, which in turn lead to myocardial contraction band necrosis and impaired coronary microcirculation.^{22,30,31} Moreover, supra-physiological levels of epinephrine may have a negative inotropic effect, as they contribute to the switching of β_2 -adrenergic receptors from canonical stimulatory G-protein-activated cardiostimulant to inhibitory G-protein-activated cardiodepressant pathways.³² The major pathophysiological aspect of altered function within the brain-heart axis in the pathogenesis of stroke-associated NSC is fur-

ther supported by experimental and clinical research evidence which indicates a correlation between the development of cardiac dysfunction and ischemic events involving the insular cortex, especially the right insular cortex.³³⁻³⁵ Indeed, current neuroimaging studies using functional magnetic resonance and positron emission tomography have confirmed the insular cortex as a cardinal regulator of the central autonomic nervous system.³⁶ The latter observation is crucial if taken into account that blood supply of insular cortex comes from the middle cerebral artery making it particularly susceptible to cerebrovascular disease.³⁷

Finally, systemic inflammation and more specifically the excessive release of interleukin (IL)-1 along with neurohormonal mechanisms seem to play a cardinal role in the development of NSC. Similarly to catecholamines, cytokine release occurs shortly after stroke; despite that elevation of their plasmatic levels lasts only few hours to days cytokine-induced myocardial remodeling can be preserved for months.^{19,22,23} Further, parasympathetic dysfunction observed in patients with severe neurological damage such as SAH, intracerebral hemorrhage, and acute ischemic stroke may lead to myocardial damage through uncontrolled release of inflammatory mediators.³⁸ Additionally, recent research suggests that microglia, the brain-resident immune cells, play a pivotal role in regulating sympathetic nervous system activity (i.e., one of the main contributors of pathological cardiac remodeling) by release of a variety of substances, including chemokines, cytokines, and growth factors.³⁹ Moreover, massive release of epinephrine and norepinephrine has also a neurohormonal basis through the activation of the HPA axis and release of adrenocorticotrophic hormone, sympathetic stimulation of the adrenal glands, elevation of plasmatic cytokines affecting the HPA axis leading to peripheral arterial vasoconstriction, sudden LV afterload increase, and elevation of LV end-systolic pressure.⁴⁰⁻⁴³ Future research regarding the development of new therapeutic targets, such as blockade of β -adrenergic receptors or IL-1-inhibition combined with the precise elucidation of the exact pathophysiological pathways, may be crucial in preventing and managing cardiac damage in patients with stroke.

Clinical presentations of NSC in patients with stroke

It is generally accepted that stroke patients develop a two-way interaction between brain and heart, which makes them highly susceptible to severe cardiovascular complications.⁴⁴ Although emotional distress⁴⁵ and SAH⁴⁶ are the major causes of NSC, cardiac dysfunction may also occur after various neurological pathologies, such as TBI, hemorrhagic and ischemic stroke, central nervous system⁴⁷ infections, and epilepsy.⁴⁸ In general, cardiac manifestations in patients with stroke and NSC are highly vari-

able and are associated with electrocardiographic (ECG) signs, such as tachyarrhythmias, QT-interval prolongation, T-wave inversion, and ST-depression; echocardiographic findings in the form of LV regional wall motion abnormalities; or troponemia and elevated N-terminal prohormone of brain natriuretic peptide (NT-proBNP).^{5,48,49}

The exact incidence of cardiac dysfunction in patients with SAH is difficult to be accurately determined given its imprecise definition and the different diagnostic criteria used in different studies. Indeed, if ECG criteria are used, its incidence seems to be in 100% of patients.⁸ If echocardiographic criteria are used, it seems that 10%–15% of patients develop global systolic dysfunction with a left ventricular ejection fraction (LVEF) less than 50%, while almost 30% of patients present wall motion abnormalities.^{50,51} Predictors of the development of NSC in patients with SAH include higher Hunt-Hess grade on presentation,⁵² older age,^{52,53} female sex,^{53–55} tobacco abuse,^{52,55} and aneurysms of the posterior circulation.⁵⁵

For patients with acute ischemic stroke, the clinical spectrum of cardiac dysfunction in the context of stroke includes ischemic and non-ischemic myocardial injury with cardiac troponin (cTn)-elevation, acute myocardial infarction, heart failure and NSC, ECG abnormalities, arrhythmias, and sudden cardiac death¹⁹ with variable overlap between the aforementioned disease entities.²¹ Recent research highlights that the frequency and severity of these functional and structural alterations peak within three days after the stroke onset^{26–28} and are proportionally associated with the severity of ischemic stroke and neurological deficits.³⁴ Moreover, experimental and clinical studies suggest that specific parameters such as location of ischemic injury influence the effect on the cardiovascular system.^{56,57} For example, cardiac injury and dysfunction occur more frequently if the ischemic injury is localized within the central autonomic network and, especially, if the ischemic injury affects the insular cortex,⁵⁸ which has been linked with the development of NSC.^{33,34}

Despite considerable inconsistencies in the reporting criteria, acute ECG alterations are more frequently associated with intracerebral hemorrhage or SAH than acute ischemic stroke being additionally more common among patients with troponemia.^{59,60} The two major categories of ECG changes during neurogenic heart disease are dysrhythmias (e.g., atrial and ventricular tachyarrhythmias, sinus bradycardia and tachycardia) and repolarization changes.^{1,2} The most common repolarization changes are seen in the anterolateral and inferolateral leads,² including prolongation of the QT-interval and changes in ST-T.⁶¹ In patients with NSC, the most frequent ECG findings are prolongation of the QT-interval and T-wave inversion.⁶²

Moreover, several studies have reported elevated levels of cTn—

specific biomarkers of myocardial injury—and NT-proBNP. These follow acute and severe brain injury, as for example SAH and acute ischemic stroke.⁶³ With highly sensitive analyses, elevations in cTn can be observed in 30%–60% of ischemic stroke patients⁶⁴ and in 20%–40% of patients after intracerebral hemorrhage and SAH with or without coronary symptoms;⁶⁵ their levels are correlated with clinical severity of aneurysmal SAH and are associated with persistent ECG and structural echocardiographic alterations, as for example QTc-prolongation, ventricular tachycardia/fibrillation, decreased ejection fraction (EF), and regional wall motion abnormalities.^{1,66} Increased troponin levels may be observed in 30% of patients with TBI, and are also associated with a worse prognosis.¹ Furthermore, a variety of studies have reported elevated levels of NT-proBNP in patients with ischemic stroke, SAH, and TBI—conditions that are also correlated with the severity of the disease.^{50,67–70} Brain natriuretic peptide (BNP) has potent natriuretic and vasodilator actions, and can decrease cerebral blood flow (CBF), cause hyponatremia, and be elevated in patients with SAH, who develop delayed ischemic injury due to vasospasm.⁷⁰ Additionally, neuroscience literature indicates that elevated levels of BNP are associated with troponemia, wall motion abnormalities, diastolic dysfunction, depressed LVEF, and pulmonary edema.⁵⁴ However, BNP is a neurohormone, that is not only secreted in the ventricles in response to volume and pressure overload, but also in the brain—specifically in the thalamus, hypothalamus, brain stem, and cerebellum.^{71–73}

Cardiac systolic impairment is not infrequent after severe neurological injury as it appears in 2% to 30% of patients with SAH,⁷⁴ in 13% to 29% of patients with ischemic stroke,⁷⁵ and in 22% of patients with TBI.⁷⁶ Furthermore, echocardiographic abnormalities include diastolic dysfunction,⁷⁶ regional wall motion abnormalities,⁷⁷ and NSC.⁷ The most commonly observed alterations in the echocardiographic patterns are of the apical type; in 25% of patients, there is right ventricular involvement⁶² and dysfunction of the basal and midventricular segments.⁶¹

Potential secondary brain alterations in patients with NSC

NSC is a specific form of TTC and a well-documented complication in patients with severe ABI.^{19,78} There is growing evidence of brain and neural changes in patients with NSC, which could exacerbate pre-existing cerebral damage. For many years, researchers have recognized that brain activation is implicated in the pathophysiology of TTC.^{24,79} Indeed, in a recent study using brain single photon emission computed tomography to assess CBF, Suzuki et al.²⁴ reported significant increases in CBF in the hippocampus, brainstem, and basal ganglia, with an associated

CBF decrease in the prefrontal cortex; these findings may partially explain the brain pathophysiology associated with TTC. As TTC may be protracted,²⁴ it is not surprising that hyperperfusion of the hippocampal and basal ganglia can progress to pathological elevation of neural activity and neuroinflammation. In addition to these findings, Klein et al.⁸⁰ have described homogenous structural and functional alterations in brain regions that are mainly associated with cardiac control, emotional processing, as well as cognitive and sensorimotor functions—suggesting communication between the heart and the brain. In a similar vein, studies used magnetic resonance imaging to compare the limbic system in TTC patients and healthy controls report structural and connective differences in patients, which may be associated with dysfunction of the autonomous nervous system, emotional processing, and cognition.^{81,82} However, it remains still unclear whether these anatomical alterations are the cause or the consequence of cardiac dysfunction. Templin et al.⁸¹ have recently compared 15 TTC patients and 39 healthy controls by means of functional MRI and found that TTC patients showed hypoconnectivity of the central brain regions of the limbic network. This may indicate that autonomic-limbic integration has a role in the pathophysiology of TTC and even in its specific variant NSC, which could result in dysfunction of emotional processing and of the autonomous nervous system.

Patients with NSC may present symptoms of acute heart failure as a result of severe myocardial dysfunction and injury that can lead to moderate to severe LV dysfunction.¹ On the other hand, there is increasing evidence that cognitive impairment is frequent in patients with heart failure.⁸³ Indeed, the term “cardiogenic dementia” has been employed to describe the co-existence of “heart and brain failure” after heart disease.⁸⁴ Nevertheless, the exact mechanisms of this relationship remain poorly understood and several pathophysiological associations have been proposed, including systemic hypotension with subsequent poor brain perfusion and occult cardiogenic emboli,^{85,86} as it is present in patients with either reduced or preserved EF.¹⁰

Pathophysiology of secondary brain injury in patients with NSC

Cerebral hypoperfusion

Despite its small size, the brain is a highly metabolic organ, that accounts for about 20% of the body's consumption of oxygen, while it receives 15% of the total cardiac output.⁸⁷⁻⁹⁰ Thus the brain is vulnerable to cardiac dysfunction, which could ultimately lead to impairment of cerebral flow. On the other hand, however, heart failure is a systemic and heterogeneous disease associated with fluctuations in blood flow and blood pressure, and may lead

to hypoperfusion of peripheral organs and systems, including the brain.⁷⁸ It is important that a growing body of evidence supports the role of cerebral hypoperfusion in the development of cognitive impairment in patients with low cardiac output.⁹¹⁻⁹³ More specifically, experimental and clinical data indicate that brain hypoperfusion results in energetic and metabolic dysregulation, loss of cholinergic receptors, abnormalities in protein synthesis, degeneration of brain capillaries, and neuronal damage that extends from the hippocampus to the temporo-parietal cortex.⁹⁴ Moreover, brain hypoperfusion also has a notorious effect on the volume of gray matter—probably due to the latter's high metabolic demands—and is associated with dysfunction of various cortical areas, resulting in cognitive deterioration.⁹⁵ These findings are consistent with the hypothesis of a “head to heart” link in patients with Alzheimer's disease and support the connection between cerebral hypoperfusion and cognitive impairment.⁹⁶

As mentioned previously, the brain receives 15% of cardiac output,^{89,90} is highly vascularized, and consumes about 20% of the total oxygen budget.⁹⁶ Under normal conditions, CBF is approximately 50 mL/min/100 g brain tissue and remains stable within a wide range of mean arterial pressure (MAP), by means of a variety of vascular and neurogenic factors, which ensure reliable autoregulation of the cerebrospinal fluid.¹⁰ However, this homeostatic mechanism is impaired or absent in severe ABI, which can lead to cerebral hypoperfusion, and this may reach ischemic levels and exacerbate primary injury.⁹⁷ A key principle of managing critically ill patients with ABI is the maintenance of brain homeostasis, which ensures oxygen delivery to the brain by stabilizing CBF.⁹⁸ Cerebral oxygenation critically depends on a close interaction between CBF, arterial content of oxygen, and cerebral oxygen metabolism.¹¹ Under non-pathological conditions, the brain autoregulates its blood flow to maintain a constant flow regardless of changes in central hemodynamics.⁹⁹ CBF is directly related to CPP, which is defined as the difference between mean arterial and intracranial pressures (MAP and ICP, respectively).⁹⁹ In healthy individuals with intact homeostatic mechanisms, the values of CPP remain stable between 50–150 mm Hg by increasing resistance when MAP and CPP increases, and decreasing cerebrovascular resistance when MAP or CPP decreases.⁹⁷ This homeostatic mechanism is often impaired in patients with ABI, where ICP can be elevated, and perfusion pressure may then change. Thus, the brain becomes vulnerable to systemic hemodynamic changes.^{11,100} In patients with ABI, brain swelling and cerebral edema, cerebrospinal fluid circulation, and neuroinflammation could additionally influence autoregulation.¹¹

It may then be concluded that it is extraordinarily important to understand how pathophysiological effects on cardiac dysfunction, including decreased cardiac output, hypotension, and ar-

rhythmias on CBF, may exacerbate primary brain injury (Figure 1).⁹

Oxygenation disturbances

In patients with ABI, it is well-documented that complications may be associated with the development of NSC² and neurogenic pulmonary edema,¹⁴ and may be associated with severe hypoxemia. As highlighted previously, arterial oxygen content is a significant regulator of CBF.¹¹ Moreover, carbon dioxide (CO₂) levels are of crucial importance in maintaining cerebral autoregulation,¹⁰¹ especially in mechanically ventilated patients with ABI.^{14,102} It has also been shown that CO₂ levels fluctuate in patients with acute heart failure, which may lead to vasoconstriction or vasodilatation of cerebral arteries.^{103,104} This point is further supported by findings showing that acute hypoxemic hypoxia (i.e., in seconds to hours) may cause cerebral vasodilation and reduce cerebral resistance, thus leading to an increase in CBF. However, vascular reactivity is additionally affected by hypoxia-induced chemoregulation, as indicated by studies showing that compensatory hyperventilation could lead to hypocapnia, cerebral vasoconstriction, and a drop in CBF.¹⁰¹ Moreover, a synergistic effect has been observed between hypoxemia and hypercapnia, which is associated with the potent vasodilating properties of CO₂.¹¹

Taken together, the pathophysiological mechanisms influence CBF through hypoxemia, and changes in blood gas are complex and multifactorial. Even though a detailed examination of all underlying mechanisms related to the role of hypoxemia and hypercapnia on brain function falls outside the scope of the present review, recent high-quality reviews are currently available on the specific topics.^{11,97,101}

Neurohormonal axis

In recent decades, it has been convincingly demonstrated that activation of the HPA axis is a key component of compensatory anti-inflammatory mechanisms in critically ill patients and involves acute elevation of free cortisol and suppression of the adrenocorticotropic hormones.^{105,106} Moreover, there is mounting evidence indicating that this neurohormonal activation plays a pivotal role in the interaction between severe heart disease, structural brain changes, and functional alterations, such as impaired cognition.¹⁰ Indeed, higher levels of cortisol have been found in patients with heart failure than in healthy controls.¹⁰⁷ This is further supported by evidence showing that the incidence of depression and cognitive impairment is significantly higher in heart failure patients than in healthy adults.¹⁰⁸ Moreover, clinical trials in healthy adults demonstrate that even transient hypercortisolism may lead to cognitive impairment, reversible decreases in verbal declarative memory, and altered spatial thinking tasks.^{109,110} On the other hand, prolonged exposure to high cortisol levels can

cause lasting alterations and atrophy of specific brain regions, through decreased neurogenesis and increased oligodendrogenesis.¹¹¹ In addition, stress-induced hypercortisolism may disrupt the plasticity and structural integrity of the hippocampus and the prefrontal cortex, and thus contribute to long-lasting impairments in memory and cognition in critically ill patients.¹⁰⁶ Lastly, high glucocorticoid levels may cause neural damage through activation of systemic inflammatory responses and exacerbation of neuroinflammation.¹⁰⁶

The role of inflammation

Systemic elevation of inflammatory cytokines and oxidative stress have been implicated in the pathophysiology of brain dysfunction after heart disease (Figure 1).^{112,113} Experimental studies in animal models of acute myocardial dysfunction indicate that neuroinflammation may develop after acute heart injury and suggest that immune activation interconnects heart and brain dysfunction.¹¹⁴ Moreover, it has been convincingly demonstrated that several heart diseases (e.g., myocarditis, myocardial infarction, and heart failure) are associated with activation of inflammatory and immune responses triggered by myocardial injury.^{95,115} Moreover, impaired cardiac function may be associated with cerebral hypoperfusion that exacerbates brain inflammatory reactions and excites the cardiac sympathetic system, and this may be related to the development of cognitive impairments in heart failure patients.¹¹⁶ These findings are further supported by several experimental studies. For example, Hong et al.¹¹⁷ reported marked cognitive impairment three months after induction of heart failure in a mouse model. This was accompanied by enhanced expression of toll-like receptor-4, tumor necrosis factor (TNF)- α , and IL-6 in the cortex and hippocampus of the animals. Moreover, additional evidence for the development of brain dysfunction in heart failure patients is provided by clinical studies which have found high levels of TNF- α and IL-6 in heart failure patients with cognitive impairment and depressive disorders in comparison to heart failure patients without cognitive dysfunction.^{118,119}

There is currently increasing interest in the pathophysiological roles of TNF- α in the central nervous system and in the development of neuroinflammation. This role is supported by findings showing that several neurological disorders are associated with marked *de novo* production of TNF- α ,¹²⁰ which is considered to be one of the main regulators of the brain's proinflammatory response, and which causes cell dysfunction and death through glutamate-mediated neurotoxicity.¹²⁰ Moreover, it is widely accepted that IL-1 β is a hallmark of neuroinflammation, which strongly contributes to cognitive impairment.¹²¹ Furthermore, TNF- α and IL-1 β can potentiate marked production of IL-6, which is also associated with neurodegenerative processes and cognitive

dysfunction.¹²² In addition, cardiac oxidative stress, activation of cascades, and release of proinflammatory cytokines (e.g., TNF- α , IL-1 β , and IL-6) have been consistently described in patients with ischemic heart disease and in heart failure patients.^{10,123} It is now known that IL-6 receptors are located in crucial areas, such as the hippocampus and the cerebral cortex. Therefore, it is not surprising that IL-6 may be an important factor in neuroinflammation, neuronal damage, and apoptosis, with subsequent development of severe neuropathological processes, so this offers a promising therapeutic target.^{10,124}

The situation is further complicated if it is born in mind that activation of inflammatory cascades and oxidative stress may induce neuropathological processes due to disruption of the BBB and activation of glial cells.^{12,13} Systemic inflammation may cause alterations in the structure and function of BBB, which may be disruptive or non-disruptive.^{13,87} Disruptive BBB alterations include changes at the histological level, such as destruction of endothelial cells, and include molecular changes leading to endothelial apoptosis, membrane abnormalities, mitochondrial damage, and modification of tight junctions.¹³ Non-disruptive alterations may be due to molecular changes.¹³ Many cytokines and inflammatory factors, for example, TNF- α and IL-1 β , have been found to mediate some of these changes (e.g., cerebral endothelial activation associated with neuroinflammation), and increased permeability of the BBB.¹²⁵ Moreover, inflammation associated with ischemic myocardial injury may promote activation of astrocytes.¹²⁶ Besides its effects on astrocytes, inflammation also plays a role in leukocyte recruitment across the BBB. Thus, migration of monocytes to the central nervous system promotes their differentiation into microglia, which has also been associated with neuroinflammation.¹²⁷ It should be remembered that activated microglia upregulate the production of proinflammatory cytokines (e.g., IL-1, IL-6, and TNF- α), thus contributing to the development and progression of neurodegeneration.¹²⁸

Finally, critical illness such as ABI may lead to excessive elevations of circulating glucocorticoids, which may generate or exacerbate inflammation-induced hippocampal and prefrontal cortical damage.¹⁰⁶

Ischemic stroke in patients with NSC

Myocardial dysfunctions, such as NSC, are well-established complications of severe brain injury.^{4,129} On the other hand, TTC is associated with acute and severe LV dysfunction, which could contribute to LV thrombus formation—with potential cardioembolic complications, such as cardioembolic stroke.¹⁵ The incidence, however, of LV thrombus formation and related embolic complications varies between studies.^{15,130,131} In a systematic review of 26 clinical studies, de Gregorio¹⁵ reported that LV thrombus

formation occurred in about 5% of patients with TTC, and that 25% of the whole study population develop major stroke or transient ischemic attack. Additionally, a recent multicenter study, which enrolled 1,676 patients with TTC, reported that intraventricular thrombus formation and/or embolism was present with a prevalence of 3.3% and that 23.2% of these patients developed ischemic stroke.¹³¹

Thromboembolic events in TTC are probably related to the depressed LV systolic dysfunction and the subsequent blood stasis during the acute phase.^{88,131-133} Moreover, endothelial dysfunction, hypercoagulability,^{131,134} activation of inflammatory pathways, and release of cytokines may provide additional pathophysiological mechanisms.^{131,135} Indeed, hypercoagulable state associated with NSC underlies the Virchow triad (e.g., endothelial cell damage, blood stasis, and platelet activation).¹³⁶ Possible pathomechanisms associated with hypercoagulability in TTC include the catecholamine storm during the acute phase of the disease with subsequent increased platelet activation and aggregation.¹³⁷ Furthermore, previous research indicates that inflammatory milieu with elevated plasmatic levels of IL-6 and IL-7 is an integral part of LV dysfunction.^{138,139} Moreover, several studies have revealed higher rates of LV thrombus formation and, thus, associated cardioembolic events, in patients with lower EF, right ventricular involvement, apical ballooning, and temporal anterior wall dysfunction in takotsubo syndrome.^{20,140,141} The hypothesis of LV thrombus formation with higher stroke rates related to apical ballooning is further supported by a recent retrospective study, which examined clinical and imaging data of 72 patients hospitalized with takotsubo syndrome, and detected apical ballooning with anterior wall motion abnormalities in 90% of the patients. Moreover, these patients showed significant recovery of LV function within two months of the event. The authors proposed that rapid changes in LV morphology might be a complementary mechanism contributing to LV thrombus formation and suggested that oral anticoagulation should be evaluated if not contraindicated until recovery of ventricular function.¹⁴²

Additionally, a variety of dysrhythmias have been observed in patients with TTC, including supraventricular tachyarrhythmias, ventricular tachycardia, ventricular fibrillation, asystole, and pulseless electrical activity.¹⁴³ More specifically, patients with TTC are at increased risk of developing various atrial arrhythmias like atrial fibrillation due to temporal left atrial dysfunction during the acute phase.¹⁴⁴ The prevalence of atrial fibrillation in patients with TTC is high, reaching a percentage of 25%^{144,145} compared to 10% in the general population.¹⁴⁶ Moreover, its presence in TTC patients serves as a powerful predictor of short-term and long-term mortality.^{144,145} However, it is conceivable that the number of unreported episodes of paroxysmal atrial fibrillation among

patients with TTC is higher due to the fact that many of them have a "silent" clinical presentation¹⁴⁷ making its diagnosis a clinical challenge.

Finally, despite that stroke in patients with TTC is primarily thought to be associated with LV thromboembolism,^{88,132,133} the question arises whether pre-existing cardiovascular risk factors are relevant to the development of stroke in patients with TTC. Moreover, several studies have shown an increased rate of common cardiovascular comorbidities and atherosclerotic risk factors in patients with TTC, including hypertension, diabetes mellitus, hyperlipidemia, and smoking.^{148,149} Indeed, a recent retrospective study of 27,970 patients hospitalized with TTC examined the trends in the incidence of ischemic stroke stratified by subtype (cardioembolic vs. thrombotic). The authors reported that thrombotic etiology of ischemic stroke was more common among patients with TTC and hypothesized that the higher rates of risk factors among these patients contribute significantly to the higher incidence of stroke of thrombotic etiology.¹⁴⁹

Despite the fact that published data are scarce, cardioembolic complications, such as ischemic stroke, could additionally develop in patients with cardiac dysfunction—after primary acute neurological injury—making the management of these patients a therapeutic challenge.

Clinical management and therapeutic implications

Screening for high-risk patients

Early detection of cardiovascular complications caused by ABI has become paramount. In fact, the therapeutic goal in patients with ABI is to ensure brain homeostasis and oxygenation, avoid intracranial hypertension, and decrease oxygen consumption^{49,150,151} in order to minimize secondary brain insults.¹⁵² Moreover, as previously discussed, ABI may be complicated by the development of NSC, which impairs LV function and systemic arterial pressure, leading to cerebral hypoperfusion, impaired brain homeostasis, and reduced oxygen supply to the brain.^{49,153,154}

Hence, close hemodynamic monitoring not only leads to improved systemic and cerebral hemodynamics, but also plays an additional fundamental role in avoiding secondary ischemic events. Mutoh et al.¹⁵³ compared the usefulness of transpulmonary thermodilution, a less invasive technique for hemodynamic monitoring, with pulmonary artery catheter monitoring in 16 SAH patients diagnosed with vasospasm. The authors then reported reduced frequencies of vasospasm and cardiopulmonary complications in patients for whom goal-directed therapy was guided by transpulmonary thermodilution. In addition, a multicenter prospective cohort study of hemodynamic variables using thermo-

dilution in patients with poor grade SAH demonstrated elevated indexes of lung water and pulmonary vascular permeability, together with a systemic vascular index corresponding to after-load mismatch and resembling heart failure.¹⁵⁴

For patients with stroke, myocardial injury is often detected and strongly associated with poor outcomes. The term "stroke-heart syndrome" involves a broad spectrum of cardiac complications after ischemic stroke, such as arrhythmias, cardiac injury and dysfunction,⁶⁴ and NSC.¹⁵⁵ Early blood pressure and continuous ECG monitoring is suggested in order to be able to identify ischemic stroke patients at high risk of cardiac complications at an early stage. Moreover, as extensive data have shown a relationship between LVEF and stroke severity and an association with neurological disability,¹⁵⁶ we strongly recommend early echocardiographic monitoring in severely affected patients (National Institutes of Health Stroke Scale [NIHSS] score >10), patients with clinical signs of myocardial dysfunction and with elevated cardiac biomarkers, and patients with disturbances during monitoring such as arrhythmias and severe hypertension.⁴⁴ Furthermore, clinicians should also take into account the likely effect of atrial fibrillation, of which incidence has been found to be statistically significantly higher in both, the group of patients with severe stroke and in patients with impaired systolic function.¹⁵⁶ Finally, prolonged cardiovascular monitoring is recommended in high-risk patients with right-located stroke, especially those with insular involvement, older patients, and patients with a history of hypertensive or coronary artery disease, or with intense emotional stress.²

Finally, in patients with NSC, a single bed-side risk score including several parameters, as for example lower LVEF, apical type, elevated troponin levels, increased inflammatory biomarkers, and prior history of vascular disease, may help identify patients at a higher risk of adverse events (e.g., intraventricular thrombus formation and embolism) and, thus improve outcome.¹³¹

Therapeutic strategies

NSC is frequently managed conservatively due to its reversible nature. In patients with classic uncomplicated TTC, the most appropriate and beneficial therapeutic intervention is the combination of β -blockers with angiotensin-converting enzyme (ACE) inhibitors.¹⁵⁷ Indeed, several studies demonstrate the therapeutic benefits of β -blockers, as they lead to rapid recovery of LV function and inhibition of the Gs-protein metabolic pathway, while ACE inhibitors prevent vasospasm and reduce hypertension.¹⁵⁸⁻¹⁶⁰ Additionally, previous research data demonstrate that pre-existing treatment with β -blockers is associated with milder stroke severity due to their sympatholytic action, anti-inflammatory properties, and reduction of norepinephrine-stimulated my-

ocytes apoptosis.^{161,162} Moreover, a classical study of Neil-Dwyer et al.¹⁶³ assessed the cardioprotective effect of propranolol plus phentolamine in patients with SAH and reported reduced myocardial necrotic lesions in patients receiving the combination propranolol/phentolamine when compared to the placebo group. These findings are in accordance with studies examining the effects of β -blocker in TBI patients with a recent meta-analysis of nine studies showing a significant reduction in in-hospital mortality in adult patients with TBI.¹⁶⁴ Similarly, a multicenter, prospective, observational study in patients with TBI found that the administration of β -blockers after TBI was associated with lower mortality and that propranolol showed better effects than other β -blockers.¹⁶⁵ In contrast, in a recent meta-analysis of three retrospective studies, which included 691 patients with SAH, no benefit of pre-admission β -blocker on mortality or on cardiac dysfunction, or vasospasm was observed.¹⁶⁶ NSC with persistent LV dysfunction should be treated according to heart failure guidelines, including ACE inhibitors, angiotensin receptor blockers, and β -blockers.¹⁶⁷ However, β -blockers are contraindicated in patients with decompensated heart failure.¹⁴² Further research is needed before β -blockers and ACE inhibitors can be routinely recommended to patients with NSC due to acute ischemic or hemorrhagic stroke.

Maintenance of systemic hemodynamics is crucial for patients with ABI, in order to ensure brain homeostasis, adequate CPP, and accurate brain oxygenation and, thus, to minimize or even avoid secondary brain injury.^{49,154,168} Despite the general agreement that vasopressors should be avoided in patients with TTC, a retrospective study reported a similar total mortality between SAH patients with NSC and SAH patients without NSC.¹⁶⁹ However, due to the pathophysiological background of the disease, it is generally recommended that adrenaline, noradrenaline, dobutamine, milrinone, and isoproterenol should be avoided in patients with TTC secondary to neurological illness.¹⁷⁰ On the other hand, if indicated, the use of the inotropes dobutamine and milrinone may improve cardiac function in SAH patients with NSC, with patients with cerebral vasospasm being especially benefited by milrinone.^{8,170} Patients with cardiogenic shock without LV outflow tract obstruction (LVOTO) usually present with hypotension. For the specific patient group, the use of dobutamine/dopamine is recommended for its treatment due to their positive inotropic effect. Inotropic support should be discontinued if the patient develops moderate or severe LVOTO. In case of refractory hypotension with impaired internal organ blood supply, vasopressor administration should be considered.^{157,171} If cardiogenic shock is accompanied by LVOTO, the treatment of choice is combined with fluid administration of β -blockers; phenylephrine administration may also be considered in preference to positive inotropic

drugs (e.g., norepinephrine).^{157,171-173} Because of its cardioprotective action and increase in blood supply to internal organs and intra-aortic balloon pump, the use of levosimendan should be evaluated in patients with refractory SAH-associated cardiogenic shock.^{8,174,175}

More clinical trials are required in order to indicate the optimal therapeutic management in patients with concurrent ABI and NSC and, consequently, improve their cardiac function and minimize secondary brain injury. More specifically, further research examining the exact role of β -blockers and ACE inhibitors is strongly recommended. Moreover, clinical trials should evaluate other therapeutic alternatives, as for example estradiol administration in postmenopausal women with NSC or medications, which are found to be effective in the management of patients with heart failure (e.g., angiotensin-nephrilysin receptor inhibitors and mineralocorticoid receptor antagonists). Finally, more experimental and controlled clinical studies are needed to determine the effectiveness of alternative therapeutic options, such as targeting inflammation and coronary microcirculation.

Conclusions

In recent years, it has been convincingly demonstrated that ABI, such as SAH, intracerebral bleeding, ischemic stroke, and TBI may cause severe cardiac complications. The pathophysiology of these brain-heart interactions is complex and involves sympathetic hyperactivity, activation of the HPA-axis as well as immune and inflammatory pathways. Patients with neurogenic cardiac dysfunction are likely to develop delayed secondary cerebral complications associated with these heart complications. Still, however, little is known about the pathophysiology of this phenomenon. The aim of the current review was to highlight the main pathophysiological mechanisms of stroke-heart-brain interactions and cardiac-associated secondary brain complications in patients with concurrent ABI and especially in patients with stroke and NSC. As was demonstrated, NSC induces severe systemic and cerebral inflammation, activation of the HPA axis, BBB dysfunction, and hemodynamic compromise, so neurocritical care patients are vulnerable to the development of delayed secondary cerebral events. The therapeutic management of neurogenic cardiac dysfunction is conservative, primarily oriented to the improvement of cardiac function, prevention of further neuroinflammation, provision of adequate oxygenation, and maintenance of brain homeostasis and CPP. Further studies are needed to investigate the complex pathophysiological interactions between heart and brain.

Disclosure

The authors have no financial conflicts of interest.

Acknowledgments

Figures created with BioRender.com.

References

- Mrozek S, Gobin J, Constantin JM, Fourcade O, Geeraerts T. Crosstalk between brain, lung and heart in critical care. *Anaesth Crit Care Pain Med* 2020;39:519-530.
- Gopinath R, Ayya SS. Neurogenic stress cardiomyopathy: what do we need to know. *Ann Card Anaesth* 2018;21:228-234.
- Chen Z, Venkat P, Seyfried D, Chopp M, Yan T, Chen J. Brain-heart interaction: cardiac complications after stroke. *Circ Res* 2017;121:451-468.
- Biso S, Wongrakpanich S, Agrawal A, Yadlapati S, Kishlyansky M, Figueredo V. A review of neurogenic stunned myocardium. *Cardiovasc Psychiatry Neurol* 2017;2017:5842182.
- Lee VH, Oh JK, Mulvagh SL, Wijdicks EF. Mechanisms in neurogenic stress cardiomyopathy after aneurysmal subarachnoid hemorrhage. *Neurocrit Care* 2006;5:243-249.
- Robba C, Bonatti G, Pelosi P, Citerio G. Extracranial complications after traumatic brain injury: targeting the brain and the body. *Curr Opin Crit Care* 2020;26:137-146.
- Dantzer R, Konsman JP, Bluthé RM, Kelley KW. Neural and humoral pathways of communication from the immune system to the brain: parallel or convergent? *Auton Neurosci* 2000;85:60-65.
- Kerro A, Woods T, Chang JJ. Neurogenic stunned myocardium in subarachnoid hemorrhage. *J Crit Care* 2017;38:27-34.
- Krishnamoorthy V, Mackensen GB, Gibbons EF, Vavilala MS. Cardiac dysfunction after neurologic injury: what do we know and where are we going? *Chest* 2016;149:1325-1331.
- Havakuk O, King KS, Grazette L, Yoon AJ, Fong M, Bregman N, et al. Heart failure-induced brain injury. *J Am Coll Cardiol* 2017;69:1609-1616.
- Claassen JAHR, Thijssen DHJ, Panerai RB, Faraci FM. Regulation of cerebral blood flow in humans: physiology and clinical implications of autoregulation. *Physiol Rev* 2021;101:1487-1559.
- Varatharaj A, Galea I. The blood-brain barrier in systemic inflammation. *Brain Behav Immun* 2017;60:1-12.
- Hoogland IC, Houbolt C, van Westerloo DJ, van Gool WA, van de Beek D. Systemic inflammation and microglial activation: systematic review of animal experiments. *J Neuroinflammation* 2015;12:114.
- Ziaka M, Exadaktylos A. Brain-lung interactions and mechanical ventilation in patients with isolated brain injury. *Crit Care* 2021;25:358.
- de Gregorio C. Cardioembolic outcomes in stress-related cardiomyopathy complicated by ventricular thrombus: a systematic review of 26 clinical studies. *Int J Cardiol* 2010;141:11-17.
- Kenigsberg BB, Barnett CF, Mai JC, Chang JJ. Neurogenic stunned myocardium in severe neurological injury. *Curr Neurol Neurosci Rep* 2019;19:90.
- Porto I, Della Bona R, Leo A, Proietti R, Pieroni M, Caltagirone C, et al. Stress cardiomyopathy (tako-tsubo) triggered by nervous system diseases: a systematic review of the reported cases. *Int J Cardiol* 2013;167:2441-2448.
- Sposato LA, Fridman S, Whitehead SN, Lopes RD. Linking stroke-induced heart injury and neurogenic atrial fibrillation: a hypothesis to be proven. *J Electrocardiol* 2018;51:430-432.
- Sposato LA, Hilz MJ, Asperg S, Murthy SB, Bahit MC, Hsieh CY, et al. Post-stroke cardiovascular complications and neurogenic cardiac injury: JACC state-of-the-art review. *J Am Coll Cardiol* 2020;76:2768-2785.
- Templin C, Ghadri JR, Diekmann J, Napp LC, Bataiosu DR, Jurguszewski M, et al. Clinical features and outcomes of takotsubo (stress) cardiomyopathy. *N Engl J Med* 2015;373:929-938.
- Scheitz JF, Nolte CH, Doehner W, Hachinski V, Endres M. Stroke-heart syndrome: clinical presentation and underlying mechanisms. *Lancet Neurol* 2018;17:1109-1120.
- Pelliccia F, Kaski JC, Crea F, Camici PG. Pathophysiology of takotsubo syndrome. *Circulation* 2017;135:2426-2441.
- Balint B, Jaremek V, Thorburn V, Whitehead SN, Sposato LA. Left atrial microvascular endothelial dysfunction, myocardial inflammation and fibrosis after selective insular cortex ischemic stroke. *Int J Cardiol* 2019;292:148-155.
- Suzuki H, Matsumoto Y, Kaneta T, Sugimura K, Takahashi J, Fukumoto Y, et al. Evidence for brain activation in patients with takotsubo cardiomyopathy. *Circ J* 2014;78:256-258.
- Masuda T, Sato K, Yamamoto S, Matsuyama N, Shimohama T, Matsunaga A, et al. Sympathetic nervous activity and myocardial damage immediately after subarachnoid hemorrhage in a unique animal model. *Stroke* 2002;33:1671-1676.
- Prosser J, MacGregor L, Lees KR, Diener HC, Hacke W, Davis S; VISTA Investigators. Predictors of early cardiac morbidity and mortality after ischemic stroke. *Stroke* 2007;38:2295-2302.
- Kallmünzer B, Breuer L, Kahl N, Bobinger T, Raaz-Schrauder D, Huttner HB, et al. Serious cardiac arrhythmias after stroke:

- incidence, time course, and predictors--a systematic, prospective analysis. *Stroke* 2012;43:2892-2897.
28. Jensen JK, Ueland T, Aukrust P, Antonsen L, Kristensen SR, Januzzi JL, et al. Highly sensitive troponin T in patients with acute ischemic stroke. *Eur Neurol* 2012;68:287-293.
 29. Lyon AR, Rees PS, Prasad S, Poole-Wilson PA, Harding SE. Stress (takotsubo) cardiomyopathy--a novel pathophysiological hypothesis to explain catecholamine-induced acute myocardial stunning. *Nat Clin Pract Cardiovasc Med* 2008;5:22-29.
 30. Ghadri JR, Wittstein IS, Prasad A, Sharkey S, Dote K, Akashi YJ, et al. International expert consensus document on takotsubo syndrome (part I): clinical characteristics, diagnostic criteria, and pathophysiology. *Eur Heart J* 2018;39:2032-2046.
 31. White M, Wiechmann RJ, Roden RL, Hagan MB, Wollmering MM, Port JD, et al. Cardiac beta-adrenergic neuroeffector systems in acute myocardial dysfunction related to brain injury. Evidence for catecholamine-mediated myocardial damage. *Circulation* 1995;92:2183-2189.
 32. Paur H, Wright PT, Sikkil MB, Tranter MH, Mansfield C, O'Gara P, et al. High levels of circulating epinephrine trigger apical cardiodepression in a β_2 -adrenergic receptor/Gi-dependent manner: a new model of takotsubo cardiomyopathy. *Circulation* 2012;126:697-706.
 33. Jung JM, Kim JG, Kim JB, Cho KH, Yu S, Oh K, et al. Takotsubo-like myocardial dysfunction in ischemic stroke: a hospital-based registry and systematic literature review. *Stroke* 2016;47:2729-2736.
 34. Yoshimura S, Toyoda K, Ohara T, Nagasawa H, Ohtani N, Kuwashiro T, et al. Takotsubo cardiomyopathy in acute ischemic stroke. *Ann Neurol* 2008;64:547-554.
 35. Bleilevens C, Roehl AB, Zoremba N, Tolba R, Rossaint R, Hein M. Insular infarct size but not levosimendan influenced myocardial injury triggered by cerebral ischemia in rats. *Exp Brain Res* 2015;233:149-156.
 36. Nagai M, Hoshida S, Kario K. The insular cortex and cardiovascular system: a new insight into the brain-heart axis. *J Am Soc Hypertens* 2010;4:174-182.
 37. Craig AD. How do you feel--now? The anterior insula and human awareness. *Nat Rev Neurosci* 2009;10:59-70.
 38. Mashaly HA, Provencio JJ. Inflammation as a link between brain injury and heart damage: the model of subarachnoid hemorrhage. *Cleve Clin J Med* 2008;75 Suppl 2:S26-S30.
 39. Wang X, Cui L, Ji X. Cognitive impairment caused by hypoxia: from clinical evidences to molecular mechanisms. *Metab Brain Dis* 2022;37:51-66.
 40. Tavazzi G, Zanierato M, Via G, Iotti GA, Procaccio F. Are neurogenic stress cardiomyopathy and takotsubo different syndromes with common pathways?: etiopathological insights on dysfunctional hearts. *JACC Heart Fail* 2017;5:940-942.
 41. Fassbender K, Schmidt R, Mössner R, Daffertshofer M, Hennerici M. Pattern of activation of the hypothalamic-pituitary-adrenal axis in acute stroke. Relation to acute confusional state, extent of brain damage, and clinical outcome. *Stroke* 1994;25:1105-1108.
 42. Sobowale OA, Parry-Jones AR, Smith CJ, Tyrrell PJ, Rothwell NJ, Allan SM. Interleukin-1 in stroke: from bench to bedside. *Stroke* 2016;47:2160-2167.
 43. Edwards AV, Jones CT. Autonomic control of adrenal function. *J Anat* 1993;183(Pt 2):291-307.
 44. Battaglini D, Robba C, Lopes da Silva A, Dos Santos Samary C, Leme Silva P, Dal Pizzol F, et al. Brain-heart interaction after acute ischemic stroke. *Crit Care* 2020;24:163.
 45. Wittstein IS, Thiemann DR, Lima JA, Baughman KL, Schulman SP, Gerstenblith G, et al. Neurohumoral features of myocardial stunning due to sudden emotional stress. *N Engl J Med* 2005;352:539-548.
 46. Kono T, Morita H, Kuroiwa T, Onaka H, Takatsuka H, Fujiwara A. Left ventricular wall motion abnormalities in patients with subarachnoid hemorrhage: neurogenic stunned myocardium. *J Am Coll Cardiol* 1994;24:636-640.
 47. Bratton SL, Chestnut RM, Ghajar J, McConnell Hammond FF, Harris OA, Hartl R, et al. XIV. Hyperventilation. *J Neurotrauma* 2007;24 Suppl 1:S87-S90.
 48. Nguyen H, Zaroff JG. Neurogenic stunned myocardium. *Curr Neurol Neurosci Rep* 2009;9:486-491.
 49. Mazzeo AT, Micalizzi A, Mascia L, Scicolone A, Siracusano L. Brain-heart crosstalk: the many faces of stress-related cardiomyopathy syndromes in anaesthesia and intensive care. *Br J Anaesth* 2014;112:803-815.
 50. Banki NM, Kopelnik A, Dae MW, Miss J, Tung P, Lawton MT, et al. Acute neurocardiogenic injury after subarachnoid hemorrhage. *Circulation* 2005;112:3314-3319.
 51. Sugimoto K, Watanabe E, Yamada A, Iwase M, Sano H, Hishida H, et al. Prognostic implications of left ventricular wall motion abnormalities associated with subarachnoid hemorrhage. *Int Heart J* 2008;49:75-85.
 52. Malik AN, Gross BA, Rosalind Lai PM, Moses ZB, Du R. Neurogenic stress cardiomyopathy after aneurysmal subarachnoid hemorrhage. *World Neurosurg* 2015;83:880-885.
 53. Kothavale A, Banki NM, Kopelnik A, Yarlagadda S, Lawton MT, Ko N, et al. Predictors of left ventricular regional wall motion abnormalities after subarachnoid hemorrhage. *Neurocrit Care* 2006;4:199-205.
 54. Tung P, Kopelnik A, Banki N, Ong K, Ko N, Lawton MT, et al. Predictors of neurocardiogenic injury after subarachnoid hem-

- orrhage. *Stroke* 2004;35:548-551.
55. Kilbourn KJ, Levy S, Staff I, Kureshi I, McCullough L. Clinical characteristics and outcomes of neurogenic stress cardiomyopathy in aneurysmal subarachnoid hemorrhage. *Clin Neurol Neurosurg* 2013;115:909-914.
 56. Colivicchi F, Bassi A, Santini M, Caltagirone C. Cardiac autonomic derangement and arrhythmias in right-sided stroke with insular involvement. *Stroke* 2004;35:2094-2098.
 57. Laowattana S, Zeger SL, Lima JA, Goodman SN, Wittstein IS, Oppenheimer SM. Left insular stroke is associated with adverse cardiac outcome. *Neurology* 2006;66:477-483.
 58. Krause T, Werner K, Fiebach JB, Villringer K, Piper SK, Haeusler KG, et al. Stroke in right dorsal anterior insular cortex is related to myocardial injury. *Ann Neurol* 2017;81:502-511.
 59. Cheung RT, Hachinski V. Cardiac effects of stroke. *Curr Treat Options Cardiovasc Med* 2004;6:199-207.
 60. Oras J, Grivans C, Bartley A, Rydenhag B, Ricksten SE, Seeman-Lodding H. Elevated high-sensitive troponin T on admission is an indicator of poor long-term outcome in patients with subarachnoid haemorrhage: a prospective observational study. *Crit Care* 2016;20:11.
 61. Murthy SB, Shah S, Rao CP, Bershad EM, Suarez JI. Neurogenic stunned myocardium following acute subarachnoid hemorrhage: pathophysiology and practical considerations. *J Intensive Care Med* 2015;30:318-325.
 62. Nasr DM, Tomasini S, Prasad A, Rabinstein AA. Acute brain diseases as triggers for stress cardiomyopathy: clinical characteristics and outcomes. *Neurocrit Care* 2017;27:356-361.
 63. Mihalovic M, Tousek P. Myocardial injury after stroke. *J Clin Med* 2022;11:2.
 64. Scheitz JF, Stengl H, Nolte CH, Landmesser U, Endres M. Neurological update: use of cardiac troponin in patients with stroke. *J Neurol* 2021;268:2284-2292.
 65. Alkhachroum AM, Miller B, Chami T, Tatsuoka C, Sila C. A troponin study on patients with ischemic stroke, intracerebral hemorrhage and subarachnoid hemorrhage: type II myocardial infarction is significantly associated with stroke severity, discharge disposition and mortality. *J Clin Neurosci* 2019;64:83-88.
 66. Hravnak M, Frangiskakis JM, Crago EA, Chang Y, Tanabe M, Gorcsan J 3rd, et al. Elevated cardiac troponin I and relationship to persistence of electrocardiographic and echocardiographic abnormalities after aneurysmal subarachnoid hemorrhage. *Stroke* 2009;40:3478-3484.
 67. Christensen H, Fogh Christensen A, Boysen G. Abnormalities on ECG and telemetry predict stroke outcome at 3 months. *J Neurol Sci* 2005;234:99-103.
 68. Iltumur K, Yavavli A, Apak I, Ariturk Z, Toprak N. Elevated plasma N-terminal pro-brain natriuretic peptide levels in acute ischemic stroke. *Am Heart J* 2006;151:1115-1122.
 69. Montaner J, García-Berrocó T, Mendioroz M, Palacios M, Perea-Gainza M, Delgado P, et al. Brain natriuretic peptide is associated with worsening and mortality in acute stroke patients but adds no prognostic value to clinical predictors of outcome. *Cerebrovasc Dis* 2012;34:240-245.
 70. Svirí GE, Soustiel JF, Zaaroor M. Alteration in brain natriuretic peptide (BNP) plasma concentration following severe traumatic brain injury. *Acta Neurochir (Wien)* 2006;148:529-533; discussion 533.
 71. Etgen T, Baum H, Sander K, Sander D. Cardiac troponins and N-terminal pro-brain natriuretic peptide in acute ischemic stroke do not relate to clinical prognosis. *Stroke* 2005;36:270-275.
 72. Takahashi K, Totsune K, Sone M, Ohneda M, Murakami O, Itoi K, et al. Human brain natriuretic peptide-like immunoreactivity in human brain. *Peptides* 1992;13:121-123.
 73. Park KI. Plasma brain-type natriuretic peptide level following seizure and syncope: pilot study. *J Epilepsy Res* 2014;4:14-17.
 74. Agewall S, Giannitsis E, Jernberg T, Katus H. Troponin elevation in coronary vs. non-coronary disease. *Eur Heart J* 2011;32:404-411.
 75. Wira CR 3rd, Rivers E, Martinez-Capolino C, Silver B, Iyer G, Sherwin R, et al. Cardiac complications in acute ischemic stroke. *West J Emerg Med* 2011;12:414-420.
 76. Krishnamoorthy V, Rowhani-Rahbar A, Gibbons EF, Rivara FP, Temkin NR, Pontius C, et al. Early systolic dysfunction following traumatic brain injury: a cohort study. *Crit Care Med* 2017;45:1028-1036.
 77. Hüttemann E, Schelenz C, Chatzinikolaou K, Reinhart K. Left ventricular dysfunction in lethal severe brain injury: impact of transesophageal echocardiography on patient management. *Intensive Care Med* 2002;28:1084-1088.
 78. Harjola VP, Mullens W, Banaszewski M, Bauersachs J, Brunner-La Rocca HP, Chioncel O, et al. Organ dysfunction, injury and failure in acute heart failure: from pathophysiology to diagnosis and management. A review on behalf of the Acute Heart Failure Committee of the Heart Failure Association (HFA) of the European Society of Cardiology (ESC). *Eur J Heart Fail* 2017;19:821-836.
 79. Wang X, Pei J, Hu X. The brain-heart connection in takotsubo syndrome: the central nervous system, sympathetic nervous system, and catecholamine overload. *Cardiol Res Pract* 2020;9:2020:4150291.
 80. Klein C, Hiestand T, Ghadri JR, Templin C, Jäncke L, Hänggi J. Takotsubo syndrome - predictable from brain imaging data. *Sci Rep* 2017;7:5434.

81. Templin C, Hänggi J, Klein C, Topka MS, Hiestand T, Levinson RA, et al. Altered limbic and autonomic processing supports brain-heart axis in takotsubo syndrome. *Eur Heart J* 2019;40:1183-1187.
82. Hiestand T, Hänggi J, Klein C, Topka MS, Jaguszewski M, Ghadri JR, et al. Takotsubo syndrome associated with structural brain alterations of the limbic system. *J Am Coll Cardiol* 2018;7:809-811.
83. Kewcharoen J, Trongtorsak A, Kanitsoraphan C, Prasitlunkum N, Mekritthikrai R, Techorueangwiwat C, et al. Cognitive impairment and 30-day rehospitalization rate in patients with acute heart failure: a systematic review and meta-analysis. *Indian Heart J* 2019;71:52-59.
84. Mccarthy ST, Wollner L, Rosenberg GA, Haaland KY. Cardiological dementia. *Lancet* 1981;318:1171.
85. Zuccalà G, Onder G, Pedone C, Carosella L, Pahor M, Bernabei R, et al. Hypotension and cognitive impairment: selective association in patients with heart failure. *Neurology* 2001;57:1986-1992.
86. Georgiadis D, Schwarz S, Baumgartner RW, Veltkamp R, Schwab S. Influence of positive end-expiratory pressure on intracranial pressure and cerebral perfusion pressure in patients with acute stroke. *Stroke* 2001;32:2088-2092.
87. Ziaka M, Exadaktylos A. ARDS associated acute brain injury: from the lung to the brain. *Eur J Med Res* 2022;27:150.
88. Kim SM, Aikat S, Bailey A, White M. Takotsubo cardiomyopathy as a source of cardioembolic cerebral infarction. *BMJ Case Rep* 2012;2012:bcr2012006835.
89. Phillips AA, Chan FH, Zheng MM, Krassioukov AV, Ainslie PN. Neurovascular coupling in humans: physiology, methodological advances and clinical implications. *J Cereb Blood Flow Metab* 2016;36:647-664.
90. Willie CK, Smith KJ. Fuelling the exercising brain: a regulatory quagmire for lactate metabolism. *J Physiol* 2011;589(Pt 4):779-780.
91. Paulson OB, Jarden JO, Godtfredsen J, Vorstrup S. Cerebral blood flow in patients with congestive heart failure treated with captopril. *Am J Med* 1984;76:91-95.
92. Choi BR, Kim JS, Yang YJ, Park KM, Lee CW, Kim YH, et al. Factors associated with decreased cerebral blood flow in congestive heart failure secondary to idiopathic dilated cardiomyopathy. *Am J Cardiol* 2006;97:1365-1369.
93. Jefferson AL, Himali JJ, Au R, Seshadri S, Decarli C, O'Donnell CJ, et al. Relation of left ventricular ejection fraction to cognitive aging (from the Framingham Heart Study). *Am J Cardiol* 2011;108:1346-1351.
94. de la Torre JC. Critically attained threshold of cerebral hypoperfusion: the CATCH hypothesis of Alzheimer's pathogenesis. *Neurobiol Aging* 2000;21:331-342.
95. Xu C, Tao X, Ma X, Zhao R, Cao Z. Cognitive dysfunction after heart disease: a manifestation of the heart-brain axis. *Oxid Med Cell Longev* 2021;2021:4899688.
96. Tini G, Scagliola R, Monacelli F, La Malfa G, Porto I, Brunelli C, et al. Alzheimer's disease and cardiovascular disease: a particular association. *Cardiol Res Pract* 2020;2020:2617970.
97. Armstead WM. Cerebral blood flow autoregulation and dysautoregulation. *Anesthesiol Clin* 2016;34:465-477.
98. Griesdale DE, Örténwall V, Norena M, Wong H, Sekhon MS, Kolmodin L, et al. Adherence to guidelines for management of cerebral perfusion pressure and outcome in patients who have severe traumatic brain injury. *J Crit Care* 2015;30:111-115.
99. Prabhakar H, Sandhu K, Bhagat H, Durga P, Chawla R. Current concepts of optimal cerebral perfusion pressure in traumatic brain injury. *J Anaesthesiol Clin Pharmacol* 2014;30:318-327.
100. Czosnyka M, Smielewski P, Piechnik S, Steiner LA, Pickard JD. Cerebral autoregulation following head injury. *J Neurosurg* 2001;95:756-763.
101. Hoiland RL, Bain AR, Rieger MG, Bailey DM, Ainslie PN. Hypoxemia, oxygen content, and the regulation of cerebral blood flow. *Am J Physiol Regul Integr Comp Physiol* 2016;310:R398-R413.
102. Ziaka M, Makris D, Fotakopoulos G, Tsilioni I, Befani C, Liakos P, et al. High-tidal-volume mechanical ventilation and lung inflammation in intensive care patients with normal lungs. *Am J Crit Care* 2020;29:15-21.
103. Derdeyn CP, Videen TO, Yundt KD, Fritsch SM, Carpenter DA, Grubb RL, et al. Variability of cerebral blood volume and oxygen extraction: stages of cerebral haemodynamic impairment revisited. *Brain* 2002;125(Pt 3):595-607.
104. Lorenzi-Filho G, Azevedo ER, Parker JD, Bradley TD. Relationship of carbon dioxide tension in arterial blood to pulmonary wedge pressure in heart failure. *Eur Respir J* 2002;19:37-40.
105. Thompson BT. Glucocorticoids and acute lung injury. *Crit Care Med* 2003;31(4 Suppl):S253-S257.
106. Hill AR, Spencer-Segal JL. Glucocorticoids and the brain after critical illness. *Endocrinology* 2021;162:bqaa242.
107. Güder G, Bauersachs J, Frantz S, Weismann D, Allolio B, Ertl G, et al. Complementary and incremental mortality risk prediction by cortisol and aldosterone in chronic heart failure. *Circulation* 2007;115:1754-1761.
108. Huffman JC, Celano CM, Beach SR, Motiwala SR, Januzzi JL. Depression and cardiac disease: epidemiology, mechanisms, and diagnosis. *Cardiovasc Psychiatry Neurol* 2013;2013:695925.
109. Newcomer JW, Selke G, Melson AK, Hershey T, Craft S, Richards K, et al. Decreased memory performance in healthy humans induced by stress-level cortisol treatment. *Arch Gen Psy-*

- chiatry* 1999;56:527-533.
110. Kirschbaum C, Wolf OT, May M, Wippich W, Hellhammer DH. Stress- and treatment-induced elevations of cortisol levels associated with impaired declarative memory in healthy adults. *Life Sci* 1996;58:1475-1483.
 111. Chetty S, Friedman AR, Taravosh-Lahn K, Kirby ED, Mirescu C, Guo F, et al. Stress and glucocorticoids promote oligodendrogenesis in the adult hippocampus. *Mol Psychiatry* 2014;19:1275-1283.
 112. Marchesi C, Paradis P, Schiffrin EL. Role of the renin-angiotensin system in vascular inflammation. *Trends Pharmacol Sci* 2008;29:367-374.
 113. Gill R, Tsung A, Billiar T. Linking oxidative stress to inflammation: Toll-like receptors. *Free Radic Biol Med* 2010;48:1121-1132.
 114. Thackeray JT, Hupe HC, Wang Y, Bankstahl JP, Berding G, Ross TL, et al. Myocardial inflammation predicts remodeling and neuroinflammation after myocardial infarction. *J Am Coll Cardiol* 2018;71:263-275.
 115. El-Menyar AA. Cytokines and myocardial dysfunction: state of the art. *J Card Fail* 2008;14:61-74.
 116. Toledo C, Andrade DC, Díaz HS, Inestrosa NC, Del Rio R. Neurocognitive disorders in heart failure: novel pathophysiological mechanisms underpinning memory loss and learning impairment. *Mol Neurobiol* 2019;56:8035-8051.
 117. Hong X, Bu L, Wang Y, Xu J, Wu J, Huang Y, et al. Increases in the risk of cognitive impairment and alterations of cerebral β -amyloid metabolism in mouse model of heart failure. *PLoS One* 2013;8:e63829.
 118. Ferketich AK, Ferguson JP, Binkley PF. Depressive symptoms and inflammation among heart failure patients. *Am Heart J* 2005;150:132-136.
 119. Parissis JT, Adamopoulos S, Rigas A, Kostakis G, Karatzas D, Venetsanou K, et al. Comparison of circulating proinflammatory cytokines and soluble apoptosis mediators in patients with chronic heart failure with versus without symptoms of depression. *Am J Cardiol* 2004;94:1326-1328.
 120. Olmos G, Lladó J. Tumor necrosis factor alpha: a link between neuroinflammation and excitotoxicity. *Mediators Inflamm* 2014;2014:861231.
 121. Gemma C, Bickford PC. Interleukin-1beta and caspase-1: players in the regulation of age-related cognitive dysfunction. *Rev Neurosci* 2007;18:137-148.
 122. Wright CB, Sacco RL, Rundek T, Delman J, Rabbani L, Elkind M. Interleukin-6 is associated with cognitive function: the Northern Manhattan study. *J Stroke Cerebrovasc Dis* 2006;15:34-38.
 123. Neri M, Fineschi V, Di Paolo M, Pomara C, Riezzo I, Turillazzi E, et al. Cardiac oxidative stress and inflammatory cytokines response after myocardial infarction. *Curr Vasc Pharmacol* 2015;13:26-36.
 124. Erta M, Quintana A, Hidalgo J. Interleukin-6, a major cytokine in the central nervous system. *Int J Biol Sci* 2012;8:1254-1266.
 125. Skelly DT, Hennessy E, Dansereau MA, Cunningham C. A systematic analysis of the peripheral and CNS effects of systemic LPS, IL-1 β , [corrected] TNF- α and IL-6 challenges in C57BL/6 mice. *PLoS One* 2013;8:e69123.
 126. Bascuñana P, Hess A, Borchert T, Wang Y, Wollert KC, Bengel FM, et al. 11C-methionine PET identifies astroglia involvement in heart-brain inflammation networking after acute myocardial infarction. *J Nucl Med* 2020;61:977-980.
 127. Heppner FL, Ransohoff RM, Becher B. Immune attack: the role of inflammation in Alzheimer disease. *Nat Rev Neurosci* 2015;16:358-372.
 128. Smith JA, Das A, Ray SK, Banik NL. Role of pro-inflammatory cytokines released from microglia in neurodegenerative diseases. *Brain Res Bull* 2012;87:10-20.
 129. Gruhl SL, Su J, Chua WC, Tay KV. Takotsubo cardiomyopathy in post-traumatic brain injury: a systematic review of diagnosis and management. *Clin Neurol Neurosurg* 2022;213:107119.
 130. Haghi D, Papavassiliu T, Heggemann F, Kaden JJ, Borggrefe M, Suselbeck T. Incidence and clinical significance of left ventricular thrombus in tako-tsubo cardiomyopathy assessed with echocardiography. *QJM* 2008;101:381-386.
 131. Ding KJ, Cammann VL, Szawan KA, Stähli BE, Wischnewsky M, Di Vece D, et al. Intraventricular thrombus formation and embolism in takotsubo syndrome: insights from the international takotsubo registry. *Arterioscler Thromb Vasc Biol* 2020;40:279-287.
 132. El-Battrawy I, Borggrefe M, Akin I. Takotsubo syndrome and embolic events. *Heart Fail Clin* 2016;12:543-550.
 133. Mitsuma W, Kodama M, Ito M, Kimura S, Tanaka K, Hoyano M, et al. Thromboembolism in takotsubo cardiomyopathy. *Int J Cardiol* 2010;139:98-100.
 134. Naegele M, Flammer AJ, Enseleit F, Roas S, Frank M, Hirt A, et al. Endothelial function and sympathetic nervous system activity in patients with takotsubo syndrome. *Int J Cardiol* 2016;224:226-230.
 135. Steffel J, Lüscher TF, Tanner FC. Tissue factor in cardiovascular diseases: molecular mechanisms and clinical implications. *Circulation* 2006;113:722-731.
 136. Mehta JL, Calcaterra G, Bassareo PP. COVID-19, thromboembolic risk, and Virchow's triad: lesson from the past. *Clin Cardiol* 2020;43:1362-1367.
 137. Pirzer R, Elmas E, Haghi D, Lippert C, Kralev S, Lang S, et al.

- Platelet and monocyte activity markers and mediators of inflammation in takotsubo cardiomyopathy. *Heart Vessels* 2012; 27:186-192.
138. Pyo MK, Yun-Choi HS, Hong YJ. Apparent heterogeneous responsiveness of human platelet rich plasma to catecholamines. *Platelets* 2003;14:171-178.
 139. Damàs JK, Waehre T, Yndestad A, Otterdal K, Hognestad A, Solum NO, et al. Interleukin-7-mediated inflammation in unstable angina: possible role of chemokines and platelets. *Circulation* 2003;107:2670-2676.
 140. Schneider B, Athanasiadis A, Schwab J, Pistner W, Gottwald U, Schoeller R, et al. Complications in the clinical course of takotsubo cardiomyopathy. *Int J Cardiol* 2014;176:199-205.
 141. Mollet NR, Dymarkowski S, Volders W, Wathiong J, Herbots L, Rademakers FE, et al. Visualization of ventricular thrombi with contrast-enhanced magnetic resonance imaging in patients with ischemic heart disease. *Circulation* 2002;106:2873-2876.
 142. Abanador-Kamper N, Kamper L, Wolfertz J, Pomjanski W, Wolf-Pütz A, Seyfarth M. Evaluation of therapy management and outcome in takotsubo syndrome. *BMC Cardiovasc Disord* 2017;17:225.
 143. Rathore SS, Iqbal K, Shafqat S, Tariq E, Tousif S, Ul-Haq ZG, et al. Meta-analysis of incidence and outcomes of life-threatening arrhythmias in takotsubo cardiomyopathy. *Indian Heart J* 2022;74:110-119.
 144. El-Battrawy I, Lang S, Ansari U, Behnes M, Hillenbrand D, Schramm K, et al. Impact of concomitant atrial fibrillation on the prognosis of takotsubo cardiomyopathy. *Europace* 2017; 19:1288-1292.
 145. Stiermaier T, Santoro F, Eitel C, Graf T, Möller C, Tarantino N, et al. Prevalence and prognostic relevance of atrial fibrillation in patients with takotsubo syndrome. *Int J Cardiol* 2017; 245:156-161.
 146. Debonnaire P, Joyce E, Hiemstra Y, Mertens BJ, Atsma DE, Schalij MJ, et al. Left atrial size and function in hypertrophic cardiomyopathy patients and risk of new-onset atrial fibrillation. *Circ Arrhythm Electrophysiol* 2017;10:e004052.
 147. Fonseca AC, Almeida AG, Santos MO, Ferro JM. Neurological complications of cardiomyopathies. *Handb Clin Neurol* 2021; 177:91-109.
 148. Dias A, Franco E, Janzer S, Koshkelashvili N, Bhalla V, Rubio M, et al. Incidence and predictors of stroke during the index event in an ethnically diverse takotsubo cardiomyopathy population. *Funct Neurol* 2016;31:157-162.
 149. Abe T, Olanipekun T, Igwe J, Khoury M, Busari O, Musonge-Effoe J, et al. Trends, predictors and outcomes of ischemic stroke among patients hospitalized with takotsubo cardiomyopathy. *J Stroke Cerebrovasc Dis* 2021;30:106005.
 150. Jaeger M, Schuhmann MU, Soehle M, Nagel C, Meixensberger J. Continuous monitoring of cerebrovascular autoregulation after subarachnoid hemorrhage by brain tissue oxygen pressure reactivity and its relation to delayed cerebral infarction. *Stroke* 2007;38:981-986.
 151. Budohoski KP, Czosnyka M, Smielewski P, Kasprócz M, Helmy A, Bulters D, et al. Impairment of cerebral autoregulation predicts delayed cerebral ischemia after subarachnoid hemorrhage: a prospective observational study. *Stroke* 2012; 43:3230-3237.
 152. van der Jagt M. Fluid management of the neurological patient: a concise review. *Crit Care* 2016;20:126.
 153. Mutoh T, Kazumata K, Ishikawa T, Terasaka S. Performance of bedside transpulmonary thermodilution monitoring for goal-directed hemodynamic management after subarachnoid hemorrhage. *Stroke* 2009;40:2368-2374.
 154. Yoneda H, Nakamura T, Shirao S, Tanaka N, Ishihara H, Suehiro E, et al. Multicenter prospective cohort study on volume management after subarachnoid hemorrhage: hemodynamic changes according to severity of subarachnoid hemorrhage and cerebral vasospasm. *Stroke* 2013;44:2155-2161.
 155. Wang TD, Wu CC, Lee YT. Myocardial stunning after cerebral infarction. *Int J Cardiol* 1997;58:308-311.
 156. Sung PH, Chen KH, Lin HS, Chu CH, Chiang JY, Yip HK. The correlation between severity of neurological impairment and left ventricular function in patients after acute ischemic stroke. *J Clin Med* 2019;8:190.
 157. Bairashevskaja AV, Belogubova SY, Kondratiuk MR, Rudnova DS, Sologova SS, Tereshkina OI, et al. Update of takotsubo cardiomyopathy: present experience and outlook for the future. *Int J Cardiol Heart Vasc* 2022;39:100990.
 158. Abu Ghalyoun B, Guragai N, Garris R, Shamooun D, Chuang A, Shamooun F. Abstract 11626: faster recovery in takotsubo with prior beta-blocker use. *Circulation* 2019;140(Suppl_1):A11626.
 159. Isogai T, Matsui H, Tanaka H, Fushimi K, Yasunaga H. Early β -blocker use and in-hospital mortality in patients with takotsubo cardiomyopathy. *Heart* 2016;102:1029-1035.
 160. Brunetti ND, Santoro F, De Gennaro L, Correale M, Gaglione A, Di Biase M. Drug treatment rates with beta-blockers and ACE-inhibitors/angiotensin receptor blockers and recurrences in takotsubo cardiomyopathy: a meta-regression analysis. *Int J Cardiol* 2016;214:340-342.
 161. Laowattana S, Oppenheimer SM. Protective effects of beta-blockers in cerebrovascular disease. *Neurology* 2007;68:509-514.
 162. Communal C, Singh K, Pimentel DR, Colucci WS. Norepinephrine stimulates apoptosis in adult rat ventricular myocytes by activation of the beta-adrenergic pathway. *Circula-*

- tion 1998;98:1329-1334.
163. Neil-Dwyer G, Walter P, Cruickshank JM, Doshi B, O'Gorman P. Effect of propranolol and phentolamine on myocardial necrosis after subarachnoid haemorrhage. *Br Med J* 1978;2: 990-992.
164. Alali AS, Mukherjee K, McCredie VA, Golan E, Shah PS, Bardes JM, et al. Beta-blockers and traumatic brain injury: a systematic review and meta-analysis. *Ann Surg* 2017;266: 952-961.
165. Ley EJ, Leonard SD, Barmparas G, Dhillon NK, Inaba K, Salim A, et al. Beta blockers in critically ill patients with traumatic brain injury: results from a multicenter, prospective, observational American Association for the Surgery of Trauma study. *J Trauma Acute Care Surg* 2018;84:234-244.
166. Luo H, Song WX, Jiang JW, Zhao JL, Rong WL, Li MH. Effects of preadmission beta-blockers on neurogenic stunned myocardium after aneurysmal subarachnoid hemorrhage: a meta-analysis. *Clin Neurol Neurosurg* 2017;158:77-81.
167. Bybee KA, Kara T, Prasad A, Lerman A, Barsness GW, Wright RS, et al. Systematic review: transient left ventricular apical ballooning: a syndrome that mimics ST-segment elevation myocardial infarction. *Ann Intern Med* 2004;141:858-865.
168. Calviello LA, Donnelly J, Zeiler FA, Thelin EP, Smielewski P, Czosnyka M. Cerebral autoregulation monitoring in acute traumatic brain injury: what's the evidence? *Minerva Anestesiol* 2017;83:844-857.
169. Talahma M, Alkhachroum AM, Alyahya M, Manjila S, Xiong W. Takotsubo cardiomyopathy in aneurysmal subarachnoid hemorrhage: institutional experience and literature review. *Clin Neurol Neurosurg* 2016;141:65-70.
170. Wagner S, Güthe T, Bhogal P, Cimpoca A, Ganslandt O, Bätzner H, et al. Aneurysmal subarachnoid hemorrhage as a trigger for takotsubo syndrome: a comprehensive review. *Rev Cardiovasc Med* 2021;22:1241-1251.
171. Omerovic E. Takotsubo syndrome—scientific basis for current treatment strategies. *Heart Fail Clin* 2016;12:577-586.
172. Nalluri N, Asti D, Anugu VR, Ibrahim U, Lafferty JC, Olkovsky Y. Cardiogenic shock secondary to takotsubo cardiomyopathy in a patient with preexisting hypertrophic obstructive cardiomyopathy. *CASE (Phila)* 2018;2:78-81.
173. Jha S, Zeijlon R, Shekka Espinosa A, Alkhoury J, Oras J, Omerovic E, et al. Clinical management in the takotsubo syndrome. *Expert Rev Cardiovasc Ther* 2019;17:83-93.
174. du Toit EF, Genis A, Opie LH, Pollesello P, Lochner A. A role for the RISK pathway and K(ATP) channels in pre- and post-conditioning induced by levosimendan in the isolated guinea pig heart. *Br J Pharmacol* 2008;154:41-50.
175. Taccone FS, Brasseur A, Vincent JL, De Backer D. Levosimendan for the treatment of subarachnoid hemorrhage-related cardiogenic shock. *Intensive Care Med* 2013;39:1497-1498.