

<https://helda.helsinki.fi>

Pharmacological fatty acid synthase inhibitors differently affect the malignant phenotype of oral cancer cells.

Boelcke, Willian Peter

2022-03

Boelcke , W P , Teixeira , I F , Aquino , I G , Mazzaro , A R , Cuadra-Zelaya , F J M , de Souza , A P , Salo , T , Della Coletta , R , Graner , E & Bastos , D C 2022 , ' Pharmacological fatty acid synthase inhibitors differently affect the malignant phenotype of oral cancer cells. ' , Archives of Oral Biology , vol. 135 , 105343 . <https://doi.org/10.1016/j.archoralbio.2021.105343>

<http://hdl.handle.net/10138/352827>

<https://doi.org/10.1016/j.archoralbio.2021.105343>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

1 **Pharmacological fatty acid synthase inhibitors differently affect the malignant**
2 **phenotype of oral cancer cells.**

3

4 Willian Peter Boelcke^{1,2*} | Isadora Ferrari Teixeira^{2*} | Iara Gonçalves Aquino² | Amanda
5 Mazzaro¹ | Florence Juana Maria Cuadra-Zelaya^{2,3} | Ana Paula de Souza¹ | Tuula
6 Salo^{4,5} | Ricardo Della Coletta² | Edgard Graner² | Débora Campanella Bastos^{1,2**}

7 * *These authors contributed equally to this work.*

8 ** *Corresponding author*

9 ¹Department of Biociences, School of Dentistry, University of Campinas, Piracicaba-SP,
10 Brazil.

11 ²Department of Oral Diagnosis, School of Dentistry, University of Campinas, Piracicaba-
12 SP, Brazil.

13 ³Department of Pathology, University of El Salvador, San Salvador, El Salvador

14 ⁴Cancer and Translational Medicine Research Unit, Faculty of Medicine and Medical
15 Research Center Oulu, Oulu University Hospital, University of Oulu, Oulu, Finland.

16 ⁵Institute of Oral and Maxillofacial Disease, University of Helsinki, and HUSLAB,
17 Department of Pathology, Helsinki University Hospital, Helsinki, Finland.

18

19 **Corresponding Author: Débora Campanella Bastos, Department of Biosciences,
20 School of Dentistry of Piracicaba, State University of Campinas (UNICAMP), Av. Limeira
21 901, Areão, Piracicaba 13414-018, São Paulo, Brazil. Phone: 55-19-2106-5315. E-mail:
22 bastosdc@unicamp.br

23

24

1 **ABSTRACT**

2 Objective: Fatty acid synthase levels are associated with aggressiveness, prognosis,
3 and risk of metastasis in oral squamous cell carcinomas. This enzyme contains seven
4 catalytic domains and its inhibition by synthetic or natural drugs has antineoplastic
5 properties such as C75, which is a synthetic inhibitor of the β - ketoacyl synthase domain,
6 the antibiotic triclosan, ligand of the enoyl reductase domain, and the antiobesity drug
7 orlistat, which inhibits the thioesterase domain. Here, we sought to investigate and
8 compare the *in vitro* effects of C75, triclosan, and orlistat on malignant phenotypes of the
9 cell line SCC-9: proliferation, cell cycle, apoptosis, adhesion, migration, and invasion.

10 Design: Half-maximal inhibitory concentration (IC_{50}) was determined using cell viability
11 assays. Cell death and cell cycle progression were analyzed by Annexin V-PE/7-ADD-
12 PerCP labeling and propidium iodide staining, respectively. Cell migration and invasion
13 were assayed by transwells assays and cell adhesion using collagen and fibronectin.

14 Results: C75 showed the lowest IC_{50} and higher inhibition of lipid droplets at low
15 concentrations and reduced cell motility. Triclosan showed the intermediate IC_{50} value,
16 excellent reduction of lipid bodies at the IC_{50} when compared with C75 and orlistat. Also,
17 triclosan reduced cell cycle progression, adhesion, migration, and invasion of SCC-9 and
18 induced the highest levels of apoptosis. Orlistat promoted cell cycle arrest, but showed
19 the lowest induction of apoptosis and did not affected invasion and adhesion of SCC-9.

20 Conclusion: Altogether, despite the particular effects of the analyzed fatty acid synthase
21 inhibitors, triclosan showed to better interfere in tumorigenic phenotypes of SCC-9 cells.

22

23 Keywords: Fatty acid synthase, FASN inhibitors, oral cancer

24

1 INTRODUCTION

2 Head and neck cancer is the sixth most common cancer worldwide and annually
3 accounts more than 650,000 cases and 330,000 deaths (Bray et al., 2018). More than
4 90% of all types of head and neck cancers are squamous cell carcinomas that affect the
5 oral mucosa (Montero & Patel, 2015). Clinically, oral squamous cell carcinoma (OSCC)
6 shows a locally aggressive behavior, frequent recurrences, and regional or distant
7 metastases (Ho et al., 2017). Despite the efforts and new modalities of treatments, the
8 mortality rates of OSCC remain unchanged for decades, with a 5-year survival rate of
9 around 50% (Cohen Goldemberg et al., 2018).

10 The reprogramming of energy metabolism plays an important role in
11 carcinogenesis and has been included as a hallmark of cancer by providing metabolic
12 targets for cancer treatment (Hanahan & Weinberg, 2011). Accordingly, several studies
13 demonstrate an aberrant expression of the enzyme fatty acid synthase (FASN) and
14 increased endogenous lipogenesis in several human cancers and its precursors,
15 including OSCC (Angeles & Hudkins, 2016; Menendez & Lupu, 2017; Silva et al., 2008;
16 Silva et al., 2009, Silva et al., 2010). FASN is the multi-enzyme protein responsible for
17 the *de novo* biosynthesis of long-chain fatty acids, especially palmitate, from acetyl-CoA
18 and malonyl-CoA (Angeles & Hudkins, 2016; Kuhajda, 2000). Structurally, FASN is
19 formed by two polypeptide chains (~270 kDa) containing seven different catalytic
20 domains sequentially organized from the N-terminal to the C-terminal: β -ketoacyl
21 synthase, acetyl-CoA transacylases and malonyl-CoA transacylases, dehydratase,
22 enoyl reductase, β -ketoacyl reductase, and thioesterase site and the acyl carrier protein
23 (Menendez & Lupu, 2017; Smith et al., 2003).

24 Under normal conditions, FASN levels in human tissues are usually low, since
25 the cells utilize circulating lipids mainly from the diet (Menendez et al., 2004). In contrast,
26 cancer cells highly express FASN to fuel membrane production, lipid-based post-
27 translational protein modifications, redox balance maintenance, and energy metabolism
28 (Menendez & Lupu, 2017; Röhrig & Schulze, 2016). In OSCC, FASN is overexpressed,
29 associated with overall survival rates, and with the histological grade, lymphatic
30 permeation, perineural infiltration, and lymph node metastasis (Silva et al., 2009; Silva
31 et al., 2008).

32 In the last decade, several natural or synthetic FASN inhibitors have been
33 described to promote anticancer effects by interfering in the lipogenic dependency
34 (Angeles & Hudkins, 2016; Buckley et al., 2017; Menendez & Lupu, 2017; Chu et al.,
35 2017). Among the most studied are cerulenin, and its synthetic derivative C75, the

1 antibiotic triclosan (TCS) and orlistat (ORL), a drug approved by FDA for the treatment
2 of obesity (Liu, 2006; Lupu & Menendez, 2006; Rendina & Cheng, 2005; Wang et al.,
3 2009). Cerulenin and C75 interact with the β -ketoacyl synthase domain and irreversibly
4 inhibit the condensation reaction and (Rendina & Cheng, 2005). TCS blocks FASN
5 activity through hydrogen bonding and hydrophobic interactions with specific amino
6 acids in the enoyl reductase domain (Liu et al., 2002; Menendez & Lupu, 2017). On the
7 other hand, ORL inhibits FASN activity through a covalent adduct with the thioesterase
8 domain (Kridel et al., 2004; Menendez & Lupu, 2017).

9 Importantly, pharmacological or genetic inhibition of FASN selectively induces
10 cancer cell death and reduction of tumor progression in mouse models (Angeles &
11 Hudkins, 2016; Jones & Infante, 2015; Menendez & Lupu, 2017). Besides the classical
12 FASN inhibitors, next-generation drugs have shown significant effects with high
13 selectivity. IPI-9119, a specific non-commercial inhibitor of the thioesterase domain of
14 FASN, is able to reduce the growth of prostate cancer xenografts and human organoids
15 by inducing substantial metabolic reprogramming (Zadra et al., 2019). TVB-3166 is an
16 imidazopyridine-based molecule which inhibits the β -ketoacyl reductase domain,
17 reduces proliferation of several cancer cell lines by interfering in oncogenic pathways,
18 especially PI3K-AKT-mTOR and β -catenin, and, when combined with paclitaxel,
19 decreases the growth of the prostate xenografts (Heuer et al., 2017; Oslob et al., 2013;
20 Ventura et al., 2015). Our group recently demonstrated that TVB-3166 decreases the
21 viability and migration and induces apoptosis and cell cycle arrest of OSCC (Aquino et
22 al., 2020). Moreover, TVB-2640 (ASC40), an analogue of TVB-3166, is the first FASN
23 inhibitor included in clinical trials for patients with solid tumors (NCT02223247), as well
24 as in phase II trials for colon (NCT02980029), KRAS mutated non-small cell lung
25 carcinomas (NCT03808558), astrocytomas (NCT03032484), ErbB2 positive breast
26 cancer (NCT03179904) and glioblastomas (NCT05118776). Finally, Omeprazole, an
27 inhibitor of proton pump, was recently reported to inhibit FASN in gastric epithelial cells
28 (Chen et al., 2020) and was safety when administrated for patients with triple negative
29 breast cancer in dosis that inhibited FASN (Sardesai et al., 2021). In fact, clinical trials
30 aiming to evaluate the response of this drug in castration resistant prostate cancer and
31 breast cancer are also being conducted (NCT04337580; NCT02595372).

32 Here we describe for the first time the effects of the pharmacological inhibition of
33 distinct catalytic domains of FASN with C75, TCS, and ORL in SCC-9 cells.

34

1 MATERIALS AND METHODS

2 Cell culture

3 Human OSCC (SCC-9) cells (American Type Culture Collection –ATCC, Manassas, VA,
4 USA) were originally isolated from a tongue squamous cell carcinoma of a 25 years-old
5 male and its tumorigenic phenotype is well documented in the literature by ours and other
6 groups (Agostini et al., 2014; de Andrade et al., 2017; Liu et al., 2002; Ramos et al.,
7 2002; Rheinwald & Beckett, 1981). SCC-9 were cultured in DMEM/F-12 medium
8 (Invitrogen, Carlsbad, CA, USA) supplemented with 2% or 10% FBS (Cultilab,
9 Campinas, Brazil), 400 ng/mL hydrocortisone (hydrocortisone sodium succinate -
10 Eurofarma, Brazil) and 1:100 antibiotic/antimycotic solution (Invitrogen) at 37°C in a
11 humidified atmosphere containing 5% CO₂. C75, TCS, ORL, and the vehicle DMSO
12 (Dimethylsulfoxide, Sigma, St Louis, MO, USA) were added to the culture medium at the
13 concentrations described in Table 1.

14 Proliferation assay and calculation of the IC₅₀

15 Cell viability was determined by seeding 8 x 10⁴ SCC-9 cells in 12-well plates with
16 DMEM/F-12 containing 10% FBS. After 24 hours, cells were serum-starved for additional
17 24 hours for cell cycle synchronization. After, the medium was replaced by 500 µL of
18 DMEM/F-12 supplemented with 2% FBS containing increasing concentrations of C75,
19 TCS or ORL (Sigma) for 24 or 48 hours. The volume of the vehicle DMSO was adjusted
20 for each concentration. Control cells were treated with the maximum volume of DMSO.
21 Cell viability was determined after the incubation with 25 mg/mL of MTT (4,5-
22 dimethylthiazol-2-yl) -2,5 diphenyltetrazolium bromide (Sigma) for 4 hours. After this
23 period, formazan crystals produced by viable cells were eluted in absolute ethanol and
24 the absorbance was determined with the aid of a microplate reader (Bio-Rad, Hercules,
25 CA, USA) at 550nm and corrected at 650 nm. Inhibitory concentration 50% (IC₅₀) values
26 were calculated by the dose-effect curve using the CompuSyn software (Cambridge, MA,
27 USA) as previously described (Chou & Martin, 2005).

28 Bodipy staining for lipid droplets

29 In order to confirm the FASN inhibition by C75, TCS, and ORL, we performed BODIPY™
30 (Invitrogen) staining for neutral lipids by plating 2 X 10³ SCC-9 cells in each well of 8-
31 well chamber slides (Lab-tek, Thermofisher-Nunc). After 24h, the medium was replaced
32 by 500 µL of DMEM-F12 without FBS and, following additional 24h, cells were incubated
33 with the concentrations corresponding to the IC_{12.5}, IC₂₅, and IC₅₀ of the studied drugs
34 diluted in DMEM/F12 with 2% FBS for 24 and 48h. After this period, lipid droplets were
35 stained by adding 3.8 µM of bodipy in the cell media (DMEMF/12 with 2% FBS) for 1

1 hour at 37°C. Cells were washed with PBS, fixed with absolute formaldehyde with 10%
2 CaCl₂ for 30 minutes, washed again, and the slides mounted in aqueous media. At least
3 10 micrographies for each condition were captured and analyzed in an epifluorescence
4 microscope (Leica DMR, Wetzlar, Germany). The number of lipid bodies was counted
5 and normalized by the corresponding cell area with the aid of the Image J software
6 (National Institute of Health, Bethesda, Maryland, USA).

7 **Cell cycle and apoptosis analysis**

8 For the flow cytometry analyses, cells were plated in 25cm² culture flasks, serum-starved
9 for 24h, and treated for 48h with three different concentrations of the FASN inhibitors.
10 Cells were then harvested by trypsinization and centrifuged at 900xg. To determine SCC-
11 9 distribution in each phase of the cell cycle, cells were fixed in cold 70% ethanol for at
12 least 16h at -20°C, and, after washing in cold PBS treated with 10 µg/ml RNase (Sigma)
13 at 37°C for 1h and incubated with 50 µg/mL propidium iodide at 4°C for 2 hours. To
14 determine the percentage of apoptotic and necrotic cells, cells were washed in PBS and
15 incubated with Annexin V-FITC (1:100) and 7-AADPerCP (1:200) in a binding buffer
16 containing 10 mM HEPES (pH 7.4), 150 mM NaCl, 5 mM KCl, 1 mM MgCl₂ and 1.8 mM
17 CaCl₂ in the dark for 20 minutes at room temperature. Ten thousand events were
18 acquired, analyzed by flow cytometry on FL-1 or FL-2 channel of a FACS Calibur (BD
19 Biosciences, San Jose, CA, USA) equipped with an argon laser for apoptosis and cell
20 cycle respectively. The distribution of cells in the cycle was evaluated with the software
21 ModFit (Verity Software House) and the percentage of apoptotic or necrotic cells
22 calculated by the software CellQuest (Becton Dickinson and Company, San Jose, CA,
23 USA).

24 **Cell adhesion assays**

25 Four wells of a 96 well plate were sensitized with 10 µg/cm² of type I collagen or
26 fibronectin diluted in 100 µL of PBS and as, negative controls, with 100 µL of PBS. The
27 plates were maintained for 24 hours at 4°C and, after this period, each well was washed
28 with PBS and blocked with 200 µL 3% BSA for 2h at 37°C. After the treatment with FASN
29 inhibitors for 48h, cells were trypsinized and resuspended in DMEM/F12 supplemented
30 with 3% BSA. For each condition, 3 X 10⁴ cells diluted in 100 µL of DMEM/F12
31 supplemented with 3% BSA were plated in each well of the previously sensitized plates
32 and incubated for 45 minutes to 1h. The period of adhesion was determined by
33 observation in a phase contrast microscope. Non-adherent cells were washed 3 times
34 with 100 µL of PBS and the adherent cells fixed with 4% formalin for 15 minutes at room
35 temperature. Cells were then stained with an aqueous solution of 1% toluidine blue and

1 1% borax for 5 minutes. The excess of dye was removed by vigorous washing in distilled
2 water. After elution in 100 μ L of a 1% SDS, the absorbances were determined at 655nm
3 in a microplate reader (Bio-Rad, Hercules, CA, USA).

4 **Transwell migration and invasion assays**

5 The effects of C75, TCS, and ORL in the migratory phenotype of SCC-9 cells were
6 evaluated by transwell migration assays. Prior to the plating, cells were serum-starved
7 for 24 hours and treated with the IC₅₀ of C75, TCS, or ORL for additional 24 hours. Briefly,
8 8×10^4 SCC-9 cells were resuspended in 200 μ L of FBS-free DMEM/ F-12 and added
9 in the upper compartment of each 8 μ M pore Transwell™ nylon filter membrane insert.
10 In the lower compartment of each transwell were added 1 mL of DMEM/F-12 containing
11 2% FBS and FASN inhibitors or vehicle. After 24 hours, migrating cells were fixed in 10%
12 formalin for 10 minutes and stained by 1% toluidine blue. The inserts were washed with
13 distilled water and the cells of the upper compartment carefully removed with distilled
14 water-soaked cotton swabs. After elution with 500 μ L of 1% SDS, the absorbances were
15 read at 655nm in a microplate reader (Bio-Rad). Invasion assays were performed as
16 previously described (Mendonça et al., 2017; Salo et al., 2015). First, 60 μ L of a 3.22
17 μ g/mL myogel solution were added in the upper chamber of 8- μ M pore Transwell™
18 inserts and solidified at 37°C for 30 minutes. After this period, serum-starved SCC-9 cells
19 (3×10^5 for each condition) were resuspended in serum-free medium containing the IC₅₀
20 concentrations of each FASN inhibitor or DMSO and seeded onto the insert with
21 solidified myogel. The lower compartments of the transwells were filled with 1 mL of
22 DMEM/F-12 supplemented with 2% FBS and the respective FASN inhibitors or the
23 vehicle. After 48h, invasive cells were fixed with 10% formalin for 10 minutes and stained
24 with 1% toluidine blue for 1 hour at room temperature. After washing the non-adherent
25 cells and elution with 500 μ L of 1% SDS, the absorbances were read at 650 nm in a
26 microplate reader (Bio-Rad).

27 **Statistical analysis**

28 All experiments were performed in technical replicates and repeated at least three times
29 independently. The appropriate statistical tests were applied according to the data
30 distribution. Normal distribution was verified using Shapiro-Wilk or D'Agostino-Pearson
31 (Bodipy quantification) tests and comparisons between groups were performed using
32 one-way ANOVA with Tukey's post-hoc analysis. The significance level was 95% ($p <$
33 0.05).

34

1 RESULTS

2

3 **C75, TCS, and ORL reduce SCC-9 proliferation and induce distinct morphologic** 4 **changes**

5 The IC₅₀ for each drug (Table 1) was calculated based on the amount of viable cells by
6 using dose-response curves with MTT assays (Figure 1). Cell death was measured by
7 the treatment with increasing concentrations of C75, TCS, and ORL for 24h (Figures 1A-
8 C) or 48h (Figure 1D-F) and, as expected, we observed a dose-dependent effect in all
9 analyzed conditions. C75 reduced the SCC-9 viability in a dose-dependent manner in
10 24h (Figure 1A) and 48h (Figure 1D) in concentrations lower than 30 μ M. On the other
11 hand, TCS was not effective in SCC-9 cells in lower concentrations but highly affected
12 the SCC-9 viability in concentrations above 36 μ M for 24h (Figure 1B) and 18 μ M for 48h
13 (Figure 1E). ORL reduced SCC-9 viability in concentrations higher than 100 μ M for 24h
14 (Figure 1C) and 50 μ M for 48h (Figure 1F). The morphology of SCC-9 treated with C75,
15 TCS, or ORL for 24h (Figure 1 G-J) or 48h (data not shown) was monitored using phase
16 contrast microscopy and compared with DMSO treated cells. Each drug promoted
17 distinct morphologic changes in SCC-9 cells, however, the morphology was not affected
18 by the period of treatment (24 or 48h). As seen in Figure 1G, SCC-9 cells incubated with
19 the vehicle showed an accentuated pleomorphism with the predominance of polygonal
20 shaped cells, large nuclei, and evident nucleoli. The treatment with C75 promoted an
21 evident cytoplasmatic enlargement and a decrease of the nucleus/cytoplasm ratio
22 (Figure 1H, arrows). SCC-9 cells treated with TCS showed an important reduction in cell
23 size and an increase of cytoplasmic granules (Figure 1I, arrows). Finally, treatment with
24 ORL promoted an evident cytoplasmic retraction and cell elongation compared to the
25 control group (Figure 1J). All compounds increased the number of detached rounded
26 cells characteristic of dead cells (Figures 1H-J, arrowheads).

27 **FASN inhibition reduce lipid droplets in SCC-9 cells**

28 In order to confirm that the effects of C75, TCS, and ORL on SCC-9 cells are associated
29 with the reduction of FASN activity, we used bodipy to stain intracellular lipid droplets.
30 These studies showed that, compared to the vehicle (Figure 2A and Figure S1A), all the
31 concentrations of C75 reduced the lipids droplets of SCC-9 cells in 24h (Figure S1) and
32 48h (Figure 2), and the concentration corresponding to its IC₅₀ had the highest effect on
33 FASN inactivation in 24h (83% of inhibition, Figures S1B, E). Similar effects were found
34 by treating SCC9 cells with C75 for 48h (Figure 2A-E). On the other hand, TCS showed
35 a dose-dependent effect and maximum FASN inhibition with the dose relative to the IC₅₀

1 when SCC9 were treated for 24h (95% of lipid droplets reduction, Figures S1C, F).
2 Similar results were obtained when the cells were treated for 48h (Figure 2F-J). ORL
3 was the less effective and reduced 28% and 38% of lipid bodies in SCC-9 cells treated
4 with its IC₂₅ and IC₅₀ for 24h, respectively (Figures S1D, G). We also did not observed
5 statistical differences when SCC9 were treated with the IC_{12.5} of ORL for 48h (Figure 2O).
6 However, the treatment with the IC₂₅ and the IC₅₀ of ORL for 48h of significantly reduced
7 the lipid bodies of SCC9 cells (Figure 2K-O).

8 **C75, TCS and ORL induces cell cycle arrest and apoptosis in SCC-9 cells**

9 In order to better characterize the effects of FASN inhibitors on SCC-9 cells, we next
10 performed cell cycle and apoptosis analysis by flow cytometry. The treatment with
11 different concentrations of C75 (Figure 3A), TCS (Figure 3B), and ORL (Figure 3C) for
12 48 h resulted in the accumulation of cells in the G0/G1 phase and significant reduction
13 of the S phase. The number of cells in the G2/M phases of the cell cycle was not affected
14 by the studied drugs (Figure 3A-C). The IC₅₀ of C75 increased in 23.19% the number of
15 cells in G0/G1 (Control: 45.05%; C75 IC₅₀: 68.25%) and reduced in 17.62% the number
16 of cells in the S phase (Control: 35.04%; C75 IC₅₀: 17.42%). The IC₅₀ of TCS resulted in
17 an accumulation of cells in G0/G1 of 35.81% (Control: 49.05%; TCS IC₅₀: 84.86%) while
18 the S phase was reduced in 26.8% (Control: 32.39%; TCS IC₅₀: 5.59%). Finally, cells in
19 G0/G1 were increased by ORL in 31.58% (Control: 52.08%; ORL IC₅₀: 83.67%) and the
20 S phase reduced in 24.05% (Control: 29.82; ORL IC₅₀: 5.77%). To compare the effects
21 of the studied drugs on proliferating SCC-9 cells, we performed statistical analyzes with
22 the number of cells in S phase (Figures 3D-F) and found that TCS and ORL similarly
23 affected cell cycle progression.

24 C75 (Figure 4A), TCS (Figure 4B), and ORL (Figure 4C) significantly reduce the
25 number of viable cells and increase the number of apoptotic cells in a dose-dependent
26 manner. As expected, the highest effect was observed in the concentration relative to
27 the IC₅₀, in which C75 (Figure 4A) and ORL (Figure 4C) increased the apoptotic levels in
28 5.5X and 4.58X, respectively (Control: 1.63; C75: 8.94 and Control: 1.13; ORL: 5.17).
29 TCS showed the highest effect with an increase of the apoptotic cells in 21.89X (Control:
30 1.08; TCS IC₅₀: 21.89) (Figure 4B). At low concentrations, TCS was also effective
31 promoting a dose dependent increase in apoptotic levels (1.75X for the IC_{12.5} and 8.42X
32 for the IC₂₅). Statistical analysis comparing the effects of the drugs using the
33 concentration relative to the IC₅₀ showed that TCS was the most effective drug in
34 reducing the number of live cells (Figure 4D) and increasing the percentage of apoptotic
35 cells (Figure 4E). Necrosis was not significantly affected by the drugs. Gate information
36 were provided in the Figures S2 (cell cycle) and S3 (apoptosis).

1 **Adhesion, migration, and invasion of SCC-9 cells were differently affected by C75,**
2 **TCS, and ORL**

3 In order to verify the effects of C75, TCS, and ORL on the adhesion of SCC-9 cells to
4 extracellular matrix components, cells were treated for 48h with different concentrations
5 of the drugs and allowed to adhere in previously sensitized plates with fibronectin or type
6 I collagen. These studies showed a slight increase, not statistically significant, in the
7 adhesion of SCC-9 cells treated with the IC_{12.5}. No significant reduction in the cell
8 adhesion to the matrix components was observed in the IC₂₅ and IC₅₀ concentrations of
9 C75 (Figure 5A). The adhesion of SCC-9 cells was significantly reduced with TCS in a
10 dose-dependent manner (Figure 5B) while ORL did not affect the adhesion to both type
11 I collagen and fibronectin (Figure 5C).

12 By using transwell cell migration (Figure 6A-D) and invasion (Figure 6F-I) assays,
13 we observed that the treatment with the IC₅₀ of C75, TCS, and ORL for 48h reduced in
14 56%, 65.7%, and 31% the migration of SCC-9 cells, respectively (Figure 6E). The
15 invasive ability of SCC-9 cells were similarly reduced by C75 and TCS and not modified
16 by ORL, when compared with the control cells (Figure 6J).

17

18

1 DISCUSSION

2

3 FASN is the key enzyme of the endogenous lipid metabolism and its activity is associated
4 with signaling pathways and maintenance of the structure and function of cancer cell
5 membranes (Buckley et al., 2017). Upregulation of FASN was demonstrated in several
6 human malignancies, including oral cancer (Agostini et al., 2014; Aquino et al., 2020;
7 Kelber et al., 2010; Ricklefs et al., 2020; Sadowski et al., 2014; Sippel et al., 2014). FASN
8 has emerged as a therapeutic target due to its differential expression in normal and
9 malignant tissues and its association with survival rates and prognosis (Röhrig &
10 Schulze, 2016; Witkiewicz et al., 2008). Importantly, the multiple catalytic domains of
11 FASN allowed the development of several FASN inhibitors through the last decade
12 (Röhrig & Schulze, 2016). The first generation of FASN inhibitors provided the basis for
13 the development and testing of new FASN-targeted drugs and clarifying the role of FASN
14 and lipid metabolism in cancer cells (Menendez & Lupu, 2017). Classic FASN inhibitors,
15 such as C75, TCS (triclosan), and ORL (orlistat), were reported to promote cell death *in*
16 *vitro* and *in vivo* in several types of cancer cells, including OSCC (Agostini et al., 2014;
17 Bastos et al., 2017; Menendez & Lupu, 2017; Zadra et al., 2019).

18 SCC-9, a model of OSCC shows a tumorigenic phenotype, documented in the
19 literature by ours and other groups (Rheinwad and Beckett, 1981; Ramos et al., 2001;
20 Liu et al., 2002; De Andrade et al., 2017; Agostini et al., 2014). Here, we demonstrated
21 that ORL has the highest IC₅₀ when compared with C75 and TCS. Our group has
22 previously shown the anti-tumoral effects of ORL in melanoma (Bastos et al., 2017;
23 Carvalho et al., 2008; Seguin et al., 2012) and OSCC cells lines (Agostini et al., 2014)
24 by using similarly high concentrations. On the other hand, a 50% reduction of the SCC-
25 9 viability was found at smaller concentrations of C75 and TCS in both, 24h and 48h.
26 There is no data in the literature regarding the effects of C75 and TCS in oral cancer
27 cells. Though, our IC₅₀ concentrations for C75 were similar to the results of previous
28 studies in cells lines derived from breast, lung, and prostate cancers (Chen et al., 2012;
29 Giró-Perafita et al., 2016; Menendez et al., 2016; Puig et al., 2008; Relat et al., 2012;
30 Sadowski et al., 2014). Also, the IC₅₀ for TCS in retinoblastoma cells is 60 µM and
31 between 4.5 to 6.9 µM in different prostate cancer cell lines (Deepa et al., 2012;
32 Sadowski et al., 2014; Vandhana et al., 2013).

33 In our work, C75 reduced lipid droplets similarly at lower and higher TCS showed
34 a dose-dependent reduction in lipogenesis. ORL also reduced the lipid bodies at its IC₂₅
35 and IC₅₀. Despite their distinct rates of FASN inhibition, all drugs reduced the S phase

1 and increased the number of SCC-9 cells in the G0/G1 phases. Accordingly, several
2 studies have demonstrated that FASN inhibitors promote a cell cycle arrest in several
3 human cancer cell lines (Veigel et al., 2015). Agostini et al. (2014) demonstrated that the
4 S phase of SCC-9 cells treated with ORL extracted from Orlistate (NeoQuímica
5 Farmacêutica, Brazil) capsules for 72 h was reduced by approximately 75%. Similarly,
6 Aquino et al. (2020) showed that TVB-3166, a new imidazopyridine-based FASN
7 inhibitor, reduces the number of cells in the S phase by 57% in a highly metastatic OSCC
8 carcinoma cell line LN-1A. The mechanisms by which FASN inhibitors affect the cell
9 cycle progression are not well described. Zadra et al. (2019) demonstrated that the
10 treatment of prostate cancer cell lines with IPI-9119 reduced the cell cycle kinetics by
11 decreasing the levels of cyclin A2. Other studies have shown that the inhibition of FASN
12 in cancer cell lines may affect several proteins involved in cell cycle control (Agostini et
13 al., 2014; Deepa et al., 2013; Sangeetha et al., 2015; Ventura et al., 2015). Scaglia et al.
14 (2014) observed with the incorporation of ¹⁴C choline that the levels of fatty acids in cell
15 membranes increase during cell cycle progression and reached the peak in G2/M.
16 According to the same authors, the presence of a lipogenic checkpoint in the G2/M phase
17 could explain, at least in part, the reduction of the cell cycle progression in FASN-
18 inhibited tumor cells.

19 We also observed here that the accumulation of SCC-9 cells in G0/G1 and
20 inhibition of S phase was accompanied by apoptosis. Although the treatment with the
21 low concentrations of C75 and ORL promotes cell cycle arrest, apoptotic levels were
22 slightly or not affected. Conversely, IC₅₀ concentration of these drugs significantly
23 increased apoptosis. In fact, TCS was the most effective FASN inhibitor in order to
24 increase apoptosis. TCS was effective even at low concentrations and produced a dose-
25 dependent increase in apoptotic levels. TCS reduces cell growth and viability of tumor
26 cell lines, such as breast cancer (MCF-7 and SKBr-3) and retinoblastoma (Y79) (Deepa
27 et al., 2012; Liu et al., 2002). According to our findings, Sadowski et al., (2014) compared
28 the inhibitory effect of TCS, C75, and ORL in normal and prostate cancer cell lines and
29 found that TCS has a greater cytotoxic effect and the lowest IC₅₀. The biological
30 mechanisms that underlie the cell cycle arrest and apoptosis promoted by the FASN
31 inhibitors need future mechanistic investigation. However, previous studies of our group
32 has shown that FASN activity is necessary for proliferation and survival of OSCC cells
33 considering that both, pharmacological or specific FASN inhibition, reduced the total lipid
34 biosynthesis in OSCC cells (by incorporation studies of ¹⁴C-acetate).

35 Here we found that while ORL did not affect cell adhesion and invasion and
36 slightly decreases migration of SCC-9 cells, C75 and TCS reduced in more than 50%

1 migration and invasion. In addition, C75 did not significantly affected the adhesion of
2 SCC-9 to type I collagen and fibronectin and TCS promoted a dose-dependent reduction
3 of cell adhesion to the studied extracellular matrix components. The role of FASN in
4 adhesion, migration, and invasion processes are still not clear. Our group has shown
5 that TVB-3166 significantly increases adhesion of SCC-9 and LN-1A cells to myogel, a
6 matrix prepared from human uterine leiomyoma composed mainly by fibronectin,
7 laminin, integrins and different collagens but reduced migration of SCC-9 cells. Zaytseva
8 et al., (2012) showed that FASN inhibition with siRNA in colon cancer cells reduces the
9 levels of MET, Akt, FAK, and paxillin, which are involved in cell adhesion and invasion.
10 Jafari et al., (2019) suggested that the inhibition of FASN by TVB-3664 decreases the
11 number of adhesion plaques in primary and metastatic colorectal cancer cells by
12 immunofluorescence reactions for p-FAK and p-paxillin. Recently, De Piano et al., (2020)
13 observed that FASN depletion using shRNA increases cell adhesion to matrigel and type
14 I collagen in prostate cancer cell lines (CPTX 1542 and PC3) by increasing the paxillin
15 levels and decreasing the palmitoylation of RhoU. Considering the controversial results
16 in the literature and the different effects of C75, TCS, and ORL it is possible to speculate
17 that cell type and the method used for the FASN inhibition can modulate the effect on
18 cell adhesion. Despite the promising results *in vitro* and/or *in vivo*, is important to
19 emphasize that C75, TCS and ORL are not available for clinical trials of cancer and
20 demonstrated limitations of pharmacological properties and side-effects (Menendez &
21 Lupu, 2017). In fact, new formulations using nanoparticles would be helpful to improve
22 the drug permeability, absorption and bioavailability of the compounds (Chu et al., 2016).

23 In summary, we found that C75, TCS, and ORL have distinct effects in the
24 tumorigenic phenotype of SCC-9 cells. C75 has the lowest IC_{50} , better inhibition of FASN
25 at low concentrations, and an important reduction of the cell motility phenotypes,
26 especially migration and invasion. TCS has the intermediate IC_{50} and an excellent
27 reduction of lipid bodies at its IC_{50} . TCS also significantly reduces cell cycle progression,
28 adhesion, migration, and invasion while induces high levels of apoptosis in SCC-9 cells.
29 ORL is effective in promoting cell cycle arrest, but induces low levels of apoptosis and
30 did not affect invasion and adhesion. These results indicate that TCS interfere in all the
31 tumorigenic phenotypes of SCC-9 cells.

32

1 **Author contributions**

2 Willian Peter Boelcke – planned and performed experiments, analyzed data, wrote and
3 approved the final version of the manuscript.

4 Isadora Ferrari Teixeira – planned and performed experiments, analyzed data, wrote and
5 approved the final version of the manuscript.

6 Iara Gonçalves de Aquino – performed experiments, revised and approved the final
7 version of the manuscript.

8 Amanda Mazzaro – analyzed data, revised and approved the final version of the
9 manuscript.

10 Florence Juana Maria Cuadra-Zelaya – analysed data, revised and approved the final
11 version of the manuscript.

12 Ana Paula Souza - analyzed data, revised and approved the final version of the
13 manuscript.

14 Tuula Salo - analyzed data, revised and approved the final version of the manuscript.

15 Ricardo Della Coletta – analyzed data, discussed the results, revised and approved the
16 final version of the manuscript.

17 Edgard Graner - analyzed data, planned the experiments, discussed the results, revised
18 and approved the final version of the manuscript.

19 Débora Campanella Bastos – conceived, planned and performed experiments, analyzed
20 data, wrote, revised and approved the final version of the manuscript.

21

22

23 **Acknowledgements**

24 This work was supported by the Fundação de Amparo à Pesquisa do Estado de São
25 Paulo (FAPESP, 2014/20832-3, 2016/07129-7 and 2016/ 24906-7) and Coordenação
26 de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES -
27 88882.306833/2018-01 and 88887.352647/2019-00).

28

1 **References**

- 2 Agostini, M., Almeida, L. Y., Bastos, D. C., Ortega, R. M., Moreira, F. S., Seguin, F., ...
3 Graner, E. (2014). The fatty acid synthase inhibitor orlistat reduces the growth and
4 metastasis of orthotopic tongue oral squamous cell carcinomas. *Molecular Cancer*
5 *Therapeutics*, 13(3), 585–595. <https://doi.org/10.1158/1535-7163.MCT-12-1136>
- 6 Angeles, T. S., & Hudkins, R. L. (2016). Recent advances in targeting the fatty acid
7 biosynthetic pathway using fatty acid synthase inhibitors. *Expert Opinion on Drug*
8 *Discovery*, 11(12), 1187–1199. <https://doi.org/10.1080/17460441.2016.1245286>
- 9 Aquino, I. G. de, Bastos, D. C., Cuadra-Zelaya, F. J. M., Teixeira, I. F., Salo, T.,
10 Coletta, R. Della, & Graner, E. (2020). Anticancer properties of the fatty acid
11 synthase inhibitor TVB-3166 on oral squamous cell carcinoma cell lines. *Archives*
12 *of Oral Biology*, 113(December 2019), 104707.
13 <https://doi.org/10.1016/j.archoralbio.2020.104707>
- 14 Bastos, D. C., Paupert, J., Maillard, C., Seguin, F., Carvalho, M. A., Agostini, M., ...
15 Graner, E. (2017). Effects of fatty acid synthase inhibitors on lymphatic vessels:
16 an in vitro and in vivo study in a melanoma model. *Laboratory Investigation; a*
17 *Journal of Technical Methods and Pathology*, 97(2), 194–206.
18 <https://doi.org/10.1038/labinvest.2016.125>
- 19 Buckley, D., Duke, G., Heuer, T. S., O'Farrell, M., Wagman, A. S., McCulloch, W., &
20 Kemble, G. (2017). Fatty acid synthase - Modern tumor cell biology insights into a
21 classical oncology target. *Pharmacology & Therapeutics*, 177, 23–31.
22 <https://doi.org/10.1016/j.pharmthera.2017.02.021>
- 23 Carvalho, M. A., Zecchin, K. G., Seguin, F., Bastos, D. C., Agostini, M., Rangel, A. L.
24 C. A., ... Graner, E. (2008). Fatty acid synthase inhibition with Orlistat promotes
25 apoptosis and reduces cell growth and lymph node metastasis in a mouse
26 melanoma model. *International Journal of Cancer*, 123(11), 2557–2565.
27 <https://doi.org/10.1002/ijc.23835>
- 28 Chen, M., Huang, J., Yang, X., Liu, B., Zhang, W., Huang, L., ... Ge, J. (2012). Serum
29 starvation induced cell cycle synchronization facilitates human somatic cells
30 reprogramming. *PLoS ONE*, 7(4). <https://doi.org/10.1371/journal.pone.0028203>
- 31 Chen, P., Li, L., Wang, H., Zhao, J., Cheng, Y., Xie, J., ... Gong, S. (2020).
32 Omeprazole, an inhibitor of proton pump, suppresses De novo lipogenesis in
33 gastric epithelial cells. *Biomedicine and Pharmacotherapy*, 130(June), 110472.
34 <https://doi.org/10.1016/j.biopha.2020.110472>
- 35 Chu, C., Deng, J., Liu, L., Cao, Y., Wei, X., Li, J., & Man, Y. (2016). Nanoparticles
36 combined with growth factors: Recent progress and applications. *RSC Advances*,
37 6(93), 90856–90872. <https://doi.org/10.1039/c6ra13636b>
- 38 Chu, C., Deng, J., Man, Y., & Qu, Y. (2017). Green Tea Extracts Epigallocatechin-3-
39 gallate for Different Treatments. *BioMed Research International*, 2017.
40 <https://doi.org/10.1155/2017/5615647>
- 41 Cohen Goldemberg, D., de Araújo, L. H. L., Antunes, H. S., de Melo, A. C., & Santos
42 Thuler, L. C. (2018). Tongue cancer epidemiology in Brazil: incidence, morbidity
43 and mortality. *Head and Neck*, 40(8), 1834–1844.
44 <https://doi.org/10.1002/hed.25166>
- 45 de Andrade, N. P., Rodrigues, M. F. S. D., Rodini, C. O., & Nunes, F. D. (2017).
46 Cancer stem cell, cytokeratins and epithelial to mesenchymal transition markers
47 expression in oral squamous cell carcinoma derived from orthotopic

- 1 xenoimplantation of CD44high cells. *Pathology Research and Practice*, 213(3),
2 235–244. <https://doi.org/10.1016/j.prp.2016.12.009>
- 3 De Piano, M., Manuelli, V., Zadra, G., Otte, J., Edqvist, P. H. D., Pontén, F., ... Wells,
4 C. M. (2020). Lipogenic signalling modulates prostate cancer cell adhesion and
5 migration via modification of Rho GTPases. *Oncogene*, 39(18), 3666–3679.
6 <https://doi.org/10.1038/s41388-020-1243-2>
- 7 Deepa, P. R., Vandhana, S., Jayanthi, U., & Krishnakumar, S. (2012). Therapeutic and
8 Toxicologic Evaluation of Anti-Lipogenic Agents in Cancer Cells Compared with
9 Non-Neoplastic Cells. *Basic and Clinical Pharmacology and Toxicology*, 110(6),
10 494–503. <https://doi.org/10.1111/j.1742-7843.2011.00844.x>
- 11 Deepa, P. R., Vandhana, S., & Krishnakumar, S. (2013). Fatty Acid Synthase Inhibition
12 Induces Differential Expression of Genes Involved in Apoptosis and Cell
13 Proliferation in Ocular Cancer Cells. *Nutrition and Cancer*, 65(2), 311–316.
14 <https://doi.org/10.1080/01635581.2013.748923>
- 15 Giró-Perafita, A., Palomeras, S., Lum, D. H., Blancafort, A., Viñas, G., Oliveras, G., ...
16 Puig, T. (2016). Preclinical evaluation of fatty acid synthase and egrinhibition in
17 triple-negative breast cancer. *Clinical Cancer Research*, 22(18), 4687–4697.
18 <https://doi.org/10.1158/1078-0432.CCR-15-3133>
- 19 Hanahan, D., & Weinberg, R. A. (2011). Hallmarks of cancer: The next generation.
20 *Cell*, 144(5), 646–674. <https://doi.org/10.1016/j.cell.2011.02.013>
- 21 Heuer, T. S., Ventura, R., Mordec, K., Lai, J., Fridlib, M., Buckley, D., & Kemble, G.
22 (2017). FASN Inhibition and Taxane Treatment Combine to Enhance Anti-tumor
23 Efficacy in Diverse Xenograft Tumor Models through Disruption of Tubulin
24 Palmitoylation and Microtubule Organization and FASN Inhibition-Mediated
25 Effects on Oncogenic Signaling and Gene Ex. *EBioMedicine*, 16, 51–62.
26 <https://doi.org/10.1016/j.ebiom.2016.12.012>
- 27 Ho, A. S., Kim, S., Tighiouart, M., Gudino, C., Mita, A., Scher, K. S., ... Zumsteg, Z. S.
28 (2017). Metastatic lymph node burden and survival in oral cavity cancer. *Journal*
29 *of Clinical Oncology*, 35(31), 3601–3609.
30 <https://doi.org/10.1200/JCO.2016.71.1176>
- 31 Jafari, N., Drury, J., Morris, A. J., Onono, F. O., Stevens, P. D., Gao, T., ... Zaytseva,
32 Y. Y. (2019). De Novo Fatty Acid Synthesis-Driven Sphingolipid Metabolism
33 Promotes Metastatic Potential of Colorectal Cancer. *Molecular Cancer Research* :
34 *MCR*, 17(1), 140–152. <https://doi.org/10.1158/1541-7786.MCR-18-0199>
- 35 Jones, S. F., & Infante, J. R. (2015). Molecular Pathways: Fatty Acid Synthase. *Clinical*
36 *Cancer Research : An Official Journal of the American Association for Cancer*
37 *Research*, 21(24), 5434–5438. <https://doi.org/10.1158/1078-0432.CCR-15-0126>
- 38 Kelber, J. A., Cao, H. S. T., Cantin, G. T., Lin, R., Wang, W., Kaushal, S., ... Yates, J.
39 R. (2010). Correction for Chiaroni et al., Y chromosome diversity, human
40 expansion, drift, and cultural evolution. *Proceedings of the National Academy of*
41 *Sciences*, 107(30), 13556–13556. <https://doi.org/10.1073/pnas.1008738107>
- 42 Kridel, S. J., Axelrod, F., Rozenkrantz, N., & Smith, J. W. (2004). Orlistat is a novel
43 inhibitor of fatty acid synthase with antitumor activity. *Cancer Research*, 64(6),
44 2070–2075. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15026345>
- 45 Kuhajda, F. P. (2000). Fatty-acid synthase and human cancer: New perspectives on its
46 role in tumor biology. *Nutrition*, 16(3), 202–208. [https://doi.org/10.1016/S0899-](https://doi.org/10.1016/S0899-9007(99)00266-X)
47 [9007\(99\)00266-X](https://doi.org/10.1016/S0899-9007(99)00266-X)

- 1 Liu, B., Wang, Y., Fillgrove, K. L., & Anderson, V. E. (2002). Triclosan inhibits enoyl-
2 reductase of type I fatty acid synthase in vitro and is cytotoxic to MCF-7 and SKBr-
3 3 breast cancer cells. *Cancer Chemotherapy and Pharmacology*, 49(3), 187–193.
4 <https://doi.org/10.1007/s00280-001-0399-x>
- 5 Liu, Y. (2006). Fatty acid oxidation is a dominant bioenergetic pathway in prostate
6 cancer. *Prostate Cancer and Prostatic Diseases*, 9(3), 230–234.
7 <https://doi.org/10.1038/sj.pcan.4500879>
- 8 Lupu, R., & Menendez, J. A. (2006). Pharmacological inhibitors of Fatty Acid Synthase
9 (FASN)--catalyzed endogenous fatty acid biogenesis: a new family of anti-cancer
10 agents? *Current Pharmaceutical Biotechnology*, 7(6), 483–493. Retrieved from
11 <http://www.ncbi.nlm.nih.gov/pubmed/17168665>
- 12 Mendonça, B. D. S., Agostini, M., Aquino, I. G., Dias, W. B., Bastos, D. C., &
13 Rumjanek, F. D. (2017). Suppression of MAGE-A10 alters the metastatic
14 phenotype of tongue squamous cell carcinoma cells. *Biochemistry and Biophysics*
15 *Reports*, 10, 267–275. <https://doi.org/10.1016/j.bbrep.2017.04.009>
- 16 Menendez, J. A., & Lupu, R. (2017). Fatty acid synthase regulates estrogen receptor- α
17 signaling in breast cancer cells. *Oncogenesis*, 6(2), e299-11.
18 <https://doi.org/10.1038/oncsis.2017.4>
- 19 Menendez, Javier A., & Lupu, R. (2017). Fatty acid synthase (FASN) as a therapeutic
20 target in breast cancer. *Expert Opinion on Therapeutic Targets*, 21(11), 1001–
21 1016. <https://doi.org/10.1080/14728222.2017.1381087>
- 22 Menendez, Javier A., Vellon, L., Espinoza, I., & Lupu, R. (2016). The metastasis
23 inducer CCN1 (CYR61) activates the fatty acid synthase (FASN)-driven lipogenic
24 phenotype in breast cancer cells. *Oncoscience*, 3(7–8), 242–257.
25 <https://doi.org/10.18632/oncoscience.314>
- 26 Menendez, Javier A., Vellon, L., Mehmi, I., Oza, B. P., Ropero, S., Colomer, R., &
27 Lupu, R. (2004). Inhibition of fatty acid synthase (FAS) suppresses HER2/neu
28 (erbB-2) oncogene overexpression in cancer cells. *Proceedings of the National*
29 *Academy of Sciences of the United States of America*, 101(29), 10715–10720.
30 <https://doi.org/10.1073/pnas.0403390101>
- 31 Montero, P. H., & Patel, S. G. (2015). Cancer of the Oral Cavity. *Surgical Oncology*
32 *Clinics of North America*, 24(3), 491–508.
33 <https://doi.org/10.1016/j.soc.2015.03.006>
- 34 Oslob, J. D., Johnson, R. J., Cai, H., Feng, S. Q., Hu, L., Kosaka, Y., ... McDowell, R.
35 S. (2013). Imidazopyridine-Based Fatty Acid Synthase Inhibitors That Show Anti-
36 HCV Activity and in Vivo Target Modulation. *ACS Medicinal Chemistry Letters*,
37 4(1), 113–117. <https://doi.org/10.1021/ml300335r>
- 38 Puig, T., Vázquez-Martín, A., Relat, J., Pétriz, J., Menéndez, J. A., Porta, R., ...
39 Colomer, R. (2008). Fatty acid metabolism in breast cancer cells: differential
40 inhibitory effects of epigallocatechin gallate (EGCG) and C75. *Breast Cancer*
41 *Research and Treatment*, 109(3), 471–479. [https://doi.org/10.1007/s10549-007-](https://doi.org/10.1007/s10549-007-9678-5)
42 [9678-5](https://doi.org/10.1007/s10549-007-9678-5)
- 43 Ramos, D. M., But, M., Regezi, J., Schmidt, B. L., Atakilit, A., Dang, D., ... Li, X.
44 (2002). Expression of integrin $\beta 6$ enhances invasive behavior in oral squamous
45 cell carcinoma. *Matrix Biology*, 21(3), 297–307. [https://doi.org/10.1016/S0945-](https://doi.org/10.1016/S0945-053X(02)00002-1)
46 [053X\(02\)00002-1](https://doi.org/10.1016/S0945-053X(02)00002-1)
- 47 Relat, J., Blancafort, A., Oliveras, G., Cufí, S., Haro, D., Marrero, P. F., & Puig, T.

- 1 (2012). Different fatty acid metabolism effects of (-)-epigallocatechin-3-gallate and
2 C75 in adenocarcinoma lung cancer. *BMC Cancer*, 12, 280.
3 <https://doi.org/10.1186/1471-2407-12-280>
- 4 Rendina, A. R., & Cheng, D. (2005). Characterization of the inactivation of rat fatty acid
5 synthase by C75: Inhibition of partial reactions and protection by substrates.
6 *Biochemical Journal*, 388(3), 895–903. <https://doi.org/10.1042/BJ20041963>
- 7 Rheinwald, J. G., & Beckett, M. A. (1981). Tumorigenic Keratinocyte Lines Requiring
8 Anchorage and Fibroblast Support Cultured from Human Squamous Cell
9 Carcinomas. *Cancer Research*, 41(5), 1657–1663.
- 10 Ricklefs, F. L., Maire, C. L., Matschke, J., Dührsen, L., Sauvigny, T., Holz, M., ...
11 Lamszus, K. (2020). Fasn is a biomarker enriched in malignant glioma-derived
12 extracellular vesicles. *International Journal of Molecular Sciences*, 21(6).
13 <https://doi.org/10.3390/ijms21061931>
- 14 Röhrig, F., & Schulze, A. (2016). The multifaceted roles of fatty acid synthesis in
15 cancer. *Nature Reviews. Cancer*, 16(11), 732–749.
16 <https://doi.org/10.1038/nrc.2016.89>
- 17 Sadowski, M. C., Pouwer, R. H., Gunter, J. H., Lubik, A. A., Quinn, R. J., & Nelson, C.
18 C. (2014). The fatty acid synthase inhibitor triclosan: repurposing an anti-microbial
19 agent for targeting prostate cancer. *Oncotarget*, 5(19), 9362–9381.
20 <https://doi.org/10.18632/oncotarget.2433>
- 21 Salo, T., Sutinen, M., Hoque Apu, E., Sundquist, E., Cervigne, N. K., de Oliveira, C. E.,
22 ... Coletta, R. D. (2015). A novel human leiomyoma tissue derived matrix for cell
23 culture studies. *BMC Cancer*, 15(1), 1. <https://doi.org/10.1186/s12885-015-1944-z>
- 24 Sangeetha, M., Deepa, P. R., Rishi, P., Khetan, V., & Krishnakumar, S. (2015). Global
25 Gene Deregulations in FASN Silenced Retinoblastoma Cancer Cells: Molecular
26 and Clinico-Pathological Correlations. *Journal of Cellular Biochemistry*, 116(11),
27 2676–2694. <https://doi.org/10.1002/jcb.25217>
- 28 Sardesai, S. D., Thomas, A., Gallagher, C., Lynce, F., Ottaviano, Y. L., Ballinger, T. J.,
29 ... Miller, K. D. (2021). Inhibiting Fatty Acid Synthase with Omeprazole to Improve
30 Efficacy of Neoadjuvant Chemotherapy in Patients with Operable TNBC. *Clinical
31 Cancer Research : An Official Journal of the American Association for Cancer
32 Research*, 27(21), 5810–5817. <https://doi.org/10.1158/1078-0432.CCR-21-0493>
- 33 Scaglia, N., Tyekucheva, S., Zadra, G., Photopoulos, C., & Loda, M. (2014). De novo
34 fatty acid synthesis at the mitotic exit is required to complete cellular division. *Cell
35 Cycle*, 13(5), 859–868. <https://doi.org/10.4161/cc.27767>
- 36 Seguin, F., Carvalho, M. A., Bastos, D. C., Agostini, M., Zecchin, K. G., Alvarez-Flores,
37 M. P., ... Graner, E. (2012). The fatty acid synthase inhibitor orlistat reduces
38 experimental metastases and angiogenesis in B16-F10 melanomas. *British
39 Journal of Cancer*, 107(6). <https://doi.org/10.1038/bjc.2012.355>
- 40 Silva, Sabrina Daniela da, Cunha, I. W., Nishimoto, I. N., Soares, F. A., Carraro, D. M.,
41 Kowalski, L. P., & Graner, E. (2009). Clinicopathological significance of ubiquitin-
42 specific protease 2a (USP2a), fatty acid synthase (FASN), and ErbB2 expression
43 in oral squamous cell carcinomas. *Oral Oncology*, 45(10), e134–e139.
44 <https://doi.org/10.1016/j.oraloncology.2009.02.004>
- 45 Silva, S D, Cunha, I. W., Younes, R. N., Soares, F. A., Kowalski, L. P., & Graner, E.
46 (2010). ErbB receptors and fatty acid synthase expression in aggressive head and
47 neck squamous cell carcinomas. *Oral Diseases*, 16(8), 774–780.

- 1 <https://doi.org/10.1111/j.1601-0825.2010.01687.x>
- 2 Silva, S D, Perez, D. E., Nishimoto, I. N., Alves, F. A., Pinto, C. A. L., Kowalski, L. P., &
3 Graner, E. (2008). Fatty acid synthase expression in squamous cell carcinoma of
4 the tongue: clinicopathological findings. *Oral Diseases*, *14*(4), 376–382.
5 <https://doi.org/10.1111/j.1601-0825.2007.01395.x>
- 6 Sippel, K. H., Vyas, N. K., Zhang, W., Sankaran, B., & Quiocho, F. A. (2014). Crystal
7 structure of the human fatty acid synthase enoyl-acyl carrier protein-reductase
8 domain complexed with triclosan reveals allosteric protein-protein interface
9 inhibition. *Journal of Biological Chemistry*, *289*(48), 33287–33295.
10 <https://doi.org/10.1074/jbc.M114.608547>
- 11 Smith, S., Witkowski, A., & Joshi, A. K. (2003). Structural and functional organization of
12 the animal fatty acid synthase. *Progress in Lipid Research*, *42*(4), 289–317.
13 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12689621>
- 14 Vandhana, S., Coral, K., Jayanthi, U., Deepa, P. R., & Krishnakumar, S. (2013).
15 Biochemical changes accompanying apoptotic cell death in retinoblastoma cancer
16 cells treated with lipogenic enzyme inhibitors. *Biochimica et Biophysica Acta -*
17 *Molecular and Cell Biology of Lipids*, *1831*(9), 1458–1466.
18 <https://doi.org/10.1016/j.bbalip.2013.06.005>
- 19 Veigel, D., Wagner, R., Stübiger, G., Wuczkowski, M., Filipits, M., Horvat, R., ... Grunt,
20 T. W. (2015). Fatty acid synthase is a metabolic marker of cell proliferation rather
21 than malignancy in ovarian cancer and its precursor cells. *International Journal of*
22 *Cancer*, *136*(9), 2078–2090. <https://doi.org/10.1002/ijc.29261>
- 23 Ventura, R., Mordec, K., Waszczuk, J., Wang, Z., Lai, J., Fridlib, M., ... Heuer, T. S.
24 (2015). Inhibition of de novo Palmitate Synthesis by Fatty Acid Synthase Induces
25 Apoptosis in Tumor Cells by Remodeling Cell Membranes, Inhibiting Signaling
26 Pathways, and Reprogramming Gene Expression. *EBioMedicine*, *2*(8), 808–824.
27 <https://doi.org/10.1016/j.ebiom.2015.06.020>
- 28 Wang, X., Lin, J., Chen, Y., Zhong, W., Zhao, G., Liu, H., ... Li, S. (2009). Novel fatty
29 acid synthase (FAS) inhibitors: Design, synthesis, biological evaluation, and
30 molecular docking studies. *Bioorganic and Medicinal Chemistry*, *17*(5), 1898–
31 1904. <https://doi.org/10.1016/j.bmc.2009.01.050>
- 32 Witkiewicz, A. K., Nguyen, K. H., Dasgupta, A., Kennedy, E. P., Yeo, C. J., Lisanti, M.
33 P., & Brody, J. R. (2008). Co-expression of fatty acid synthase and caveolin-1 in
34 pancreatic ductal adenocarcinoma: Implications for tumor progression and clinical
35 outcome. *Cell Cycle*, *7*(19), 3021–3025. <https://doi.org/10.4161/cc.7.19.6719>
- 36 Zadra, G., & Loda, M. (2019). When fat goes down, prostate cancer is on the ropes.
37 *Molecular and Cellular Oncology*, *6*(3), 1–3.
38 <https://doi.org/10.1080/23723556.2019.1595308>
- 39 Zadra, G., Ribeiro, C. F., Chetta, P., Ho, Y., Cacciatore, S., Gao, X., ... Loda, M. (n.d.).
40 Inhibition of de novo lipogenesis targets androgen receptor signaling in castration-
41 resistant prostate cancer. <https://doi.org/10.1073/pnas.1808834116>
- 42 Zaytseva, Y. Y., Rychahou, P. G., Gulhati, P., Elliott, V. A., Mustain, W. C., O'Connor,
43 K., ... Evers, B. M. (2012). Inhibition of fatty acid synthase attenuates CD44-
44 associated signaling and reduces metastasis in colorectal cancer. *Cancer*
45 *Research*, *72*(6), 1504–1517. <https://doi.org/10.1158/0008-5472.CAN-11-4057>

1 **Figure Legends:**

2 **Figure 1 - FASN inhibition with C75, TCS, and ORL reduces proliferation and**
3 **induces distinct morphologic changes in SCC-9 cells.** The IC₅₀ was calculated using
4 MTT viability assays after treating SCC-9 cells with FASN inhibitors for 24h (A-C) and
5 48h (D-F). C75 (A and D), TCS (B and E) and ORL (C and F) induced a dose-dependent
6 SCC-9 cell death, allowing the determination of IC₅₀. Compared to control cells (G),
7 treatment with the IC₅₀ of C75 (H) induced cytoplasmic retraction (arrow) and
8 predominantly polygonal cells with abundant cytoplasm (arrow). (I) TCS reduced the cell
9 size and increased the number of rounded cells (arrow) and the amount of cytoplasmic
10 granules (arrow). ORL (J) induced a spindle pattern in SCC-9 cells (arrow). Highlighted
11 rounded cells, characteristics of dead cells (arrowheads). Error bar = mean ± SD of at
12 least 3 independent experiments. * p <0.05 ANOVA and Tukey tests. Original 200X
13 magnification.

14

15 **Figure 2 – C75, TCS, and ORL reduces the amount of lipid bodies in SCC-9 cells.**
16 Compared with the control cells (A, F and K), the treatment with C75 (B-D), TCS (F-I)
17 and ORL (L-N) significantly reduce the number of lipid bodies normalized by the cell area
18 in comparison with to control cells. (E,J and O) number of lipid droplets stained by bodipy
19 were normalized by the area of each cell. Error bars = mean ± SD of at least 50 cells in
20 10 microscopic fields (original magnification 400X). # different from all, * p <0.05,
21 **p<0.0001 ANOVA and Tukey tests (ORL) and Kruskal-Wallis (C75 and TCS).

22

23 **Figure 3 – FASN inhibition promotes cell cycle arrest in G0/G1 and increases the**
24 **S phase in SCC-9 cells.** Treatment of SCC-9 cells with IC_{12.5}, IC₂₅, and IC₅₀ for 48 hours
25 of C75 (A) and ORL (C) induces a cell cycle arrest in the G0/G1 phase and reduces the
26 number of cells in the S phase in a dose dependent manner. The treatment of SCC-9
27 cells with TCS (B) reduces S phase and increases the percentage of cells in the G0/G1 at
28 the IC₂₅ and IC₅₀. The compounds equally affected the S-phase progression at the IC_{12.5}
29 (D) but at the IC₂₅ (E) and IC₅₀ (F), TCS and ORL show higher inhibitory effect on S-phase,
30 when compared to the C75. Error bars = mean ± SD of the experimental triplicate. * p
31 <0.05, ANOVA and Tukey tests.

32

1 **Figure 4 – Apoptosis is induced by FASN inhibitors.** The treatment of SCC-9 with
2 the IC₅₀ of C75 (A) and the IC₂₅ and IC₅₀ of TCS (B) reduces the number of viable cells
3 and increases apoptotic cells. All tested concentrations of ORL (C) increase the number
4 of apoptotic cells. TCS was the most effective when compared with C75 and ORL in
5 order to reduce the number of living cells (D) and induce apoptosis (E). Error bar = mean
6 ± SD of the experimental triplicate. * p<0.05, ** p<0.005, *** p <0.0001, ANOVA and
7 Tukey tests.

8

9 **Figure 5 - Cell adhesion is differently modulated by pharmacological FASN**
10 **inhibitors.** The treatment with C75 (A) resulted in a slight increase, not statistically
11 significant, in SCC-9 adhesion at the IC_{12.5} and IC₅₀. TCS significantly reduces SCC-9
12 adhesion to fibronectin and type I collagen in all the analyzed concentrations (B) and
13 ORL (C) did not significantly affect the adhesion. Error bar = mean ± SD of the
14 experimental triplicate. # different from all, * p <0.05, ANOVA and Tukey tests

15

16 **Figure 6 – Migration and invasion phenotypes of SCC-9 cells are affected by the**
17 **inhibition of FASN.** Transwell migration (A-E) and invasion (F-J) assays with SCC-9
18 cells treated with the IC₅₀ of C75, TCS and ORL. C75 (B) and TCS (C) showed a
19 significant inhibition (E) of cell migration when compared to the control (A) and to the
20 ORL (D). The invasiveness of the cells were significantly affected by C75 (G) and TCS
21 (H) but not modulated by ORL (I-J). Original magnification 5X. Error bar = mean ± SD of
22 the experimental triplicate. ANOVA and Tukey tests, # different from all, * p <0.05, ** p
23 <0.005.

24

25 **Supplementary file**

26 **Figure S1: C75, TCS, and ORL reduces the amount of lipid bodies in SCC-9 cells**
27 **in 48h.** The treatment with C75 (A-E), TCS (F-J) and ORL (K-O) significantly reduce the
28 number of lipid bodies normalized by the cell area in comparison with to control cells.
29 (E,J and O) number of lipid droplets satained by bodipy were normalized by the area of
30 each cell. Error bars = mean ± SD of at least 50 cells in 10 microscopic fields (original
31 magnification 400X). # different from all, * p <0.05, **p<0.0001 ANOVA and Tukey tests
32 (ORL) and Kruskall-Wallis (C75 and TCS).

33 **Figure S2: Cell cycle gate information.**

34 **Figure S3: Apoptosis gate information.**