



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Heat stimulation as a modulatory tool for the histaminergic and non-histaminergic itch

Riccio, Daniele

DOI (link to publication from Publisher):  
[10.54337/aau510569253](https://doi.org/10.54337/aau510569253)

Creative Commons License  
Unspecified

Publication date:  
2022

Document Version  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):  
Riccio, D. (2022). *Heat stimulation as a modulatory tool for the histaminergic and non-histaminergic itch*. Aalborg Universitetsforlag. Aalborg Universitet. Det Sundhedsvidenskabelige Fakultet. Ph.D.-Serien  
<https://doi.org/10.54337/aau510569253>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

### Take down policy

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.



**HEAT STIMULATION AS A MODULATORY TOOL  
FOR THE HISTAMINERGIC AND  
NON-HISTAMINERGIC ITCH**

**BY  
DANIELE RICCIO**

DISSERTATION SUBMITTED 2022



**AALBORG UNIVERSITY**  
DENMARK



**HEAT STIMULATION AS A  
MODULATORY TOOL FOR THE  
HISTAMINERGIC AND NON-  
HISTAMINERGIC ITCH**

Dissertation submitted: October 2022

PhD supervisor: Prof. Lars Arendt-Nielsen, Dr. Med. Sci., PhD,  
Aalborg University

PhD committee: Associate Professor Eugen Romulus Lontis (chair)  
Aalborg University, Denmark

Associate Professor Romina Nassini  
University of Florence, Italy

Associate Professor Sara Invitto  
General and Experimental Psychology, Italy

PhD Series: Faculty of Medicine, Aalborg University

Department: Department of Health Science and Technology

ISSN (online): 2246-1302

ISBN (online): 978-87-7573-806-9

Published by:  
Aalborg University Press  
Krogstræde 3  
DK – 9220 Aalborg Ø  
Phone: +45 99407140  
aauf@forlag.aau.dk  
forlag.aau.dk

© Copyright: Daniele Riccio

Printed in Denmark by Stibo Complete, 2022



## CV

Daniele obtained his B.Sc and M.Sc in pharmaceutical biotechnology from the University of Milan (Italy).

Subsequently he worked as research assistant in Prof. Abbracchio's lab in Milan with focus on trigeminal pain and the role of glial cells in this sensory modality.

In 2016 he enrolled at Aalborg University as a PhD fellow at the Center for Neuroplasticity and Pain under the supervision of Lars Arendt-Nielsen.

During his research experiences he develop a solid scientific knowledge and expertise in the biological sciences, from a completely in-vitro approach to human preclinical studies.

He is interested in neurosciences in particular the mechanism underlying pain and itch perception as well as how these two relate with thermosensation.

He focused his research on the evaluation of the effects that various heat stimuli have on different itch modalities.

Daniele had the opportunity to present his work at different international conferences and school both as poster and oral presentation.

# ENGLISH SUMMARY

Itch is a sensation characterized by the innate desire to scratch. Although acute itch is generally tolerable, chronic itch (itch occurring for more than 6 weeks) can severely affect the patients' quality of life and have a prevalence up to 25%. In the last decades, many therapies and treatments have been developed, but to date, their efficacies are scarce and may present with side effects. If efficient non-pharmacological treatment options could be developed this would have a major clinical potential.

Scratching itchy skin is ultimately a form of self-inflicted mechanical pain that can help in relieving itch to some extent. Therefore, many studies have focused on the interaction between itch and pain.

Regarding heat stimulations, it has been shown that long noxious heat stimuli can inhibit itch, but on the other hand, such stimuli may cause hyperalgesia and skin damage.

Many clinical observations indicate that patients can experience increased itch in warm environments, suggesting some relationships between heat and itch sensations.

A 70 years old study showed that a paradoxical itch sensation could be achieved when anaesthetised skin was stimulated with what was defined as a "burning" sensation.

Taken together these independent observations, seem to indicate that thermal perception may have a role in modulating different aspects of itch sensation, and therefore it is of interest to understand better the mechanism underlining these modulatory effects.

Within this context, this PhD project aimed at investing these aspects: 1) the effects of very short heat stimuli (ranging from innocuous to noxious) on various itch modalities; 2) how continuous skin warming affects itch intensity in human surrogate itch models; 3) investigate and possibly confirm with a more modern and systematic approach the observed dysesthesia of paradoxical itch sensation when a "burning" stimulus is applied on top of anaesthetised skin.

The first study showed that homotopical transient noxious heat stimuli were able to inhibit histaminergic itch and, to a lesser extent, non-histaminergic itch induced by cowhage. When the stimuli were applied heterotopically there was no significant inhibition suggesting that, at least for very transient stimuli, the observed inhibitory effect relies on peripheral mechanisms.

The second study showed that mild skin heating selectively aggravates serotonergic and histaminergic itch but not cowhage-induced itch. These



results partially agree with what is observed in animals (increased serotonergic itch intensity) and parts are in direct contrast with them (unchanged histaminergic itch intensity).

The third study failed to confirm what was previously in a 70 years old observation. Applying a heat ramp (ranging from warm to noxious heat) on anaesthetised skin did not evoke a paradoxical itch sensation at any of the time points analysed.

In conclusion, this dissertation unveils a potential role of thermal stimulation to modulate itch. Further studies are needed to investigate this in a clinical context.

# DANSK RESUME

Kløe er en følelse, som karakteriseres ved et instinktivt ønske om at krads sig. Selvom akut kløe generelt er udholdelig, så kan kronisk kløe (dvs. kløe som varer ved i mere end seks uger) påvirke patienternes livskvalitet alvorligt, og kronisk kløe har en prævalens på op til 25%. I de seneste årtier er der blevet søgt udviklet mange terapiformer og behandlinger til lindring af kløe, men virkningen af disse har indtil nu vist sig at være mangelfuld og forbundet med bivirkninger. Hvis der kunne udvikles ikke-farmakologiske behandlingsmuligheder til kløe, ville disse have et betydeligt klinisk potentiale. Når man krads i kløende hud, påfører man ultimativt sig selv en slags selvforskyldt mekanisk smerte, som til en vis grad kan hjælpe med at lindre kløen. Derfor har mange studier fokuseret på interaktionen mellem kløe og smerte. For varmestimulationer har det vist sig, at langvarige smertefulde varmestimulationer kan hæmme smerte, men samtidig kan sådanne stimuli også forårsage hyperalgesi og hudskader.

Mange kliniske observationer peger på, at patienter kan opleve forøget kløe i varme miljøer, hvilket indikerer en forbindelse mellem varme og følelse af kløe.

Et 70 år gammelt studie viste, at der kunne opnås en paradoksal følelse af kløe, når bedøvet hud blev stimuleret med hvad, der blev defineret som en "brændende" følelse.

Overordnet synes disse uafhængige observationer at indikere, at termisk perception kan spille en rolle i moduleringen af forskellige aspekter af kløe, og derfor er det interessant at opnå en bedre forståelse af de mekanismer, der ligger til grund for disse modulatoriske effekter.

Med baggrund heri var formålet med dette ph.d.-projekt at undersøge følgende aspekter: 1) Effekterne af meget korte varmestimuli (fra ikke-smertefulde til smertefulde) på forskellige kløe-modaliteter; 2) Hvordan kontinuerlig opvarmning af huden påvirker kløeintensiteten i humane surrugatmodeller for kløe; 3) Ved hjælp af en mere moderne og systematisk tilgang at undersøge og eventuelt bekræfte den observerede dysæstesi med paradoksal kløe, når der påføres en "brændende" stimulation på bedøvet hud. Det første studie viste, at homotopiske, kortvarige smertefulde varmestimulationer var i stand til at hæmme histaminerg kløe og i mindre grad ikke-histaminerg kløe fremkaldt ved hjælp af nåle fra planten mucuna pruriens. Når stimulationerne blev påført heterotopisk, var der ingen signifikant hæmning, hvilket antyder, at den observerede hæmmende effekt - i hvert fald for meget kortvarige stimuli - er afhængig af perifere mekanismer.

Det andet studie viste, at let opvarmning af huden selektivt forværrer serotonerg og histaminerg kløe men ikke kløe fremkaldt ved hjælp af nåle fra mucuna pruriens. Disse resultater er til dels i overensstemmelse med observationer hos dyr (forøget serotonerg kløeintensitet) og andre dele er i direkte kontrast med disse (uændret histaminerg kløeintensitet).

Det tredje studie kunne ikke bekræfte det, som tidligere var en 70 år gammel observation. Påføring af en varmerampe (fra varm til smertefuld varme) på bedøvet hud fremkaldte ikke en paradoksal kløefølelse på nogen af de analyserede tidspunkter.

Denne afhandling præsenterer dermed varmestimulationers potentielle rolle til modulation af kløe. Yderligere studier er nødvendige for at undersøge denne rolle i klinisk kontekst.

# ACKNOWLEDGEMENTS

The first person to whom I would like to express my most sincere gratitude is my supervisor Lars Arendt-Nielsen, for allowing me to work in such an amazing environment. His vast scientific knowledge together with patience and advice vastly helped me in getting this work done.

Thanks to Prof. Thomas Graven Nielsen, who let me be part of CNAP and be surrounded by a high-level scientific environment.

Thanks to Hjalte, for his scientific support, when I had to start projects on new and unexplored terrain for me.

Thanks to all the CNAP family that went along with me during these years and gave me tips and tricks on how to survive the PhD life. In particular, I would like to thank Dennis, Line, Megan and Sinead for always being ready to share a laugh, gossip, some tips or a beer.

Thanks to Giulia, even though our lives took different paths, we helped each other at the start of this new adventure.

A very special thanks to my first mentor Marco. Your teachings are still rooted in me and thanks to them I developed curiosity and passion for science.

Thanks to Elisa and Veronica always there when I needed to laugh or just a trip down memory lane.

Thanks to Maurizio, your constant support in the last 10-15 years is priceless. Thanks for the patience and advice, thanks for the tips on series and music, thanks for the calls, and thanks for being there when I need you. I could write 3 books focused on thanking you, I hope I will be able to give you back at least part of what you are giving me.

Thanks to Masiello and Heriwa for the trips, dinners, gossip, concerts, and experiences. It is always a pleasure to be around you.

Thanks to Federica, you massively helped me in getting things “back” in order in my life.

Thanks to Iga, even though our lives crossed for a small time, your support helped me complete this document.

Thanks to Silvia and Simone for (among other things) exploring together the restaurants in Aalborg and giving me a way out when the carnival is in town.

Thanks to Rocco and Thea for the parties, the food, the weekends, and the fun, we had together. In particular Rocco for being there since my second and your very first day in Denmark.

Thanks to Alessandro and Giulia for adopting me. Ale thanks to you I can add “foosball proficiency” to my CV (clearly a highly required skill). Giulia we will plan soon our next HP marathon.

Thanks to Marco for always having your office door open for me. For teaching me how to make pasta, for the football struggles, for the support and the stress-free attitude you are trying to teach me.

Thanks to Davide, Federica, Timothy, and Elektra for the amazing food, the various tastings, the support, and the roof during the most difficult time of my life.

Thanks to all the Italians in Aalborg, we are way too many (Carolina, Enrico Davide, Anna, Samuel, Alessia, Sticchi, Carlo... etc), it is like I never left Italy.

Last but not for importance thanks to my family. Sarebbe scontato dire che non potrei essere qui se non fosse per voi, ma rimane un’assoluta verità. Non avrebbe senso elencare tutto quello per cui vi sono grato, la lista sarebbe infinita. Grazie per esserci sempre, spero ci sarete anche nel futuro.

*"All right," said the computer, and settled into silence again. The two men fidgeted. The tension was unbearable.*

*"You're really not going to like it," observed Deep Thought.*

*"Tell us!"*

*"All right," said Deep Thought. "The Answer to the Great Question..."*

*"Yes..!"*

*"Of Life, the Universe and Everything..." said Deep Thought.*

*"Yes...!"*

*"Is..." said Deep Thought, and paused.*

*"Yes...!"*

*"Is..."*

*"Yes...!!!...?"*

*"Forty-two," said Deep Thought, with infinite majesty and calm."*

**Douglas Adams**

# Table of Contents

Chapter 1. Introduction.....	13
1.1. Itch at a glance .....	13
1.2. Similarities and differences between itch and pain.....	14
1.2.1. Itch coding hypotheses.....	14
1.3. Itch human models.....	15
1.4. Heat perception.....	18
1.4.1. TRPs in heat and itch perception.....	18
1.5. Topical cutaneous anaesthesia .....	19
1.6. Aims of the PhD project .....	20
1.6.1. Papers and dissertation overview .....	21
Chapter 2. The effects of noxious heat on itch .....	23
2.1 Noxious heat reverse histaminergic itch.....	23
2.2 Noxious heat partially inhibits non-histaminergic itch .....	24
2.3 Heterotopic noxious heat does not affect itch sensation.....	24
Chapter 3. The effects of innocuous heat on itch .....	27
3.1 Selective warmth hyperknesis for histaminergic and serotonergic itch.....	27
3.2 Cowhage-induced itch is not affected by innocuous skin heating.....	28
3.3 Enhanced superficial blood perfusion due to innocuous skin heating .....	28
Chapter 4. The effects of escalating heat on top of anaesthetized skin.....	31
4.1 Pain intensity.....	32
4.2 Itch intensity.....	32
Chapter 5. General discussion.....	33
5.1 Limitations .....	35
Chapter 6. Conclusions.....	37
Literature list .....	39





# CHAPTER 1. INTRODUCTION

## 1.1. ITCH AT A GLANCE

Itch is considered an uncomfortable sensation that causes the desire to scratch. It can be caused by different plants, animals or clinical conditions giving rise to acute or chronic itch<sup>1-4</sup>.

Acute itch can occur in many instances with moderate to no effects on day-to-day life whereas chronic itch can be very burdensome.

Itch is primarily a skin condition, but the aetiology could originate from central neurobiological mechanisms or as secondary symptoms too eg. vascular conditions, kidney conditions, pharmacological treatments, etc...<sup>5-7</sup>.

The desire to scratch is a primordial instinct and it is basically the urge to use mechanical pain to abolish the itch sensation. This consideration led the researchers to focus extensively on the interaction and the relationship between itch and pain transmission. When the pruritus sensation lasts for more than 6 weeks it is clinically referred to as chronic itch. To understand better the extent of such a condition, it has to be noticed that one every 5 people will experience chronic itch in their life<sup>8,9</sup>. In the most recent study on the global burden of disease (2017), "Skin and subcutaneous diseases" ranked ninth when measuring the "years lived with disability" parameter<sup>10</sup>. And among these diseases, the first four positions were held by atopic dermatitis, psoriasis, urticaria and scabies all of which are heavily characterized by the presence of itch<sup>11</sup>. It must be noticed that these results were before to the recent global pandemic that most likely would have skewed the results. Furthermore, chronic itch has been associated with a decrease in Health-Related Quality of Life (HRQoL) and can involve psychiatric comorbidities and sleep disorders<sup>9,12-16</sup>.

Since 1990 there has been an increase in itch prevalence globally, from 812 to 932 in 2017 (out of 100000 people)<sup>8</sup>. Highlighting itch as a growing problem in our society. Interestingly, data indicates also being female and older as risk factors linked to itch<sup>8,9,17-19</sup>.

The impact that itch diseases have on society can be also measured in health care expenditures. Only in the United States, itch has been associated with a yearly cost of more than \$90 billion for the all population<sup>8,20</sup>. Altogether, these data highlight a clinical challenge that can affect a very large part of the population. Historically itch perception has been divided into a histaminergic and non-histaminergic itch, based on which receptors and pathways are involved.

## 1.2. SIMILARITIES AND DIFFERENCES BETWEEN ITCH AND PAIN

Traditionally, it is known that painful counter stimuli can inhibit itch. The same definition of itch recalls the desire to scratch which is substantially self-inflicted mechanical pain, that aims at removing the source of itch. Whereas the natural reflex for acute pain is a withdrawal response, intended to escape from the source that causes this sensation<sup>21,22</sup>.

One of the main differences between these two sensory modalities is that pain can occur in various tissues (cutaneous, musculoskeletal, etc) whereas itch is present solely in the cutaneous tissue.

Interestingly, the known inhibitory effect of  $\mu$ -opioids on pain does not affect itch sensation whereas  $\mu$ -antagonists can inhibit itch and have no effects on pain sensation<sup>23,24</sup>. Whereas,  $\kappa$ -receptor's agonists showed to have analgesic and antipruritic effects, leading to the development of new potential drugs<sup>25-27</sup>. This line of evidence suggests a sort of specificity in the itch and pain transmission, in opposition to what has been widely demonstrated in the literature, where these two distinct sensory modalities have many receptors<sup>28</sup> and pathways<sup>29-31</sup> in common. Therefore, many theories have been postulated to evaluate and describe the relationship between these two sensory modalities.

### 1.2.1. ITCH CODING HYPOTHESES

Understanding how itch and pain are transmitted is important to identify new targets for a selective and effective therapy able to help the many patients suffering from chronic itch and chronic pain conditions.

Historically, itch has been considered a submodality of pain, giving rise to the so-called intensity theory according to which, some nociceptors can be activated by both algogens and pruritogens (hence the name polymodal fibres) and are connected to the same neurons' population at the central level. Depending on the intensity at which the neurons fire these can signal itch (weak activation) or pain (strong activation)<sup>32-34</sup>.

The specificity theory (also referred to as labelled line theory) assumes that itch and pain are transmitted by fibres creating a distinct and selective pathway up to the central nervous system<sup>32-34</sup>. Many studies have identified various markers expressed in neurons attributed to specifically signal itch sensation<sup>35-41</sup>.

The selectivity theory expands what is already said by the specificity theory by adding the presence of spinal interneurons activated by painful stimuli, that exerts an inhibitory effect on itch sensation<sup>42</sup>.

The spatial contrast theory postulates that itch and pain sensation share the same set of neurons and nociceptors and the discrimination between these two sensory modalities is given by the number of nociceptors recruited in the epidermis. When there is a focal stimulation few nerve endings are activated, and this translate into an itchy sensation. When the area stimulated is larger, more nociceptors are excited, and this gives rise to a pain sensation<sup>43,44</sup>.

Finally, more recently a leaky gate theory has been presented<sup>45</sup>. The latter has some similarities with the selectivity theory and shows a set of spinal neurons that are activated by both nociceptive and pruriceptive stimuli and can evoke simultaneously itch and pain sensations. Upon strong nociceptive stimulus, endogenous opioids inhibit what would be an overwhelming pain sensation.

Many studies in the last years have explored several aspects of itch and pain signalling and interaction, but to date, an unambiguous theory has not been presented. Further studies using both human and animal models are needed to understand how the brain and the nervous system can discriminate between itch and pain.

### **1.3. ITCH HUMAN MODELS**

It is of utmost importance to study and understand itch mechanisms to develop new devices and therapies able to help to improve the patients' quality of life. To do so, many models have been developed both for animal and human experiments. Animal models could potentially provide a lot of information but unfortunately, they are not reliable when it comes to establishing the intensity of itch experienced, and they rely on behavioural responses such as scratching bouts and licking (itch and pain behaviour respectively)<sup>46,47</sup>. However, animal models remain the first choice when it comes to investigating new molecular mechanisms both at the central and peripheral levels. Several animal itch models have been developed and used in many studies, these will not be treated in detail here and further information can be found in literature<sup>3,34,46,48</sup>.

A good way to study and investigate itch mechanisms is to use human surrogate models of itch. The current itch models, rely on the intracutaneous application of various substances, called pruritogens, producing a mild-to-moderate itch sensation. Except for cowhage-induced itch (where itch disappear as soon as the spicules are removed; more details below),not counter stimulated itch sensations will slowly fade when the substances in the

skin have been cleared physiologically. Although not optimal, these models helped in progressing toward a better understanding of the basic mechanisms of itch.

The most used pruritogen is histamine, able to evoke substantial itch, lasting between 5 to 20 minutes, coupled with a well-documented wheal and neurogenic inflammation (also referred to as neurogenic flare)<sup>49–52</sup>. The wheal appears as a small oedema (approximately 5 mm in diameter) in the exact provocation site resembling to the wheal present after a mosquitoes bite<sup>49,52–56</sup>. The neurogenic flare is characterised by a short increase in superficial blood perfusion due to the activation of peptidergic nerve fibers<sup>57–59</sup>. This latter cutaneous response, can be measured by full-field laser perfusion imaging (FLPI), a modern technique that can quantify the superficial blood perfusion by analysing the haemoglobin content<sup>60–65</sup>.

Histamine activates histamine 1 receptor (H1) peripherally, inducing an increase in intracellular  $\text{Ca}^{2+}$  influx through intracellular cooperation with phospholipase  $\text{C}\beta$ <sup>66</sup> or phospholipase  $\text{A}2$ <sup>67</sup>. Interestingly, it has been shown recently that selective and systematic inhibition of histamine 4 receptor (H4) lead to decreased itch intensity in rodents, highlighting a potential new role for H4<sup>68</sup>. Nonetheless, in clinic most chronic itch conditions are unaffected by the treatment with antihistamines, demonstrating how chronic itch does not rely solely on histaminergic pathways<sup>1,69–72</sup>.

Histamine has to be introduced into the skin, to achieve an itch sensation. This can be done using a skin prick test, iontophoresis, intradermal injection and inactivated cowhage spicules soaked in a histamine solution<sup>50,52,73–76</sup>.

Another itch human model is achieved by the use of cowhage spicules already characterized in 1955<sup>77,78</sup>. These are collected from the pod of *Mucuna pruriens* and applied onto the skin by gently rubbing with the fingertip until a light pricking sensation is felt on the treated skin. In recent years, cowhage received more attention as a pruritogen due to the unique sensory qualities that evokes<sup>43,79</sup>. The itch sensation evoked by cowhage is stronger compared to histamine, furthermore is coupled with nociceptive sensory qualities like pricking/stinging<sup>43,80</sup>. Cowhage-evoked itch lasts up to 5-15 minutes and rapidly disappears when the spicules are carefully removed<sup>73,74</sup>. The molecule believed to be responsible for its action is mucunain, an enzyme present on the spicules<sup>78</sup>. The mechanism that leads to the itch sensation is still not fully clear, two pathways have been suggested so far: the activation of proteinase-activated receptor 2 and 4 (PAR2/4) and it has been suggested the contribution of mas-related G protein-coupled receptor X1 and 2 (MRGPRX1/2)<sup>35,81</sup>. The activation of the MRGPRX2 also lead to mast cell degranulation, a mechanism that characterize histamine-dependent itch

pathways, suggesting that cowhage-induced itch pathways could also recruit histaminergic itch pathways<sup>82-84</sup>. Nonetheless, it has been shown that cowhage do not evoke any wheal typical of mast cells degranulation<sup>50</sup>. More molecular studies are needed to fully elucidate cowhage-evoked itch pathway. It is worth mentioning that all the receptors activated by mucunain are expressed and activated on superficial epidermal C-fibres<sup>35,81</sup>.

A third model used for inducing itch in healthy subjects relies on the use of serotonin. Serotonin can be introduced to the skin by iontophoresis and induces a mild itch sensation<sup>85,86</sup>. A study explored itch induction by intradermal injection achieving slightly weaker results<sup>87</sup>, overall iontophoresis remain the preferred induction modality. Serotonin in the epidermis activates both mechanosensitive, also called polymodal C-fibres (PmC-fibres), and mechano-insensitive C-fibres (Cmi-fibres), the latter a specific group that can be activated by histamine as well<sup>88</sup>. Notably, pre-treatment with antihistamines does not affect itch intensity following iontophoresis with serotonin<sup>86</sup>, indicating that serotonergic itch does not rely on histamine-releasing mechanisms to develop itch. However, treatment with tropisetron, a selective serotonin type 3 receptor (5-HT<sub>3</sub>) antagonist, showed a marked serotonergic itch inhibition, suggesting that 5-HT<sub>3</sub> may be the responsible for serotonergic itch sensation<sup>89</sup>.

Other itch models rely on the application of various pruritogens. Bovine adrenal medulla 8-22 (BAM8-22) can be applied to the skin to evoke an itch sensation similar to the one achieved with cowhage spicules<sup>52,90-92</sup>. BAM8-22 is a direct agonist of MRGPRX1 and can be introduced by the use of intradermal injection or inactivated cowhage spicules soaked in a solution of this compound<sup>90</sup>.

The use of  $\beta$ -alanine as an agonist of MRGPRD has been used as itch model<sup>93</sup>. Qualitatively, it evokes a less intense itch compared to histamine- or cowhage-evoked itch. Furthermore,  $\beta$ -alanine is not able to provoke itch sensitization<sup>94</sup>.

Transient Receptor Potential Ankyrin 1 (TRPA1) activation with trans-cinnamaldehyde lead to mild cutaneous pain as well as itch in some participants, but further investigation is needed to make it a reliable itch model.<sup>95,96</sup> These latter three models were not used in the studies on which the dissertation is based, more detailed information can be found in the literature.

Thanks to these human models it is possible to analyse and investigate how itch interacts with other sensory modalities, as well as to develop new therapies.

For this thesis, the attention will be set on histamine-, cowhage- and serotonin-induced itch models.

## 1.4. HEAT PERCEPTION

Numerous studies have shown the effect of various thermal stimuli on top of the skin. These studies investigated the temperature at which generally warm detection and pain perception occurs.

Normal skin temperature is set around 32°C<sup>95,97-100</sup>, and the Heat Pain Threshold (HPT) is ≈45°C<sup>101</sup>. These temperatures can vary based on different body regions stimulated, gender and method of stimulation<sup>102-105</sup>.

Heat stimuli can be applied to the skin using various methods: radiation (infrared, microwave, laser, etc)<sup>74,102,106</sup>, convection (through a fluid, i.e. air and liquid)<sup>107,108</sup> and conduction (by putting the skin in contact with a hot object)<sup>73,97,109-111</sup> or a combination of these. This dissertation will focus mainly on conductive heat transfer (using a controlled Peltier element; Study I and III) and radiant heat transfer (by the use of infrared radiation; Study II).

With regards to the interaction between itch and thermal stimulation, it has been shown how noxious heat and cold stimulation can inhibit itch sensation<sup>73,97,110,112</sup>. Specifically, these studies explored the effect of 30-seconds long stimulation at 49°C, showing that this was able to inhibit itch when applied homotopically and/or heterotopically to a previous itch provocation<sup>97,109,110</sup>. These results suggest that the inhibitory mechanism relied on spinal or supraspinal pathways, because the effect was observed also when the stimuli were applied heterotopically i.e. in a cutaneous area differently innervated compared to the itch provocation area<sup>97</sup>. Nonetheless, it is still unclear how this known mechanism can be exploited for the development of new therapies.

### 1.4.1. TRPs IN HEAT AND ITCH PERCEPTION

Heat perception resides mainly on a specific class of calcium-permeable ion channel receptors, of the TRPs family called Transient Receptor Potential Vanilloid (TRPV).

Two decades ago TRPV1 was indicated as the receptor able to sense noxious heat, and to respond to temperatures above 48°C; since then the research around these ion channels and their relation with thermal sensation has been widely explored<sup>113,114</sup>. Further studies showed that this receptor is activated by temperatures above 42°C<sup>115,116</sup>. Its central role as a primary noxious heat detector was confirmed also by genetic ablation experiments showing inhibition in heat pain sensation that was not completely abolished, suggesting

the involvement of other receptors as well<sup>115,117,118</sup>. TRPV1 is markedly expressed by small-diameter sensory neurons<sup>113,117</sup> as well as in keratinocytes<sup>119,120</sup>.

Although direct activation of this receptor is known to cause pain and not itch<sup>121</sup>, following itch induction with histamine, sensory neurons can intracellularly activate TRPV1<sup>122</sup>. Similarly, it has been shown how TRPV1 can help mediate itch when induced by PAR2 activation or IL-31<sup>122,123</sup>.

TRPV2 activation starts at a higher temperature (>52°C), to date no studies are showing a connection between itch sensation and this receptor<sup>115,124</sup>.

Warm innocuous stimuli activate TRPV3, specifically above 33°C<sup>115,125</sup>. This receptor is capsaicin-insensitive<sup>125</sup> and its expression is mainly in keratinocytes<sup>126,127</sup> rather than neuronal cells. Interestingly, a specific mutation of this receptor, characterized by a gain of function has been associated with pruritus and dermatitis<sup>128,129</sup>. Moreover, itch elicited by PAR2 activation has been linked with TRPV3<sup>130,131</sup>.

Also, TRPV4 is activated by innocuous stimuli in the range of 27-42°C, the higher the temperature goes the more active this receptor is, and with repeated or sustained stimulation >42°C desensitization is achieved<sup>115,132</sup>. Furthermore, recently it has been shown its involvement both in serotonergic and histaminergic itch<sup>133-135</sup>.

Another TRP channel receptor is somehow involved in the heat perception, TRP Ankyrin 1 (TRPA1). TRPA1 is widely known for his role as chemosensor and in inflammation, but some studies also showed its involvement in thermal and mechanical sensation<sup>136-139</sup>. Although it has been recognised as a cold stimuli detector<sup>115,140,141</sup>, it has been shown how its genetic deletion or pharmacological blockade is necessary in order to achieve a complete loss to noxious heat sensitivity<sup>142</sup>. One could think that these two sets of evidence contradict each other, nonetheless it has been shown that TRPA1 could in theory be able to exert this double sensitivity<sup>143</sup> and moreover some evidence in vitro showed activation with cold and heat stimuli<sup>144</sup>. Further investigations are needed to fully verify TRPA1 thermal sensitivity.

## 1.5. TOPICAL CUTANEOUS ANAESTHESIA

Local anaesthetic exerts their anaesthetic effect by blocking the sodium channels (Nav) and consequently blocking the majority of sensory fibres from firing any signals<sup>145,146</sup>. They have been extensively used in clinic for dermatologic procedures<sup>147,148</sup>, in certain case they can also be used as treatments for particular neuropathic pain conditions in the form of plasters<sup>149,150</sup>.

A recent study investigating the RNA expression profile of all the cutaneous sensory fibres showed that Nav channels are expressed by the so-called peptidergic (PEP) and non-peptidergic (NP) sensory fibres' endings in the epidermis<sup>151</sup>. According to this study, these fibres are involved in the pain and itch perception<sup>151</sup>. Therefore, topical Nav blockers are a potent tool to modulate the sensorial cutaneous inputs.

The use of topical anaesthesia, besides the well-known analgesic effect, has been reported to partly alleviate itch, usually in the form of anaesthetic creams<sup>25,152,153</sup>. These creams supposedly achieve anaesthesia in a large section of the skin, meaning that they can block fibres in the upper as well as the deeper cutaneous layers.

This distinction is important because it has been shown how the different fibres PEPs and NPs, involved in itch and pain transduction have their terminals ending in different cutaneous layers. Specifically, NPs terminals are in the upmost layer of the skin (superficial epidermis), whereas, PEPs endings are closer to the dermal-epidermal junction (deep epidermis)<sup>151,154,155</sup>.

Interestingly, a 70 years old study investigated the effect of heat stimuli following topical anaesthetization due to subcutaneous injection of procaine (a local anaesthetic known to block the Nav channels)<sup>156</sup>. In this study, a stimulation intended to achieve a “decided sensation of burning” was applied on top of the anaesthetized skin and led to the development of a paradoxical itch sensation when anaesthetization started to subside<sup>156</sup>. This observation highlights a potential interaction between heat perception and itch pathways that needs to be further investigated to be fully understood.

## **1.6. AIMS OF THE PHD PROJECT**

Given these premises, this PhD project aimed to explore and investigate how heat perception interacts with itch sensation and development. Furthermore, to understand if it can be used as a tool to modulate itch. For these purposes there were three main focuses:

1. To evaluate the effects of short noxious heat stimuli, homotopically and heterotopically to itch induction.
2. To investigate the pruritic effects of continuous warm innocuous stimulation pre- and post-itch induction in various human itch models.
3. To explore the short- and long-term effects of escalating heat stimulation on top of anaesthetised skin.
- 4.



### 1.6.1. PAPERS AND DISSERTATION OVERVIEW

To this end, the thesis is based on two published papers, and one submitted to peer-reviewed journals, each paper exploring one of the above bullet points.

- **Study I<sup>73</sup>**: **Riccio D.**, Andersen HH. & Arendt-Nielsen L. Antipruritic effects of transient heat stimulation on histaminergic and non-histaminergic itch. *Br. J. Dermatol.* **181**, 786–795 2019.
- **Study II<sup>74</sup>**: **Riccio D.**, Andersen HH. & Arendt-Nielsen L. Mild skin heating evokes warmth hyperknesis selectively for histaminergic and serotonergic itch in humans. *Acta Derm. Venereol.* **102**, adv00649 2022.
- **Study III<sup>111</sup>**: **Riccio D.**, Lo Vecchio S., & Arendt-Nielsen L., The effect of escalating heat stimulation on top of anaesthetized skin. Itch 2022. (SUBMITTED).

All the studies were conducted on healthy human participants with the approval of the regional ethical committee (N-20180035). The full methodologies will not be discussed in this dissertation, a detailed version of them is available in the papers.

The figure 1 (Fig. 1) shows an overview of how the different studies analyse different aspects of the interaction between heat perception and itch.




Figure 1. Schematic overview of the dissertation studies with the temperature ranges, models applied and outcomes measured: Study I and II investigated the effects of various heat stimuli on well-established itch models (histamine, cowhage and serotonin). Study III investigated the interaction between heat perception and anaesthesia (lidocaine).

# CHAPTER 2. THE EFFECTS OF NOXIOUS HEAT ON ITCH

The content of this chapter is related to Study I<sup>73</sup>.

In chapter 1.2 it was briefly discussed how counter stimuli can inhibit, partially or completely the itch sensation<sup>5,157,158</sup>. Particularly, it has been shown that long noxious heat stimulation can reverse itch in healthy human participants<sup>97,109</sup>. These long stimulations usually induce also primary and secondary hyperalgesia on the skin area stimulated and therefore they are not considered a suitable treatment option<sup>159,160</sup>. Nonetheless, it is important to investigate the mechanisms behind the observed effect as well as try a different approach to exploit this known outcome.

In Study I, the effects of very transient (5 seconds) heat stimuli in healthy participants after itch induction with histamine or cowhage has been analysed. The stimuli were applied, exactly 2 minutes after the itch provocation, at the same site as the itch induction (experiment 1, homotopically; Study I Fig. 1A) or in a different site (experiment 2, heterotopically; Study I Fig. 1B) either next to the provocation site (intra-segmentally) or on the other side of the forearm (extra-segmentally; Study I Fig. 1C).

Itch intensity was measured starting from the itch induction up to 10 minutes after. Notably, the heat stimuli applied in Study I did not evoke a long-lasting pain sensation in any of the conditions, mainly due to their transient nature.

## 2.1 NOXIOUS HEAT REVERSE HISTAMINERGIC ITCH

Fifteen years ago Yosipovitch et al., started investigating the effects of noxious heat stimuli on histamine-evoked itch both in healthy participants<sup>97,109</sup> and atopic dermatitis patients<sup>110</sup>. As mentioned before, these focused on long noxious stimuli (30 seconds stimulations), applied homotopically or heterotopically. Little is known about the effects of transient and very short heat stimuli.

The results in Study I show that a very transient homotopical noxious heat stimulation was able to greatly inhibit histaminergic itch sensation. Specifically, a significant reduction in itch sensation was achieved with stimulations of 45°C (-41.27%,  $p < 0.01$ ) and 50°C (-76.66%,  $p < 0.0001$ ; Study I Fig. 2A)<sup>73</sup>.

Interestingly, as explained in section 1.4, these stimuli, able to inhibit itch, directly activate TRPV1<sup>115,161</sup>, known to hold an important role in signalling histaminergic itch<sup>162–166</sup>. Following histamine receptor activation, intracellular pathways lead to the co-activation of TRPV1<sup>91,167</sup>.

It can be speculated that, due to the external activation of TRPV1 with transient heat counter stimulation, histaminergic intracellular pathways are not able to co-opt it and therefore an inhibition in itch sensation is observed. This latter hypothesis is yet to be confirmed, with further investigations and possibly with pharmacological tools.

## **2.2 NOXIOUS HEAT PARTIALLY INHIBITS NON-HISTAMINERGIC ITCH**

With regards to cowhage-induced itch, Study I shows that a homotopic transient noxious stimulation (50°C) was able to achieve a partial itch inhibition (-43.6%,  $p=0.011$ ; Study I Fig. 2B).

As explained in section 1.3, cowhage spicules lead to the activation of PAR2/4 and/or MRGPX1/2<sup>81</sup>. Rather than recruiting TRPV1 as for histaminergic itch, these pathways are known to recruit TRPA1<sup>168,169</sup>, which is substantially less activated by heat noxious stimuli<sup>142,170,171</sup>. As mentioned in the section 1.4, TRPA1 historically has been considered as a cold detector but some evidence showed a possible activation by heat stimuli, that needs further data to be confirmed<sup>144</sup>.

Since cowhage-induced itch was inhibited to a lower extent (compared to histaminergic itch; Study I Fig. 2B and Fig. 3) and there was no significant inhibition by stimuli lower than 50°C, one can assume that the different inhibition pattern derives from the different TRP receptors involved, specifically cowhage-induced itch was less inhibited due to the less prominent response to heat stimuli by TRPA1<sup>73</sup>.

## **2.3 HETEROTOPIC NOXIOUS HEAT DOES NOT AFFECT ITCH SENSATION**

The second experiment conducted in Study I, explored the effect of heterotopic counter stimuli to itch induction.

For both histaminergic and non-histaminergic itch, heterotopic stimuli were not able to inhibit itch sensation (Study I Fig.5).

The effect known as “diffuse noxious inhibitory control” was for the first time described in rodents in 1979<sup>172,173</sup>. This psychophysical mechanism refers to the reduced pain perception following a painful stimulus applied to a

heterotopic area. Later on, the human behavioural correlate was investigated and named conditioned pain modulation (CPM)<sup>174,175</sup>. Studies have shown that CPM signal starts at central level in the brainstem<sup>175–178</sup>. Thus, heterotopic counter-stimuli to the itch induction site were explored to rule out the presence of a similar mechanism called conditioned itch modulation<sup>179,180</sup>.

As showed in Study I, both histaminergic and non-histaminergic itch were inhibited only by homotopical counter-stimuli (Study I Fig. 5). This is in sharp contrast with pieces of evidence showing pain-evoked itch inhibition being mainly a central mechanism because itch inhibition by intra- and extrasegmental counterstimuli would have meant the central nervous system involvement<sup>42,181</sup>.

Furthermore, some previous studies presented opposite results to what showed in Study I. Specifically, itch inhibition was achieved by intrasegmental stimuli in the noxious range<sup>97,109,110,182</sup>. It has to be notice that in all of these studies the counter stimulations were long (3 minutes<sup>182</sup> and 30 seconds<sup>97,109</sup>) in opposition to the present study in which the stimuli lasted 5 seconds to avoid any possibility of hyperalgesia (both primary and secondary). The transient and short nature of the applied stimuli could be the reason for lack of involvement of the descending inhibitory pathways on the itch sensation.

Taken together the results of this study show that depending on the type and the pathways involved, itch can be differently modulated by counter-stimuli.



## **CHAPTER 3. THE EFFECTS OF INNOCUOUS HEAT ON ITCH**

The considerations made in this chapter are based on Study II<sup>74</sup>.

Itch is defined as chronic when it persists for more than 6 weeks and generally comes with two characteristic symptoms: alloknesis (when a non-pruritogen stimulus causes itch sensation) and hyperknesis (when a pruritogen stimulus causes a stronger and/or longer itch sensation)<sup>183–185</sup>. Many studies have focused on a specific type of dysesthesia called warmth hyperknesis, the mechanism where skin warming lead to an exacerbation in itch intensity, both in animal models<sup>108</sup> and in patients<sup>186–188</sup>. Nonetheless, a clear understanding of the mechanism characterizing warmth hyperknesis is lacking and therefore further investigations are needed.

Study II aimed to explore the effect of mild skin warming on various human itch models. To do so, the skin of the volar forearms of the healthy subject was heated either by +4 or +7°C above their measured and recorded individual baseline ( $\approx 32^\circ\text{C}$ ) using an infrared lamp. Once the target temperature was achieved, itch was provoked either by histamine, serotonin or cowhage, and itch intensity was assessed for up to 10 minutes, with constant and precise skin warming by the infrared lamp (Study II Fig1).

Furthermore, it was also investigated the superficial blood perfusion at two different time points after the skin warming and the itch intensity recording were stopped (Study II Fig1).

### **3.1 SELECTIVE WARMTH HYPERKNESIS FOR HISTAMINERGIC AND SEROTONINERGIC ITCH**

Histaminergic itch intensity was significantly increased when the skin was warmed by 7°C ( $\approx 39^\circ\text{C}$ ), the same effect was not observed when the skin temperature was increased by 4°C ( $\approx 36^\circ\text{C}$ ; Study II Fig.3). This result is in direct contrast with what was previously reported in animals, where histaminergic itch was not enhanced at any of the analysed temperatures<sup>108</sup>. Interestingly, the warmth hyperknesis phenomenon was present only in the first 2 minutes after itch induction, meaning the onset phase, and not at later stages.

Serotonin-evoked itch intensity was enhanced when the skin temperature was increased by 4°C ( $\approx 36^\circ\text{C}$ ) and not by 7°C ( $\approx 39^\circ\text{C}$ ; Study II Fig. 4). Animal preclinical findings confirm these data, where serotonergic itch was

enhanced when the skin temperature was  $\approx 36^{\circ}\text{C}$ <sup>108</sup>. Also, in this case, the increased itch sensation was present in the early part after the itch provocation, specifically in the first 4 minutes (Study II Fig.4).

As explained in section 1.4.1 of this dissertation, TRPVs have a major role in transmitting heat perception from innocuous to noxious<sup>115</sup>. The temperatures used in Study II are mainly sensed by TRPV3 and TRPV4 ( $\geq 33^{\circ}\text{C}$  and  $27-42^{\circ}\text{C}$  respectively). TRPV4 has been recently linked to histaminergic itch<sup>133</sup> and serotonergic itch<sup>134,135</sup>, whereas a direct interaction between TRPV3 and these itch modalities has yet to be shown. Therefore, it can be speculated that there is the direct involvement of TRPV4 in the hyperknesis effect hereby observed, but other key players cannot be ruled out.

Of particular interest is the observation that hyperknesis occurred only in the initial phase of the itch sensation (Study II Fig.3-4), suggesting that the itch modulation by activation of TRPVs is more relevant in the onset phase and at later stages is less important (in regard to the itch intensity).

## **3.2 COWHAGE-INDUCED ITCH IS NOT AFFECTED BY INNOCUOUS SKIN HEATING**

Cowhage-evoked itch intensity was not modified by the increase in skin temperature both by  $4$  and  $7^{\circ}\text{C}$  (Study II Fig.5). Interestingly, even though these results showed a lack of itch modulation following TRPV3 activation (via temperature), it has been shown how TRPV3 is important in the itch transmission when evoked by cowhage<sup>130</sup>. Specifically, when keratinocytes lacked TRPV3, there was a decreased itch behaviour in mice, when treated with PAR2 agonists (compared with wild-type animals)<sup>130</sup>. One could assume that, despite the well-documented interaction, prior activation of TRPV3 is not able to exacerbate itch intensity. It could be that cowhage spicules activate PAR2 which in turn recruits TRPV3 to exert its pruritic effect, but further or greater TRPV3 activation is unable to increase itch intensity.

Another reason for the lack of warmth hyperknesis in this condition could be that pre-activation of TRPV3 does not facilitate itch.

But we cannot exclude the possibility of increased warmth sensation given that this was not recorded in the experiment.

## **3.3 ENHANCED SUPERFICIAL BLOOD PERFUSION DUE TO INNOCUOUS SKIN HEATING**

Other than itch intensity, Study II explored the combined effect of different individual itch provocation and mild skin warming onto superficial blood



perfusion. The results from this analysis showed that superficial blood perfusion following histaminergic itch induction was unaffected by the continuous increase in skin temperature (Study II Fig. 6). It is known that histamine-evoked itch alone can generate a substantial neurogenic inflammation with subsequent increased blood perfusion together with cutaneous redness and wheal<sup>51</sup>. Therefore, one could suggest that higher skin temperature does not increase this state any further.

Superficial blood perfusion was significantly enhanced when skin temperature was increased by 7°C for serotonergic itch, while for cowhage-evoked itch it was increased both at +4 and +7°C above the resting skin temperature. This line of evidence confirm, what previously observed, that neurogenic inflammation is not necessary for the development of itch sensation because the first was increased exactly when the latter was not different from the control condition<sup>49,80,92,189,190</sup>. A very captivating explanation could be that increased superficial blood perfusion leads to increased tissue clearance and therefore this would counteract what would be an increased itch intensity, but this study does not provide enough data to make such an assumption. Further investigations in this regard are needed.

It has been shown that TRPV3 and PAR2 participate in the inflammatory response<sup>127,191–195</sup>. Therefore, the increased blood perfusion could be due to the probable co-activation of these two receptors that would lead to a more pronounced inflammatory response.

Regarding serotonin-evoked itch, increased blood perfusion was achieved only when the skin was heated by 7°C. As we said earlier, to date there is no evidence suggesting an interaction between serotonergic itch and TRPV3, rather with TRPV4<sup>134,135</sup>. Temperature-selective increased neurogenic inflammation in this setup could be due to a marked activation of TRPV4 (i.e. skin temperature ≈39°C); once again this is just a speculation that needs to be confirmed.



## CHAPTER 4. THE EFFECTS OF ESCALATING HEAT ON TOP OF ANAESTHETIZED SKIN

This chapter is based on the results shown in the manuscript submitted, Study III<sup>111</sup>.

As discussed throughout the introduction and the previous studies of this dissertation, the relationship between heat and itch is complicated and has many aspects to be analysed. In section 1.5 it was mentioned how topical cutaneous anaesthetisation is able to inhibit both pain and itch by blocking Nav channel transmission<sup>25,145,146,152,153,196</sup>. There is some evidence regarding the effectiveness of topical anaesthetics in various itch conditions, particularly neuropathic itch<sup>25,152,197,198</sup>. However, the use of these topical anaesthetics is not a feasible clinical solution to itch and pain, especially when these are chronic.

Study III investigated whether escalating heat on top of anaesthetized skin can provoke a paradoxical itch sensation when the anaesthesia is subsiding. This was done to explore, using a more modern and systematic approach, the results achieved by a 70 years old study<sup>156</sup>. In this study, the skin was anaesthetised by 2% procaine with subcutaneous injections. Once the efficacy of the anaesthesia was verified by pinprick sensibility, the area was heated by an infrared lamp to 41°C. After 15 minutes from the anaesthetisation, a paradoxical itch sensation started to develop, while simultaneously numbness was still present but pinprick sensibility was recovered<sup>156</sup>. To date, we are not aware of any further study investigating this peculiar mechanism, that could be useful to develop new sensitive targets.

In Study III, healthy skin was superficially anaesthetized by using intradermal injection with lidocaine 2% together with adrenalin as vasoconstrictor to achieve a qualitative better anaesthetisation<sup>199</sup>. Subsequently, on top of the same spot, an escalating heat ramp, lasting 1 minute in total and ranging from 38 to 50 °C, was applied every five minutes. To the participants it was asked to continuously rate their itch and pain sensation simultaneously during every heat stimulation (Study III Fig.S1). The results were compared with saline treatment as internal control.

To have a better idea of the different modulations involved in this study, the analysis of the itch and pain sensation was divided into 3 subgroups according to the temperature at which the intensity was rated. Innocuous range i.e. 38-42°C, painful range i.e. 42.2-46°C and finally noxious range i.e. 46.2-50°C.

## 4.1 PAIN INTENSITY

Pain sensation was reduced in the area treated with lidocaine compared to saline, specifically in the noxious range in the first 15 minutes (Study III Fig.1). This result suggests that, after this period, the anaesthetisation is subsided enough to not be able to inhibit strong nociceptive stimulation.

Interestingly, a previous study showed that an intradermal injection of lidocaine led to anaesthetisation being present after 20 minutes and subsiding after 120 minutes, this was tested by various heat stimuli ranging from 40 to 52°C<sup>200</sup>. Whereas, Cormia and Kuykendall<sup>156</sup> started to see the aforementioned “paradoxical itch sensation” exactly 15 minutes after the anaesthetic treatment.

When the escalating stimulation was in the painful and noxious range, 40 minutes after the anaesthetisation, it was observed a decreased pain intensity. To the best of our knowledge, this pain inhibition at this specific time point does not have a clear explanation and further studies are needed to assess whether it was an artefact or a specific unexplored new mechanism.

## 4.2 ITCH INTENSITY

The results in Study III showed no significant differences between saline and lidocaine treatment in itch intensity, in any of the temperature ranges and time points analysed. This is in sharp contrast with what was observed 70 years ago<sup>156</sup>.

As explained in section 1.5, a recent study characterized the different sensory fibres by discriminating them according to their transcriptome<sup>151</sup>. The leading hypothesis prior to Study III was that after about 15 minutes (as previously showed<sup>156</sup>) the fibres of the uppermost layer (superficial epidermis<sup>154,155</sup>) (i.e. NP fibres) would recover faster from the anaesthetisation and therefore would transmit a paradoxical itch sensation, as a consequence to noxious heat stimulation. For this reason, intradermal injections were preferred to a more common and modern method such as anaesthetic cream (e.g. EMLA), which would achieve a deeper, longer and less specific anaesthetisation.

It has to be pointed out that in the previous study, the temperature at which it was achieved an itch sensation was constant and developed by a radiant heat source, in opposition to Study III in which there was a heat ramp by direct contact with the skin. This methodological difference could have been what led to the development of paradoxical itch sensation due to the prolonged stimulation. Further studies are needed to explore different heat modalities and more specific temperatures in this setup to understand better the mechanisms at play.

## CHAPTER 5. GENERAL DISCUSSION

There is a need for new targets and therapies able to help chronic itch patients. The three studies presented in this thesis explore how heat from innocuous to noxious range can modulate itch. The ultimate goal of these studies was to understand which receptors and/or pathways involved in the processing of thermal sensation can be exploited toward the development of new treatments.

In brief the take home messages are:

- 1) Short noxious heat stimuli can greatly inhibit itch.
- 2) Innocuous skin warming can exacerbate itch.
- 3) Heat stimuli onto anaesthetized skin are not able to evoke itch sensation.

1) Very transient (5 seconds) noxious homotopical heat stimuli can inhibit different modalities of itch (histaminergic and cowhage-induced itch) to a different extent. Specifically, histaminergic itch showed a greater inhibition compared to cowhage-evoked itch (up to 76% compared to 43%, respectively<sup>73</sup>).

Recently, some tentative device has been developed and studied in a clinical context, consisting of the application of 51°C for 3 or 6 seconds on top of insect bites<sup>201</sup>. This latter study, showed how swelling, pain intensity and itch intensity, coming from wasps, bees or mosquitoes bites, were greatly inhibited, demonstrating that transient noxious heat stimuli can help in modulating itch intensity<sup>201</sup>.

Although, it is widely known that itch sensation due to insect bites is primarily histaminergic and this study can provide some knowledge about the intricate interaction between itch and pain, further systematic investigations are needed to better understand the mechanisms at play.

It has to be kept in mind that this approach would be scarcely applicable in chronic itch patients due to the large section of skin on which noxious heat should be applied as well as probably repeated exposure during the day.

Moreover, most of the itch symptomatic conditions (except for urticaria and mastocytosis) are unaffected by the treatment with antihistamines<sup>25,152</sup> therefore it is important to find mechanisms able to inhibit mostly non-histaminergic itch.

For this reason, Study I focused more on the potential mechanism underlining such an effect and the involvement of TRPV1 (important for the development of histaminergic itch<sup>91,162-167</sup>) and TRPA1 (recruited during cowhage-induced itch<sup>168,169</sup>) was hypothesized.

Capsaicin (a potent TRPV1 agonist) is used in clinical setup as an 8% cream able to achieve on some occasions long-lasting itch relief<sup>153,202–206</sup>. Whereas, to the best of our knowledge compounds acting on TRPA1 are yet to be explored as potential drugs for treating itch.

The data presented in Study I together with these clinical observations strongly suggest continuing the research toward TRPV1 and/or TRPA1 activation as a valuable targets to inhibit chronic itch.

2) The known mechanism of warmth hyperknesis<sup>108,186–188</sup>, was observed when itch was evoked by histamine and serotonin and not by cowhage. It can be assumed that the enhanced itch sensation could originate from the activation of TRPV4 rather than TRPV3, because of the selective documented interaction that this receptor has with both histaminergic<sup>133</sup> and serotonergic itch<sup>134,135</sup> pathways. Moreover, despite the known interaction between cowhage-induced itch and TRPV3<sup>130</sup>, the pre-activation by skin warming of this latter receptor did not result in enhanced itch intensity.

TRPV4 is generally expressed by both sensory neurons and keratinocytes<sup>207</sup>. In keratinocytes, other than participate in itch development of histaminergic and serotonergic itch, TRPV4 activation leads to the release of endothelin 1, known for its pruritogen activity<sup>67,133,208</sup>. Interestingly, this evidence shows also that keratinocytes are able to initiate itch sensation without the involvement of neuronal cells<sup>34</sup>.

Given these premises, a potential new target for therapies could be the antagonism of TRPV4. To the best of our knowledge, such a mechanism has yet to be explored for clinical purposes.

3) By using a more systematic and modern approach to investigate a previously observed effect<sup>156</sup> data presented in Study III did not confirm the results from an old study, however, few considerations can be done.

Firstly, if the Study III hypothesis would have been confirmed, could have given a new potential target by retro-analysing at what exact temperature itch sensation occur and therefore which among the thermal receptor was most likely involved in this mechanism.

Secondly, in the paper published by Cormia and Kuykendall, the stimulation able to evoke a paradoxical itch sensation was achieved by irradiation with an infrared lamp. The same methodology was used in Study II to mildly warm the skin which resulted in hyperknesis. Together these pieces of evidence could suggest that radiant heat stimulation, specifically using an infrared lamp, can facilitate itch sensation in certain conditions. This would be a novel and

unexplored role for such a stimulation but need further investigations to be verified and understood.

All things considered, the results and speculations highlighted in this dissertation point out that heat perception and TRP receptors as key player in itch sensation, that need more attention in the development of new potential therapies.

## 5.1 LIMITATIONS

The studies hereby analysed present some limitations.

In Study I<sup>73</sup>, only cowhage was used as a non-histaminergic itch model. Whereas, in Study II it was used also serotonin-evoked itch. This was done for practical reasons, serotonergic itch is induced by iontophoresis which requires more space on the skin due to the presence of the conductive pad and ion chamber (compared to histamine- and cowhage evoked itch, which requires only the treated area). Therefore having 5 different areas on the forearm (as for histamine- and cowhage-evoked itch) would have been not feasible. Moreover, the second experiment of Study I would have been impossible to be performed in a single session without applying itch provocation on the same area already treated previously (Study I Fig.1B-C). We could hypothesize that serotonergic itch would have been inhibited by noxious heat as it happened for the other 2 itch modalities, to understand to what extent further investigations are needed.

The analysis of the skin blood perfusion in Study II<sup>74</sup> was not adopted in the other two studies. This was due to the intensity of the heat stimulation applied that most likely would have resulted in an oversaturated picture, to be expected due to the cutaneous redness seen after the heat stimulations (data not shown).

The heat stimulations applied in the three studies differ substantially from the qualitative point of view. In Study I and III contact heat stimuli were used, whereas in Study II it was used a source of radiant heat. The latter was chosen to better simulate what was observed in clinic, where patients have exacerbated itch in warm environments. Moreover, by applying a contact form of stimulation the results could have been invalidated due to mechanical itch inhibition.

In Study I and II more intermediate temperatures could have been explored, however, due to limited space on the forearms (Study I Fig.1 and Study II Fig.1) this was not possible.

On the other hand, in Study III it could have been more helpful to use very specific heat stimuli (i.e. few chosen temperatures) instead of using a heat

ramp. However, our purpose was to understand at exactly what temperature the potential paradoxical itch sensation would occur, to have a systematic view of the effects at play, for this reason, a heat ramp was preferred.



## CHAPTER 6. CONCLUSIONS

This dissertation aimed to evaluate 1) the effects of transient noxious heat stimuli on itch, 2) investigate how various itch models interact with continuous warm innocuous heat stimulation and 3) explore old results showing a “paradoxical itch sensation” when anaesthetised skin receives a “burning” heat stimulation<sup>156</sup>.

Study I<sup>73</sup> showed that transient homotopical noxious heat counter stimulations can inhibit itch showing a dose-response effect, where the higher the temperature the higher the inhibitory effect is. These are more effective on histaminergic compared with non-histaminergic itch. Moreover, heterotopic counter stimuli failed to inhibit itch, suggesting that transient heat stimuli effect is mainly due to peripheral mechanisms.

Study II<sup>74</sup> partially confirmed what was already observed in animal<sup>108</sup> where serotonergic itch was enhanced by innocuous skin warming. In contrast with data in rodents, it also showed exacerbation of histaminergic itch due to skin warming. These data may give a more detailed explanation of what has been observed in clinic<sup>186–188</sup>, the so-called warmth hyperknesis and open up a new avenue in the development of new therapies by antagonising TRPV4, the main suspect in the observed itch exacerbation.

Interestingly, cowhage-induced itch was not aggravated, however, there was an increased skin blood perfusion. This was probably due to the co-activation of PAR2 (by cowhage spicules) and TRPV3 (by increased temperature) known to interact<sup>130</sup> and probably cause an increased inflammatory response. Study III failed to confirm what was observed in an old study<sup>156</sup>, since noxious heat applied on top of previously anaesthetised skin did not evoke any paradoxical itch sensation. As briefly explained in section 5.1, it could be useful to investigate the effect of specifically chosen temperature to better match the methodology used in the previous study.

In summary, these studies highlight a unique role in the interaction between heat and itch perception and intensity, suggesting how fine-tuning and modulation of TRPVs receptors can translate into inhibition or increased itch intensity. Particularly, following up on the use and the development of TRPV1 agonists and exploring the effect of endogenous and synthetic TRPV4 antagonists.

Further proof-of-concept studies in patient populations are needed to validate the experimental data and the initial step toward a better understanding of new non-pharmacological targets to treat chronic itch.



## LITERATURE LIST

1. Yosipovitch, G. & Bernhard, J. D. Chronic Pruritus. *N. Engl. J. Med.* **368**, 1625–1634 (2013).
2. Yosipovitch, G., Rosen, J. D. & Hashimoto, T. Itch: From mechanism to (novel) therapeutic approaches. *Journal of Allergy and Clinical Immunology* **142**, 1375–1390 (2018).
3. Yosipovitch, G., Greaves, M. W. & Schmelz, M. Itch. *Lancet (London, England)* **361**, 690–4 (2003).
4. Lay, M. & Dong, X. Neural Mechanisms of Itch. *Annual Review of Neuroscience* **43**, 187–205 (2020).
5. Dong, X. & Dong, X. Peripheral and Central Mechanisms of Itch. *Neuron* **98**, 482–494 (2018).
6. Weisshaar, E. Epidemiology of Itch. *Curr. Probl. Dermatology* **50**, 5–10 (2016).
7. Roh, Y. S., Choi, J., Sutaria, N. & Kwatra, S. G. Itch: Epidemiology, clinical presentation, and diagnostic workup. *J. Am. Acad. Dermatol.* **86**, 1–14 (2022).
8. Kulkarni, V., Liu, D., Aggarwal, P., Gamsarian, V. & Fleischer, A. B. Global epidemiology of itch from 1990 to 2017: gender, age, sanitation, and air pollution as risk factors. *Itch* **7**, e60–e60 (2022).
9. Weisshaar, E. *et al.* European S2k Guideline on Chronic Pruritus. *Acta Derm. Venereol.* **99**, 469–506 (2019).
10. Global Health Data Exchange (GBDE). *Global Burden of Disease Study 2017 (GBDE 2017)*. (The Lancet, 2017).
11. Brenaut, E., Garlantezec, R., Talour, K. & Misery, L. Itch characteristics in five dermatoses: Non-atopic eczema, atopic dermatitis, urticaria, psoriasis and scabies. *Acta Dermato-Venereologica* **93**, 573–574 (2013).
12. Zachariae, R., Lei, U., Haedersdal, M. & Zachariae, C. Itch severity and quality of life in patients with pruritus: Preliminary validity of a danish adaptation of the itch severity scale. *Acta Derm. Venereol.* **92**, 508–514 (2012).

13. Schneider, G. *et al.* Psychosomatic cofactors and psychiatric comorbidity in patients with chronic itch. *Clin. Exp. Dermatol.* **31**, 762–767 (2006).
14. Tessari, G. *et al.* The impact of pruritus on quality of life of patients undergoing dialysis: A single-center cohort study. *J. Nephrol.* **22**, 241–248 (2009).
15. Chen, S. C. Pruritus. *Dermatologic Clinics* **30**, 309–321 (2012).
16. Leader, B., Carr, C. W. & Chen, S. C. Pruritus epidemiology and quality of life. *Handb. Exp. Pharmacol.* **226**, 15–38 (2015).
17. Matteredne, U., Apfelbacher, C. J., Vogelgsang, L., Loerbroks, A. & Weisshaar, E. Incidence and determinants of chronic pruritus: A population-based cohort study. *Acta Derm. Venereol.* **93**, 532–537 (2013).
18. Holm, E. A., Esmann, S. & Jemec, G. B. E. Does visible atopic dermatitis affect quality of life more in women than in men? *Gend. Med.* **1**, 125–130 (2004).
19. Stander, S. *et al.* Gender differences in chronic pruritus: women present different morbidity, more scratch lesions and higher burden. *Br J Dermatol* **168**, 1273–1280 (2013).
20. Tripathi, R., Knusel, K. D., Ezaldein, H. H., Bordeaux, J. S. & Scott, J. F. The cost of an itch: A nationally representative retrospective cohort study of pruritus-associated health care expenditure in the United States. *J. Am. Acad. Dermatol.* **80**, 810–813 (2019).
21. Yosipovitch, G., Carstens, E. & McGlone, F. Chronic itch and chronic pain: Analogous mechanisms. *Pain* **131**, 4–7 (2007).
22. Liu, X., Miao, X. H. & Liu, T. More than Scratching the Surface: Recent Progress in Brain Mechanisms Underlying Itch and Scratch. *Neurosci. Bull.* **36**, 85–88 (2020).
23. Akiyama, T., Carstens, M. I. & Carstens, E. Differential itch- and pain-related behavioral responses and  $\mu$ -opioid modulation in mice. *Acta Derm. Venereol.* **90**, 575–581 (2010).
24. Dawn, A. G. & Yosipovitch, G. Butorphanol for treatment of intractable pruritus. *J. Am. Acad. Dermatol.* **54**, 527–531 (2006).
25. Leslie, T. A., Greaves, M. W. & Yosipovitch, G. Current topical and

- systemic therapies for itch. *Handb. Exp. Pharmacol.* **226**, 337–356 (2015).
26. Marsella, R., Ahrens, K., Wilkes, R., Soeberdt, M. & Abels, C. Topical  $\kappa$ -opioid receptor agonist asimadoline improves dermatitis in a canine model of atopic dermatitis. *Exp. Dermatol.* **31**, 628–632 (2022).
  27. Ádám, D. *et al.* Opioidergic Signaling—A Neglected, Yet Potentially Important Player in Atopic Dermatitis. *International Journal of Molecular Sciences* **23**, 4140 (2022).
  28. Han, L. *et al.* A subpopulation of nociceptors specifically linked to itch. *Nat. Neurosci.* **16**, 174–182 (2013).
  29. Davidson, S. *et al.* Pruriceptive spinothalamic tract neurons: physiological properties and projection targets in the primate. *J. Neurophysiol.* **108**, 1711–23 (2012).
  30. Jansen, N. A. & Giesler, G. J. Response characteristics of pruriceptive and nociceptive trigeminoparabrachial tract neurons in the rat. *J. Neurophysiol.* **113**, 58–70 (2015).
  31. Moser, H. R. & Giesler, G. J. Characterization of pruriceptive trigeminothalamic tract neurons in rats. *J. Neurophysiol.* **111**, 1574–1589 (2014).
  32. Schmelz, M. How Do Neurons Signal Itch? *Frontiers in Medicine* **8**, 284 (2021).
  33. Carstens, E., Follansbee, T. & Iodi Carstens, M. The challenge of basic itch research. *Acta Dermato-Venereologica* **100**, 3–9 (2020).
  34. Yosipovitch, G., Andersen, H. H., Arendt-Nielsen, L. & Andersen, H. H. *Itch and Pain: similarities, Interactions, and Differences*. (Wolters Kluwer, 2020).
  35. Reddy, V. B., Iuga, A. O., Shimada, S. G., LaMotte, R. H. & Lerner, E. A. Cowhage-evoked itch is mediated by a novel cysteine protease: A ligand of protease-activated receptors. *J. Neurosci.* **28**, 4331–4335 (2008).
  36. Sun, Y.-G. G. *et al.* Cellular basis of itch sensation. *Science* **325**, 1531–4 (2009).
  37. Buddenkotte, J. & Steinhoff, M. Pathophysiology and therapy of pruritus in allergic and atopic diseases. *Allergy: European Journal of*

- Allergy and Clinical Immunology* **65**, 805–821 (2010).
38. Mishra, S. K. & Hoon, M. A. The cells and circuitry for itch responses in mice. *Science* **340**, 968–71 (2013).
  39. Bader, M., Alenina, N., Andrade-Navarro, M. A. & Santos, R. A. Mas and its related G protein-coupled receptors, Mrgprs. *Pharmacol. Rev.* **66**, 1080–1105 (2014).
  40. Aresh, B. *et al.* Spinal cord interneurons expressing the gastrin-releasing peptide receptor convey itch through VGLUT2-mediated signaling. *Pain* **158**, 945–961 (2017).
  41. Mu, D. *et al.* A central neural circuit for itch sensation. *Science* (80-. ). **357**, 695–699 (2017).
  42. Ross, S. E. *et al.* Loss of Inhibitory Interneurons in the Dorsal Spinal Cord and Elevated Itch in Bhlhb5 Mutant Mice. *Neuron* **65**, 886–898 (2010).
  43. LaMotte, R. H., Shimada, S. G., Green, B. G. & Zeltzman, D. Pruritic and nociceptive sensations and dysesthesias from a spicule of cowhage. *J. Neurophysiol.* **101**, 1430–1443 (2009).
  44. Steinhoff, M., Oaklander, A. L., Szabó, I. L., Ständer, S. & Schmelz, M. Neuropathic itch. *Pain* **160**, S11–S16 (2019).
  45. Sun, S. *et al.* Leaky Gate Model: Intensity-Dependent Coding of Pain and Itch in the Spinal Cord. *Neuron* **93**, 840-853.e5 (2017).
  46. LaMotte, R. H., Shimada, S. G. & Sikand, P. Mouse models of acute, chemical itch and pain in humans. *Experimental Dermatology* **20**, 778–782 (2011).
  47. Yuan, T. *et al.* Assessment of itch and pain in animal models and human subjects. *Adv. Exp. Med. Biol.* **904**, 1–22 (2016).
  48. Akiyama, T., Nagamine, M., Carstens, M. I. & Carstens, E. Behavioral model of itch, allodynia, pain and allodynia in the lower hindlimb and correlative responses of lumbar dorsal horn neurons in the mouse. *Neuroscience* **266**, 38–46 (2014).
  49. Darsow, U., Ring, J., Scharein, E. & Bromm, B. Correlations between histamine-induced wheal, flare and itch. *Arch. Dermatol. Res.* **288**, 436–441 (1996).

50. Sikand, P., Shimada, S. G., Green, B. G. & LaMotte, R. H. Similar itch and nociceptive sensations evoked by punctate cutaneous application of capsaicin, histamine and cowhage. *Pain* **144**, 66–75 (2009).
51. Sikand, P., Shimada, S. G., Green, B. G. & LaMotte, R. H. Sensory responses to injection and punctate application of capsaicin and histamine to the skin. *Pain* **152**, 2485–2494 (2011).
52. Andersen, H. H., Elberling, J. & Arendt-Nielsen, L. Human Surrogate Models of Histaminergic and Non-histaminergic Itch. *Acta Derm. Venereol.* **95**, 771–7 (2015).
53. Bjerring, P. & Arendt-Nielsen, L. A quantitative comparison of the effect of local analgesics on argon laser induced cutaneous pain and on histamine induced wheal, flare and itch. *Acta Derm. Venereol.* **70**, 126–131 (1990).
54. Schmelz, M., Luz, O., Averbek, B. & Bickel, A. Plasma extravasation and neuropeptide release in human skin as measured by intradermal microdialysis. *Neurosci. Lett.* **230**, 117–120 (1997).
55. Rukwied, R., Lischetzki, G., McGlone, F., Heyer, G. & Schmelz, M. Mast cell mediators other than histamine induce pruritus in atopic dermatitis patients: A dermal microdialysis study. *Br. J. Dermatol.* **142**, 1114–1120 (2000).
56. Heinzerling, L. *et al.* The skin prick test - European standards. *Clin. Transl. Allergy* **3**, 3 (2013).
57. Lewis, T. The blood vessels of the human skin. *Br. Med. J.* **2**, 61–62 (1926).
58. Schmelz, M. & Petersen, L. J. Neurogenic inflammation in human and rodent skin. *News Physiol. Sci.* **16**, 33–37 (2001).
59. Steinhoff, M. *et al.* Modern Aspects of Cutaneous Neurogenic Inflammation. *Archives of Dermatology* **139**, 1479–1488 (2003).
60. Boas, D. A. & Dunn, A. K. Laser speckle contrast imaging in biomedical optics. *J. Biomed. Opt.* **15**, 011109 (2010).
61. Eriksson, S., Nilsson, J. & Stureson, C. Non-invasive imaging of microcirculation: A technology review. *Med. Devices Evid. Res.* **7**, 445–452 (2014).
62. Lindahl, F., Tesselaar, E. & Sjöberg, F. Assessing paediatric scald

- injuries using laser speckle contrast imaging. *Burns* **39**, 662–666 (2013).
63. Video capability revives interest in laser method. *Optics & Laser Europe magazine* Available at: <https://optics.org/article/27263>.
64. Stewart, C. J. *et al.* A comparison of two laser-based methods for determination of burn scar perfusion: Laser Doppler versus laser speckle imaging. *Burns* **31**, 744–752 (2005).
65. Nomura, S. *et al.* Reliability of laser speckle flow imaging for intraoperative monitoring of cerebral blood flow during cerebrovascular surgery: Comparison with cerebral blood flow measurement by single photon emission computed tomography. *World Neurosurgery* **82**, E753–E757 (2014).
66. Han, S. K., Mancino, V. & Simon, M. I. Phospholipase C $\beta$  3 Mediates the Scratching Response Activated by the Histamine H1 Receptor on C-Fiber Nociceptive Neurons. *Neuron* **52**, 691–703 (2006).
67. Kim, B. M., Lee, S. H., Shim, W. S. & Oh, U. Histamine-induced Ca<sup>2+</sup> influx via the PLA2/lipoxygenase/TRPV1 pathway in rat sensory neurons. *Neurosci. Lett.* **361**, 159–162 (2004).
68. Ko, K. *et al.* Discovery of a Novel Highly Selective Histamine H4 Receptor Antagonist for the Treatment of Atopic Dermatitis. *J. Med. Chem.* **61**, 2949–2961 (2018).
69. Mollanazar, N. K., Smith, P. K. & Yosipovitch, G. Mediators of Chronic Pruritus in Atopic Dermatitis: Getting the Itch Out? *Clin. Rev. Allergy Immunol.* **51**, 263–292 (2016).
70. Wahlgren, C. F. Itch and atopic dermatitis: An overview. *J. Dermatol.* **26**, 770–779 (1999).
71. Patel, T. & Yosipovitch, G. Therapy of pruritus. *Expert Opin. Pharmacother.* **11**, 1673–1682 (2010).
72. Twycross, R. *et al.* Itch: scratching more than the surface. *QJM An Int. J. Med.* **96**, 7–26 (2003).
73. Riccio, D., Andersen, H. H. & Arendt-Nielsen, L. Antipruritic effects of transient heat stimulation on histaminergic and nonhistaminergic itch. *Br. J. Dermatol.* **181**, 786–795 (2019).
74. Riccio, D., Andersen, H. H. & Arendt-Nielsen, L. Mild Skin Heating



- Evokes Warmth Hyperknesis Selectively for Histaminergic and Serotonergic Itch in Humans. *Acta Derm. Venereol.* **102**, adv00649 (2022).
75. Wahlgren, C. F. & Ekblom, A. Perception of histamine-induced itch elicited in three different skin regions. *Acta Derm. Venereol.* **71**, 205–208 (1991).
  76. Marzell, R., Reichwein, G., Gieler, U., Kupfer, J. & Schut, C. Itch induction by audiovisual stimuli and histamine iontophoresis: a randomized, controlled noninferiority study. *Br. J. Dermatol.* **182**, 1253–1261 (2020).
  77. Shelley, W. B. & Arthur, R. P. Studies on Cowhage (*Mucuna Pruriens*) and Its Pruritogenic Proteinase, Mucunain. *A. M. A. Arch. Dermatology* **72**, 399–406 (1955).
  78. Shelley, W. B. & Arthur, R. P. Mucunain, the active pruritogenic proteinase of cowhage. *Science (80- )*. **122**, 469–470 (1955).
  79. Johaneck, L. M. *et al.* Psychophysical and Physiological Evidence for Parallel Afferent Pathways Mediating the Sensation of Itch. *J. Neurosci.* **27**, 7490–7497 (2007).
  80. Andersen, H. H., Elberling, J., Vecchio, S. Lo & Arendt-Nielsen, L. Topography of itch: evidence of distinct coding for pruriception in the trigeminal nerve. *Itch (Philadelphia, Pa.)* **2**, e2 (2017).
  81. Reddy, V. B., Azimi, E., Chu, L. & Lerner, E. A. Mas-Related G-Protein Coupled Receptors and Cowhage-Induced Itch. *J. Invest. Dermatol.* **138**, 461–464 (2018).
  82. McNeil, B. D. *et al.* Identification of a mast-cell-specific receptor crucial for pseudo-allergic drug reactions. *Nature* **519**, 237–241 (2015).
  83. McNeil, B. & Dong, X. Mrgprs as Itch Receptors. in *Itch: Mechanisms and Treatment* (eds. Carstens, E. & Akiyama, T.) (CRC Press/Taylor & Francis, Boca Raton (FL), US, 2014).
  84. Azimi, E. *et al.* Dual action of neurokinin-1 antagonists on Mas-related GPCRs. *JCI Insight* **1**, 89362 (2016).
  85. Weisshaar, E., Ziethen, B. & Gollnick, H. Lack of efficacy of topical capsaicin in serotonin-induced itch. *Skin Pharmacol. Appl. Skin Physiol.* **13**, 1–8 (2000).

86. Hosogi, M., Schmelz, M., Miyachi, Y. & Ikoma, A. Bradykinin is a potent pruritogen in atopic dermatitis: A switch from pain to itch. *Pain* **126**, 16–23 (2006).
87. Rasul, A., Nordlind, K. & Wahlgren, C. F. Pruritic and vascular responses induced by serotonin in patients with atopic dermatitis and in healthy controls. *Acta Derm. Venereol.* **93**, 277–280 (2013).
88. Schmelz, M. *et al.* Chemical response pattern of different classes of C-nociceptors to pruritogens and algogens. *J. Neurophysiol.* **89**, 2441–2448 (2003).
89. Weisshaar, E., Ziethen, B. & Gollnick, H. Can a serotonin type 3 (5-HT<sub>3</sub>) receptor antagonist reduce experimentally-induced itch? *Inflamm. Res.* **46**, 412–416 (1997).
90. Sikand, P., Dong, X. & LaMotte, R. H. BAM8-22 peptide produces itch and nociceptive sensations in humans independent of histamine release. *J. Neurosci.* **31**, 7563–7 (2011).
91. Lamotte, R. H., Dong, X. & Ringkamp, M. Sensory neurons and circuits mediating itch. *Nat. Rev. Neurosci.* **15**, 19–31 (2014).
92. Aliotta, G. E., Vecchio, S. Lo, Elberling, J. & Arendt-Nielsen, L. Evaluation of itch and pain induced by bovine adrenal medulla (BAM)8–22, a new human model of non-histaminergic itch. *Exp. Dermatol.* (2022). doi:10.1111/EXD.14611
93. Liu, Q. *et al.* Mechanisms of itch evoked by  $\beta$ -alanine. *J. Neurosci.* **32**, 14532–7 (2012).
94. Christensen, J. D., Vecchio, S. L. O., Elberling, J., Arendt-Nielsen, L. & Andersen, H. H. Assessing punctate administration of beta-alanine as a potential human model of non-histaminergic itch. *Acta Derm. Venereol.* **99**, 222–223 (2019).
95. Olsen, R. V., Andersen, H. H., Møller, H. G., Eskelund, P. W. & Arendt-Nielsen, L. Somatosensory and vasomotor manifestations of individual and combined stimulation of TRPM8 and TRPA1 using topical L-menthol and trans-cinnamaldehyde in healthy volunteers. *Eur. J. Pain (United Kingdom)* **18**, 1333–1342 (2014).
96. Namer, B., Seifert, F., Handwerker, H. O. & Maihöfner, C. TRPA1 and TRPM8 activation in humans: Effects of cinnamaldehyde and menthol. *Neuroreport* **16**, 955–959 (2005).

97. Yosipovitch, G., Duque, M. I., Fast, K., Dawn, A. G. & Coghill, R. C. Scratching and noxious heat stimuli inhibit itch in humans: a psychophysical study. *Br. J. Dermatol.* **156**, 629–634 (2007).
98. Werner, Y. & Lindberg, M. Transepidermal water loss in dry and clinically normal skin in patients with atopic dermatitis. *Acta Derm. Venereol.* **65**, 102–105 (1985).
99. Geber, C. *et al.* Test-retest and interobserver reliability of quantitative sensory testing according to the protocol of the German Research Network on Neuropathic Pain (DFNS): A multi-centre study. *Pain* **152**, 548–556 (2011).
100. Rolke, R. *et al.* Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): standardized protocol and reference values. *Pain* **123**, 231–43 (2006).
101. Lowenstein, E. & Dallenbach, K. M. The Critical Temperatures for Heat and for Burning Heat. *Am. J. Psychol.* **42**, 423 (1930).
102. Jones, L. A. & Berris, M. The psychophysics of temperature perception and thermal-interface design. in *Proceedings - 10th Symposium on Haptic Interfaces for Virtual Environment and Teleoperator Systems, HAPTICS 2002* 137–142 (IEEE Comput. Soc, 2002). doi:10.1109/HAPTIC.2002.998951
103. Fillingim, R. B. *et al.* Sex, Gender, and Pain: A Review of Recent Clinical and Experimental Findings. *J. Pain* **10**, 447 (2009).
104. Park, S., Roh, S. H. & Lee, J. Y. Body regional heat pain thresholds using the method of limit and level: a comparative study. *Eur. J. Appl. Physiol.* **119**, 771–780 (2019).
105. Dyck, P. J. *et al.* Cool, warm, and heat-pain detection thresholds: testing methods and inferences about anatomic distribution of receptors. *Neurology* **43**, 1500–8 (1993).
106. Frahm, K. S., Gervasio, S., Arguissain, F. & Mouraux, A. New Insights into Cutaneous Laser Stimulation – Dependency on Skin and Laser Type. *Neuroscience* **448**, 71–84 (2020).
107. Hardy, J. D., Stolwijk, J. A., Hammel, H. T. & Murgatroyd, D. Skin temperature and cutaneous pain during warm water immersion. <https://doi.org/10.1152/jappl.1965.20.5.1014> **20**, 1014–1021 (1965).
108. Akiyama, T. *et al.* Innocuous warming enhances peripheral

- serotonergic itch signaling and evokes enhanced responses in serotonin-responsive dorsal horn neurons in the mouse. *J. Neurophysiol.* **117**, 251–259 (2017).
109. Yosipovitch, G., Fast, K. & Bernhard, J. D. Noxious heat and scratching decrease histamine-induced itch and skin blood flow. *J. Invest. Dermatol.* **125**, 1268–1272 (2005).
  110. Ishiiji, Y. *et al.* Repetitive scratching and noxious heat do not inhibit histamine-induced itch in atopic dermatitis. *Br. J. Dermatol.* **158**, 78–83 (2008).
  111. Riccio, D., Lo Vecchio, S. & Arendt-Nielsen, L. The effect of escalating heat stimulation on top of anaesthetized skin (SUBMITTED). (2022).
  112. Murray, F. S. & Weaver, M. M. Effects of ipsilateral and contralateral counterirritation on experimentally produced itch in human beings. *J. Comp. Physiol. Psychol.* **89**, 819–826 (1975).
  113. Caterina, M. J. *et al.* The capsaicin receptor: A heat-activated ion channel in the pain pathway. *Nature* **389**, 816–824 (1997).
  114. Voets, T. *et al.* The principle of temperature-dependent gating in cold- and heat-sensitive TRP channels. *Nat. 2004 4307001* **430**, 748–754 (2004).
  115. Dhaka, A., Viswanath, V. & Patapoutian, A. Trp Ion Channels and Temperature Sensation. *Annu. Rev. Neurosci.* **29**, 135–161 (2006).
  116. Tóth, B. I., Oláh, A., Szöllösi, A. G. & Bíró, T. TRP channels in the skin. *British Journal of Pharmacology* **171**, 2568–2581 (2014).
  117. Tominaga, M. & Caterina, M. J. Thermosensation and pain. *J. Neurobiol.* **61**, 3–12 (2004).
  118. Caterina, M. J. *et al.* Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science (80- )*. **288**, 306–313 (2000).
  119. Inoue, K. *et al.* Functional vanilloid receptors in cultured normal human epidermal keratinocytes. *Biochem. Biophys. Res. Commun.* **291**, 124–129 (2002).
  120. Tóth, B. I. *et al.* Endocannabinoids modulate human epidermal keratinocyte proliferation and survival via the sequential engagement of cannabinoid receptor-1 and transient receptor potential vanilloid-1. *J. Invest. Dermatol.* **131**, 1095–1104 (2011).

121. Petersen, K. L. & Rowbotham, M. C. A new human experimental pain model: The heat/capsaicin sensitization model. *Neuroreport* **10**, 1511–1516 (1999).
122. Kittaka, H. & Tominaga, M. The molecular and cellular mechanisms of itch and the involvement of TRP channels in the peripheral sensory nervous system and skin. *Allergology International* **66**, 22–30 (2017).
123. Tóth, B. I., Szallasi, A. & Bíró, T. Transient receptor potential channels and itch: How deep should we scratch? *Handb. Exp. Pharmacol.* **226**, 89–133 (2015).
124. Kojima, I. & Nagasawa, M. TRPV2. *Handb. Exp. Pharmacol.* **222**, 247–272 (2014).
125. Smith, G. D. *et al.* TRPV3 is a temperature-sensitive vanilloid receptor-like protein. *Nat. 2002 4186894* **418**, 186–190 (2002).
126. Peier, A. M. *et al.* A heat-sensitive TRP channel expressed in keratinocytes. *Science (80-. )*. **296**, 2046–2049 (2002).
127. Szöllösi, A. G. *et al.* Activation of TRPV3 Regulates Inflammatory Actions of Human Epidermal Keratinocytes. *J. Invest. Dermatol.* **138**, 365–374 (2018).
128. Asakawa, M. *et al.* Association of a mutation in TRPV3 with defective hair growth in rodents. *J. Invest. Dermatol.* **126**, 2664–2672 (2006).
129. Lin, Z. *et al.* Exome sequencing reveals mutations in TRPV3 as a cause of Olmsted syndrome. *Am. J. Hum. Genet.* **90**, 558–564 (2012).
130. Zhao, J. *et al.* PAR2 Mediates Itch via TRPV3 Signaling in Keratinocytes. *J. Invest. Dermatol.* **140**, 1524–1532 (2020).
131. Park, C. *et al.* TRPV3 Channel in Keratinocytes in Scars with Post-Burn Pruritus. *Int. J. Mol. Sci.* **18**, 2425 (2017).
132. Güler, A. D. *et al.* Heat-evoked activation of the ion channel, TRPV4. *J. Neurosci.* **22**, 6408–6414 (2002).
133. Chen, Y. *et al.* Transient Receptor Potential Vanilloid 4 Ion Channel Functions as a Pruriceptor in Epidermal Keratinocytes to Evoke Histaminergic Itch \*. *J. Biol. Chem.* **291**, 10252–10262 (2016).
134. Akiyama, T. *et al.* Involvement of TRPV4 in Serotonin-Evoked Scratching. *J. Invest. Dermatol.* **136**, 154–160 (2016).

135. Snyder, L. M., Kuzirian, M. S. & Ross, S. E. An unexpected role for TRPV4 in serotonin-mediated itch. *J. Invest. Dermatol.* **136**, 7 (2016).
136. Julius, D. TRP channels and pain. *Annual Review of Cell and Developmental Biology* **29**, 355–384 (2013).
137. Zygmunt, P. M. & Högesätt, E. D. TRPA1. *Handb. Exp. Pharmacol.* **222**, 583–630 (2014).
138. Trevisan, G. *et al.* TRPA1 mediates trigeminal neuropathic pain in mice downstream of monocytes/macrophages and oxidative stress. *Brain* aww038- (2016). doi:10.1093/brain/aww038
139. Landini, L. *et al.* TRPA1 Role in Inflammatory Disorders: What Is Known So Far? *International Journal of Molecular Sciences* **23**, 4529 (2022).
140. Karashima, Y. *et al.* TRPA1 acts as a cold sensor in vitro and in vivo. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 1273–1278 (2009).
141. Camino, D. Del *et al.* TRPA1 Contributes to Cold Hypersensitivity. *J. Neurosci.* **30**, 15165–15174 (2010).
142. Vandewauw, I. *et al.* A TRP channel trio mediates acute noxious heat sensing. *Nature* **555**, 662–666 (2018).
143. Clapham, D. E. & Miller, C. A thermodynamic framework for understanding temperature sensing by transient receptor potential (TRP) channels. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 19492–19497 (2011).
144. Moparthy, L. *et al.* Human TRPA1 is a heat sensor displaying intrinsic U-shaped thermosensitivity. *Sci. Rep.* **6**, 1–10 (2016).
145. Arcisio-Miranda, M., Muroi, Y., Chowdhury, S. & Chanda, B. Molecular mechanism of allosteric modification of voltage-dependent sodium channels by local anesthetics. *J. Gen. Physiol.* **136**, 541–554 (2010).
146. Ahern, C. A., Payandeh, J., Bosmans, F. & Chanda, B. *The hitchhiker's guide to the voltage-gated sodium channel galaxy.* *Journal of General Physiology* **147**, 1–24 (The Rockefeller University Press, 2016).
147. Sobanko, J. F., Miller, C. J. & Alster, T. S. Topical anesthetics for dermatologic procedures: A review. *Dermatologic Surgery* **38**, 709–721 (2012).

148. Friedman, P. M., Mafong, E. A., Friedman, E. S. & Geronemus, R. G. Topical anesthetics update: EMLA and beyond. *Dermatologic Surg.* **27**, 1019–1026 (2001).
149. Mick, G. & Correa-Illanes, G. Topical pain management with the 5 lidocaine medicated plaster a review. *Current Medical Research and Opinion* **28**, 937–951 (2012).
150. Derry, S., Wiffen, P. J., Moore, R. A. & Quinlan, J. Topical lidocaine for neuropathic pain in adults. *Cochrane Database of Systematic Reviews* **2017**, (2014).
151. Usoskin, D. *et al.* Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nat. Neurosci.* **18**, 145–153 (2015).
152. Weisshaar, E. *et al.* European guideline on chronic pruritus: In cooperation with the European dermatology forum (EDF) and the European academy of dermatology and venereology (EADV). *Acta Derm. Venereol.* **92**, 563–581 (2012).
153. Rosen, J. D., Fostini, A. C. & Yosipovitch, G. Diagnosis and Management of Neuropathic Itch. *Dermatologic Clinics* **36**, 213–224 (2018).
154. Zylka, M. J., Rice, F. L. & Anderson, D. J. Topographically distinct epidermal nociceptive circuits revealed by axonal tracers targeted to Mrgprd. *Neuron* **45**, 17–25 (2005).
155. Lowy, D. B., Makker, P. G. S. & Moalem-Taylor, G. Cutaneous Neuroimmune Interactions in Peripheral Neuropathic Pain States. *Front. Immunol.* **12**, 1064 (2021).
156. Cormia, F. E. & Kuykendall, V. Experimental histamine pruritus. II. Nature; physical and environmental factors influencing development and severity. *J. Invest. Dermatol.* **20**, 429–446 (1953).
157. Braz, J., Solorzano, C., Wang, X. & Basbaum, A. I. Transmitting Pain and Itch Messages: A Contemporary View of the Spinal Cord Circuits that Generate Gate Control. *Neuron* **82**, 522–536 (2014).
158. Wallengren, J. & Sundler, F. Cutaneous field stimulation in the treatment of severe itch. *Arch. Dermatol.* **137**, 1323–1325 (2001).
159. Pedersen, J. L. & Kehlet, H. Secondary hyperalgesia to heat stimuli after burn injury in man. *Pain* **76**, 377–84 (1998).

160. Brull, S. J., Atanassoff, P. G., Silverman, D. G., Zhang, J. & LaMotte, R. H. Attenuation of experimental pruritus and mechanically evoked dysesthesiae in an area of cutaneous allodynia. *Somatosens. Mot. Res.* **16**, 299–303 (1999).
161. Patapoutian, A., Peier, A. M., Story, G. M. & Viswanath, V. Sensory systems: ThermoTRP channels and beyond: mechanisms of temperature sensation. *Nat. Rev. Neurosci.* **4**, 529–539 (2003).
162. Imamachi, N. *et al.* TRPV1-expressing primary afferents generate behavioral responses to pruritogens via multiple mechanisms. *Proc. Natl. Acad. Sci.* **106**, 11330–11335 (2009).
163. Jian, T. *et al.* TRPV1 and PLC Participate in Histamine H4 Receptor-Induced Itch. *Neural Plast.* **2016**, 1–9 (2016).
164. Shim, W.-S. *et al.* TRPV1 mediates histamine-induced itching via the activation of phospholipase A2 and 12-lipoxygenase. *J. Neurosci.* **27**, 2331–7 (2007).
165. Min, H. *et al.* TLR4 enhances histamine-mediated pruritus by potentiating TRPV1 activity. *Mol. Brain* **7**, 59 (2014).
166. Gibson, R. A. *et al.* A randomised trial evaluating the effects of the TRPV1 antagonist SB705498 on pruritus induced by histamine, and cowhage challenge in healthy volunteers. *PLoS One* **9**, e100610 (2014).
167. Davidson, S. & Giesler, G. J. The multiple pathways for itch and their interactions with pain. *Trends Neurosci.* **33**, 550–558 (2010).
168. Wilson, S. R. *et al.* TRPA1 is required for histamine-independent, Mas-related G protein-coupled receptor-mediated itch. *Nat. Neurosci.* **14**, 595–603 (2011).
169. Wilson, S. R. *et al.* The ion channel TRPA1 is required for chronic itch. *J. Neurosci.* **33**, 9283–9294 (2013).
170. Bautista, D. M. *et al.* TRPA1 Mediates the Inflammatory Actions of Environmental Irritants and Proalgesic Agents. *Cell* **124**, 1269–1282 (2006).
171. Hoffmann, T. *et al.* TRPA1 and TRPV1 are differentially involved in heat nociception of mice. *Eur. J. Pain* **17**, 1472–1482 (2013).
172. Le Bars, D., Dickenson, A. H. & Besson, J.-M. Diffuse noxious



- inhibitory controls (DNIC). I. Effects on dorsal horn convergent neurones in the rat. *Pain* **6**, 283–304 (1979).
173. Yarnitsky, D. *et al.* Recommendations on terminology and practice of psychophysical DNIC testing. *Eur. J. Pain* **14**, 339 (2010).
  174. Kennedy, D. L., Kemp, H. I., Ridout, D., Yarnitsky, D. & Rice, A. S. C. Reliability of conditioned pain modulation: A systematic review. *Pain* **157**, 2410–2419 (2016).
  175. Yarnitsky, D. Conditioned pain modulation (the diffuse noxious inhibitory control-like effect): its relevance for acute and chronic pain states. *Curr. Opin. Anaesthesiol.* **23**, 611–615 (2010).
  176. Youssef, A. M., Macefield, V. G. & Henderson, L. A. Cortical influences on brainstem circuitry responsible for conditioned pain modulation in humans. *Hum. Brain Mapp.* **37**, 2630–2644 (2016).
  177. Youssef, A. M., Macefield, V. G. & Henderson, L. A. Pain inhibits pain; human brainstem mechanisms. *Neuroimage* **124**, 54–62 (2016).
  178. Ramaswamy, S. & Wodehouse, T. Conditioned pain modulation—A comprehensive review. *Neurophysiologie Clinique* **51**, 197–208 (2021).
  179. Andersen, H. H., van Laarhoven, A. I. M., Elberling, J. & Arendt-Nielsen, L. Modulation of Itch by Conditioning Itch and Pain Stimulation in Healthy Humans. *J. Pain* **18**, 1437–1450 (2017).
  180. van Laarhoven, A. I. M., Kraaimaat, F. W., Wilder-Smith, O. H., van de Kerkhof, P. C. M. & Evers, A. W. M. Heterotopic pruritic conditioning and itch--analogous to DNIC in pain? *Pain* **149**, 332–7 (2010).
  181. Snyder, L. M. & Ross, S. E. Itch and its inhibition by counter stimuli. *Handb. Exp. Pharmacol.* **226**, 191–206 (2015).
  182. Bickford, R. Experiments relating to the itch sensation, its peripheral mechanism, and central pathways. *Clin Sci* **3**, 377–386 (1938).
  183. Andersen, H. H. *et al.* Alloknese and hyperknese—mechanisms, assessment methodology, and clinical implications of itch sensitization. *Pain* **159**, 1185–1197 (2018).
  184. LaMotte, R. H. Subpopulations of 'nocifensor neurons' contributing to pain and allodynia, itch and alloknese. *APS J.* **1**, 115–126 (1992).

185. Simone, D. A., Alreja, M. & Lamotte, R. H. Psychophysical studies of the itch sensation and itchy skin ('alloknesis' produced by intracutaneous injection of histamine. *Somatosens. Mot. Res.* **8**, 271–279 (1991).
186. Goon, A. T. J., Yosipovitch, G., Chan, Y. H. & Goh, C. L. Clinical characteristics of generalized idiopathic pruritus in patients from a tertiary referral center in Singapore. *Int. J. Dermatol.* **46**, 1023–1026 (2007).
187. Darsow, U. *et al.* New aspects of itch pathophysiology: component analysis of atopic itch using the 'Eppendorf Itch Questionnaire'. *Int. Arch. Allergy Immunol.* **124**, 326–31 (2001).
188. Dawn, A. *et al.* Itch characteristics in atopic dermatitis: Results of a web-based questionnaire. *Br. J. Dermatol.* **160**, 642–644 (2009).
189. Andersen, H. H., Lo Vecchio, S., Elberling, J., Yosipovitch, G. & Arendt-Nielsen, L. UVB- and NGF-induced cutaneous sensitization in humans selectively augments cowhage- and histamine-induced pain and evokes mechanical hyperknesis. *Exp. Dermatol.* **27**, 258–267 (2018).
190. Andersen, H. H., Marker, J. B., Hoeck, E. A., Elberling, J. & Arendt-Nielsen, L. Antipruritic effect of pretreatment with topical capsaicin 8% on histamine- and cowhage-evoked itch in healthy volunteers: a randomized, vehicle-controlled, proof-of-concept trial. *Br. J. Dermatol.* **177**, 107–116 (2017).
191. Dai, Y. *et al.* Sensitization of TRPA1 by PAR2 contributes to the sensation of inflammatory pain. *J. Clin. Invest.* **117**, 1979–1987 (2007).
192. Vergnolle, N., Hollenberg, M. D., Sharkey, K. A. & Wallace, J. L. Characterization of the inflammatory response to proteinase-activated receptor-2 (PAR2)-activating peptides in the rat paw. *Br. J. Pharmacol.* **127**, 1083–1090 (1999).
193. Vergnolle, N., Wallace, J. L., Bunnett, N. W. & Hollenberg, M. D. Protease-activated receptors in inflammation, neuronal signaling and pain. *Trends in Pharmacological Sciences* **22**, 146–152 (2001).
194. Seeliger, S. *et al.* Proinflammatory role of proteinase-activated receptor-2 in humans and mice during cutaneous inflammation in vivo. *FASEB J.* **17**, 1871–1885 (2003).
195. Saifeddine, M., Al-ani, B., Cheng, C. H., Wang, L. & Hollenberg, M. D.

- Rat proteinase-activated receptor-2 (PAR-2): cDNA sequence and activity of receptor-derived peptides in gastric and vascular tissue. *Br. J. Pharmacol.* **118**, 521–530 (1996).
196. Davidson S, Moser H, G. G. Itch Mechanisms and Treatments. in *Itch Mechanisms and Treatment* (CRC Press, Taylor & Francis group, 2014).
197. Freitag, G. & Höppner, T. Results of a postmarketing drug monitoring survey with a polidocanolurea preparation for dry, itching skin. *Curr. Med. Res. Opin.* **13**, 529–537 (1997).
198. Young, T. A. *et al.* A pramoxine-based anti-itch lotion is more effective than a control lotion for the treatment of uremic pruritus in adult hemodialysis patients. *J. Dermatolog. Treat.* **20**, 76–81 (2009).
199. Sisk, A. L. Vasoconstrictors in local anesthesia for dentistry. *Anesthesia Progress* **39**, 187–193 (1992).
200. Atanassoff, P. G., Brull, S. J., Printsev, Y. & Silverman, D. G. The effect of intradermal administration of lidocaine and morphine on the response to thermal stimulation. *Anesth. Analg.* **84**, 1340–1343 (1997).
201. Müller, C., Großjohann, B. & Fischer, L. The use of concentrated heat after insect bites/ stings as an alternative to reduce swelling, pain, and pruritus: An open cohort-study at German beaches and bathing-lakes. *Clin. Cosmet. Investig. Dermatol.* **4**, 191–196 (2015).
202. Papoiu, A. D. & Yosipovitch, G. Topical capsaicin. the fire of a ‘hot’ medicine is reignited. *Expert Opinion on Pharmacotherapy* **11**, 1359–1371 (2010).
203. Andersen, H. H., Sand, C. & Elberling, J. Considerable variability in the efficacy of 8% capsaicin topical patches in the treatment of chronic pruritus in 3 patients with notalgia paresthetica. *Ann. Dermatol.* **28**, 86–89 (2016).
204. Zeidler, C. *et al.* Capsaicin 8% cutaneous patch: A promising treatment for brachioradial pruritus? *British Journal of Dermatology* **172**, 1669–1671 (2015).
205. Andersen, H. H., Arendt-Nielsen, L. & Elberling, J. Topical capsaicin 8% for the treatment of neuropathic itch conditions. *Clin. Exp. Dermatol.* **42**, 596–598 (2017).
206. Misery, L. *et al.* Successful treatment of refractory neuropathic pruritus

with capsaicin 8% patch: A bicentric retrospective study with long-term follow-up. *Acta Dermato-Venereologica* **95**, 864–865 (2015).

207. Chung, M. K., Lee, H. & Caterina, M. J. Warm temperatures activate TRPV4 in mouse 308 keratinocytes. *J. Biol. Chem.* **278**, 32037–32046 (2003).
208. Moore, C. *et al.* UVB radiation generates sunburn pain and affects skin by activating epidermal TRPV4 ion channels and triggering endothelin-1 signaling. *Proc. Natl. Acad. Sci. U. S. A.* **110**, (2013).



ISSN (online): 2246-1302  
ISBN (online): 978-87-7573-806-9

**AALBORG UNIVERSITY PRESS**