

ANNALES DE L'INSTITUT FOURIER

DMITRI I. PANYUSHEV

On spherical nilpotent orbits and beyond

Annales de l'institut Fourier, tome 49, n° 5 (1999), p. 1453-1476

http://www.numdam.org/item?id=AIF_1999__49_5_1453_0

© Annales de l'institut Fourier, 1999, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON SPHERICAL NILPOTENT ORBITS AND BEYOND

by Dmitri I. PANYUSHEV

Introduction.

Let \mathfrak{g} be a semisimple Lie algebra and G its adjoint group. The ground field \mathbf{k} is algebraically closed and of characteristic zero. We continue investigations started in [Pa94], which are primarily concerned with the complexity of nilpotent G -orbits (conjugacy classes) in \mathfrak{g} . Let $\mathcal{N} \subset \mathfrak{g}$ be the nilpotent cone and $\mathcal{O} \subset \mathcal{N}$ an orbit. We gave in [Pa94] a formula for the complexity of nilpotent orbits and proved that \mathcal{O} is spherical (i.e., of complexity 0) if and only if $\text{ht}(\mathcal{O}) \leq 3$. Here $\text{ht}(\mathcal{O})$ is the height of \mathcal{O} , which can be defined as $\max\{n \in \mathbb{N} \mid (\text{ad } e)^n \neq 0, e \in \mathcal{O}\}$. In this article we give yet another characterization of spherical nilpotent orbits in terms of minimal Levi subalgebras intersecting them, see (3.2). This yields a kind of canonical form for such orbits, see (3.4):

an orbit $\mathcal{O} \subset \mathcal{N}$ is spherical if and only if it contains a representative which is a sum of root vectors corresponding to orthogonal simple roots.

Along the way, in Sections 2 and 3, we prove several auxiliary results about the height and the type of \mathcal{O} . The minimal non-spherical orbits in the simple Lie algebras are described in Section 4. These are of complexity 1 for SL_N and of complexity 2 for all other simple groups. In Section 5, we study the complexity of nilpotent orbits for Vinberg's θ -groups. Recall that associated with a finite order automorphism θ of \mathfrak{g} , one has the periodic

Keywords: Semisimple Lie algebra – Nilpotent orbit – Spherical variety.
Math classification: 14L30 – 14M17 – 17B45.

grading $\mathfrak{g} = \bigoplus_{j \in \mathbb{Z}_m} \mathfrak{g}_j$ and the connected reductive group G_0 acting linearly on \mathfrak{g}_1 . In this situation we are interested in the complexity of G_0 -orbits in $\mathcal{N} \cap \mathfrak{g}_1$. Our main results are:

- a monotonicity result for the complexity of Ge and G_0e ($e \in \mathcal{N} \cap \mathfrak{g}_1$), see (5.1);
- a formula for the complexity of G_0e in terms of a bi-grading of \mathfrak{g} , see (5.4);
- in case θ is of order 2, an almost complete description of spherical G_0 -orbits is found, see (5.7).

The situation for θ -groups is not however so simple, as it could have been: By [Vi76], the irreducible components of $Ge \cap \mathfrak{g}_1$ are just G_0 -orbits. If θ is of order 2, these components have the same dimension [KR71]. But it may happen that these have different complexity, see (5.10). Finally, Section 6 is a collection of observations and questions related to spherical nilpotent orbits. In particular, we show that theory of spherical orbits has some relationship with the index of Borel subalgebras.

As usual, algebraic groups are denoted by capital Roman letters, and their Lie algebras by the corresponding small Gothic letters. For $x \in \mathfrak{g}$, we write Gx in place of $(\text{ad } G)x$.

Acknowledgements. I would like to thank A.G. Elashvili and E.B. Vinberg for illuminating conversations about nilpotent orbits. Thanks are also due to D. Shmel'kin for questions reviving my interest to the subject and to the referee for several helpful suggestions. This research was carried out while I was visiting University of Poitiers and MPI (Bonn). I am grateful to both institutions for hospitality and support. This research was supported in part by RFFI Grant No. 98-01-00598.

1. Recollections on nilpotent orbits and the complexity.

Let \mathfrak{g} be a semisimple Lie algebra with a fixed triangular decomposition $\mathfrak{g} = \mathfrak{u}_- \oplus \mathfrak{t} \oplus \mathfrak{u}_+$, Δ the corresponding root system, and $\Pi = \{\alpha_1, \dots, \alpha_p\}$ the set of simple roots. Let $\mathcal{N} \subset \mathfrak{g}$ be the nilpotent cone. By the Morozov-Jacobson theorem, any nonzero element $e \in \mathcal{N}$ can be included in an \mathfrak{sl}_2 -triple $\{e, h, f\}$ (i.e., $[e, f] = h$, $[h, e] = 2e$, $[h, f] = -2f$). The semisimple

element h defines the \mathbb{Z} -grading in \mathfrak{g} :

$$\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{g}(i),$$

where $\mathfrak{g}(i) = \{x \in \mathfrak{g} \mid [h, x] = ix\}$. It is well known that all \mathfrak{sl}_2 -triples containing e are G_e -conjugate. Therefore the structure of this \mathbb{Z} -grading does not depend on a particular choice of h .

Following E.B. Dynkin, we shall say that h is a *characteristic* of e . The orbit Gh contains a unique element h_+ such that $h_+ \in \mathfrak{t}$ and $\alpha(h_+) \geq 0$ for all $\alpha \in \Pi$. The Dynkin diagram of \mathfrak{g} equipped with the numerical labels $\alpha_i(h_+)$, $\alpha_i \in \Pi$, at the corresponding nodes is called the *weighted Dynkin diagram* of e . After Dynkin and Kostant, it is known (see e.g. [SpSt]) that

(a) $\alpha_i(h_+) \in \{0, 1, 2\}$;

(b) elements $e, e' \in \mathcal{N}$ are G -conjugate if and only if their characteristics h and h' are G -conjugate if and only if their weighted Dynkin diagrams coincide.

We shall need the following standard results on the structure of the stabilizer $G_e \subset G$ and the centralizer $\mathfrak{g}_e \subset \mathfrak{g}$ (see [SpSt, ch. III]).

1.1. PROPOSITION. — (i) *The Lie algebra \mathfrak{g}_e (resp. \mathfrak{g}_f) is positively (resp. negatively) graded; $\mathfrak{g}_e = \bigoplus_{i \geq 0} \mathfrak{g}_e(i)$, where $\mathfrak{g}_e(i) = \mathfrak{g}_e \cap \mathfrak{g}(i)$, and likewise for \mathfrak{g}_f ;*

(ii) *Let $L := G(0)$ be the connected subgroup in G corresponding to $\mathfrak{g}(0)$ and $K := L_e^{(1)}$. Then $K = G_e \cap G_f$ and it is a maximal reductive subgroup in both G_e and G_f ;*

(iii) *For any i , there are K -stable decompositions:*

$$\mathfrak{g}(i) = \mathfrak{g}_e(i) \oplus [f, \mathfrak{g}(i+2)], \quad \mathfrak{g}(i) = \mathfrak{g}_f(i) \oplus [e, \mathfrak{g}(i-2)].$$

In particular, $\text{ad } f : \mathfrak{g}(i) \rightarrow \mathfrak{g}(i-2)$ is injective when $i \geq 1$ and surjective when $i \leq 1$;

(iv) *$(\text{ad } f)^i : \mathfrak{g}(i) \rightarrow \mathfrak{g}(-i)$ is one-to-one.*

The notation related to the \mathbb{Z} -grading associated with a nilpotent orbit will be used throughout the paper.

1.2. PROPOSITION. — *For i even (resp. odd), $\mathfrak{g}(i)$ is an orthogonal (resp. symplectic) K -module. In particular, $\dim \mathfrak{g}(i)$ is even for i odd.*

(1) K can be disconnected.

Proof. — For $i \geq 0$, consider the bilinear form Ψ_i on $\mathfrak{g}(i)$ given by $(x, y) \mapsto \langle (\text{ad } f)^i x, y \rangle$, where $\langle \cdot, \cdot \rangle$ is a G -invariant inner product on \mathfrak{g} . By Proposition 1.1(ii),(iv), Ψ_i is nondegenerate and K -invariant. It follows from G -invariance of $\langle \cdot, \cdot \rangle$ that Ψ_i is symmetric for i even and alternate for i odd. □

Recall that e or Ge is called

- even whenever $\mathfrak{g}(i) = 0$ for i odd or, equivalently, if all $\alpha_i(h_+) \in \{0, 2\}$;
- distinguished, if \mathfrak{g}_e contains no semisimple elements, i.e., K is finite.

For a reductive group R , we let B_R denote a Borel subgroup of R . If X is an irreducible R -variety, then X is called (R -)spherical whenever B_R has an open orbit in X . The complexity of X relative to R , which is denoted by $c_R(X)$, is equal to the minimal codimension of B_R -orbits in X . Clearly, $c_R(X) = c_{R^o}(X)$, where R^o stands for the identity component of R .

2. The height of a nilpotent orbit.

DEFINITION. — The integer $\max\{i \mid \mathfrak{g}(i) \neq 0\}$ is called the height of e or the orbit $\mathcal{O} = Ge$ and is denoted by $\text{ht}(e)$ or $\text{ht}(\mathcal{O})$.

Since $e \in \mathfrak{g}(2)$, we have $\text{ht}(e) \geq 2$ for any $e \in \mathcal{N} \setminus \{0\}$. Let $\Lambda \in \Delta_+$ be the highest root, $\Lambda = \sum_{i=1}^p n_i \alpha_i$. Clearly, we then have

$$(2.1) \quad \text{ht}(e) = \Lambda(h_+) = \sum_{i=1}^p \alpha_i(h_+) n_i.$$

An immediate consequence of (1.1) is an intrinsic characterization of the height

$$(2.2) \quad \text{ht}(e) = \max\{n \in \mathbb{N} \mid (\text{ad } e)^n \neq 0\}.$$

For the classical Lie algebras $\mathfrak{sl}(V)$, $\mathfrak{sp}(V)$, and $\mathfrak{so}(V)$, it is sometimes more convenient to describe nilpotent orbits by the sizes of blocks in the Jordan normal form, i.e., in terms of partitions (a_1, \dots, a_t) , where $a_1 \geq a_2 \geq \dots \geq a_t$ and $\sum_{i=1}^t a_i = \dim V$. As is well known, this correspondence is one-to-one in case of $\mathfrak{sl}(V)$. For $\mathfrak{so}(V)$ and $\mathfrak{sp}(V)$, there is a correspondence

between the nilpotent orbits and partitions satisfying a special condition. That is, in the symplectic (resp. orthogonal) case, one considers the partitions whose *odd* (resp. *even*) parts occur pairwise. This correspondence turns out to be a bijection, the only exception being that for $\mathfrak{so}(V)$ with $\dim V \equiv 0 \pmod{4}$, each partition whose all parts are even (“a very even partition”) arises from two $SO(V)$ -orbits, see [CM93, 5.1]. Since these two $SO(V)$ -orbits form a single $O(V)$ -orbit, these have the same height and complexity. In the sequel, we shall identify “classical” nilpotent orbits with corresponding partitions, keeping in mind this exception.

Let us give simple formulas for the height of nilpotent orbits in the classical Lie algebras.

2.3. THEOREM. — *Let $\mathcal{O} = (a_1, \dots, a_t)$ be a nilpotent orbit in a classical Lie algebra \mathfrak{g} ($a_1 \geq a_2 \geq \dots \geq a_t$).*

1. *If $\mathfrak{g} = \mathfrak{sl}(V)$ or $\mathfrak{sp}(V)$, then $\text{ht}(\mathcal{O}) = 2(a_1 - 1)$;*
2. *If $\mathfrak{g} = \mathfrak{so}(V)$, then $\text{ht}(\mathcal{O}) = \begin{cases} a_1 + a_2 - 2, & \text{if } a_2 \geq a_1 - 1 \\ 2a_1 - 4, & \text{if } a_2 \leq a_1 - 2. \end{cases}$*

In particular, either $\text{ht}(\mathcal{O})$ is even or $\text{ht}(\mathcal{O}) \equiv 3 \pmod{4}$.

Proof. — Let $\mathfrak{a} \subset \mathfrak{g}$ be a simple 3-dimensional subalgebra containing $e \in \mathcal{O}$. Denote by $R(n)$ a simple \mathfrak{a} -module of dimension $n + 1$. Considering \mathfrak{g} as an \mathfrak{a} -module, say $\mathfrak{g} = \bigoplus_i R(n_i)$, one sees that $\text{ht}(e) = \max\{n_i\}$. On the other hand, $V = \bigoplus_{i=1}^t R(a_i - 1)$ as \mathfrak{a} -module. The relationship between V and the adjoint representation is well known:

$$\mathfrak{g} = \begin{cases} V \otimes V^* \oplus \mathbf{I} & \text{for } \mathfrak{sl}(V) \\ S^2V & \text{for } \mathfrak{sp}(V) \\ \wedge^2V & \text{for } \mathfrak{so}(V). \end{cases}$$

Combining these relations with the Clebsch–Gordan formula $R(n) \otimes R(m) = R(n + m) \oplus R(n - 1) \otimes R(m - 1)$ and with the decomposition of $S^2R(n_i)$ and $\wedge^2R(n_i)$, one easily detects the biggest \mathfrak{a} -submodule in \mathfrak{g} . Whence the formulas for the height. In the orthogonal case, the constraint on parity must be satisfied. That is, the equality $a_2 = a_1 - 1$ is only possible, if a_1 is odd. □

Remark. — The above relationship between the adjoint and the simplest representations of classical algebras was used in [El85] for obtaining a quick classification of distinguished nilpotent orbits.

The orbits with odd height, in all simple Lie algebras, are not numerous and my feeling is that these ought to have some interesting properties. The following is a simple observation for them:

2.4. PROPOSITION. — *Suppose $\text{ht}(e)$ is odd. Then the weighted Dynkin diagram of e contains no 2's.*

Proof. — By (2.3), such elements e do not exist in $\mathfrak{sl}(V)$ and $\mathfrak{sp}(V)$. For $\mathfrak{so}(V)$, we must then have $a_2 = a_1 - 1$. Then a formula for the weighted Dynkin diagram (see [SpSt, 2.32] or [CM93, 5.3]) shows that $\alpha_i(h_+) \in \{0, 1\}$. In the exceptional cases, one can consult Dynkin's tables [Dy52, Tables 16–20] of the weighted Dynkin diagrams (see also [El75] or [CM93, ch. 8]). \square

The \mathbb{Z} -gradings associated with \mathfrak{sl}_2 -triples form only a small part among all possible \mathbb{Z} -gradings. Many interesting features of the former were described in [Ka80]. The following assertion concerns the same subject.

2.5. PROPOSITION. — *Suppose that $\mathfrak{g}(2n) = 0$ for the \mathbb{Z} -grading associated with a nilpotent element e . Then $\text{ht}(e) \leq 2n - 1$.*

Proof. — It suffices to prove that $\mathfrak{g}(2n + 1) = 0$. Assume not. Then $\mathfrak{g}(2n - 1) \neq 0$ as well. The space $\mathfrak{g}(2n + 1)$ is a sum of root spaces. Because each positive root is a sum of simple roots and $\mathfrak{g}(2n) = 0$, to reach $\mathfrak{g}(2n + 1)$ from $\mathfrak{g}(2n - 1)$, we must have a root vector $e_{\alpha_i} \in \mathfrak{g}(2)$ for some $\alpha_i \in \Pi$. Thus, the weighted Dynkin diagram must contain a label “2”. Then (2.4) says that $\text{ht}(e)$ is even, which is not the case, if $\mathfrak{g}(2n + 1) \neq 0$. \square

A relationship between the complexity and the height of nilpotent orbits is given by the following theorem proved in [Pa94]:

2.6. THEOREM. — *A nilpotent orbit Ge is spherical if and only if $\text{ht}(e) \leq 3$.* \square

3. The type of a nilpotent orbit and sphericity.

In this section we characterize the spherical nilpotent orbits in terms of minimal Levi subalgebras intersecting them. For any $e \in \mathcal{N}$, there exists a unique, up to conjugation, minimal Levi subalgebra \mathfrak{z} intersecting Ge and, moreover, the orbit $Ze \subset \mathfrak{z}' := [\mathfrak{z}, \mathfrak{z}]$ is distinguished. This fact is usually attributed to Bala and Carter [BC76]. Not everybody has observed

that a much more general assertion, in the context of graded Lie algebras, has independently been proved by E.B. Vinberg (see [Vi75] and [Vi79]). The construction itself is quite simple. Let \mathfrak{h} be a Cartan subalgebra in $\mathfrak{k} = \mathfrak{g}_e(0)$. Then the centralizer $\mathfrak{z} := \mathfrak{z}_{\mathfrak{g}}(\mathfrak{h})$ is the desired Levi subalgebra. Put $\mathfrak{q} := \mathfrak{z}_{\mathfrak{g}}(\mathfrak{h})'$. Obviously, the elements e , h , and f lie in \mathfrak{z} and hence in \mathfrak{q} . We also have $\mathfrak{z} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{z}(i)$ and $\mathfrak{q} = \bigoplus_{i \in \mathbb{Z}} \mathfrak{q}(i)$.

- 3.1. LEMMA.** — 1. $\mathfrak{q}(i) = \mathfrak{z}(i)$ for $i \neq 0$;
 2. $\mathfrak{q}(i) = 0$ whenever i is odd.

Proof. — 1. Since \mathfrak{z} contains h , the centre of \mathfrak{z} is contained in $\mathfrak{g}(0) \cap \mathfrak{z} = \mathfrak{z}(0)$.

2. Obviously, $\mathfrak{q}(0)_e = \mathfrak{q}_e(0) = 0$, i.e., e is distinguished in \mathfrak{q} . By a result of Bala-Carter [BC76] and Vinberg [Vi79], any distinguished nilpotent element is even⁽²⁾. □

The Cartan label of the semisimple subalgebra $\mathfrak{q} \subset \mathfrak{g}$ is called the *type* of Ge . Indication of the type forms a part of the notation for the nilpotent orbits in the exceptional Lie algebras used in [BC76]⁽³⁾. In case of two root lengths, if a simple component of \mathfrak{q} involves only short roots, then one places tilde over its Cartan label. The type does not determine the orbit uniquely. For instance, if e is distinguished in \mathfrak{g} , then $\mathfrak{h} = \{0\}$ and $\mathfrak{q} = \mathfrak{g}$. In order to distinguish different distinguished orbits and different conjugacy classes of Levi subalgebras, the Cartan label is accompanied by additional symbols, see [BC76] or [CM93, 8.4] for more details.

The notion of type applies to the classical Lie algebras as well and it is worth to write down explicit formulas for the type of a nilpotent orbit in this case. The partitions corresponding to the distinguished orbits was pointed out in [Vi75], but the general formulas, though being known to experts, seem not to be in print.

Let $\mathcal{O} = (a_1, \dots, a_t)$ be a ‘classical’ nilpotent orbit.

For $SL(V)$: By the theory of Jordan normal form, we have $\mathfrak{q} = \mathbf{A}_{a_1-1} + \dots + \mathbf{A}_{a_t-1}$.

For $Sp(V)$: If all the a_i ’s are distinct and even, then e is distinguished and $\mathfrak{q} = \mathbf{C}_n$, where $n = (\dim V)/2$. In general, each pair of equal parts

⁽²⁾ Both proofs were case-by-case. An elegant *a priori* proof was found by Jantzen, see [Ka80, Note added in proof].

⁽³⁾ Actually, this is a truncation of the data that were already used by E.B. Dynkin in [Dy52].

(equal Jordan blocks) $a_i = a_{i+1}$ gives rise to a summand $\tilde{\mathbf{A}}_{a_i-1}$. After deleting all equal pairs, we obtain a partition with distinct even parts. This little partition determines the last summand in \mathfrak{q} , a smaller symplectic algebra. Because there are two root lengths, one has to distinguish between \mathbf{A}_1 and $\tilde{\mathbf{A}}_1$. The answer is that \mathbf{A}_1 occurs if and only if one obtains at the very end the partition (2). That is, formally $\mathbf{C}_1 = \mathbf{A}_1$.

For $SO(V)$: The procedure is similar. Each pair of equal Jordan blocks gives rise to a summand \mathbf{A}_{a_i-1} in \mathfrak{q} . After deleting all equal pairs, we obtain a partition with distinct odd parts. This partition determines the last summand in \mathfrak{q} , a smaller orthogonal algebra. We have again to distinguish between \mathbf{A}_1 and $\tilde{\mathbf{A}}_1$. The answer is that $\tilde{\mathbf{A}}_1$ occurs if and only if one obtains at the very end the partition (3). That is, formally $\mathbf{B}_1 = \tilde{\mathbf{A}}_1$.

Modulo the description of distinguished orbits, the proof immediately amounts to the claim that the orbit(s) corresponding to the partition (n, n) is of type \mathbf{A}_{n-1} in \mathfrak{so}_{2n} and of type $\tilde{\mathbf{A}}_{n-1}$ in \mathfrak{sp}_{2n} .

- Examples. 1. $\mathcal{O} = (4, 4, 4, 3, 3, 1, 1) \in \mathfrak{sp}_{20}$. Then $\mathfrak{q} = \tilde{\mathbf{A}}_3 + \tilde{\mathbf{A}}_2 + \mathbf{C}_2$.
- 2. $\mathcal{O} = (3, 3, 3, 2, 2, 2, 2) \in \mathfrak{so}_{17}$. Then $\mathfrak{q} = \mathbf{A}_2 + 2\mathbf{A}_1 + \tilde{\mathbf{A}}_1$.

3.2. THEOREM. — *Let $e \in \mathcal{N} \setminus \{0\}$. The following conditions are equivalent:*

- 1. $\text{ht}(e) \leq 3$;
- 2. $\mathfrak{g}(4) = 0$;
- 3. *the type of Ge is $r\mathbf{A}_1 + l\tilde{\mathbf{A}}_1$;*
- 4. *there exist pairwise orthogonal simple roots β_1, \dots, β_t such that Ge contains an element of the form $\sum_{i=1}^t e_i$, where $e_i \in \mathfrak{g}_{\beta_i} \setminus \{0\}$. (Then $t = r + l$ and there are r long and l short roots among the β_i 's.)*

Proof. — $1 \Rightarrow 2$ — Obvious.

$2 \Rightarrow 3$. It follows from the definition of \mathfrak{q} and (3.1) that $\mathfrak{q} = \mathfrak{q}(-2) \oplus \mathfrak{q}(0) \oplus \mathfrak{q}(2)$, $e \in \mathfrak{q}(2)$, and $h \in \mathfrak{q}(0)$. Since $\mathfrak{q}_e(0) = \{0\}$, we have $\dim \mathfrak{q}(0) = \dim \mathfrak{q}(2)$. On the other hand, $\mathfrak{q}(2)$ is a spherical $Q(0)$ -module by [Pa94, 3.2]. Hence $Q(0)$ is a torus. Clearly, a semisimple Lie algebra having such a grading is just a sum of several 3-dimensional simple algebras.

$3 \Leftrightarrow 4$. This follows at once from the definition of the type.

$4 \Rightarrow 1$. Consider the Levi subalgebra \mathfrak{f} corresponding to the roots β_1, \dots, β_t and the corresponding connected subgroup $F \subset G$. We may

assume that $e = \sum_{i=1}^t e_i$ and $\mathfrak{a} = \langle e, h, f \rangle$ is embedded diagonally in $\mathfrak{f}' \simeq (\mathfrak{sl}_2)^t$. Take the unique F -stable decomposition $\mathfrak{g} = \mathfrak{f} \oplus \mathfrak{m}$ and consider an arbitrary irreducible F -submodule $V \subset \mathfrak{m}$. I claim that F has finitely many orbits in V . Indeed, as F is a Levi subgroup of G , there is a \mathbb{Z}^{p-t} -grading of \mathfrak{g} whose ‘zero’-part is \mathfrak{f} . (Recall that $p = \text{rk } \mathfrak{g}$.) The F -module V is contained in some homogeneous subspace of this polygrading (it will actually be equal to some homogeneous subspace, but we do not need this fact). Now, finiteness follows by famous Vinberg’s lemma, see [Vi76, Lemma in §2]. Let s be the number of simple factors of F acting non-trivially on V . Then $V \simeq R(d_1) \otimes \dots \otimes R(d_s)$, where all $d_i \geq 1$. Since F has a dense orbit in V , we have $\dim((SL_2)^s \times \mathbf{k}^*) \geq \dim V$, i.e. $3s+1 \geq (d_1+1) \dots (d_s+1) \geq 2^s$. This inequality has exactly three solutions:

$$\begin{aligned} s = 3, \quad d_1 = d_2 = d_3 = 1; \\ s = 2, \quad d_1 \leq 2, d_2 = 1; \\ s = 1, \quad d_1 \leq 3. \end{aligned}$$

In each case we have $\sum_i d_i \leq 3$. Together with the Clebsch–Gordan formula, this shows that the biggest irreducible \mathfrak{a} -module that can occur in V is $R(3)$. Since $\mathfrak{f}|_{\mathfrak{a}} \simeq tR(2) + (p-t)R(0)$, the same is true for the whole Lie algebra \mathfrak{g} . But, this means precisely that $\text{ht}(e) \leq 3$. □

Equivalence of conditions (1) and (2) is a particular case of Proposition 2.5. But the last proof does not appeal to case-by-case considerations as in 2.4.

3.3. COROLLARY (of the proof). — *For any $\mu \in \Delta$, let $\{\alpha_{i_1}, \dots, \alpha_{i_s}\} \subset \Pi$ be a set of pairwise orthogonal roots such that $\langle \alpha_{i_j}, \mu \rangle \neq 0, j = 1, \dots, s$. Then $s \leq 3$ (even $s \leq 2$, if an addition μ is long and one of the α_{i_j} ’s is short).*

Proof. — Replacing μ by a suitable root $\mu' = \mu + \sum_j n_j \alpha_{i_j}$, one can achieve that $\langle \alpha_{i_j}, \mu' \rangle > 0$ for all j . Then μ' is the highest weight of an irreducible $(SL_2)^s$ -submodule V of \mathfrak{g} . Since all copies of SL_2 act non-trivially on V , the proof of $4 \Rightarrow 1$ shows that $s \leq 3$ (even $s \leq 2$, if $d_1 \geq 2$). □

Remarks. — 1. In case $\mu \in \Pi$, we obtain the well-known assertion that the number of other simple roots that are not orthogonal to μ is at most 3. That is, we have given an invariant-theoretic proof of it.

2. If we drop the assumption that the α_{i_j} 's are pairwise orthogonal, then it is easy to give an example with $s = 5$ (at least).

Combining (2.6) and (3.2) yields a kind of “normal form” for spherical nilpotent orbits:

3.4. THEOREM. — *Suppose $e \in \mathcal{N}$. Then the orbit Ge is spherical if and only if it contains an element of the form $\sum_{i=1}^t e_i$, where $e_i \in \mathfrak{g}_{\beta_i}$ and β_1, \dots, β_t are pairwise orthogonal simple roots.* □

Obviously, this normal form is not unique in general. For instance, if Ge is the orbit of highest weight vectors, then $t = 1$ and β_1 can be any long simple root.

Examples. — 1. The spherical nilpotent orbits in \mathfrak{sp}_{2p} are $\mathcal{O}_i = (2^i, 1^{2p-2i})^{(4)}$ ($0 \leq i \leq p$). Then the type of \mathcal{O}_i is $\begin{cases} l\tilde{\mathbf{A}}_1, & \text{if } i = 2l, \\ l\tilde{\mathbf{A}}_1 + \mathbf{A}_1, & \text{if } i = 2l + 1. \end{cases}$

In particular for $i = p = 2l + 1$, the respective set of simple roots is $\alpha_1, \alpha_3, \dots, \alpha_{2l+1}$. (The last of them is long.)

2. The type of spherical orbit $(3, 2^2, 1^l)$ in $\mathfrak{g} = \mathfrak{so}(V)$ is $\begin{cases} 3\mathbf{A}_1, & \text{if } l \text{ is odd,} \\ \mathbf{A}_1 + \tilde{\mathbf{A}}_1, & \text{if } l \text{ is even.} \end{cases}$

4. Orbits of small complexity.

Recall, with some variations, a formula for the complexity of Ge obtained in [Pa94]. The following assertion which is implicit in [Pa94, 1.2] was suggested by the referee.

4.1. LEMMA. — *Let P be a parabolic subgroup of G and Y a P -variety. If L is a Levi subgroup of P , then $c_G(G *_P Y) = c_L(Y)$.*

Proof. — Consider the canonical projection $G *_P Y \rightarrow G/P$. Since B_G has a dense orbit in G/P , the minimal codimension of B_G -orbits in $G *_P Y$ is equal to the minimal codimension of $(B_G)_*$ -orbits in Y , where $(B_G)_*$ is the stabilizer of a point in the dense orbit in G/P . For a suitable choice of B_G , we obtain $(B_G)_* = B_G \cap P = B_L$. □

Maintain the notation of sect. 1. Let S be a *stabilizer in general position* (= s.g.p.) for the K -action on $\mathfrak{g}(2)$. We shall use the notation

(4) As usual in the theory of partitions, $a^j := a, \dots, a$ (j times).

$S = \text{s.g.p.}(K, \mathfrak{g}(2))$ as a shorthand. The reader is referred to [VP89, §7] for the basic facts on s.g.p. As $\mathfrak{g}(2)$ is an orthogonal K -module, a result of D. Luna [Lu72] asserts that S is reductive. Set $\mathfrak{g}(\geq j) = \bigoplus_{i \geq j} \mathfrak{g}(i)$.

4.2. THEOREM. — 1. $c_G(Ge) = c_L(\mathfrak{g}(2)) + c_S(\mathfrak{g}(\geq 3))$;

2. $c_G(Ge) = c_L(\mathfrak{g}(\geq 2))$.

Proof. — 1. Since $Le \simeq L/K$ is the dense orbit in $\mathfrak{g}(2)$, we have $c_L(\mathfrak{g}(2)) = c_L(L/K)$. Hence (1) is nothing but the first formula in [Pa94,2.3].

2. Let P be the parabolic subgroup corresponding to $\mathfrak{p} := \mathfrak{g}(\geq 0)$. Then L is a Levi subgroup of P . Since $G_e = P_e$ and $\overline{P_e} = \mathfrak{g}(\geq 2)$ (see 1.1), the homogeneous vector bundle $G *_P \mathfrak{g}(\geq 2)$ is birationally isomorphic to G_e . (Actually, the collapsing $G *_P \mathfrak{g}(\geq 2) \rightarrow G \cdot \mathfrak{g}(\geq 2) = \overline{G_e}$ is an equivariant resolution of $\overline{G_e}$.) Hence $c_G(Ge) = c_G(G *_P \mathfrak{g}(\geq 2))$. We conclude by Lemma 4.1. □

Remark. — The first formula in (4.2) is convenient for theoretical arguments, while the second one is sometimes better suited for practical computations. The significance of these formulas is that computing of the complexity of Ge is reduced to that for a representation space. In case of representations, there is an explicit algorithm for doing this [Pa87]. Actually, given a representation $R \rightarrow GL(V)$ of a reductive group R , the algorithm says how to find $\text{s.g.p.}(R, V \oplus V^*) =: R_*$. The group R_* is reductive and has some other nice properties. Then $c_R(V) = \dim V - \dim B_R + \dim B_{R_*}$.

4.3. PROPOSITION. — *If $\dim \mathfrak{g}(4) \geq 2$ or $\text{ht}(e) \geq 5$, then $c_G(Ge) \geq 2$.*

Proof. — In view of Theorem 4.2(1), it suffices to show that $c_S(\mathfrak{g}(\geq 3)) \geq 2$. We are to find at least two algebraically independent B_S -invariant rational functions on $\mathfrak{g}(\geq 3)$.

It follows from Theorem 3.2 that in both cases $\mathfrak{g}(4) \neq 0$. Since $\text{ad } f : \mathfrak{g}(4) \rightarrow \mathfrak{g}(2)$ is injective, there is a K -module W_1 such that $\mathfrak{g}(2) \simeq \mathfrak{g}(4) \oplus W_1$. Because $S = \text{s.g.p.}(K, \mathfrak{g}(2))$, we have $K(\mathfrak{g}(2)^S)$ is dense in $\mathfrak{g}(2)$. It follows that $\mathfrak{g}(4)^S \neq 0$. Write $\mathfrak{g}(4) = \mathfrak{g}(4)^S \oplus W_2$, where W_2 is an S -module. Because $\mathfrak{g}(4)$ is an orthogonal K -module (1.2), the same holds for W_2 . Hence $\mathbf{k}[\mathfrak{g}(4)]^S$ contains at least two algebraically independent functions whenever $\dim \mathfrak{g}(4) \geq 2$. If $\dim \mathfrak{g}(4) = 1$, we examine the following possibilities:

(a) Assume $\mathfrak{g}(6) \neq 0$. Then $\mathfrak{g}(6) = (\text{ad } e)\mathfrak{g}(4)$ and $\mathfrak{g}(6) = \mathfrak{g}(6)^S$. This yields another S -invariant function in $k[\mathfrak{g}(\geq 3)]$.

(b) Assume $\mathfrak{g}(5) \neq 0$. Then $\mathfrak{g}(5)$ and $[f, \mathfrak{g}(5)] \subset \mathfrak{g}(3)$ are two different isomorphic S -submodules in $\mathfrak{g}(\geq 3)$. Obviously, this produces a non-constant B_S -invariant rational function on $\mathfrak{g}(\geq 3)$. \square

Remark. — Similar arguments prove that if $\text{ht}(e) \geq 2n + 1$ ($n \geq 2$), then $c_G(Ge) \geq n$. But I think there ought to exist a quadratic polynomial $n \mapsto \phi(n)$ such that $c_G(Ge) \geq \phi(n)$.

By (2.6) and (4.3), the nilpotent orbits of complexity 1 are contained among those with height 4 and $\dim \mathfrak{g}(4) = 1$. Such orbits exist in all simple Lie algebras. However, routine computations lead to the following conclusion:

4.4. THEOREM. — 1. Nilpotent orbits of complexity 1 exist only for $G = SL_n$. For each $n \geq 3$ there exist a unique such orbit. Its weighted Dynkin diagram is $2-0-\dots-0-2$ and the partition is $(3, 1^{n-3})$. Moreover, this orbit is the unique minimal non-spherical one.

2. For all other simple groups, the minimal non-spherical orbits are of complexity 2.

Proof. — For the classical groups, the classification of the minimal non-spherical (= m.n.s.) orbits follows from (2.3), (2.6), and an explicit description, due to Gerstenhaber and Hesselink, of the closure ordering, see [CM93, 6.2]. For the exceptional groups, one uses 3.2(3) and the Hasse diagrams for the closure ordering, see [Spal, IV.2].

1. For SL_n , there is a unique m.n.s. orbit $\mathcal{O} = (3, 1^{n-3})$. It is even and of height 4. For $n = 3$, \mathcal{O} is the regular nilpotent orbit and the assertion follows from counting dimensions. Let $n \geq 4$. Then $L = SL_{n-2}(\mathbf{k}^*)^2$ and the L -modules $\mathfrak{g}(2)$ and $\mathfrak{g}(4)$ are as follows: $\mathfrak{g}(2) = R(\varphi_1) \otimes \varepsilon + R(\varphi_1)^* \otimes \mu$, $\mathfrak{g}(4) = R(0) \otimes \varepsilon \mu$. Here $R(\lambda)$ stands for the irreducible representation of the semisimple part with highest weight λ , while ε and μ are basic characters of the central torus; $\dim \mathfrak{g}(2) = 2n - 4$ and $\dim \mathfrak{g}(4) = 1$. Obviously, $\mathfrak{g}(2)$ is a spherical L -module. It is not hard to compute that $K \simeq SL_{n-3} \cdot \mathbf{k}^*$, $S \simeq SL_{n-4} \cdot \mathbf{k}^*$, and S acts trivially on $\mathfrak{g}(4)$. Applying then (4.2), we get $c_G(\mathcal{O}) = 1$.

The minimal nilpotent orbit lying 'over' \mathcal{O} is

$\mathcal{O}' = \begin{cases} (4), & \text{if } n = 4 \\ (3, 2, 1^{n-5}), & \text{if } n \geq 5 \end{cases}$. Then it already turns out that $c_G(\mathcal{O}') = 2$.

2. For SO_n ($n \geq 7$) and E_n ($n = 6, 7, 8$), the unique m.n.s. orbit is of type A_2 ; F_4 has two m.n.s. orbits of types A_2 and \tilde{A}_2 ; Sp_{2n} ($n \geq 2$) has the m.n.s. orbits of types \tilde{A}_2 (for $n \geq 3$) and C_2 . The corresponding partitions are $(3, 3, 1^{2n-6})$ and $(4, 1^{2n-4})$, respectively. Finally, the m.n.s. orbit for G_2 is 10-dimensional (and distinguished). All these orbits are of complexity 2.

For instance, consider the orbit of type \tilde{A}_2 for $G = F_4$. Here $L = Spin_7 \cdot k^*$, $\mathfrak{g}(2) = R(\varphi_3) \otimes \varepsilon$, and $\mathfrak{g}(4) = R(\varphi_1) \otimes \varepsilon^2$; $\dim \mathfrak{g}(2) = 8$, $\dim \mathfrak{g}(4) = 7$. This information is easily being extracted from the weighted Dynkin diagram given in Table 1 in Section 5. Because $\dim \mathfrak{g}(4) \geq 2$, Proposition 4.3 implies that $c_G(\mathcal{O}) \geq 2$. Let us find the exact value. Here $K \simeq G_2$ and $c_L(L/K) = 0$. Next, $\mathfrak{g}(2)$ affords the sum of the simplest (7-dimensional) and the trivial 1-dimensional representation of G_2 . Therefore $S \simeq SL_3$, the long root subgroup of G_2 . It is easily seen that $\mathfrak{g}(4)$ affords the simplest representation of G_2 and that $\mathfrak{g}(4)|_{SL_3} = R(\varphi'_1) + R(\varphi'_2) + R(0)$. Whence $c_S(\mathfrak{g}(4)) = 2$ and $c_G(\mathcal{O}) = 2$. □

This result confirms a claim in [Vi86, n. 9] concerning orbits of complexity 1 in the universe that “it appears there should be few of them”.

5. The complexity of nilpotent orbits of θ -groups.

Let θ be an automorphism of \mathfrak{g} , of finite order m . Fix a primitive m -th root of unity ζ . Consider the periodic grading

$$\mathfrak{g} = \bigoplus_{j \in \mathbb{Z}_m} \mathfrak{g}_j,$$

where \mathfrak{g}_j is the θ -eigenspace of \mathfrak{g} corresponding to ζ^j . Following Vinberg, we shall say that the connected reductive group G_0 acting linearly on \mathfrak{g}_1 is a θ -group. The main references on θ -groups are [Vi75], [Vi76], [Vi79]. One of the basic results is that $\overline{G_0 e} \ni 0$ ($e \in \mathfrak{g}_1$) if and only if $e \in \mathcal{N}$. Such G_0 -orbits are called nilpotent, too. Our aim is to study the complexity of them. Throughout this section, it is assumed that $e \in \mathcal{N} \cap \mathfrak{g}_1$. The first result is:

5.1. THEOREM. — $c_G(Ge) \geq c_{G_0}(G_0e)$.

To demonstrate the theorem, we need the following variation on Vinberg’s themes:

5.2. LEMMA. — *Let $G \rightarrow GL(V)$ be a representation of a reductive group, $H \subset G$ a reductive subgroup, and $W \subset V$ an H -stable subspace. Choose Borel subgroups in H and G such that $B_H \subset B_G$. Suppose that*

$$(*) \quad \mathfrak{b}_H v = \mathfrak{b}_G v \cap W \quad \text{for all } v \in W.$$

For any G -stable locally-closed subvariety $X \subset V$ and each irreducible component Y of $X \cap W$, we then have

$$c_G(X) \geq c_H(Y).$$

Proof. — By [Vi86], the complexity $c_G(X)$ is equal to the modality of B_G -action on X , i.e., to $\max_{X' \subset X} \text{trdeg } k(X')^{B_G}$, where X' runs through the irreducible B_G -stable subvarieties of X . On the other side, it follows from (*) and [Vi76, §2] that $B_G v \cap W$ is a union of finitely many B_H -orbits for all $v \in W$, each B_H -orbit being an irreducible component of $B_G v \cap W$. Therefore, if $Y' \subset Y$ is B_H -stable and irreducible, then $\text{trdeg } k(Y')^{B_H} = \text{trdeg } k(\overline{B_G \cdot Y'})^{B_G}$. □

Proof of 5.1. — The lemma applies to $V = \mathfrak{g}$, $H = G_0$, and $W = \mathfrak{g}_1$. The condition (*) follows from presence of periodic grading. As (*) holds also for \mathfrak{g}_0 and \mathfrak{g} in place of \mathfrak{b}_H and \mathfrak{b}_G , Vinberg’s lemma [Vi76, §2] implies that each irreducible component of $Ge \cap \mathfrak{g}_1$ is a G_0 -orbit. In particular, one of the components is $G_0 e$. □

It follows that, given a spherical orbit Ge , each irreducible component of $Ge \cap \mathfrak{g}_1$ is a spherical G_0 -orbit, too. But a naive hope for the converse fails to be true. For, any simple Lie algebras has a periodic grading such that G_0 is a torus. (Indeed, if $x \in \mathfrak{g}$ is regular semisimple, with integral eigenvalues, then $\theta = \exp\left(\frac{2\pi\sqrt{-1}}{n}x\right)$ yields such a grading for n large enough.) Then all G_0 -orbits in \mathfrak{g}_1 are spherical, while this is not always the case for the G -orbits in $G\mathfrak{g}_1$. To develop a technique for dealing with the complexity of G_0 -orbits and, in particular, for classifying the spherical ones, we need some preparations.

By a modification of the Morozov-Jacobson theorem, one may assume that $\{e, h, f\}$ is adapted to θ , i.e., $h \in \mathfrak{g}_0$ and $f \in \mathfrak{g}_{-1}$. Then \mathfrak{g} gains a $\mathbb{Z} \times \mathbb{Z}_m$ -grading

$$\mathfrak{g} = \bigoplus_{i \in \mathbb{Z}} \bigoplus_{j \in \mathbb{Z}_m} \mathfrak{g}^{(i)_j}.$$

We have $e \in \mathfrak{g}(2)_1$, $h \in \mathfrak{g}(0)_0$, and $f \in \mathfrak{g}(-2)_{-1}$. By [Vi79, Th. 1], all adapted triples containing e are $(G_0)_e$ -conjugate. Therefore the structure

of this bi-grading does not depend on choice of an adapted triple. One may say that G_0e determines a refinement $\mathfrak{g}(i) = \bigoplus_{j \in \mathbb{Z}_m} \mathfrak{g}(i)_j$ of the \mathbb{Z} -grading associated with Ge . However, it is worth noting that another irreducible component of $Ge \cap \mathfrak{g}_1$ may and usually does determine another refinement of the same \mathbb{Z} -grading. We defer discussing of this and related problems until (5.9) and (5.10).

Denote by L_0 and K_0 the identity components of the θ -fixed subgroups in L and K respectively. Then $\mathfrak{l}_0 = \mathfrak{g}(0)_0$ and it follows from (1.1) that $[\mathfrak{l}_0, e] = \mathfrak{g}(2)_1$ and $(\mathfrak{l}_0)_e = \mathfrak{k}_0$. Clearly, each $\mathfrak{g}(i)_j$ is a K_0 -module and $\langle \mathfrak{g}(i)_j, \mathfrak{g}(i')_{j'} \rangle = 0$ unless $i + i' = 0$ and $j + j' = 0$. An extension of Proposition 1.2 to the bi-grading is given by

5.3. PROPOSITION. — *For all $i, j \in \mathbb{Z}$ we have*

1. $\mathfrak{g}(2i)_i$ and $\mathfrak{g}(2i)_j \oplus \mathfrak{g}(2i)_{2i-j}$ are orthogonal K_0 -modules;
2. $\mathfrak{g}(2i + 1)_j$ and $\mathfrak{g}(2i + 1)_{2i+1-j}$ are dual K_0 -modules.

(The subscripts are being considered as elements of $\mathbb{Z}/m\mathbb{Z}$.)

Proof. — Let $i \geq 0$. Recall from (1.2) the bilinear form Ψ_i . Since $(\text{ad } f)^{2i} : \mathfrak{g}(2i)_i \rightarrow \mathfrak{g}(-2i)_{-i}$ is bijective, the restriction of the symmetric form Ψ_{2i} to $\mathfrak{g}(2i)_i$ is non-degenerate. The other cases are treated similarly. □

Denote by M an s.g.p. for the K_0 -action on $\mathfrak{g}(2)_1$. Again, M is reductive by Luna’s result, since $\mathfrak{g}(2)_1$ is orthogonal. Note that there is no relation in general between $M = \text{s.g.p.}(K_0, \mathfrak{g}(2)_1)$ and $S = \text{s.g.p.}(K, \mathfrak{g}(2))$.

- 5.4. THEOREM.** — 1. $c_{G_0}(G_0e) = c_{L_0}(\mathfrak{g}(2)_1) + c_M(\mathfrak{g}(\geq 3)_1)$;
 2. $c_{G_0}(G_0e) = c_{L_0}(\mathfrak{g}(\geq 2)_1)$.

Proof. — 1. As well as Theorem 4.2(1), it will be a consequence of [Pa94, 1.2]. Namely, to derive a formula for the complexity of $G_0e \simeq G_0/(G_0)_e$, we exploit an embedding of $(G_0)_e$ into some parabolic subgroup in G_0 . Recall that $\mathfrak{p} = \mathfrak{g}(\geq 0)$ and $\mathfrak{l} = \mathfrak{g}(0)$ is a Levi subalgebra in it. Obviously, then $\mathfrak{p}_0 := \bigoplus_{i \geq 0} \mathfrak{g}(i)_0$ is parabolic in \mathfrak{g}_0 and \mathfrak{l}_0 is a Levi subalgebra in it. Let A^u denote the unipotent radical of an algebraic group A . By Proposition 1.1, $G_e \subset P$ and $(G_e)^u \subset P^u$. Set $N = (G_e)^u \cap G_0$. Then the identity component of $(G_0)_e$ is K_0N and $N = (K_0N)^u$. Since $N \subset (P_0)^u$, the embedding $K_0N \subset P_0$ is *right* in terminology of [Pa94]. Because the component group of the stabilizer does not affect the complexity of an orbit,

we may apply [Pa94, 1.2] to conclude

$$c_{G_0}(G_0e) = c_{G_0}(G_0/K_0N) = c_{L_0}(L_0/K_0) + c_M((\mathfrak{p}_0)^u/\mathfrak{n}).$$

Since K_0 is the identity component of $(L_0)_e$ and L_0e is dense in $\mathfrak{g}(2)_1$, we have $c_{L_0}(L_0/K_0) = c_{L_0}(\mathfrak{g}(2)_1)$. It follows from Proposition 1.1(iii) that

$$\mathfrak{p}^u = \mathfrak{g}(\geq 1) = (\mathfrak{g}_e)^u \oplus \text{ad } f \cdot \mathfrak{g}(\geq 3).$$

Whence

$$(\mathfrak{p}_0)^u = (\mathfrak{p}^u)_0 = (\mathfrak{g}_e)_0^u \oplus \text{ad } f \cdot \mathfrak{g}(\geq 3)_1 = \mathfrak{n} \oplus \text{ad } f \cdot \mathfrak{g}(\geq 3)_1.$$

Thus, $(\mathfrak{p}_0)^u/\mathfrak{n}$ is isomorphic to $\mathfrak{g}(\geq 3)_1$ as K_0 - and hence M -module.

2. As is explained in the first part of the proof, the identity component of $(G_0)_e$ lies in P_0 . It follows that the collapsing

$$G_0 *_{P_0} \mathfrak{g}(\geq 2)_1 \rightarrow G_0 \cdot \mathfrak{g}(\geq 2)_1 = \overline{G_0e}$$

is generically finite-to-one. Hence these varieties have the same G_0 -complexity. Again, we conclude by Lemma 4.1. □

Remarks. — 1. All the previous results hold without changes if $\mathfrak{g} = \oplus_j \mathfrak{g}_j$ is a \mathbb{Z} -grading, i.e., formally $m = \infty$.

2. The paper [Pa94] has dealt not only with the complexity, but also with the rank of nilpotent orbits. Although the notions are quite different, the formulas for the both and the proofs turned out to be the same. This also holds in case of θ -groups. For instance, making use of the above embedding $K_0N \subset P_0$ and Theorem 1.2 in [loc cit.], one proves the formula for the rank of G_0e : $r_{G_0}(G_0e) = r_{L_0}(\mathfrak{g}(2)_1) + r_M(\mathfrak{g}(\geq 3)_1)$.

Although it is already possible to give some estimates for $c_{G_0}(G_0e)$, these are isolated and do not enable us to achieve attractive results. For this reason, we stick to the case where θ is involutory. That is, from now on $m = 2$ and we intend to describe spherical nilpotent orbits for the isotropy representation of a symmetric variety G/G_0 . Now, an adapted \mathfrak{sl}_2 -triple yields a splitting of $\mathfrak{g}(i)$ ($i \in \mathbb{Z}$) in two K_0 -submodules and as an immediate corollary of Proposition 5.3 we have

5.5. LEMMA. — (i) For i odd, $\mathfrak{g}(i)_0$ and $\mathfrak{g}(i)_1$ are dual K_0 -modules;

(ii) for i even, $\mathfrak{g}(i)_0$ and $\mathfrak{g}(i)_1$ are orthogonal K_0 -modules. □

Our idea is to characterize spherical G_0 -orbits in terms of the G -orbits in \mathfrak{g} these generate. In view of (5.1), one has to realize which nonspherical G -orbits may arise in this way. To this end, our main tool is Theorem 5.4.

5.6. THEOREM. — Suppose θ is involutory and $e \in \mathfrak{g}_1 \cap \mathcal{N}$. If $\text{ht}(e) \geq 5$ or $\mathfrak{g}(4)_1 \neq 0$, then $c_{G_0}(G_0e) > 0$.

Proof. — Actually, we shall prove that $\mathbf{k}[\mathfrak{g}(\geq 3)_1]^M \neq \mathbf{k}$, which certainly implies that $c_M(\mathfrak{g}(\geq 3)_1) > 0$, cf. (4.3). The situation splits into 3 cases.

(a) Assume $\mathfrak{g}(5) \neq 0$. By (5.5), $\mathfrak{g}(5)_1$ and $\mathfrak{g}(5)_0$ are dual K_0 -modules and hence dual M -modules. Then $(\text{ad } f)\mathfrak{g}(5)_0$ and $\mathfrak{g}(5)_1$ are dual M -modules in $\mathfrak{g}(\geq 3)_1$.

(b) Assume $\mathfrak{g}(4)_1 \neq 0$. By (5.5), it is an orthogonal K_0 -module and hence M -module. Thus, $\mathbf{k}[\mathfrak{g}(\geq 3)_1]^M \neq \mathbf{k}$.

(c) If $\text{ht}(e) \geq 5$, $\mathfrak{g}(5) = 0$, and $\mathfrak{g}(4) \subset \mathfrak{g}_0$, then $0 \neq (\text{ad } e)\mathfrak{g}(4) = \mathfrak{g}(6) = \mathfrak{g}(6)_1$. Consider $(\text{ad } f)^2\mathfrak{g}(6)_1 = (\text{ad } f)\mathfrak{g}(4) \subset \mathfrak{g}(2)_1$. It is a K_0 -submodule. Since $M = \text{s.g.p.}(K_0, \mathfrak{g}(2)_1)$, we have $K_0(\mathfrak{g}(2)_1^M)$ is dense in $\mathfrak{g}(2)_1$. Whence $((\text{ad } f)^2\mathfrak{g}(6)_1)^M \neq 0$ and, finally, $\mathfrak{g}(6)_1^M \neq 0$. Thus, $\mathbf{k}[\mathfrak{g}(\geq 3)_1]^M$ contains a linear function. \square

Combining (5.1) and (5.6) gives a complete coherent description for all G -orbits except for the orbits of height 4:

Let θ be involutory and \mathcal{O} a nilpotent orbit. If $\text{ht}(\mathcal{O}) \leq 3$, then each irreducible component of $\mathcal{O} \cap \mathfrak{g}_1$ is G_0 -spherical. If $\text{ht}(\mathcal{O}) \geq 5$, then none of the irreducible components is G_0 -spherical. This holds regardless of θ provided only that $\mathcal{O} \cap \mathfrak{g}_1 \neq \emptyset$.

(5.7) *If $\text{ht}(\mathcal{O}) = 4$, the answer already depends on relationship between θ and \mathcal{O} . The complexity of the irreducible components of $\mathcal{O} \cap \mathfrak{g}_1$ may or may not be equal to zero, depending on θ (see 5.10). Given $e \in \mathcal{O} \cap \mathfrak{g}_1$, Theorem 5.6 yields a necessary condition for sphericity of G_0e : $\mathfrak{g}(4) = \text{Im}(\text{ad } e)^4 \subset \mathfrak{g}_0$. It is not however sufficient.*

Note that, given e and θ , it is easy to realize when $Ge \cap \mathfrak{g}_1 \neq \emptyset$. By a result of L. Antonyan [An82, Th. 1], $Ge \cap \mathfrak{g}_1 \neq \emptyset$ if and only if $Gh \cap \mathfrak{g}_1 \neq \emptyset$. The last condition is immediately being verified by comparing the weighted Dynkin diagram of e and the Satake diagram of θ :

$Gh \cap \mathfrak{g}_1 \neq \emptyset$ if and only if the weighted Dynkin diagram has zero labels on the black nodes of the Satake diagram and equal labels on the pairs of nodes connected by arrow.

5.8. Example. — Using (5.4), one can give a recipe for producing triples $(\mathfrak{g}, \theta, e)$ such that $e \in \mathfrak{g}_1$, $c_G(Ge) > 0$, and $c_{G_0}(G_0e) = 0$. Let e be an even nilpotent element with $\text{ht}(e) = 4$. Then $c_G(Ge) > 0$.

Define a \mathbb{Z}_2 -grading of \mathfrak{g} by the formulas $\mathfrak{g}_0 = \mathfrak{g}(-4) \oplus \mathfrak{g}(0) \oplus \mathfrak{g}(4)$, $\mathfrak{g}_1 = \mathfrak{g}(-2) \oplus \mathfrak{g}(2)$. Then $L_0 = L$, $\mathfrak{g}(2) = \mathfrak{g}(2)_1$, and $\mathfrak{g}(\geq 3)_1 = 0$. Therefore $c_{G_0}(G_0e) = c_L(\mathfrak{g}(2))$. All such orbits with spherical L -module $\mathfrak{g}(2)$ are listed in Table 1.

Table 1

\mathfrak{g}	characteristic	type of \mathcal{O}	L	$\dim \mathfrak{g}(2)$	\mathfrak{g}_0
\mathfrak{sl}_n ($n \geq 3$)	2-0-...-0-2	A_2	$SL_{n-2} \times (\mathfrak{k}^*)^2$	$2n - 4$	$\mathfrak{sl}_{n-2} \oplus \mathfrak{sl}_2 \oplus \mathfrak{k}$
\mathfrak{sl}_6	0-2-0-2-0	$2A_2$	$(SL_2)^3 \times (\mathfrak{k}^*)^2$	8	$\mathfrak{sl}_4 \oplus \mathfrak{sl}_2 \oplus \mathfrak{k}$
\mathfrak{sp}_{2n} ($n \geq 3$)	0-2-0-...-0	A_2	$SL_2 \times Sp_{2n-4} \times \mathfrak{k}^*$	$4n - 8$	$\mathfrak{sp}_{2n-4} \oplus \mathfrak{sp}_4$
\mathfrak{sp}_{12}	0-0-0-2-0-0	$2A_2$	$SL_4 \times Sp_4 \times \mathfrak{k}^*$	16	$\mathfrak{sp}_8 \oplus \mathfrak{sp}_4$
\mathfrak{f}_4	2-0-0-0-0	\tilde{A}_2	$Spin_7 \times \mathfrak{k}^*$	8	\mathfrak{so}_9
\mathfrak{e}_6	2-0-0-0-2	$2A_2$	$Spin_8 \times (\mathfrak{k}^*)^2$	16	$\mathfrak{so}_{10} \oplus \mathfrak{k}$
	0				

In the column “ L ”, we indicate the simply connected group with Lie algebra $\mathfrak{g}(0)$.

5.9. On behaviour of irreducible components. — In general, different irreducible components of $\mathcal{O} \cap \mathfrak{g}_1$ determine non-isomorphic bi-gradings of \mathfrak{g} ; in particular, the groups L_0 can be different. One may address the following questions in this regard:

Is it possible that these components have different complexity relative to G_0 ?

Is it possible that spherical and non-spherical components occur together?

The answer to the first question is “yes” and we present below an example of triple $(\mathfrak{g}, \theta, e)$ such that the complexity of irreducible components of $\mathcal{O} \cap \mathfrak{g}_1$ takes three values. As for the second question, it seems that the answer is “no”. Such a situation might only occur if $\text{ht}(\mathcal{O}) = 4$. But our computations based on an explicit classification of the G_0 -orbits confirm the negative answer. For instance, to compute the complexity of the irreducible components of $Ge \cap \mathfrak{g}_1$ in the exceptional case, we have used Djoković’s tables [Dj88]. This will be published elsewhere. It is however desirable to have a classification-free proof.

5.10. Example. — Let $\mathfrak{g} = \mathfrak{sl}_N$. Denote by θ_n ($1 \leq n \leq N/2$) an inner involution of \mathfrak{g} such that $\mathfrak{g}_0 \simeq \mathfrak{sl}_n \oplus \mathfrak{sl}_{N-n} \oplus \mathfrak{k}$. To emphasize the dependence on n , we shall write $G_0^{(n)}$ and $\mathfrak{g}_1^{(n)}$. The elements of $\mathfrak{g}_1^{(n)}$ can be thought of as the pairs of counter operators (A, B) , where $A \in \mathbf{Hom}(V, W)$,

$B \in \mathbf{Hom}(W, V)$, $\dim V = n$, and $\dim W = N - n$. The orbit classification in this case was first obtained in [DP65]. (From the modern point of view, this is a special case of the quiver theory.) We explain this classification using the language of ab -diagrams introduced in [KP79, sect. 4]. Given $\mathcal{O} = (a_1, \dots, a_t)$, a corresponding ab -diagram is obtained if one writes a string of consecutive symbols “ a ” and “ b ”, of length a_i , in place of part “ a_i ”. Two ab -diagrams are proclaimed to be equivalent if these are obtained from each other by reordering ab -strings of equal length. There is a bijection between the irreducible components of $\mathcal{O} \cap \mathfrak{g}_1^{(n)}$ and the classes of equivalent ab -diagrams such that the total number of a ’s is n .

For $N \geq 6$, consider $\mathcal{O} = (3, 3, 1^{N-6})$. Its weighted Dynkin diagram is $0-2-0-\dots-0-2-0$. Comparing with the Satake diagram of θ_n , one finds that $\mathcal{O} \cap \mathfrak{g}_1^{(n)} \neq \emptyset$ if and only if $n \geq 2$. Here the different irreducible components of $\mathcal{O} \cap \mathfrak{g}_1^{(n)}$ correspond to the following ab -diagrams:

I : (bab, bab, \dots) ; II : (aba, aba, \dots) ; III : (aba, bab, \dots) .

The strings of length 1 are uniquely determined by the constraint that the symbol “ a ” appears exactly n times. Note that case I (resp. II) occurs if and only if $n \geq 2$, $N - n \geq 4$ (resp. $n \geq 4$, $N - n \geq 2$) and case III occurs if and only if $3 \leq n \leq N - 3$. This again shows that $\mathcal{O} \cap \mathfrak{g}_1^{(n)} \neq \emptyset$ only for $2 \leq n \leq N - 2$. Furthermore, this intersection contains at most 3 irreducible components. Making use of these ab -diagrams, one can write explicitly an \mathfrak{sl}_2 -triple adapted to θ_n . An explicit matrix form of the latter enables us to determine the decompositions $\mathfrak{g}(i)_0 \oplus \mathfrak{g}(i)_1$ and then to compute the complexity. The answer is that the complexity relative to $G_0^{(n)}$ of the irreducible components of $\mathcal{O} \cap \mathfrak{g}_1^{(n)}$ is equal to: $n - 2$ in case I; $N - n - 2$ in case II; 1 in case III. For instance, if $N = 9$ and $n = 4$ then there are irreducible components of complexity 1, 2, and 3. Since $c_G(\mathcal{O}) = N - 2$, one obtains a nice illustration to Theorem 5.1 as well. It also follows from above formulas that, for $n = 2$, $\mathcal{O} \cap \mathfrak{g}_1^{(2)}$ is irreducible and is a spherical $G_0^{(2)}$ -orbit.

6. Questions, observations, remarks.

6.1. Irreducibility.— Theorem 2.6 says that \mathcal{N}^{sph} , the union of all nilpotent spherical G -orbits, is determined set-theoretically by the equations $(\text{ad } e)^4 = 0$. Hence \mathcal{N}^{sph} is closed⁽⁵⁾. A direct verification shows

⁽⁵⁾ It is not hard to prove that X^{sph} is closed for any G -variety X .

that \mathcal{N}^{sph} is irreducible. The explicit expression for the maximal spherical orbit in the classical case is derived from [Pa94, sect. 4]; in the exceptional case, look at the tables in [Spal, IV.2]. It might be interesting to give an *a priori* proof. However, irreducibility fails for arbitrary θ -groups. For instance, $(\mathcal{N} \cap \mathfrak{g}_1)^{\text{sph}}$ has 2 irreducible components for an involution of SL_2 . A more exotic example is an involution of \mathbf{F}_4 of maximal rank.

Making use of the known normality results (due to Broer and Kraft & Procesi), one sees that \mathcal{N}^{sph} is normal for $G = \mathbf{A}_p, \mathbf{B}_{2p}, \mathbf{C}_p, \mathbf{D}_p, \mathbf{F}_4$; and not normal for \mathbf{B}_{2p+1} and \mathbf{G}_2 . It is likely \mathcal{N}^{sph} is normal for \mathbf{E}_p , and I think there ought to be a unified proof for **A-D-E**.

6.2. The defining ideal. — It would be interesting to describe conceptually the defining ideal of $(\mathcal{N} \cap \mathfrak{g}_1)^{\text{sph}}$.

In case $m = 1$, i.e., for the adjoint representation, Theorem 2.6 shows that \mathcal{N}^{sph} is determined set-theoretically by polynomials of degree 4. In the classical case, it is however clear that the matrix coefficients of $(\text{ad } e)^4$ do not generate a radical ideal. Indeed, for the “multiplicity-free” reason, the covariants of type \mathfrak{g} of degree > 1 must vanish on \mathcal{N}^{sph} . But there exist such covariant of degree 2 (resp. 3) for $\mathfrak{g} = \mathfrak{sl}(V)$ (resp. $\mathfrak{g} = \mathfrak{so}(V)$ or $\mathfrak{sp}(V)$). However, I see no obstructions to that the coefficients of $(\text{ad } e)^4$ would generate a radical ideal in the exceptional case.

6.3. Dimension. — By definition, $\dim \mathcal{O} \leq \dim B_G$ for $\mathcal{O} \subset \mathcal{N}^{\text{sph}}$. Our aim here is a bit sharper inequality. Without loss of generality, we may assume that \mathfrak{g} is simple. Denote by k_0 (resp. k_1) the number of even (resp. odd) exponents of \mathfrak{g} . Then $k_0 + k_1 = \text{rk } \mathfrak{g} =: p$.

PROPOSITION. — *If $\mathcal{O} \subset \mathcal{N}^{\text{sph}}$, then $\dim \mathcal{O} \leq \dim B_G - k_0$.*

Proof. — This is easily verified case-by-case, but we give a conceptual proof. Let $\mathfrak{g} = \mathfrak{g}_0^{\text{max}} \oplus \mathfrak{g}_1^{\text{max}}$ be the decomposition corresponding to an involution of maximal rank θ^{max} . This means $\mathfrak{g}_1^{\text{max}}$ contains a Cartan subalgebra (= C.s.a.) of \mathfrak{g} . By [An82, th. 2], we have $\mathcal{O} \cap \mathfrak{g}_1^{\text{max}} \neq \emptyset$. It then follows from [KR71, prop. 5] that $\dim(\mathcal{O} \cap \mathfrak{g}_1^{\text{max}}) = \frac{1}{2} \dim \mathcal{O}$. Since each irreducible component of $\mathcal{O} \cap \mathfrak{g}_1^{\text{max}}$ is G_0^{max} -spherical (5.1), we have

$$\frac{1}{2} \dim \mathcal{O} = \dim(\mathcal{O} \cap \mathfrak{g}_1^{\text{max}}) \leq \dim B_0^{\text{max}}.$$

Here B_0^{max} is a Borel subgroup of G_0^{max} . Thus, it suffices to demonstrate that $\dim B_0^{\text{max}} = \frac{1}{2}(\dim B_G - k_0)$. This is equivalent to the equality in the next Lemma, since $\dim G_0^{\text{max}} = \dim B_G - p$. □

LEMMA. — $\text{rk } G_0^{\max} = k_1$.

Proof. — Let \mathfrak{h} be a θ^{\max} -stable C.s.a. of \mathfrak{g} such that \mathfrak{h}_0 is a C.s.a. of \mathfrak{g}_0^{\max} . Set $\sigma := \theta^{\max}|_{\mathfrak{h}}$. Since the G -orbit of characteristics of regular nilpotent elements intersects \mathfrak{g}_0^{\max} (consider an adapted \mathfrak{sl}_2 -triple!), \mathfrak{h}_0 contains regular semisimple elements. By [Sp74, 6.5], the eigenvalues of σ are ε_i^{-1} ($1 \leq i \leq p$), where the ε_i 's are the "eigenvalues of θ^{\max} on the set of basic invariants". The latter means that homogeneous generators F_1, \dots, F_p of the polynomial algebra $\mathbf{k}[\mathfrak{g}]^G$ can be chosen so that $\theta^{\max} \cdot F_i = \varepsilon_i F_i$. (Of course, $\varepsilon_i \in \{1, -1\}$.) On the other hand, there is a C.s.a. $\mathfrak{h}' \subset \mathfrak{g}_1^{\max}$, i.e., $\sigma' := \theta^{\max}|_{\mathfrak{h}'} = -\text{id}$. By [Sp74, 6.4(v)], the eigenvalues of σ' are $\varepsilon_i^{-1}(-1)^{m_i}$ ($1 \leq i \leq p$), where m_1, \dots, m_p are the exponents of \mathfrak{g} . Whence $\varepsilon_i = 1$ if m_i is odd and $\varepsilon_i = -1$ if m_i is even. Finally, $\text{rk } G_0^{\max} = \dim \mathfrak{h}_0 = \#\{i \mid \varepsilon_i = 1\} = k_1$. □

We have proven that $\dim \mathcal{N}^{\text{sph}} \leq \dim B_G - k_0$. Actually, the equality holds, but I do not know a unified proof. A promising approach is discussed in the next subsection.

There is another curious coincidence related to another involution. By [An82, Th. 3(2)], there is a unique, up to G -conjugation, *inner* involution θ^{int} of \mathfrak{g} such that *all* nilpotent G -orbits intersect $\mathfrak{g}_1^{\text{int}}$. ($\theta^{\text{int}} \neq \theta^{\max}$ for $\mathbf{A}_p, \mathbf{D}_{2p+1}$, and \mathbf{E}_6 .) One can conceptually prove that $\dim \mathfrak{g}_0^{\text{int}} - \dim \mathfrak{g}_1^{\text{int}} = k_0 - k_1$. Since $\text{rk } G_0^{\text{int}} = \text{rk } G$, this implies $\dim \mathfrak{g}_1^{\text{int}} = \dim B_G - k_0$. That is, $\dim \mathcal{N}^{\text{sph}} = \dim \mathfrak{g}_1^{\text{int}}$.

6.4. A relationship with the index of B_G . — The index of an algebraic group (or its Lie algebra) is the minimal codimension of its orbits in the coadjoint representation. The index $\text{ind } B_G$ of B_G for all simple groups was computed in [Tr83, §4]. This result can be stated as: $\text{ind } B_G = k_0$. It seems however that no explanation of this equality is known. Our observation is:

PROPOSITION. — *If $\mathcal{O} \subset \mathcal{N}^{\text{sph}}$, then $\dim \mathcal{O} \leq \dim B_G - \text{ind } B_G$.*

Proof. — By the very definition of sphericity, there is $x \in \mathcal{O}$ such that $\mathfrak{g}_x + \mathfrak{b}_G = \mathfrak{g}$. Taking the orthocomplements, we obtain

$$(\heartsuit) \quad [\mathfrak{g}, x] \cap [\mathfrak{b}_G, \mathfrak{b}_G] = \{0\}.$$

Let \bar{x} be the image of x in $\mathfrak{g}/[\mathfrak{b}_G, \mathfrak{b}_G]$. The latter is identified with the B_G -module $(\mathfrak{b}_G)^*$. Since $x \in [\mathfrak{g}, x]$, we have $\bar{x} \neq 0$. Because $[\mathfrak{g}, x] = [\mathfrak{b}_G, x]$, equality (\heartsuit) implies that $\dim B_G - \text{ind } B_G \geq \dim B_G \bar{x} = \dim B_G x = \dim \mathcal{O}$.

□

Given an $\xi \in (\mathfrak{b}_G)^* \simeq \mathfrak{g}/[\mathfrak{b}_G, \mathfrak{b}_G]$, it would be interesting to realize, is there $x \in \mathfrak{g}$ over ξ such that Gx is spherical?

6.5. In search of a uniform proof. — Recently, Fan and Stembridge found a relationship in the *simply laced* case between the spherical nilpotent orbits and the “commutative” elements in the Weyl group W . They proved (see [FS97, Th. 3.1(a)]) that, for a natural map $\phi : W \rightarrow \mathcal{N}/G$, the set W_c of commutative elements is just the preimage of \mathcal{N}_4/G , where $\mathcal{N}_4 = \{e \in \mathcal{N} \mid \text{ht}(e) < 4\}$. But their hope [FS97, 3.4(d)] that better understanding of the fibres of ϕ may lead to a uniform proof of (2.6) seems to me groundless. For: (1) the proof of Theorem 3.1(a) in [loc.cit.] also uses classification results; (2) while the concept of a spherical orbit is quite general, the above relationship breaks in the non simply laced case, even if one replaces “commutative elements” in W by “fully commutative” or “short-braid avoiding” ones. To see this, it suffices to consider the groups Sp_4 and G_2 .

Actually, case-by-case (classification) arguments in proving (2.6) were used only for the orbits of height 3. My opinion is that a right way consists in a better understanding of the orbits of height 3, as special case of orbits of odd height.

BIBLIOGRAPHY

- [An82] L.V. ANTONYAN, On classification of homogeneous elements of \mathbb{Z}_2 -graded semisimple Lie algebras, Vestnik Mosk. Un-ta, Ser. Matem. & Mech. No. 2 (1982), 29–34 (Russian). English translation: Moscow Univ. Math. Bulletin, 37, No. 2 (1982), 36–43.
- [BC76] P. BALA, R.W. CARTER, Classes of unipotent elements in simple algebraic groups, II, Math. Proc. Cambridge Philos. Soc., 80 (1976), 1–18.
- [CM93] D.H. COLLINGWOOD, W.M. MCGOVERN, Nilpotent orbits in semisimple Lie algebras, New York: Van Nostrand Reinhold, 1993.
- [Dj88] D. DJOKOVIĆ, Classification of nilpotent elements in simple exceptional real Lie algebras of inner type and description of their centralizers, J. Alg., 112 (1988), 503–524.
- [DP65] N.M. DOBROVOLSKAYA, V.A. PONOMAREV, Pairs of counter operators, Uspekhi Matem. Nauk, 20, No. 6 (1965), 81–86 (Russian).

- [Dy52] E.B. DYNKIN, Semisimple subalgebras of semisimple Lie algebras, *Matem. Sbornik*, 30, No. 2 (1952), 349–462 (Russian). English translation: *Amer. Math. Soc. Transl. II, Ser.*, 6 (1957), 111–244.
- [El75] A.G. ELASHVILI, The centralizers of nilpotent elements in semisimple Lie algebras, *Trudy Tbiliss. Matem. Inst. Akad. Nauk Gruzin. SSR*, 46 (1975), 109–132 (Russian).
- [El85] A.G. ELASHVILI, Frobenius Lie algebras II, *Trudy Tbiliss. Matem. Inst. Akad. Nauk Gruzin. SSR*, 77 (1985), 127–137 (Russian).
- [FS97] C.K. FAN, J.R. STEMBRIDGE, Nilpotent orbits and commutative elements, *J. Algebra*, 196 (1997), 490–498.
- [Ka80] V.G. KAC, Some remarks on nilpotent orbits, *J. Algebra*, 64 (1980), 190–213.
- [KR71] B. KOSTANT, S. RALLIS, Orbits and representations associated with symmetric spaces, *Amer. J. Math.*, 93 (1971), 753–809.
- [KP79] H. KRAFT, C. PROCESI, Closures of conjugacy classes of matrices are normal, *Invent. Math.*, 53 (1979), 227–247.
- [Lu72] D. LUNA, Sur les orbites fermées des groupes algébriques réductifs, *Invent. Math.*, 16 (1972), 1–5.
- [Pa87] D. PANYUSHEV, Orbits of maximal dimension of solvable subgroups of reductive algebraic groups and reduction for U -invariants, *Matem. Sb.*, 132, No. 3 (1987), 371–382 (Russian). English translation: *Math. USSR-Sb.*, 60 (1988), 365–375.
- [Pa94] D. PANYUSHEV, Complexity and nilpotent orbits, *Manuscripta Math.*, 83 (1994), 223–237.
- [Spal] N. SPALTENSTEIN, “Classes Unipotentes et Sous-groupes de Borel”, *Lecture notes in Math.*, 946, Berlin Heidelberg New York: Springer 1982.
- [Sp74] T.A. SPRINGER, Regular elements in finite reflection groups, *Invent. Math.*, 25 (1974), 159–198.
- [SpSt] T.A. SPRINGER, R. STEINBERG, Conjugacy classes, In: “Seminar on algebraic groups and related finite groups”. *Lecture notes in Math.*, 131, pp. 167–266, Berlin-Heidelberg-New York, Springer, 1970.
- [Tr83] V.V. TROFIMOV, Semi-invariants of the coadjoint representation of Borel subalgebras of simple Lie algebras, In: “Trudy seminar po vect. i tenz. analizu”, vol. 21, pp. 84–105. Moscow: MGU 1983 (Russian). English translation: *Selecta Math. Sovietica*, 8 (1989), 31–56.
- [Vi75] E.B. VINBERG, On the classification of nilpotent elements of graded Lie algebras, *Dokl. Akad. Nauk SSSR*, 225 (1975), No. 4, 745–748 (Russian). English translation: *Soviet Math. Dokl.*, 16 (1975), 1517–1520.
- [Vi76] E.B. VINBERG, The Weyl group of a graded Lie algebra, *Izv. Akad. Nauk SSSR, Ser. Mat.*, 40 (1976), No. 3, 488–526 (Russian). English translation: *Math USSR-Izv.*, 10 (1976), 463–495.
- [Vi79] E.B. VINBERG, Classification of homogeneous nilpotent elements of a semisimple graded Lie algebra, In: “Trudy seminar po vect. i tenz. analizu”, vol. 19, pp. 155–177. Moscow: MGU 1979 (Russian). English translation: *Selecta Math. Sovietica*, 6 (1987), 15–35.

- [Vi86] E.B. VINBERG, Complexity of actions of reductive groups, *Funkt. Anal. i Prilozhen*, 20, No. 1 (1986), 1–13 (Russian). English translation: *Funct. Anal. Appl.*, 20 (1986), 1–11.
- [VP89] E.B. VINBERG, V.L. POPOV, Invariant theory, In: *Sovremennye problemy matematiki. Fundamentalnye napravleniya*, t. 55, pp. 137–309. Moscow: VINITI 1989 (Russian). English translation in: *Algebraic Geometry IV (Encyclopaedia Math. Sci.*, vol. 55, pp. 123–284) Berlin-Heidelberg-New York, Springer, 1994.

Manuscrit reçu le 2 février 1999,
accepté le 19 mars 1999.

Dmitri I. PANYUSHEV,
ul. Akad. Anokhina, d.30, kor.1, kv.7
Moscow 117602 (Russia).
panyush@dpa.msk.ru