

ALGEBRAIC COMBINATORICS

Valentin Bonzom, Guillaume Chapuy & Maciej Dołęga

Enumeration of non-oriented maps via integrability

Volume 5, issue 6 (2022), p. 1363-1390.

<https://doi.org/10.5802/alco.268>

© The author(s), 2022.



This article is licensed under the
CREATIVE COMMONS ATTRIBUTION 4.0 INTERNATIONAL LICENSE.
<http://creativecommons.org/licenses/by/4.0/>




*Algebraic Combinatorics is published by The Combinatorics Consortium
and is a member of the Centre Mersenne for Open Scientific Publishing*

www.tccpublishing.org www.centre-mersenne.org

e-ISSN: 2589-5486





Enumeration of non-oriented maps via integrability

Valentin Bonzom, Guillaume Chapuy & Maciej Dołęga

ABSTRACT In this note, we examine how the BKP structure of the generating series of several models of maps on non-oriented surfaces can be used to obtain explicit and/or efficient recurrence formulas for their enumeration according to the genus and size parameters.

Using techniques already known in the orientable case (elimination of variables via Virasoro constraints or Tutte equations), we naturally obtain recurrence formulas with non-polynomial coefficients. This non-polynomiality reflects the presence of shifts of the charge parameter in the BKP equation. Nevertheless, we show that it is possible to obtain non-shifted versions, meaning pure ODEs for the associated generating functions, from which recurrence relations with polynomial coefficients can be extracted. We treat the cases of triangulations, general maps, and bipartite maps.

These recurrences with polynomial coefficients are conceptually interesting but bigger to write than those with non-polynomial coefficients. However they are relatively nice-looking in the case of one-face maps. In particular we show that Ledoux's recurrence for non-oriented one-face maps can be recovered in this way, and we obtain the analogous statement for the (bivariate) bipartite case.

1. INTRODUCTION AND MAIN RESULTS

In this note, we are interested in obtaining simple, or at least efficient, recurrence formulas to count maps on surfaces according to their genus and size parameters. For us, a *map* is the 2-cell embedding of a connected multigraph in a compact connected surface, considered up to homeomorphism. Our surfaces are not necessarily orientable, and we call *genus* of a surface the number $g \in \frac{1}{2}\mathbb{N}$ such that its Euler characteristic is $2 - 2g$. The sphere has genus 0, the projective plane has genus $\frac{1}{2}$, the torus and Klein bottle have genus 1, etc.

Perhaps one of the nicest-looking formulas in the field of map enumeration is the Goulden–Jackson recurrence formula for orientable *triangulations*, i.e. maps in which all faces are incident to three edge-sides. The Goulden–Jackson recurrence [29], in fact also discovered in an equivalent form in [32, Eq. (B.6)], asserts that the number $t_{n,g}$ of rooted⁽¹⁾ triangulations

Manuscript received 25th October 2021, revised 9th September 2022, accepted 14th September 2022.

KEYWORDS. combinatorial maps, nonorientable surfaces, enumeration, BKP hierarchy, zonal polynomials, matrix integrals.

ACKNOWLEDGEMENTS. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. ERC-2016-STG 716083 "CombiTop"). GC is also supported by the grants ANR-18-CE40-0033 "Dimers" and ANR-19-CE48-0011 "Combiné". MD is supported from *Narodowe Centrum Nauki*, grant UMO-2017/26/D/ST1/00186. VB is partially supported by the ANR-20-CE48-0018 3DMaps.

⁽¹⁾precise definitions of all terms used in this introduction are given in the later sections.

with n faces on an *orientable* surface of genus g is solution of the equation

$$(1) \quad (n + 1)t_{n,g} = 4n(3n - 2)(3n - 4)t_{n-1,g-1} + 4 \sum_{\substack{i+j=n-2 \\ h+k=g}} (3i + 2)(3j + 2)t_{i,h}t_{j,k}.$$

This formula was immediately recognized as a breakthrough in the field, because it gives a much better access to these numbers (computational or theoretical) than the classical techniques.

Indeed, in the classical approach, one introduces generating functions of maps of genus g with a certain number of additional boundaries, and one shows that a combinatorial operation of root-deletion on the maps (the “Tutte decomposition”) implies a functional equation for these functions. This approach has been very successful in the planar case since the work of Tutte, see e.g. [39, 40, 41, 8, 11, 10]. In higher genus, it was pioneered by Lehman and Walsh [43] and later Bender and Canfield, who showed that the generating functions of maps of fixed genus and number of boundaries can be computed inductively on the Euler characteristic, thus revealing their particularly nice algebraic structure as well as their singular behaviour [6, 7]. Bender and Canfield’s inductive technique can be seen as a predecessor of the Chekhov–Eynard–Orantin topological recursion [24, 27], a powerful theory invented in the context of matrix integrals [26] which has now been applied to study the structure of fixed-genus generating functions of many models of maps or in enumerative geometry [28, 4, 19, 5, 14].

The need to introduce additional boundaries (and “catalytic” variables to mark their sizes) makes these approaches ineffective for large values of g . There seems to be no hope to obtain control on the bivariate numbers $t_{n,g}$ for non-fixed g in this way, a striking contrast with the recurrence (1). For example (1) also gives access to the so-called *double-scaling limit* of the numbers $t_{n,g}$ [32, 9], and it is also crucially used in the recent Budzinski-Louf breakthrough on large genus asymptotics [18]. Perhaps we should insist on the fact that we mean no harm to the “classical approach”. The study of the rational parametrization it gives rise to is a fascinating subject, including purely bijective combinatorics [23, 22, 35, 3, 25], with probable link to the study of random geometries [16, 17]. On the other hand, as of today, the bijective interpretation of the Goulden–Jackson recurrence (1) is wide open.

One reason why the recurrence (1) gives access to different results is because it comes from a completely different technique. It is based on the fact that the generating function of maps on orientable surfaces, with an infinite number of variables $p_i, i \geq 1$ (p_i marking faces of degree i), is a solution of the KP hierarchy – an infinite sequence of PDEs originating from the theory of integrable systems, with deep connections to infinite dimensional Lie algebras and algebraic combinatorics [31, 37]. The first equation of the hierarchy (*the KP equation*) reads

$$(2) \quad F_{2^2} - F_{3,1} + \frac{1}{2}F_{1^2}^2 + \frac{1}{12}F_{1^4} = 0,$$

where each i -index indicates a partial derivative with respect to p_i . In order to go from the KP equation (2) to the recurrence (1), Goulden and Jackson use the fact that the generating function $F(p_1, p_2, p_3, 0, \dots)$ of maps having only faces of sizes 1, 2, 3 can in fact be expressed in terms of the series $F(0, 0, t, 0, \dots)$ of triangulations only. This enables one to set $p_i = t\delta_{i,3}$ in (2) and obtain an *ODE* for the generating function of triangulations (in this paper we use the Kronecker δ symbol). The fact that the variables p_1 and p_2 can be *eliminated* in this way relies on local surgery operations that can, in fact, be interpreted as first cases of the classical Tutte decomposition.

A similar elimination technique has since been used to obtain similar results for other models of maps. In [21], Carrell and the second author use the fact that the generating function of bipartite maps solves the KP hierarchy, and local operations related to the first Tutte equations for bipartite quadrangulations, to obtain a recurrence formula similar to (1) to count

maps by vertices and faces. In [33], Kazarian and Zograf use a slightly different elimination procedure, using the so-called Virasoro constraints (which are also related to Tutte decompositions) to recover the recurrence of [21] and to obtain an analogue for bipartite maps. These three works only use the first KP equation. Finally, Louf [36] uses a different integrable hierarchy (the Toda hierarchy) to obtain a remarkable recurrence counting bipartite maps of arbitrary genus with control on all face degrees, using a different elimination technique inspired by Okounkov’s work on Hurwitz numbers [38].

In this paper, we are interested in obtaining variants of these results for non-oriented surfaces. Our starting point is the fact that generating functions of maps (or bipartite maps) are solutions of the BKP hierarchy of Kac and Van De Leur [30] (see also the appendix of [15]). An important difference between the KP and BKP hierarchy is that the function F which is a solution of this hierarchy also involves a so-called *charge parameter* N , which in our context will always be a variable marking faces or vertices of a certain kind. The first BKP equation reads

$$(3) \quad F_{2^2}(N) - F_{3,1}(N) + \frac{1}{2}F_{1^2}(N)^2 + \frac{1}{12}F_{1^4}(N) = S_2(N)\tau(N-2)\tau(N+2)\tau(N)^{-2},$$

where $\tau(N) = e^{F(N)}$ and where $S_2(N)$ is a model-dependant normalizing factor that will always be an explicit rational function in our case. In [20], Carrell used the fact that the generating function of non-oriented maps satisfies this equation, together with the elimination techniques developed by Goulden and Jackson in the orientable case, to obtain a functional equation for the case of triangulations (this technique leads to an explicit recurrence, see Theorem 4.10 below).

The first task we perform in this paper, somehow unsurprisingly, is to apply the elimination of variables from the papers [21, 33] to the BKP equation, to obtain recurrences of the same kind to count maps (by vertices and edges) and bipartite maps (by edges, and vertices of each colour) on non-oriented surfaces. The Virasoro constraints for these models are known (e.g. [15]) and our main task here is to make sure that the elimination procedure indeed works, i.e. that these equations indeed enable to reduce all derivatives appearing in (3) to differential polynomials in a single variable. For completeness, we also treat Carrell’s case of triangulations explicitly. All these recurrences are larger than (1), but incredibly short compared to any alternative, and it is not unreasonable to believe that they could have a combinatorial interpretation. For example, we obtain in Section 5 the following recurrence formula. Everywhere in the paper, the symbol \sum denotes a sum over elements of $\frac{1}{2}\mathbb{N}$.

THEOREM 1.1 (Counting maps by edges and genus). *The number h_n^g of rooted maps of genus g with n edges, orientable or not, can be computed from the following recurrence formula:*

$$(4) \quad h_n^g = \frac{2}{(n+1)(n-2)} \left(n(2n-1)(2h_{n-1}^g + h_{n-1}^{g-1/2}) + \frac{(2n-3)(2n-2)(2n-1)(2n)}{2} h_{n-2}^{g-1} \right. \\ + 12 \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n \\ n_1+n_2=n}} \frac{(2n_2-1)(2n_1-1)n_1}{2} h_{n_2-1}^{g_2} h_{n_1-1}^{g_1} \\ - \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n-1 \\ n_1+n_2=n}} \sum_{\substack{g_0=0..g_1 \\ g_1-g_0 \in \mathbb{N}}} \binom{n_1+2-2g_0}{n_1-2g_1} 2^{2(1+g_1-g_0)} h_{n_1}^{g_0} \\ \left. \left(\frac{(2n_2-1)(2n_2-2)(2n_2-3)}{2} h_{n_2-2}^{g_2-1} - \delta_{(n_2, g_2) \neq (n, g)} \frac{n_2+1}{4} h_{n_2}^{g_2} + \frac{2n_2-1}{2} (2h_{n_2-1}^{g_2} + h_{n_2-1}^{g_2-1/2}) \right) \right. \\ \left. + 6 \sum_{\substack{g_3=0..g_2 \\ g_3+g_4=g_2}} \sum_{\substack{n_3=0..n_2 \\ n_3+n_4=n_2}} \frac{(2n_3-1)(2n_4-1)}{4} h_{n_3-1}^{g_3} h_{n_4-1}^{g_4} \right) \Bigg)$$

for $n > 2$, with the initial conditions $h_0^0 = 1, h_1^0 = 2, h_2^0 = 9, h_1^{1/2} = 1, h_2^{1/2} = 10, h_2^1 = 5$, and $h_n^g = 0$ if $n < 2g$.

We will obtain similar theorems for other models, in particular one with control on vertices and faces (Theorem 5.2 or Theorem 3.7 depending on the elimination technique), one for bipartite maps (Theorem 4.4), and one for triangulations which is implicit in Carrel’s work (Theorem 4.10).

The crucial fact that the BKP equation (3) involves not only the function $F(N)$ but also its shifts $F(N + 2)$ and $F(N - 2)$ has an important effect on the recurrence formulas we obtain. The functional equations corresponding to these recurrences, which involve derivatives but also shifts of variables, are *not* ODE in their main variable. In return, the recurrences obtained do not have polynomial coefficients (for example (4) contains binomial coefficients, which are not polynomials in the summation variables). This is a deep structural difference between the recurrence (1) and recurrences such as (4).

It is natural to ask if one could instead obtain formulas in which the shifts are not involved, i.e. true polynomial recurrence formulas, corresponding to nonlinear ODEs with polynomial coefficients for the associated generating functions. This would be much more satisfying, at least at the conceptual level. Maybe surprisingly, we will see that the answer to this question is *yes*. To see this, we will have to use several (in fact, three) equations of the BKP hierarchy. Using additional derivations and manipulations, we will be able to eliminate the shifts from equations, and obtain equations at fixed N , at the price of having to consider higher derivatives. It is not obvious, but it will be true, that a finite number of Virasoro constraints will still be sufficient to perform the elimination of variables in this context.

Due to the use of higher BKP equations and additional manipulations involved, the equations thus obtained are bigger than the previous ones.⁽²⁾ We will only state them here in a non-explicit form. The reader eager to see them at work may access these equations, and use them to compute numbers of maps, in the accompanying Maple worksheet [13]. A typical statement we obtain from these methods is the following.

THEOREM 1.2 (Counting maps by edges and genus – unshifted recurrence). *The number h_n^g of rooted maps of genus g with n edges, orientable or not, is solution of an explicit recurrence relation of the form*

$$(5) \quad h_n^g = \sum_{a=0}^{K_1} \sum_{b=0}^{K_2} \sum_{k=1}^{K_3} \sum_{\substack{n_1, \dots, n_k \geq 1 \\ n_1 + \dots + n_k = n - a}} \sum_{\substack{g_1, \dots, g_k \geq 0 \\ g_1 + \dots + g_k = g - b}} P_{a,b,k}(n_1, \dots, n_k) h_{n_1}^{g_1} h_{n_2}^{g_2} \dots h_{n_k}^{g_k},$$

where the $P_{a,b,k}$ are rational functions with $P_{0,0,1} = 0$, and $K_1, K_2, K_3 < \infty$.

We will obtain similar theorems for other models, in particular one for bipartite maps (Theorem 4.5), and for triangulations (Theorem 4.11). Moreover, we will in fact prove a version of Theorem 1.2 with control on the number of faces, from which we obtain a closed recurrence formula enumerating one-face maps, small enough to be explicitly written.

THEOREM 1.3 (Ledoux’s recursion for non-oriented one-face maps). *The number u_n^g of rooted non-oriented maps of genus g with n edges and only one face (or equivalently with n*

⁽²⁾They are however not gigantic, see the accompanying worksheet [13]. The main ODE for maps fits in slightly more than a page in \tiny LaTeX print, we have however chosen not to reproduce it here.

edges and only one vertex) is given by the recursion

$$(6) \quad \begin{aligned} (n+1)u_n^g &= (8n-2)u_{n-1}^g - (4n-1)u_{n-1}^{g-1/2} + n(2n-3)(10n-9)u_{n-2}^{g-1} - 8(2n-3)u_{n-2}^g \\ &\quad - 10(2n-3)(2n-4)(2n-5)u_{n-3}^{g-1} + 5(2n-3)(2n-4)(2n-5)u_{n-3}^{g-3/2} \\ &\quad + 8(2n-3)u_{n-2}^{g-1/2} - 2(2n-3)(2n-4)(2n-5)(2n-6)(2n-7)u_{n-4}^{g-2} \end{aligned}$$

with the convention that $u_n^g = 0$ for $g < 0$ and $g > \frac{n}{2}$ and with the initial condition $u_1^{1/2} = 1$, $u_2^1 = u_2^{1/2} = 5$, $u_3^{3/2} = 41$, $u_3^1 = 52$, $u_3^{1/2} = 22$.

The recurrence (6) was first obtained by Ledoux [34] using matrix integral techniques unrelated (as far as we know) to the BKP equation. It is remarkable to see that it is, in fact, the shadow of bigger nonlinear recurrence giving access to an arbitrary number of faces. The Ledoux recurrence can be viewed as a non-oriented version of the infamous Harer–Zagier recurrence, a similar (but smaller) formula which covers the case of orientable one-face maps (and which is itself a special case of the recurrence of [21]). The Harer–Zagier recurrence has a nice analogue in the bipartite case due to Adrianov [2], and it is natural to ask if our non-shifted recursion in the bipartite case implies a non-oriented version of Adrianov’s result. The answer is yes.

THEOREM 1.4 (A recurrence for non-oriented bipartite one-face maps). *The number $b_n^{i,j}$ of rooted one-face maps with n edges, i white and j black vertices, orientable or not, is given by the recursion:*

$$(7) \quad \begin{aligned} (n+1)b_n^{i,j} &= (4n-1)(b_{n-1}^{i-1,j} + b_{n-1}^{i,j-1} - b_{n-1}^{i,j}) + (5n^3 - 16n^2 + 13n - 1)b_{n-2}^{i,j} \\ &\quad + (2n-3)(4b_{n-2}^{i-1,j} + 4b_{n-2}^{i,j-1} - 3b_{n-2}^{i-2,j} + 3b_{n-2}^{i,j-2} - 2b_{n-2}^{i-1,j-1}) \\ &\quad + (10n^3 - 68n^2 + 150n - 107)(b_{n-3}^{i,j} - b_{n-3}^{i-1,j} - b_{n-3}^{i,j-1}) \\ &\quad + (4n-11)(b_{n-3}^{i-3,j} + b_{n-3}^{i,j-3} - b_{n-3}^{i-2,j-1} - b_{n-3}^{i-1,j-2} - b_{n-3}^{i-2,j} - b_{n-3}^{i,j-2} + 2b_{n-3}^{i-1,j-1}) \\ &\quad + (4-n)((2n-7)^2(n-2)^2 b_{n-4}^{i,j} + (5n^2 - 32n + 53)(b_{n-4}^{i-2,j} + b_{n-4}^{i,j-2} - 2b_{n-4}^{i-1,j-1}) \\ &\quad + b_{n-4}^{i-4,j} + b_{n-4}^{i,j-4} - 4b_{n-4}^{i-3,j-1} + 4b_{n-4}^{i-1,j-3} + 6b_{n-4}^{i-2,j-2})) \end{aligned}$$

with the convention that $b_n^{i,j} = 0$ for $i + j > n + 1$, and $b_n^{i,0} = b_n^{0,j} = 0$ and the initial conditions $b_1^{1,1} = b_2^{2,1} = b_2^{1,2} = b_2^{1,1} = 1$, $b_3^{3,1} = b_3^{1,3} = 1$, $b_3^{2,2} = b_3^{2,1} = b_3^{1,2} = 3$, $b_3^{1,1} = 4$.

To conclude this introduction, it is natural to ask if our techniques of shift elimination are specific to the case of maps or apply to general solutions of the BKP hierarchy. The latter is in fact true, and any function $F(N)$ which solves the BKP hierarchy is in fact solution of an explicit (but big) PDE involving only the function $F(N)$ and its derivatives, with no shifts (Theorem B.1 in the appendix). We are not aware of any in-depth study of such “fixed charge” BKP equations, which might be worth considering in the future.

STRUCTURE OF THE PAPER. In Section 2, we will recall what we need about the first BKP equations, directing the reader to other sources for the depth of the BKP theory. In Section 3, we will address the case of maps, taking the time to explain the main ideas and techniques. We will write the Virasoro constraints, and show how to use them to express some derivatives of a specialization of the main BKP tau function as univariate differential polynomials. This will give us “shifted” equations. We will also show how to eliminate the shifts appearing in the BKP equation using instead the first three BKP equations to obtain non-shifted ODEs. In Section 4.1 and Section 4.2, we will address the cases of bipartite maps and triangulations. The main steps are similar to the case of maps and we will give fewer details than in the

previous section. In Section 5, we apply the technique of elimination of variables of the paper [21] to obtain slightly different recurrence formulas than in Section 3 to count maps.

Appendix A contains tables of the numbers of rooted maps and bipartite maps of genus g with n edges and of rooted triangulations of genus g with $2n$ faces, generated with our recurrences. Appendix B derives the fixed charge equation for BKP solutions (Theorem B.1), which we do not use directly in this paper.

Throughout the paper, the notation $\mathbf{R}[\cdot]$, $\mathbf{R}(\cdot)$, $\mathbf{R}[[\cdot]]$ denote respectively polynomials, rational functions, and formal power series with coefficients in the ring \mathbf{R} .

ACCOMPANYING MAPLE WORKSHEET. A Maple worksheet containing an implementation of the recurrences of this paper, together with automated calculations of the bigger ODEs for the different cases (as well as certain proofs regarding their top coefficients) is available in both Maple and html form in [13]. The worksheet also contains recursive programs obtained from these ODEs, as well tables for small genus and consistency checks against existing formulas of the literature.

2. A FEW WORDS ON THE BKP HIERARCHY

In this paper, we will use the BKP hierarchy as a black box, and only recall the statements and equations needed for our purposes. We refer the reader to [30, 42] for the general theory, and to the appendix of our previous paper [15] for details about the applications to maps and bipartite maps.

The BKP hierarchy is an infinite set of partial differential equations (PDEs) for a sequence of functions $\tau(N)_{N \in \mathbb{Z}}$ depending on “time parameters” (formal variables) p_1, p_2, \dots . For our combinatorial purposes, it will be convenient to think of the symbol N as a formal variable rather than an integer, and this turns out to be possible under technical conditions, formalized in the notion of “formal- N ” BKP tau function in [15].

A formal- N BKP tau function is in fact a pair, consisting of a formal power series $\tau(N) \in \mathbb{Q}(N)[[p_1, p_2, \dots]]$, together with a normalizing sequence $(\beta_N)_{N \in \mathbb{Z}}$ which is such that

$$(8) \quad \frac{\beta_{N-1}\beta_{N+k-1}}{\beta_N\beta_{N+k-2}} = R_k(N) \quad , \quad \frac{\beta_{N-2}\beta_{N+k}}{\beta_N\beta_{N+k-2}} = S_k(N)$$

for $N \geq 0$, and for respectively every odd positive integer k and every positive integer k , for some *rational* functions $R_k(N), S_k(N) \in \mathbb{Q}(N)$. These conditions may seem technical but they are crucial to stating the equations of the BKP hierarchy in a formal way as we will do here. In the context of enumeration, the field \mathbb{Q} will often be promoted to a field of rational functions or formal Laurent series involving additional variables, for example $\mathbb{Q}(t)$ so that $R_k(N), S_k(N) \in \mathbb{Q}(t)(N)$.

The typical definition of a (formal or not) BKP tau function makes use of the infinite wedge formalism. It is the image of the orbit of the exponential of an infinite-dimensional Lie algebra, often denoted $b(\infty)$, via the boson-fermion correspondence. We refer the reader to the references mentioned above. For the purposes of this paper, we will admit the PDEs of the BKP hierarchy as a definition:

DEFINITION 2.1. A pair $(\tau(N), \beta_N)$ as above is a formal- N BKP tau function if for $k \in \mathbb{N}, k \geq 1$ the following bilinear identity holds in $\mathbb{C}(N)[\mathbf{p}, \mathbf{q}][[t]]$,

$$(9) \quad \frac{1}{2} \left((-1)^k - 1 \right) R_k(N) U(\mathbf{q}) \tau(N-1) \cdot \tau(N+k-1) \\ + S_k(N) \sum_{j \geq k+1} h_j(2\mathbf{q}) h_{j-k-1}(-\check{\mathbf{D}}) U(\mathbf{q}) \tau(N-2) \cdot \tau(N+k) \\ + \sum_{j \geq 0} h_j(-2\mathbf{q}) h_{j+k-1}(\check{\mathbf{D}}) U(\mathbf{q}) \tau(N) \cdot \tau(N+k-2) = 0$$

where $\mathbf{q} = (q_1, q_2, \dots)$ is a vector of formal indeterminates. Here, h_j denotes the complete homogeneous symmetric function of degree j , and we define $U(\mathbf{q}) = e^{\sum_{r \geq 1} q_r D_r}$ and $\check{\mathbf{D}} = (kD_k)_{k \geq 1}$, where D_r is the Hirota derivative with respect to p_r ,

$$(10) \quad D_r f \cdot g = \frac{\partial}{\partial s_r} f(p_r + s_r)g(p_r - s_r)|_{s_r=0}.$$

By extracting coefficients in the variables q_1, q_2, \dots in (9), one obtains explicit PDEs for the function $\tau(N)$, which altogether form the BKP hierarchy. For example, by setting $k = 2$ and extracting the coefficient of q_3 , we obtain the BKP equation (3) stated in the introduction, where we recall the notation

$$\tau(N) = \exp F(N),$$

in the sense of formal power series, and where indices indicate partial derivatives,

$$f_i := \frac{\partial}{\partial p_i} f.$$

We will only need two other equations of the hierarchy, namely the following bilinear identities valid in $\mathbb{C}(N)[\mathbf{p}, \mathbf{q}][[t]]$:

$$(11) \quad -2F_{4,1}(N) + 2F_{3,2}(N) + 2F_{2,1}(N)F_{1^2}(N) + \frac{1}{3}F_{2,1^3}(N) \\ = S_2(N) \frac{\tau(N-2)\tau(N+2)}{\tau(N)^2} (F_1(N+2) - F_1(N-2)).$$

$$(12) \quad -6F_{5,1}(N) + 4F_{4,2}(N) + 2F_{3^2}(N) + 4F_{3,1}(N)F_{1^2}(N) + \frac{2}{3}F_{3,1^3}(N) + 4F_{2,1}(N)^2 \\ + 2F_{2^2}(N)F_{1^2}(N) + F_{2^2,1^2}(N) + \frac{1}{3}F_{1^2}(N)^3 + \frac{1}{6}F_{1^4}(N)F_{1^2}(N) + \frac{1}{180}F_{1^6}(N) \\ = S_2(N) \frac{\tau(N-2)\tau(N+2)}{\tau(N)^2} (F_{1^2}(N+2) + F_{1^2}(N-2) + 2F_2(N+2) - 2F_2(N-2) \\ + (F_1(N+2) - F_1(N-2))^2),$$

obtained respectively by extracting the coefficient of q_4 and q_5 , again with $k = 2$.

We now proceed with map enumeration.

3. THE CASE OF MAPS

3.1. GENERATING FUNCTIONS OF MAPS. For us a *surface* is a non-oriented two-dimensional real manifold without boundary. A surface of Euler characteristic $2 - 2g$ has *genus* g . A *map* is a graph (with loops and multiple edges allowed) embedded in a surface such that the complement of the embedding is a disjoint collection of contractible components, called *faces*. The *genus* of the map is the one of the underlying surface. A *corner* of a map is a small angular sector around a vertex delimited by two consecutive edge-sides; a corner is *oriented* if an orientation of it (among the two possible ones) is distinguished; the degree of a face/vertex is the number of corners belonging to it/adjacent to it, respectively. In this paper orientable surfaces do not play a particular role, however in some places we will explicitly use the terminology *non-oriented maps* to emphasize that our surfaces can be orientable or not.

We will be interested in enumeration of *rooted maps*, i.e. maps with a distinguished and oriented corner called the *root corner*. Rooted maps are considered up to homeomorphisms preserving the root corner. Define the generating function

$$(13) \quad F(t, \mathbf{p}, u) := \sum_M \frac{t^{2e(M)}}{4e(M)} u^{v(M)} \prod_{f \in F(M)} p_{\deg(f_i)},$$

where we sum over all rooted non-oriented maps, where $F(M)$, $E(M)$, $V(M)$ denote the set of faces, edges and vertices of M , and $f(M)$, $e(M)$, $v(M)$ denote their cardinalities. Note

that since the sum of face degrees in a map is equal to twice the number of edges, F satisfies the *homogeneity relation*

$$(14) \quad t \frac{\partial}{\partial t} F = \sum_{k \geq 1} k p_k \frac{\partial}{\partial p_k} F.$$

The following specialization operator plays a crucial role throughout the paper.

DEFINITION 3.1 (Specialization θ). *We let θ be the operator that specializes all variables p_i to the variable z , namely $\theta(p_i) = z$ for every $i \geq 1$.*

We define the formal power series $\Theta(t, z, u) \in \mathbb{Q}[u, z][[t]]$

$$(15) \quad \Theta(t, z, u) := \theta F(t, \mathbf{p}, u) = \sum_M \frac{t^{2e(M)}}{4e(M)} u^{v(M)} z^{f(M)} = \sum_{n \geq 1} \sum_{i, j \geq 1} \frac{H_n^{i, j}}{4n} t^{2n} u^i z^j$$

which is the bivariate generating function of rooted non-oriented maps M with variables t, u, z marking respectively twice the number of edges, the numbers of vertices, and faces, i.e. $H_n^{i, j}$ denotes the number of rooted non-oriented maps with n edges, i vertices and j faces. It is important to note that a map with n edges, i vertices, and j faces, has Euler characteristic $i - n + j = 2 - 2g$, so the genus is implicitly controlled in this generating function and $\Theta(t, z, u)$ can be rewritten as

$$\Theta(t, z, u) := \sum_{n \geq 1} \sum_{g \geq 0} \frac{H_n^g(u, z)}{4n} t^{2n}, \quad \text{where } H_n^g(u, z) := \sum_{i+j=n+2-2g} H_n^{i, j} u^i z^j.$$

We additionally set $h_n^g := H_n^g(1, 1)$ for the number of rooted, non-oriented maps of genus g with n edges and $u_n^g := H_n^{n+1-2g, 1}$ for the number of rooted, non-oriented maps of genus g with n edges and only one face.

The main goal of this section is to obtain functional equations on the function $\Theta(t, z, u)$, allowing us to compute its coefficients. For this, we start from the fact that the “bigger” function F has a deep structure inherited from the BKP hierarchy, which was proved by [42] using a connection with matrix integrals (see also [15, Appendix] for details on the connection with maps). Here we use the notation $2\mathbf{p} = (2p_1, 2p_2, 2p_3, \dots)$, and Γ denotes the usual gamma function.

PROPOSITION 3.2 ([42]). *Let $\beta_N := 2^{\lfloor \frac{N}{2} \rfloor} \frac{(2\pi)^{\frac{N}{2}} (2t^2)^{\frac{N(N+1)}{4}}}{N! \Gamma(\frac{3}{2})^N} \prod_{j=1}^N \Gamma(1 + \frac{j}{2})$. Then the pair $(\tau(t, 2\mathbf{p}, N) := \exp F(t, 2\mathbf{p}, N), \beta_N)$ is a formal- N tau function of the BKP hierarchy. The function β_N satisfies (8) with in particular $S_2(N) = t^4 N(N - 1)$.*

Proposition 3.2 implies that $F(t, 2\mathbf{p}, N)$ satisfies the BKP equation (3). It is tempting to apply the operator θ to this equation in order to get information on the function $\Theta(t, z, u)$, however because partial derivatives with respect to the p_i do not commute with θ , it is not obvious that such an approach will succeed. For a sequence of non-negative integers $\lambda = (i_1, \dots, i_k)$, we introduce the quantity

$$F_\lambda^\theta \equiv F_\lambda^\theta(t, z, u) := \theta(F_\lambda) = \theta \left(\prod_{j=1}^k \frac{\partial}{\partial p_{i_j}} F(t, \mathbf{p}, u) \right).$$

The F_λ^θ are the quantities naturally appearing when applying θ to the BKP equation (3).

In order to obtain information on the F_λ^θ , we use the fact that $\tau(t, \mathbf{p}, u)$ satisfies the following Virasoro constraints.

PROPOSITION 3.3. [15, Proposition A.1] We have $L_i\tau(t, \mathbf{p}, u) = 0$ for $i \geq -1$, where $(L_i)_{i \geq -1}$ are given by

(16)

$$L_i = \frac{p_{i+2}^*}{t^2} - \left(2 \sum_{\substack{a,b \geq 1 \\ a+b=i}} p_a^* p_b^* + \sum_{a \geq 1} p_a p_{a+i}^* + ((i+1) + 2u)p_i^* + \frac{\delta_{i,-1} u p_1 + u(u+1)\delta_{i,0}}{2} \right)$$

and $p_i^* := \frac{i\partial}{\partial p_i}$ for $i > 0$ and $p_i^* := 0$ for $i < 1$.

Equation (16) has a simple combinatorial interpretation corresponding to the deletion of the root edge in a (non necessarily connected) map whose root face has degree $i + 2$. It is thus closely related to the Tutte/Lehman–Walsh equations. The term “Virasoro constraints” comes from the fact that the operators L_i satisfy the commutation relations of the Virasoro algebra with central charge $c = -2$ [1] – a fact that we will not use here. The Virasoro constraints imply the following proposition:

PROPOSITION 3.4. For $i \geq -1$ and $n_1, n_2, n_3 \geq 0$, one has the recurrence relation

$$(17) \quad \frac{(i+2)F_{i+2,3^{n_3},2^{n_2},1^{n_1}}^\theta}{t^2} = \left(\delta_{i,-1}(\delta_{n_1,1} + z\delta_{n_1,0}) + (u+1)\delta_{i,0}\delta_{n_1,0} \right) \frac{u}{2} \delta_{n_2,0} \delta_{n_3,0} \\ + 2 \sum_{\substack{a+b=i \\ a,b \geq 1}} \sum_{\substack{l_i=0 \\ i=1,2,3}}^{n_i} ab \binom{n_1}{l_1} \binom{n_2}{l_2} \binom{n_3}{l_3} F_{a,3^{l_3},2^{l_2},1^{l_1}}^\theta F_{b,3^{n_3-l_3},2^{n_2-l_2},1^{n_1-l_1}}^\theta \\ + 2 \sum_{\substack{a+b=i \\ a,b \geq 1}} ab F_{a,b,3^{n_3},2^{n_2},1^{n_1}}^\theta + \sum_{j=1}^3 n_j(i+j) F_{i+j,3^{n_3-\delta_{3,j}},2^{n_2-\delta_{2,j}},1^{n_1-\delta_{1,j}}}^\theta + t \frac{\partial}{\partial t} F_{3^{n_3},2^{n_2},1^{n_1}}^\theta \\ - (n_1 + 2n_2 + 3n_3) F_{3^{n_3},2^{n_2},1^{n_1}}^\theta - z \sum_{a=1}^i a F_{a,3^{n_3},2^{n_2},1^{n_1}}^\theta + \delta_{i \neq -1} (2u+i+1) i F_{i,3^{n_3},2^{n_2},1^{n_1}}^\theta,$$

with the initial condition that $F_{3^0 2^0 1^0}^\theta = F_\emptyset^\theta = \Theta(t, z, u)$.

Consequently, for any integer vector λ of the form $\lambda = [\ell, 3^{n_3}, 2^{n_2}, 1^{n_1}]$ with $\ell \leq 9$ and of size $|\lambda| = \ell + n_1 + 2n_2 + 3n_3$, there exists a polynomial P_λ in $|\lambda|$ variables, with coefficients in $\mathbb{Q}[t, u, z]$, such that

$$(18) \quad F_\lambda^\theta = P_\lambda \left(\frac{\partial}{\partial t} \Theta(t, z, u), \dots, \frac{\partial^{|\lambda|}}{\partial t^{|\lambda|}} \Theta(t, z, u) \right).$$

This polynomial is linear for $\ell \leq 3$ and $|\lambda| \leq 5$, and quadratic for $4 \leq \ell \leq 6$ and $|\lambda| \leq 6$.

Proof. The constraints read explicitly

$$(19) \quad (i+2)F_{i+2} = t^2 \left(2 \sum_{a+b=i} ab(F_{a,b} + F_a F_b) + \sum_{a \geq 1} p_a(i+a)F_{i+a} \right) \\ + t^2(2u+i+1)iF_i + t^2 \frac{u(u+1)}{2} \delta_{i,0} + t^2 \delta_{i,-1} \frac{p_1 u}{2}.$$

We act on both sides of (19) with $\frac{\partial^{n_1+n_2+n_3}}{\partial p_1^{n_1} \partial p_2^{n_2} \partial p_3^{n_3}}$ and apply θ . The action on $\sum_{a \geq 1} p_a(i+a)F_{i+a}$ can be re-written using the following equation:

$$\theta \sum_{a \geq 1} p_a(i+a)F_{i+a,3^{n_3},2^{n_2},1^{n_1}} = z \sum_{a \geq 1} (i+a)F_{i+a,3^{n_3},2^{n_2},1^{n_1}}^\theta \\ = t \frac{\partial F_{3^{n_3},2^{n_2},1^{n_1}}^\theta}{\partial t} - (n_1 + 2n_2 + 3n_3) F_{3^{n_3},2^{n_2},1^{n_1}}^\theta - z \sum_{a=1}^i a F_{a,3^{n_3},2^{n_2},1^{n_1}}^\theta.$$

It itself comes from the homogeneity relation (14) by acting with $\frac{\partial^{n_1+n_2+n_3}}{\partial p_1^{n_1} \partial p_2^{n_2} \partial p_3^{n_3}}$ and applying θ . This produces (17).

Note that the size of all vectors indexing F^θ s appearing in the RHS of (17) are strictly smaller than the size $i + 2 + n_1 + 2n_2 + 3n_3$ in the LHS. In order to be able to iterate this equation on all these terms, we need all vectors appearing in the RHS to have at most one part larger than 3. The only term on the RHS that could have two parts larger than 3 is of the form $F_{a,b,3n_3,2n_2,1n_1}^\theta$. Since $a + b = i$, this does not happen unless $i + 2 > 9$. We thus obtain (18). The last statement is a direct check. \square

REMARK 3.5. The Virasoro constraints in fact imply a more general result: F_λ^θ for any λ is a differential polynomial of Θ . This is proved by applying $\prod_{j=1}^k \frac{\partial}{\partial p_{i_j}}$ to (19) and performing an induction on $|\lambda|$. We will refrain from writing it since we will not need it in full generality. Instead, the previous proposition is enough to cover the cases we need with explicit formulas, involving vectors of size $|\lambda| \leq 6$.

We insist on the fact that the recurrence given in Proposition 3.4 can be fully automated to compute the polynomials P_λ , and this is done in the accompanying Maple worksheet [13].

It is now immediate to see that applying the operator θ to the BKP equation (3) produces a functional equation on $\Theta(t, z, u)$.

THEOREM 3.6. *The generating function $\Theta(t, z, u)$ satisfies a functional equation of the following form:*

$$\begin{aligned} \left(\frac{\partial}{\partial t} (\Theta(t, z, u + 2) + \Theta(t, z, u - 2) - 2\Theta(t, z, u)) \right) P \left(\frac{\partial}{\partial t} \Theta(t, z, u), \dots, \frac{\partial^5}{\partial t^5} \Theta(t, z, u) \right) \\ = Q \left(\frac{\partial}{\partial t} \Theta(t, z, u), \dots, \frac{\partial^5}{\partial t^5} \Theta(t, z, u) \right), \end{aligned}$$

where P and Q are quadratic polynomials with coefficients in $\mathbb{Q}[t, u, z]$.

Proof. Consider (3) for $\tau(t, 2\mathbf{p}, u)$. Applying θ to both sides we get:

$$4F_{2^2}^\theta - 4F_{3,1}^\theta + \frac{4}{3}(6(F_{1^2}^\theta)^2 + F_{1^4}^\theta) = S_2(u)e^{\Theta(t,z,u+2)+\Theta(t,z,u-2)-2\Theta(t,z,u)}.$$

The difference between the LHS in the equation above and the LHS in (3) is due to the fact that $\tau(N)$ is a formal- N tau function of the BKP hierarchy after rescaling the variables $\mathbf{p} \rightarrow 2\mathbf{p}$. Taking the derivative with respect to t and substituting the initial equation (3) back into it to eliminate exponentials, we obtain

$$\begin{aligned} (20) \quad \frac{\partial}{\partial t} \left(4F_{2^2}^\theta - 4F_{3,1}^\theta + \frac{4}{3}(6(F_{1^2}^\theta)^2 + F_{1^4}^\theta) \right) - \frac{4}{t} \left(4F_{2^2}^\theta - 4F_{3,1}^\theta + \frac{4}{3}(6(F_{1^2}^\theta)^2 + F_{1^4}^\theta) \right) = \\ \left(\frac{\partial}{\partial t} (\Theta(t, z, u + 2) + \Theta(t, z, u - 2) - 2\Theta(t, z, u)) \right) \left(4F_{2^2}^\theta - 4F_{3,1}^\theta + \frac{4}{3}(6(F_{1^2}^\theta)^2 + F_{1^4}^\theta) \right) \end{aligned}$$

thanks to the identity $\frac{\partial}{\partial t} S_2(u) = \frac{4}{t} S_2(u)$ implied by $S_2(u) = t^4 u(u - 1)$. Proposition 3.4 immediately concludes the proof. \square

The functional equation above can be transformed into a recurrence to compute coefficients. It has the following relatively compact form:

THEOREM 3.7 (Counting maps by vertices, faces, and genus). *The generating polynomial*

$$H_n^g \equiv H_n^g(u, z) = \sum_{i+j=n+2-2g} H_n^{i,j} u^i z^j$$

of rooted non-oriented maps of genus g with n edges, with weight u per vertex and z per face, can be computed from the following recurrence formula:

$$\begin{aligned}
 H_n^g + \frac{3u^2 \partial^2}{n(n+1) \partial u^2} H_n^g &= \frac{1}{n(n+1)} \times \left(n \left(2(2n-1) \left((4u+z) H_{n-1}^g - 2H_{n-1}^{g-1/2} \right) + 4(2n-3) \left(3uz H_{n-2}^g \right. \right. \right. \\
 &+ (2n-1)(n-1) H_{n-2}^{g-1} \left. \left. \left. + 6 \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n \\ n_1+n_2=n}} (2n_1-1)(2n_2-1) H_{n_2-1}^{g_2} H_{n_1-1}^{g_1} \right) \right) \\
 &- \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=1..n \\ n_1+n_2=n}} \sum_{\substack{g_0=0..g_1 \\ g_1-g_0 \in \mathbb{N}}} \sum_{p+j=n_1+2-2g_0} 2^{2(1+g_1-g_0)} \binom{p}{2(1+g_1-g_0)} u^{n_1-2g_1-j} z^j H_{n_1}^{p,j} \\
 &\left(-\frac{n_2+1}{2} H_{n_2}^{g_2} + (2n_2-1) \left((4u+z) H_{n_2-1}^{g_2} - 2H_{n_2-1}^{g_2-1/2} \right) + 2(2n_2-3) \left((2n_2-1)(n_2-1) H_{n_2-2}^{g_2-1} \right. \right. \\
 &+ 3uz H_{n_2-2}^{g_2} \left. \left. + \frac{3}{2} \delta_{g_0 \neq g} \delta_{n_1, n} u (\delta_{g_1, g} u - \delta_{g_1, g-1/2}) + \delta_{n_1, n-1} uz (\delta_{g_1, g} (4u+z) - 2\delta_{g_1, g-1/2}) \right) \right) \\
 &+ 3uz \delta_{n_1, n-2} (\delta_{g_1, g} uz + 2\delta_{g_1, g-1}) + 3 \sum_{\substack{g_3=0..g_2 \\ g_3+g_4=g_2}} \sum_{\substack{n_3=0..n_2 \\ n_3+n_4=n_2}} (2n_3-1)(2n_4-1) H_{n_4-1}^{g_4} H_{n_3-1}^{g_3} \left. \right) \Big)
 \end{aligned}$$

for $n > 2$, with the initial conditions $H_0^g = 0$, $H_1^0 = uz(u+z)$, $H_2^0 = uz(2u^2 + 5uz + 2z^2)$, $H_1^{1/2} = uz$, $H_2^{1/2} = 5uz(u+z)$, $H_2^1 = 5uz$, and $H_n^g = 0$ if $n < 2g$.

Proof. We extract the coefficient of $[t^{2n+4} r^{n+2-2g}]$ in (20) after substitution $u \rightarrow ur, z \rightarrow zr$. We obtain from Proposition 3.4

$$\begin{aligned}
 &n \left(-(n+1) H_n^g + 2(2n-1) \left((4u+z) H_{n-1}^g - 2H_{n-1}^{g-1/2} \right) + 4(2n-3) \left((2n-1)(n-1) H_{n-2}^{g-1} \right. \right. \\
 &\quad \left. \left. + 3uz H_{n-2}^g \right) + 6 \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n \\ n_1+n_2=n}} (2n_1-1)(2n_2-1) H_{n_2-1}^{g_2} H_{n_1-1}^{g_1} \right) \\
 &= \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=1..n \\ n_1+n_2=n}} \sum_{\substack{g_0=0..g_1 \\ g_1-g_0 \in \mathbb{N}}} \sum_{p+j=n_1+2-2g_0} 2^{2(1+g_1-g_0)} \binom{p}{2(1+g_1-g_0)} u^{n_1-2g_1-j} z^j H_{n_1}^{p,j} \\
 &\left(-\frac{n_2+1}{2} H_{n_2}^{g_2} + (2n_2-1) \left((4u+z) H_{n_2-1}^{g_2} - 2H_{n_2-1}^{g_2-1/2} \right) + 2(2n_2-3) \left((2n_2-1)(n_2-1) H_{n_2-2}^{g_2-1} \right. \right. \\
 &\quad \left. \left. + 3uz H_{n_2-2}^{g_2} \right) + \frac{3}{2} \delta_{n_1, n} u (\delta_{g_1, g} u - \delta_{g_1, g-1/2}) + \delta_{n_1, n-1} uz (\delta_{g_1, g} (4u+z) - 2\delta_{g_1, g-1/2}) \right) \\
 &+ 3uz \delta_{n_1, n-2} (\delta_{g_1, g} uz + 2\delta_{g_1, g-1}) + 3 \sum_{\substack{g_3=0..g_2 \\ g_3+g_4=g_2}} \sum_{\substack{n_3=0..n_2 \\ n_3+n_4=n_2}} (2n_3-1)(2n_4-1) H_{n_4-1}^{g_4} H_{n_3-1}^{g_3} \Big).
 \end{aligned}$$

Here we have extracted, respectively, in (20) (after substitution $u \rightarrow ur, z \rightarrow zr$), the coefficient of $[t^{2n+4} r^{n+2-2g}]$, of $[t^{2n_1-1} r^{n_1-2g_1}]$, and of $[t^{2n_2+5} r^{n_2+2-2g_2}]$, in the RHS, in the

first factor of the LHS, and in the second factor of the LHS. Moreover we have used

$$\begin{aligned}
 & [r^m](f(zr, ur + 2) + f(zr, ur - 2) - 2f(zr, ur)) \\
 &= \sum_{i+j=m} u^i z^j \sum_{p>i} \binom{p}{i} (2^{p+j-m} + (-2)^{p+j-m}) [u^p z^j] f(z, u) \\
 &= \sum_{i+j=m} u^i z^j \sum_{\substack{p \geq i+2, \\ p+j-m \in 2\mathbb{N}}} \binom{p}{i} 2^{p+j-m+1} [u^p z^j] f(z, u) \\
 &= \sum_{\substack{k \geq m+2 \\ k-m \in 2\mathbb{N}}} 2^{1+k-m} \sum_{p+j=k} \binom{p}{m-j} u^{m-j} z^j [u^p z^j] f(z, u).
 \end{aligned}$$

In our case $m = n_1 - 2g_1$, and we parametrized k as $k = n_1 + 2 - 2g_0$ (the condition $k \geq m + 2$ translates into $g_0 \leq g_1$, and the summand is null when $g_0 < 0$).

It now only remains to group the terms of the form $H_n^{i,j}$ with $i + j = n + 2 - 2g$. In the LHS they contribute to the first term H_n^g and in the RHS they appear as the terms $H_{n_1}^{p,j}$ when $n_1 = n, n_2 = 0, g_1 = g, g_2 = 0, g_0 = g$. Collecting these terms on the RHS gives

$$\begin{aligned}
 \frac{3}{2} \delta_{n_1, n} \delta_{g_1, g} \delta_{g_0, g} u^2 \sum_{p+j=n_1+2-2g_0} 2^{2(1+g_1-g_0)} \binom{p}{2(1+g_1-g_0)} u^{n_1-2g_1-j} z^j H_{n_1}^{p,j} \\
 = 6 \sum_{p+j=n+2-2g} \binom{p}{2} u^p z^j H_n^{p,j} = 3 \frac{u^2 \partial^2}{\partial u^2} H_n^g,
 \end{aligned}$$

which leads to the main equation of the theorem. The identification of the initial conditions for $n \geq 2$ can be done, either: by hand drawing, or from the OEIS, or from explicit expansions in small genera using the equations of this paper, or from the expansion in Zonal polynomials up to order $n = 2$. □

3.2. REMOVING THE SHIFTS. We now proceed with the task of obtaining a functional equation on the function $\Theta(t, z, u)$ which does not involve any shift on the variable u . We will do this by using the three equations (3), (11), (12) to eliminate the shifts, and apply the operator θ . This will make terms of the form F_λ^θ appear, with larger partitions λ than in the previous section, but fortunately they are still in the range covered by Proposition 3.4. We have

THEOREM 3.8. *There exists a polynomial $P \in \mathbb{Q}[t, u, z][x_1, \dots, x_6]$ of degree 5 such that*

$$P \left(\frac{\partial}{\partial t} \Theta(t, z, u), \dots, \frac{\partial^6}{\partial t^6} \Theta(t, z, u) \right) \equiv 0.$$

An explicit form of P can be obtained by applying Proposition 3.4 to the following equation

$$\begin{aligned}
 (21) \quad & t^6 \left(\frac{\partial}{\partial t} \text{KP1} \right)^2 - \left(\text{KP2} \right)^2 + \text{KP1} \left(\text{KP3} - \frac{1}{2} \text{KP2} - \right. \\
 & \left. - \left(t^6 \frac{\partial^2}{\partial t^2} + 2t^5 \frac{\partial}{\partial t} + 2t^6 \frac{\partial^2}{\partial t^2} \Theta + 4t^5 \frac{\partial}{\partial t} \Theta + t^4(uz - 4) + t^2(3u + 1 - z) \right) \text{KP1} \right) \equiv 0,
 \end{aligned}$$

where

$$\begin{aligned} \text{KP1} &= -4F_{3,1}^\theta + 4F_{2^2}^\theta + \frac{4}{3}(6(F_{1^2}^\theta)^2 + F_{1^4}^\theta), \\ \text{KP2} &= -4F_{4,1}^\theta + 4F_{3,2}^\theta + \frac{8}{3}(6F_{2,1}^\theta F_{1^2}^\theta + F_{2,1^3}^\theta), \\ \text{KP3} &= -6F_{5,1}^\theta + 4F_{4,2}^\theta + 2F_{3,3}^\theta + \frac{8}{3}(6F_{3,1}^\theta F_{1^2}^\theta + F_{3,1^3}^\theta) + 4(4(F_{2,1}^\theta)^2 + 2F_{2^2}^\theta F_{1^2}^\theta + F_{2^2,1^2}^\theta) \\ &\quad + \frac{4}{45}(60(F_{1^2}^\theta)^3 + 30F_{1^4}^\theta F_{1^2}^\theta + F_{1^6}^\theta), \end{aligned}$$

and $\Theta \equiv \Theta(t, z, u)$. An explicit form of this ODE can be found in the accompanying Maple worksheet [13].

Proof. Denote $\Delta f(u) = f(u + 2) - f(u - 2)$ and $\nabla f = f(u + 2) + f(u - 2)$, and

$$E = S_2(u)e^{\nabla\Theta(t,z,u) - 2\Theta(t,z,u)}.$$

Using (3), (11), (12) and applying θ we obtain the following equations

$$(22) \quad E = \text{KP1}, \quad E\Delta(F_1^\theta) = \text{KP2}, \quad E(\nabla(F_{1^2}^\theta) + \Delta(F_2^\theta) + (\Delta F_1^\theta)^2) = \text{KP3},$$

where KP1, KP2, KP3 are given in the statement of theorem. The difference between KP1, KP2, KP3 and the LHS in (3), (11), (12) is due to the fact that $\tau(N)$ is a formal- N tau function of the BKP hierarchy after rescaling the variables $\mathbf{p} \rightarrow 2\mathbf{p}$. Using (17) and the identity $S_2(u) = t^4u(u - 1)$ we have

$$\Delta(F_1^\theta) = t^3\Delta\frac{\partial}{\partial t}\Theta(t, z, u) + 2t^2z, \quad \Delta(F_2^\theta) = \frac{t^3}{2}\Delta\frac{\partial}{\partial t}\Theta(t, z, u) + t^2(2u + 1)$$

and

$$\nabla(F_{1^2}^\theta) = (t^6\frac{\partial^2}{\partial t^2} + 2t^5\frac{\partial}{\partial t})\nabla\Theta(t, z, u) + t^4uz + t^2u$$

so that the third BKP equation reads

$$(23) \quad E\left(\left(t^6\frac{\partial^2}{\partial t^2} + 2t^5\frac{\partial}{\partial t}\right)\nabla\Theta(t, z, u) + \frac{t^3}{2}\Delta\frac{\partial}{\partial t}\Theta(t, z, u) + t^4uz + t^2(3u + 1) + \left(t^3\Delta\frac{\partial}{\partial t}\Theta(t, z, u) + 2t^2z\right)^2\right) = \text{KP3}.$$

We now use the first two BKP equations to express $\nabla\frac{\partial}{\partial t}\Theta(t, z, u)$, $\nabla\frac{\partial^2}{\partial t^2}\Theta(t, z, u)$ and $\Delta\frac{\partial}{\partial t}\Theta(t, z, u)$ in terms of $\Theta(t, z, u)$ and its t -derivatives. Taking the t -derivative of the first BKP equation, we have

$$E\nabla\frac{\partial}{\partial t}\Theta(t, z, u) = \frac{\partial}{\partial t}\text{KP1} + \left(2\frac{\partial}{\partial t}\Theta(t, z, u) - \frac{4}{t}\right)\text{KP1},$$

and another derivative gives

$$\begin{aligned} E\nabla\frac{\partial^2}{\partial t^2}\Theta(t, z, u) &= \frac{\partial^2}{\partial t^2}\text{KP1} + \left(2\frac{\partial}{\partial t}\Theta(t, z, u) - \frac{4}{t}\right)\frac{\partial}{\partial t}\text{KP1} + \left(2\frac{\partial^2}{\partial t^2}\Theta(t, z, u) + \frac{4}{t^2}\right)\text{KP1} \\ &\quad - \frac{\partial}{\partial t}\text{KP1}\nabla\frac{\partial}{\partial t}\Theta(t, z, u) \end{aligned}$$

and further

$$E^2\nabla\frac{\partial^2}{\partial t^2}\Theta(t, z, u) = \text{KP1}\frac{\partial^2}{\partial t^2}\text{KP1} + \left(2\frac{\partial^2}{\partial t^2}\Theta(t, z, u) + \frac{4}{t^2}\right)\text{KP1}^2 - \left(\frac{\partial}{\partial t}\text{KP1}\right)^2.$$

From the second BKP equation,

$$Et^3\Delta\frac{\partial}{\partial t}\Theta(t, z, u) = -2t^2z\text{KP1} + \text{KP2}.$$

Those expressions can then be substituted into (23), which gives (21). The statement about the form of polynomial P is a direct consequence of Proposition 3.4. \square

Theorem 1.2 is an immediate consequence of Theorem 3.8.

Proof of Theorem 1.2. It is enough to make the change of variables $u \rightarrow ur, z \rightarrow zr$ in (21) and extract the coefficient of $[t^{2n}r^{n+2g-2}]$. This substitution allows to track the genus of the underlying maps. Extracting the coefficient gives a recursion for the bivariate version of h_n^g which additionally tracks the number of vertices and faces via u and z . Specializing $u = z = 1$ gives the recursion for h_n^g of the form (5) with a, b depending on the specific form of ODE given by (21). A direct examination of the highest degree terms of this recurrence implemented in [13] shows that

$$\begin{aligned}
 h_n^g &= h_n^{g-1/2} - \sum_{\substack{n_1=1..n-1 \\ n_1+n_2=n}} \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \frac{(n_1+1)(n_2+1)}{42(n+1)} h_{n_1}^{g_1} h_{n_2}^{g_2} \\
 &+ \sum_{a=1}^{K_1} \sum_{b=0}^{K_2} \sum_{k=1}^{K_3} \sum_{\substack{n_1, \dots, n_k \geq 1 \\ n_1 + \dots + n_k = n-a}} \sum_{\substack{g_1, \dots, g_k \geq 0 \\ g_1 + \dots + g_k = g-b}} P_{a,b,k}(n_1, \dots, n_k) h_{n_1}^{g_1} h_{n_2}^{g_2} \dots h_{n_k}^{g_k},
 \end{aligned}$$

which finishes the proof. \square

The coefficient of z^1 in $\Theta(t, z, u)$ is the generating function of maps having only one face, with control on the number of edges and vertices (equivalently, edges and genus). Extracting the bottom coefficient in z in (21), we obtain a linear ODE for this generating function. It is equivalent to Ledoux’s recurrence (6) stated in the introduction.

COROLLARY 3.9. *The generating function*

$$u(t, u) := [z]\Theta(t, z, u) = \sum_{n \geq 1} \sum_{g \geq 0} \frac{u_n^g}{4n} t^{2n} u^{n+1-2g}$$

of rooted non-oriented maps with only one face satisfies the following linear ODE

$$\begin{aligned}
 (24) \quad &(32t^4(u^2 - u - 5) + 240t^6(2u - 1) + t^2(10 - 20u) + 2880t^8 + 3) \frac{\partial}{\partial t} u \\
 &+ t(2t^4(8u^2 - 8u - 109) + 360t^6(2u - 1) + t^2(4 - 8u) + 7200t^8 + 1) \frac{\partial^2}{\partial t^2} u \\
 &+ 6t^6(20t^2(2u - 1) + 800t^4 - 11) \frac{\partial^3}{\partial t^3} u + 5t^7(-1 + 2t^2(2u - 1) + 240t^4) \frac{\partial^4}{\partial t^4} u \\
 &+ 120t^{12} \frac{\partial^5}{\partial t^5} u + 4t^{13} \frac{\partial^6}{\partial t^6} u + 240t^7 u + 30t^5(2u^2 - u) + 2t^3(4u^3 - 4u^2 - 11u) - 2t(u^2 + u) \equiv 0.
 \end{aligned}$$

4. RECURRENCES FOR BIPARTITE MAPS AND TRIANGULATIONS

4.1. NON-ORIENTED BIPARTITE MAPS. Consider the generating function

$$(25) \quad G(t, \mathbf{p}, u, v) := \sum_M \frac{t^{e(M)}}{2e(M)} u^{v_o(M)} v^{v_\bullet(M)} \prod_{f \in F(M)} p_{\deg(f_i)},$$

where we sum over all rooted non-oriented bipartite maps, and $v_o(M), v_\bullet(M)$ denote the number of white and black vertices, respectively. Similarly, as in the case of general maps, the function G inherits a deep structure from the BKP hierarchy. This result can be derived directly from Van de Leur’s work [42], even though it is not stated explicitly there (see [15, Appendix] for additional details on the connection with maps).

PROPOSITION 4.1 ([42]). Let $\beta_N = \frac{2^{\lfloor \frac{N}{2} \rfloor} (2t)^{\frac{N(N+\delta)}{2}}}{N! \Gamma(3/2)^N} \prod_{j=1}^N \Gamma(1 + \frac{j}{2}) \Gamma(\frac{\delta+j}{2})$. Then the pair $(\tau(t, 2\mathbf{p}, \delta, N) := \exp G(t, 2\mathbf{p}, N, N+\delta), \beta_N)$ is a formal- N tau function of the BKP hierarchy. The function β_N satisfies (8) with in particular $S_2(N) = t^4 N(N+\delta)(N-1)(N+\delta-1)$.

We recall that $\theta(p_i) = z$ for $i \geq 1$. Define the power series $\eta(t, z, u, v) \in \mathbb{Q}[u, v, z][[t]]$

$$(26) \quad \eta(t, z, u, v) := \theta G(t, \mathbf{p}, u, v) = \sum_M \frac{t^{e(M)}}{2e(M)} u^{v \bullet (M)} v^{v \circ (M)} z^{f(M)} = \sum_{n, i, j, k \geq 1} \frac{K_n^{i, j, k}}{2n} t^n u^i v^j z^k,$$

which is the generating function of rooted, non-oriented bipartite maps M . The variables t, u, v, z mark the number of edges, black vertices, white vertices and faces, respectively so that $K_n^{i, j, k}$ denotes the number of rooted non-oriented bipartite maps with n edges, i black vertices, j white vertices and k faces (the root vertex is black by convention). Note that due to the Euler relation we can rewrite $\eta(t, z, u, v)$ so that it is parametrized by the number of edges and genus:

$$\eta(t, z, u, v) := \sum_{n \geq 1} \sum_{g \geq 0} \frac{K_n^g(u, v, z)}{2n} t^n, \text{ where } K_n^g(u, v, z) := \sum_{i+j+k=n+2-2g} K_n^{i, j, k} u^i v^j z^k.$$

We additionally set $\mathfrak{k}_n^g := K_n^g(1, 1, 1)$ for the number of rooted non-oriented bipartite maps of genus g with n edges and $\mathfrak{b}_n^{i, j} := K_n^{i, j, 1}$ for the number of rooted non-oriented bipartite maps of genus g with n edges, i black and j white vertices, and only one face.

In analogy with Proposition 3.4 we express G_λ^θ in terms of $\frac{\partial^i}{\partial t^i} \eta(t, z, u, v)$, where

$$G_\lambda^\theta \equiv G_\lambda^\theta(t, z, u, v) := \theta(G_\lambda) = \theta \left(\prod_{j=1}^k \frac{\partial}{\partial p_{i_j}} G(t, \mathbf{p}, u, v) \right)$$

for a sequence of non-negative integers $\lambda = (i_1, \dots, i_k)$.

PROPOSITION 4.2. For $i \geq 0$ and $n_1, n_2, n_3 \geq 0$, one has the recurrence relation

$$(27) \quad \frac{(i+1)G_{i+1, 3^{n_3}, 2^{n_2}, 1^{n_1}}^\theta}{t} = \frac{uv}{2} \delta_{i,0} \delta_{n_1,0} \delta_{n_2,0} \delta_{n_3,0} \\ + 2 \sum_{\substack{a+b=i \\ a, b \geq 1}} \sum_{\substack{n_i \\ l_i=0 \\ i=1,2,3}} ab \binom{n_1}{l_1} \binom{n_2}{l_2} \binom{n_3}{l_3} G_{a, 3^{l_3}, 2^{l_2}, 1^{l_1}}^\theta G_{b, 3^{n_3-l_3}, 2^{n_2-l_2}, 1^{n_1-l_1}}^\theta \\ + 2 \sum_{\substack{a+b=i \\ a, b \geq 1}} ab G_{a, b, 3^{n_3}, 2^{n_2}, 1^{n_1}}^\theta + \sum_{j=1}^3 n_j (i+j) G_{i+j, 3^{n_3-\delta_{3,j}}, 2^{n_2-\delta_{2,j}}, 1^{n_1-\delta_{1,j}}}^\theta + t \frac{\partial}{\partial t} G_{3^{n_3}, 2^{n_2}, 1^{n_1}}^\theta \\ - (n_1 + 2n_2 + 3n_3) G_{3^{n_3}, 2^{n_2}, 1^{n_1}}^\theta - z \sum_{a=1}^i a G_{a, 3^{n_3}, 2^{n_2}, 1^{n_1}}^\theta + (u+v+i) i G_{i, 3^{n_3}, 2^{n_2}, 1^{n_1}}^\theta,$$

with the convention that $G_{3^0, 2^0, 1^0}^\theta = G_\emptyset^\theta = \eta(t, z, u, v)$.

Consequently, for any partition λ of the form $\lambda = [\ell, 3^{n_3}, 2^{n_2}, 1^{n_1}]$ and of size $|\lambda| = \ell + n_1 + 2n_2 + 3n_3$, there exists a polynomial Q_λ in $|\lambda|$ variables, with coefficients in $\mathbb{Q}[t, u, v, z]$ which is linear for $\ell \leq 2$ and $|\lambda| \leq 5$, and quadratic for $\ell \geq 3, 4 \leq |\lambda| \leq 6$ and satisfies

$$(28) \quad G_\lambda^\theta = Q_\lambda \left(\frac{\partial}{\partial t} \eta(t, z, u, v), \dots, \frac{\partial^{|\lambda|}}{\partial t^{|\lambda|}} \eta(t, z, u, v) \right).$$

The proof is identical to the proof of Proposition 3.4 (and left to the reader) with the only difference being in replacing Proposition 3.3 by its bipartite counterpart:

PROPOSITION 4.3. [15, Proposition A.1] *We have $L_i\tau(t, \mathbf{p}, u, v) = 0$ for $i \geq 0$, where $(L_i)_{i \geq 0}$ are given by*

$$L_i = \frac{p_{i+1}^*}{t} - \left(2 \sum_{\substack{a,b \geq 1 \\ a+b=i}} p_a^* p_b^* + \sum_{a \geq 1} p_a p_{a+i}^* + (i+u+v)p_i^* + \frac{uv\delta_{i,0}}{2} \right).$$

where $p_i^* := \frac{i\partial}{\partial p_i}$ for $i > 0$ and $p_i^* := 0$ for $i < 1$.

THEOREM 4.4 (Counting bipartite maps by black/white vertices, faces, and genus). *The generating polynomial*

$$K_n^g \equiv K_n^g(u, v, z) = \sum_{i+j+k=n+2-2g} K_n^{i,j,k} u^i v^j z^k$$

of rooted non-oriented bipartite maps of genus g with n edges, with weight u per black vertex, v per white vertex and z per face, can be computed from the following recurrence formula:

$$\begin{aligned} K_n^g &= \frac{1}{(n+1)} \left((2n-1) \left((u+v+z)K_{n-1}^g - K_{n-1}^{g-1/2} \right) - \psi_n(u, v, z)K_{n-2}^g + (2n-1)(2n-3)nK_{n-2}^{g-1} \right. \\ &\quad \left. - 6(n-1)(u+v-z)K_{n-2}^{g-1/2} + 2 \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n \\ n_1+n_2=n}} (6n_1n_2 - 2(n_1+n_2) + 1)K_{n_2-1}^{g_2}K_{n_1-1}^{g_1} \right) \\ &\quad - \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=1..n-1 \\ n_1+n_2=n}} \sum_{\substack{g_0=0..g_1 \\ g_1-g_0 \in \mathbb{N}}} \frac{2^{2(1+g_1-g_0)}}{(n-2)(n+1)} \sum_{\substack{p+q+k= \\ n_1+2-2g_0}} \sum_{i=0}^{n_1-2g_1-k} \binom{p}{i} \binom{q}{n_1-2g_1-k-i} u^i v^{n_1-2g_1-k-i} z^k K_{n_1}^{p,q,k} \\ &\quad \left(-(n_2+1)K_{n_2}^{g_2} + (2n_2-1) \left((u+v+z)K_{n_2-1}^{g_2} - K_{n_2-1}^{g_2-1/2} \right) + (2n_2-1)(2n_2-3)n_2K_{n_2-2}^{g_2-1} \right. \\ &\quad \left. - 6(n_2-1)(u+v-z)K_{n_2-2}^{g_2-1/2} + \delta_{n_2,1}\delta_{g_2,0}2uvz + 6uv\delta_{n_2,2}(\delta_{g_2,0}uv - \delta_{g_2,1/2}(u+v-z) + \delta_{g_2,1}) \right. \\ &\quad \left. - \psi_{n_2}(u, v, z)K_{n_2-2}^{g_2} + 2 \sum_{\substack{g_3=0..g \\ g_3+g_4=g_2}} \sum_{\substack{n_3=0..n_2 \\ n_3+n_4=n_2}} (6n_3n_4 - 2(n_3+n_4) + 1)K_{n_3-1}^{g_3}K_{n_4-1}^{g_4} \right) \end{aligned}$$

for $n > 2$, with

$$\psi_n(u, v, z) := (n-2)(u^2 + v^2 + z^2 - 14uv - 2uz - 2vz) - 12uv$$

and the initial conditions $K_0^g = 0$, $K_1^0 = uvz$, $K_2^0 = uvz(u+v+z)$, $K_1^{1/2} = 0$, $K_2^{1/2} = uvz$, $K_2^1 = 0$, and $K_n^g = 0$ if $n < 2g$.

Proof. The proof is almost identical to the proof of Theorem 3.7. The only difference is that (20) should be replaced by

$$\begin{aligned} &\left(\frac{\partial}{\partial t} \left(\eta(t, z, u+2, v+2) + \eta(t, z, u-2, v-2) - 2\eta(t, z, u, v) \right) \right) \left(G_{22}^\theta - G_{3,1}^\theta + \frac{1}{3}(6(G_{12}^\theta)^2 + G_{14}^\theta) \right) \\ &= \frac{\partial}{\partial t} \left(G_{22}^\theta - G_{3,1}^\theta + \frac{1}{3}(6(G_{12}^\theta)^2 + G_{14}^\theta) \right) - \frac{1}{t} \left(4G_{22}^\theta - 4G_{3,1}^\theta + \frac{4}{3}(6(G_{12}^\theta)^2 + G_{14}^\theta) \right). \end{aligned}$$

The computational details are left to the reader. □

As in the case of maps, it is possible to manipulate the first three BKP equations and obtain an ordinary differential equation on $\eta(t, z, u, v)$. In particular, it does not involve any shifts on the variables u and v .

THEOREM 4.5. *There exists a polynomial $Q \in \mathbb{Q}[t, u, v, z][x_1, \dots, x_6]$ of degree 5 such that*

$$Q \left(\frac{\partial}{\partial t} \eta(t, z, u, v), \dots, \frac{\partial^6}{\partial t^6} \eta(t, z, u, v) \right) \equiv 0.$$

An explicit form of Q can be obtained by applying Proposition 4.2 to the following equation

$$(29) \quad t^4 \left(\frac{\partial}{\partial t} \text{KP1} \right)^2 - (\text{KP2})^2 + \text{KP1} \left(\text{KP3} - \frac{1}{2} (t(u+v+1-z) + 1) \theta \text{KP2} - \left(t^4 \frac{\partial^2}{\partial t^2} + 4t^3 \frac{\partial}{\partial t} + 2t^4 \frac{\partial^2}{\partial t^2} \eta + 8t^3 \frac{\partial}{\partial t} \eta + 3uvt^2 - (u+v)t \right) \text{KP1} \right) \equiv 0$$

where

$$\begin{aligned} \text{KP1} &= -4G_{3,1}^\theta + 4G_{2,2}^\theta + \frac{4}{3}(6(G_{1,2}^\theta)^2 + G_{1,4}^\theta), \\ \text{KP2} &= -4G_{4,1}^\theta + 4G_{3,2}^\theta + \frac{8}{3}(6G_{2,1}^\theta G_{1,2}^\theta + G_{2,1,3}^\theta), \\ \text{KP3} &= -6G_{5,1}^\theta + 4G_{4,2}^\theta + 2G_{3,3}^\theta + \frac{8}{3}(6G_{3,1}^\theta G_{1,2}^\theta + G_{3,1,3}^\theta) \\ &\quad + 4(4(G_{2,1}^\theta)^2 + 2G_{2,2}^\theta G_{1,2}^\theta + G_{2,2,1,2}^\theta) + \frac{4}{45}(60(G_{1,2}^\theta)^3 + 30G_{1,4}^\theta G_{1,2}^\theta + G_{1,6}^\theta), \end{aligned}$$

and $\eta \equiv \eta(t, z, u, v)$. An explicit form of this ODE can be found in the accompanying Maple worksheet [13].

The proof is analogous to the proof of Theorem 3.8 and left to the reader. We have two immediate corollaries, Theorem 4.6 which is analogous to Theorem 1.2 for bipartite maps, and Theorem 4.7 which is a bipartite analogue of Ledoux’s recurrence and a non-oriented analogue of Adrianov’s.

THEOREM 4.6 (Counting bipartite maps by edges and genus – unshifted recurrence). *The number \mathfrak{b}_n^g of rooted bipartite maps of genus g with n edges, orientable or not, is solution of an explicit recurrence relation of the form*

$$(30) \quad \begin{aligned} \mathfrak{b}_n^g &= \frac{1}{2(n+1)} \left(\sum_{\substack{n_1=1..n \\ n_1+n_2=n+1}} \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \delta_{(n_1, g_1) \neq (n, g)} \delta_{(n_2, g_2) \neq (n, g)} (n_1+1)(n_2+1) \mathfrak{b}_{n_1}^{g_1} \mathfrak{b}_{n_2}^{g_2} \right. \\ &\quad + \sum_{\substack{n_1=1..n-1 \\ n_1+n_2=n}} (6n_1n_2 + 5(n_1+n_2) + 4) \left(\sum_{\substack{g_1=0..g-1 \\ g_1+g_2=g-1}} \mathfrak{b}_{n_1}^{g_1} \mathfrak{b}_{n_2}^{g_2} - 3 \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \mathfrak{b}_{n_1}^{g_1} \mathfrak{b}_{n_2}^{g_2} \right) \\ &\quad \left. + \sum_{a=1}^{K_1} \sum_{b=0}^{K_2} \sum_{k=1}^{K_3} \sum_{\substack{n_1, \dots, n_k \geq 1 \\ n_1 + \dots + n_k = n-a}} \sum_{\substack{g_1, \dots, g_k \geq 0 \\ g_1 + \dots + g_k = g-b}} Q_{a,b,k}(n_1, \dots, n_k) \mathfrak{b}_{n_1}^{g_1} \mathfrak{b}_{n_2}^{g_2} \dots \mathfrak{b}_{n_k}^{g_k}, \right) \end{aligned}$$

where the $Q_{a,b,k}$ are rational functions and $K_1, K_2, K_3 < \infty$.

THEOREM 4.7 (A recurrence for non-oriented bipartite one-face maps). *The number $\mathfrak{b}_n^{i,j}$ of rooted one-face maps with n edges, i white and j black vertices, orientable or not, is given by the recursion:*

$$(31) \quad \begin{aligned} (n+1)\mathfrak{b}_n^{i,j} &= (4n-1)(\mathfrak{b}_{n-1}^{i-1,j} + \mathfrak{b}_{n-1}^{i,j-1} - \mathfrak{b}_{n-1}^{i,j}) + (5n^3 - 16n^2 + 13n - 1)\mathfrak{b}_{n-2}^{i,j} \\ &\quad + (2n-3)(4\mathfrak{b}_{n-2}^{i-1,j} + 4\mathfrak{b}_{n-2}^{i,j-1} - 3\mathfrak{b}_{n-2}^{i-2,j} + 3\mathfrak{b}_{n-2}^{i,j-2} - 2\mathfrak{b}_{n-2}^{i-1,j-1}) \\ &\quad + (10n^3 - 68n^2 + 150n - 107)(\mathfrak{b}_{n-3}^{i,j} - \mathfrak{b}_{n-3}^{i-1,j} - \mathfrak{b}_{n-3}^{i,j-1}) \\ &\quad + (4n-11)(\mathfrak{b}_{n-3}^{i-3,j} + \mathfrak{b}_{n-3}^{i,j-3} - \mathfrak{b}_{n-3}^{i-2,j-1} - \mathfrak{b}_{n-3}^{i-1,j-2} - \mathfrak{b}_{n-3}^{i-2,j} - \mathfrak{b}_{n-3}^{i,j-2} + 2\mathfrak{b}_{n-3}^{i-1,j-1}) \\ &\quad + (4-n)((2n-7)^2(n-2)^2 \mathfrak{b}_{n-4}^{i,j} + (5n^2 - 32n + 53)(\mathfrak{b}_{n-4}^{i-2,j} + \mathfrak{b}_{n-4}^{i,j-2} - 2\mathfrak{b}_{n-4}^{i-1,j-1}) \\ &\quad + \mathfrak{b}_{n-4}^{i-4,j} + \mathfrak{b}_{n-4}^{i,j-4} - 4\mathfrak{b}_{n-4}^{i-3,j-1} + 4\mathfrak{b}_{n-4}^{i-1,j-3} + 6\mathfrak{b}_{n-4}^{i-2,j-2}) \end{aligned}$$

with the convention that $\mathfrak{b}_n^{i,j} = 0$ for $i + j > n + 1$, and $\mathfrak{b}_n^{i,0} = \mathfrak{b}_n^{0,j} = 0$ and the initial conditions $\mathfrak{b}_1^{1,1} = \mathfrak{b}_2^{2,1} = \mathfrak{b}_2^{1,2} = \mathfrak{b}_2^{1,1} = 1$, $\mathfrak{b}_3^{3,1} = \mathfrak{b}_3^{1,3} = 1$, $\mathfrak{b}_3^{2,2} = \mathfrak{b}_3^{2,1} = \mathfrak{b}_3^{1,2} = 3$, $\mathfrak{b}_3^{1,1} = 4$.

Proof of Theorem 4.7. To obtain a linear ODE for the generating function

$$\mathfrak{b} \equiv \mathfrak{b}(t, u, v) := [z]\eta(t, z, u, v) = \sum_{n,i,j} \frac{\mathfrak{b}_n^{i,j}}{2n} t^n u^i v^j$$

of rooted non-oriented bipartite maps with only one face, we extract the coefficient of z^1 in (29), and multiply by $\frac{45}{14t^7uv(u-1)(v-1)}$. This gives

$$(32) \quad \begin{aligned} &(-uv + 2\frac{\partial}{\partial t}\mathfrak{b}) + t(2u^2v + 2uv^2 - 5uv + 7\frac{\partial}{\partial t}\mathfrak{b}(1-u-v) + \frac{\partial^2}{\partial t^2}\mathfrak{b}) + t^2(-uv((u-v)^2 - 1) \\ &\quad + (3(3u^2 + 3v^2 + 2uv) - 12(u+v) - 29)\frac{\partial}{\partial t}\mathfrak{b} + 4(1-u-v)\frac{\partial^2}{\partial t^2}\mathfrak{b}) \\ &\quad + t^3(5(-u^3 + u^2v + uv^2 - v^3 + (u-v)^2 + 7(u+v-1))\frac{\partial}{\partial t}\mathfrak{b} + (2(3u^2 + 3v^2 + 2uv) \\ &\quad - 8(u+v) - 86)\frac{\partial^2}{\partial t^2}\mathfrak{b}) + t^4(((u-v)^4 - 18(u-v)^2 + 81)\frac{\partial}{\partial t}\mathfrak{b} - 4(u^3 - u^2v - uv^2 + v^3 \\ &\quad - (u-v)^2 - 37(u+v-1))\frac{\partial^2}{\partial t^2}\mathfrak{b} - 44\frac{\partial^3}{\partial t^3}\mathfrak{b}) + t^5(((u-v)^4 - 64(u-v)^2 + 719)\frac{\partial^2}{\partial t^2}\mathfrak{b} \\ &\quad + 82(u+v-1)\frac{\partial^3}{\partial t^3}\mathfrak{b} - 5\frac{\partial^4}{\partial t^4}\mathfrak{b}) + t^6((-38(u-v)^2 + 1078)\frac{\partial^3}{\partial t^3}\mathfrak{b} \\ &\quad + 10(u+v-1)\frac{\partial^4}{\partial t^4}\mathfrak{b}) + t^7(-5(u-v)^2 + 493)\frac{\partial^4}{\partial t^4}\mathfrak{b} + 80t^8\frac{\partial^5}{\partial t^5}\mathfrak{b} + 4t^9\frac{\partial^6}{\partial t^6}\mathfrak{b} = 0. \end{aligned}$$

Extracting the coefficient of $[t^n u^i v^j]$ produces the desired recursion. □

4.2. NON-ORIENTED TRIANGULATIONS. The generating series of triangulations can be obtained from $F(t, \mathbf{p}, u)$ (given by (13)) by applying another specialization instead of θ . Indeed, define the specialization operator θ_3 by $\theta_3(p_i) := z\delta_{3,i}$. This operator enforces that all faces must be of degree 3, and

$$\Xi(t, z, u) := \theta_3 F(t, \mathbf{p}, u) = \sum_{\substack{M: \forall f \in F(M) \\ \deg(f)=3}} \frac{t^{2e(M)}}{4e(M)} z^{f(M)} u^{v(M)} = \sum_{n \geq 1} \sum_{g \geq 0} \frac{t_n^g}{12n} t^{6n} z^{2n} u^{n+2-2g},$$

is the generating function of rooted, non-oriented triangulations M . Of course, triangulations satisfy $2e(M) = 3f(M)$. By using Euler's relation, one can expand $\Xi(t, z, u)$ by the genus and the number of edges, and here t_n^g denotes the number of rooted, non-oriented triangulations with $3n$ edges (or equivalently $2n$ faces) and genus g .

Similarly as in the previous sections, we want to express $F_\lambda^{\theta_3}$ as a polynomial in $\Xi(t, z, u)$ and its derivatives with respect to t , where

$$F_\lambda^{\theta_3} \equiv F_\lambda^{\theta_3}(t, z, u) := \theta_3(F_\lambda) = \theta_3 \left(\prod_{j=1}^k \frac{\partial}{\partial p_{i_j}} F(t, \mathbf{p}, u) \right).$$

for a sequence of non-negative integers $\lambda = (i_1, \dots, i_k)$.

PROPOSITION 4.8. For $i \geq -1$ and $n_1, n_2 \geq 0$, one has the recurrence relations

$$(33) \quad t^2 z(i+3)F_{i+3,2^{n_2},1^{n_1}}^{\theta_3} = -2t^2 \sum_{\substack{a+b=i \\ a,b \geq 1}} \sum_{\substack{l_1=0 \\ i=1,2}}^{n_1} ab \binom{n_1}{l_1} \binom{n_2}{l_2} F_{a,2^{l_2},1^{l_1}}^{\theta_3} F_{b,2^{n_2-l_2},1^{n_1-l_1}}^{\theta_3} \\ - 2t^2 \sum_{\substack{a+b=i \\ a,b \geq 1}} ab F_{a,b,2^{n_2},1^{n_1}}^{\theta_3} - t^2 \sum_{j=1}^2 n_j(i+j)F_{i+j,2^{n_2-\delta_{2,j}},1^{n_1-\delta_{1,j}}}^{\theta_3} + (i+2)F_{i+2,2^{n_2},1^{n_1}}^{\theta_3} \\ - t^2 \delta_{i \neq -1} (2u+i+1)iF_{i,2^{n_2},1^{n_1}}^{\theta_3} - t^2 \left(\delta_{i,-1} \delta_{n_1,1} + (u+1)\delta_{i,0} \delta_{n_1,0} \right) \frac{u}{2} \delta_{n_2,0},$$

$$(34) \quad F_{3,i_1,\dots,i_k}^{\theta_3} = \frac{t}{3z} \frac{\partial}{\partial t} F_{i_1,\dots,i_k}^{\theta_3} - \frac{i_1 + \dots + i_k}{3z} F_{i_1,\dots,i_k}^{\theta_3}$$

$$(35) \quad F_{1^i}^{\theta_3} = t^5 z \frac{\partial}{\partial t} F_{1^{i-1}}^{\theta_3} + \frac{t^4 z(u^2+u)}{2} \delta_{i,1} + \frac{t^2 u}{2} \delta_{i,2},$$

with the convention that $F_{3^0,2^0,1^0}^{\theta_3} = \Xi(t, z, u)$.

Consequently, for any sequence of integers of the form $\lambda = [\ell, 3^{n_3}, 2^{n_2}, 1^{n_1}]$ and of size $|\lambda| = \ell + n_1 + 2n_2 + 3n_3$, there exists a polynomial R_λ in $|\lambda|$ variables, with coefficients in $\mathbb{Q}[t, t^{-1}, z, z^{-1}, u]$ which is linear for $\ell \leq 4$, and quadratic for $5 \leq \ell \leq 8$ and satisfies

$$(36) \quad F_\lambda^{\theta_3} = R_\lambda \left(\frac{\partial}{\partial t} \Xi(t, z, u), \dots, \frac{\partial^{|\lambda|}}{\partial t^{|\lambda|}} \Xi(t, z, u) \right).$$

Proof. Similarly to the proof of Proposition 3.4, we act with $\frac{\partial^{n_1+n_2}}{\partial p_1^{n_1} \partial p_2^{n_2}}$ on both sides of (19) and apply θ_3 to obtain

$$(37) \quad (i+2)F_{i+2,2^{n_2},1^{n_1}}^{\theta_3} = t^2 z(i+3)F_{i+3,2^{n_2},1^{n_1}}^{\theta_3} + \delta_{i \neq -1} t^2 i(i+1+2u)F_{i,2^{n_2},1^{n_1}}^{\theta_3} \\ + 2t^2 \sum_{a+b=i} ab \left(F_{a,b,2^{n_2},1^{n_1}}^{\theta_3} + \sum_{l_1=0}^{n_1} \sum_{l_2=0}^{n_2} \binom{n_1}{l_1} \binom{n_2}{l_2} F_{a,2^{l_2},1^{l_1}}^{\theta_3} F_{b,2^{n_2-l_2},1^{n_1-l_1}}^{\theta_3} \right) \\ + t^2 n_1(i+1)F_{i+1,2^{n_2},1^{n_1-1}}^{\theta_3} + t^2 n_2(i+2)F_{i+2,2^{n_2-1},1^{n_1}}^{\theta_3} \\ + t^2 \left(\delta_{i,-1} \delta_{n_1,1} + (u+1)\delta_{i,0} \delta_{n_1,0} \right) \frac{u}{2} \delta_{n_2,0}.$$

Equation (33) is obtained from (37) by moving $t^2 z(i+3)F_{i+3,2^{n_2},1^{n_1}}^{\theta_3}$ to the left of the equality and $(i+2)F_{i+2,2^{n_2},1^{n_1}}^{\theta_3}$ to the right.

To get Equation (34), one acts on both sides of the homogeneity relation $\sum_{i \geq 1} p_i p_i^* F = t \frac{\partial F}{\partial t}$ with $\frac{\partial^k}{\partial p_1 \dots \partial p_k}$ and applies θ_3 .

Furthermore, specializing $i = -1, n_2 = 0, n_1 = l - 1$ and then $i = 0, n_2 = 0, n_1 = l - 1$ in (37), we get

$$F_{1^i}^{\theta_3} = t^2 z 2 F_{2,1^{i-1}}^{\theta_3} + t^2 \frac{u}{2} \delta_{i,2} = t^2 z \left(t^2 z 3 F_{3,1^{i-1}}^{\theta_3} + t^2 (l-1) F_{1^{i-1}}^{\theta_3} + t^2 \frac{u(u+1)}{2} \delta_{i=1} \right) + t^2 \frac{u}{2} \delta_{i,2}.$$

To get (35), one substitutes (34) into the above for $i_1, \dots, i_k = 1$ and $k = l - 1$.

Since the sizes of the vectors indexing F^{θ_3} s appearing in the RHS of (33) are strictly smaller than $|\lambda|$, one computes $F_\lambda^{\theta_3}$ recursively for vectors of the form $\lambda = [\ell, 3^{n_3}, 2^{n_2}, 1^{n_1}]$, where $\ell \leq 10$ (by eliminating all the parts of length 3 thanks to (34), and reducing the sizes of the indexing vectors thanks to (33) and finally using the recurrence (35) for the parts of the form $F_{1^i}^{\theta_3}$). The last statement follows by induction on ℓ . \square

REMARK 4.9. We want to highlight the fact that the above computations are possible because we are working with the specific model of triangulations. Replacing the specialization θ_3 by $\theta_l : p_i \mapsto \delta_{i,l}$ with $l \geq 4$ makes the above technique fail to even compute $F_1^{\theta_l}$.

The following theorem gives a recurrence formula with non-polynomial coefficients in the case of triangulations. It is essentially equivalent to the functional equation for the associated generating function given in [20, Corollary 3.2], which is obtained through (essentially) the same method.

THEOREM 4.10 (Counting triangulations by faces, and genus). *The number t_n^g of rooted non-oriented triangulations of genus g with $2n$ faces (or, equivalently $3n$ edges) can be computed from the following recurrence formula:*

$$\begin{aligned}
 t_n^g &= \frac{2}{2n^2+(3-2g)n+(1-g)(1-2g)} \times \left(n \left(6(3n-1)t_{n-1}^g + 12(3n-4)((3n-2)nt_{n-2}^{g-1} \right. \right. \\
 &\quad \left. \left. + 2(t_{n-2}^{g-1/2} + t_{n-2}^g) \right) + 6 \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n \\ n_1+n_2=n}} (3n_1-1)(3n_2-1)t_{n_2-1}^{g_2}t_{n_1-1}^{g_1} \right) \\
 &- \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=1..n \\ n_1+n_2=n}} \left(\sum_{\substack{g_0=0..g_1 \\ g_1-g_0 \in \mathbb{N}}} \binom{n_1+2-2g_0}{n_1-2g_1} 2^{2(1+g_1-g_0)} t_{n_1}^{g_0} \right) \left(-\frac{n_2+1}{8} t_{n_2}^{g_2} + (3n_2-1)t_{n_2-1}^{g_2} \right. \\
 &\quad \left. + 2(3n_2-4)((3n_2-2)n_2 t_{n_2-2}^{g_2-1} + 2(t_{n_2-2}^{g_2-1/2} + t_{n_2-2}^{g_2})) + \frac{1}{8} \delta_{g_0 \neq g} \delta_{n_1, n} (\delta_{g_1, g} - \delta_{g_1, g-1/2}) \right. \\
 &\quad \left. + \delta_{n_1, n-1} (2\delta_{g_1, g} + 2\delta_{g_1, g-1/2} + \delta_{g_1, g-1}) + 4\delta_{n_1, n-2} (\delta_{g_1, g} + 2\delta_{g_1, g-1/2} \right. \\
 &\quad \left. + 9\delta_{g_1, g-1} + 8\delta_{g_1, g-3/2}) + \sum_{\substack{g_3=0..g_2 \\ g_3+g_4=g_2}} \sum_{\substack{n_3=0..n_2 \\ n_3+n_4=n_2}} (3n_3-1)(3n_4-1)t_{n_3-1}^{g_3}t_{n_4-1}^{g_4} \right) \Big)
 \end{aligned}$$

for $n > 2$, with the initial conditions $t_0^g = 0$, $t_1^0 = 4$, $t_2^0 = 32$, $t_1^{1/2} = 9$, $t_2^{1/2} = 118$, $t_1^1 = 7$, $t_2^1 = 202$, $t_2^{3/2} = 128$ and $t_n^g = 0$ if $n < 2g - 1$.

Proof. As in the case of bipartite maps, the proof is almost identical to the proof of Theorem 3.7 and we leave the details for the interested reader. The only difference is that (20) should be replaced by

$$\begin{aligned}
 &\left(\frac{\partial}{\partial t} (\Xi(t, z, u + 2) + \Xi(t, z, u - 2) - 2\Xi(t, z, u)) \right) \left(4F_{2^2}^{\theta_3} - 4F_{3,1}^{\theta_3} + \frac{4}{3}(6(F_{1^2}^{\theta_3})^2 + F_{1^4}^{\theta_3}) \right) \\
 &= \frac{\partial}{\partial t} \left(4F_{2^2}^{\theta_3} - 4F_{3,1}^{\theta_3} + \frac{4}{3}(6(F_{1^2}^{\theta_3})^2 + F_{1^4}^{\theta_3}) \right) - \frac{4}{t} \left(4F_{2^2}^{\theta_3} - 4F_{3,1}^{\theta_3} + \frac{4}{3}(6(F_{1^2}^{\theta_3})^2 + F_{1^4}^{\theta_3}) \right).
 \end{aligned}$$

□

THEOREM 4.11. *There exists a polynomial $R \in \mathbb{Q}[t, u, z][x_1, \dots, x_6]$ of degree 5 such that*

$$R \left(\frac{\partial}{\partial t} \Xi(t, z, u), \dots, \frac{\partial^6}{\partial t^6} \Xi(t, z, u) \right) \equiv 0.$$

An explicit form of R can be obtained by applying Proposition 4.8 to the following equation

$$\begin{aligned}
 (38) \quad &t^{10}z^2 \left(\frac{\partial}{\partial t} \text{KP1} \right)^2 - (\text{KP2})^2 + \text{KP1} \left(\text{KP3} - \frac{1}{2t^2z} \text{KP2} \right. \\
 &\left. - \left(t^{10}z^2 \frac{\partial^2}{\partial t^2} + 5t^9z^2 \frac{\partial}{\partial t} + 2t^{10}z^2 \frac{\partial^2}{\partial t^2} \Xi + 10t^9z^2 \frac{\partial}{\partial t} \Xi + 4t^8z^2(u^2 + u) + t^2u \right) \text{KP1} \right) \equiv 0
 \end{aligned}$$

where

$$\begin{aligned} \text{KP1} &= -4F_{3,1}^{\theta_3} + 4F_{2^2}^{\theta_3} + \frac{4}{3}(6F_{1^2}^{\theta_3^2} + F_{1^4}^{\theta_3}), \\ \text{KP2} &= -4F_{4,1}^{\theta_3} + 4F_{3,2}^{\theta_3} + \frac{8}{3}(6F_{2,1}^{\theta_3}F_{1^2}^{\theta_3} + F_{2,1^3}^{\theta_3}), \\ \text{KP3} &= -6F_{5,1}^{\theta_3} + 4F_{4,2}^{\theta_3} + 2F_{3,3}^{\theta_3} + \frac{8}{3}(6F_{3,1}^{\theta_3}F_{1^2}^{\theta_3} + F_{3,1^3}^{\theta_3}) + 4(4F_{2,1}^{\theta_3^2} + 2F_{2^2}^{\theta_3}F_{1^2}^{\theta_3} + F_{2^2,1^2}^{\theta_3}) \\ &\quad + \frac{4}{45}(60F_{1^2}^{\theta_3^3} + 30F_{1^4}^{\theta_3}F_{1^2}^{\theta_3} + F_{1^6}^{\theta_3}) \end{aligned}$$

where $\Xi \equiv \Xi(t, z, u)$. An explicit form of this ODE can be found in the accompanying Maple worksheet [13].

The proof is the same as in the previous cases, so we leave it as an exercise. As a standard consequence we have:

THEOREM 4.12 (Counting triangulations by edges and genus – unshifted recurrence). *The number t_n^g of rooted triangulations of genus g with $3n$ edges, orientable or not, is solution of an explicit recurrence relation of the form*

$$\begin{aligned} (39) \quad t_n^g &= t_n^{g-1/2} - \sum_{\substack{n_1=1..n-1 \\ n_1+n_2=n}} \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \frac{(n_1+1)(n_2+1)}{14(n+1)} t_{n_1}^{g_1} t_{n_2}^{g_2} \\ &\quad + \sum_{a=1}^{K_1} \sum_{b=0}^{K_2} \sum_{k=1}^{K_3} \sum_{\substack{n_1, \dots, n_k \geq 1 \\ n_1 + \dots + n_k = n-a}} \sum_{\substack{g_1, \dots, g_k \geq 0 \\ g_1 + \dots + g_k = g-b}} R_{a,b,k}(n_1, \dots, n_k) t_{n_1}^{g_1} t_{n_2}^{g_2} \dots t_{n_k}^{g_k}, \end{aligned}$$

where the $R_{a,b,k}$ are rational functions and $K_1, K_2, K_3 < \infty$.

In analogy to what we did for maps and bipartite maps, it would be natural to study now the case of triangulations with only one vertex (or by duality, cubic one-face maps). However, there exist very explicit and simple formulas in this case, obtained from bijective methods [12] so we prefer not to go into such calculations here.

5. ANOTHER METHOD IN THE CASE OF MAPS

In this section, we quickly address the case of maps treated in Section 3 with another method, which actually leads to different recurrence relations. The situation is similar to the orientable case, where the approaches used in [21] and [33] differ. In Section 3 (non-oriented analogue of [33]) we started from the fact that the generating function F of maps is a BKP tau function, and applied the substitution operator $\theta : p_i \mapsto z$. In this section, we will instead start from the fact that the generating function G of bipartite maps is a BKP tau function, and apply the different substitution operator $\theta_2 : p_i \mapsto \delta_{i,2}$. We will only treat the equations with shifts, our main motivation being that they are relatively nice looking – for example we will prove here Theorem 1.1.

Our starting point is the well-known fact that, from a famous bijection due to Tutte and valid on all surfaces, the number of rooted maps with n edges on a surface is equal to the number of rooted bipartite quadrangulations on the same surface, with vertices and faces of the map corresponding respectively to black and white vertices of the quadrangulation (see e.g. [21]). Therefore, the generating function $\Theta(t, z, u)$ of maps defined in Section 3 and $G(t, \mathbf{p}, u, v)$ of bipartite maps defined in Section 4.1 satisfy the relation

$$\theta_2 G(t, \mathbf{p}, u, z) = \Theta(t, z, u) = \sum_{n \geq 1} \sum_{g \geq 0} \frac{H_n^g}{4n},$$

where $H_n^g \in \mathbb{N}[u, z]$ is the generating polynomial of rooted maps of genus g with n edges, with u and z marking respectively vertices and faces. Let us write $G^{u,z} \equiv G(t, \mathbf{p}, u, z)$, so that the BKP equation (3) for the function G can be rewritten

(40)

$$G_{1^4}^{u,z} + 3G_{2^2}^{u,z} - 3G_{3,1}^{u,z} + 6(G_{1^2}^{u,z})^2 = C(u, z) \exp \left(G^{u+2;z+2} - 2G^{u,z} + G^{u-2;z-2} \right),$$

where the prefactor $C(u, z)$ (which is equal to $S_2(N)$ after the substitution $u = N, z = N + \delta$), independent of the variables $(p_i)_{i \geq 1}$, does not play any role in what follows, and where as before the indices indicate derivatives with respect to the \mathbf{p} -variables. By hitting (40) with the operator $\frac{\partial}{\partial p_1}$, and using (40) again to eliminate the factor $C(u, z) \exp(\dots)$, we obtain an equation which involves no exponential anymore, and in which the prefactor has disappeared. Namely:

$$(41) \quad G_{1^5}^{u,z} + 3G_{2^2,1}^{u,z} - 3G_{3,1^2}^{u,z} + 12G_{1^2}^{u,z} G_{1^3}^{u,z} = \left(G_1^{u+2;z+2} - 2G_1^{u,z} + G_1^{u-2;z-2} \right) \left(G_{1^4}^{u,z} + 3G_{2^2}^{u,z} - 3G_{3,1}^{u,z} + 6(G_{1^2}^{u,z})^2 \right).$$

To extract coefficients in this equation, we will use the following lemma. Here we use an additional variable r which will be convenient to track the genus parameter. In this section, it is convenient to use the convention $H_0^g = uz$.

LEMMA 5.1. *We have, for $(n, g) \in \mathbb{N} \times (\frac{1}{2}\mathbb{N})$, $n \geq 1$,*

$$\begin{aligned} [t^{2n+1}r^{n+2-2g}] \theta_2 \quad & \partial_1 G^{ru;rz} = \frac{1}{2} H_n^g, \\ [t^{2n}r^{n+2-2g}] \theta_2 \quad & \partial_{2^2} G^{ru;rz} = \frac{n-1}{4} H_n^g, \\ [t^{2n}r^{n+1-2g}] \theta_2 \quad & \partial_{1^2} G^{ru;rz} = \frac{2n-1}{2} H_{n-1}^g, \\ [t^{2n}r^{n+2-2g}] \theta_2 \quad & \partial_{1^4} G^{ru;rz} = \frac{(2n-1)(2n-2)(2n-3)}{2} H_{n-2}^{g-1}, \\ [t^{2n}r^{n+2-2g}] \theta_2 \quad & \partial_{1,3} G^{ru;rz} = \frac{(2n-1)}{6} \left(H_n^g - (u+z)H_{n-1}^g - H_{n-1}^{g-1/2} \right), \\ [t^{2n+1}r^{n+2-2g}] \theta_2 \quad & \partial_{2^2,1} G^{ru;rz} = \frac{n(n-1)}{2} H_n^g, \\ [t^{2n+1}r^{n+1-2g}] \theta_2 \quad & \partial_{1^3} G^{ru;rz} = \frac{2n(2n-1)}{2} H_{n-1}^g, \\ [t^{2n+1}r^{n+2-2g}] \theta_2 \quad & \partial_{1^5} G^{ru;rz} = \frac{2n(2n-1)(2n-2)(2n-3)}{2} H_{n-2}^{g-1}, \\ [t^{2n+1}r^{n+2-2g}] \theta_2 \quad & \partial_{3,1^2} G^{ru;rz} = \frac{n(2n-1)}{3} \left(H_n^g - (u+z)H_{n-1}^g - H_{n-1}^{g-1/2} \right). \end{aligned}$$

For $n = 0$, the first equality remains valid, while all other quantities in left-hand sides vanish.

Proof. The lemma can easily be proved with Virasoro constraints in the same manner as Proposition 4.8 and details are left to the reader. However, a calculation-free proof based on digon contraction and elementary combinatorial map operation is also easily doable. The proof is completely similar to [21, Lemma 7], the only difference is the extra term of genus $g - 1/2$ in the two equations involving a hexagonal default (i.e. a p_3 -derivative). This term comes from the possibility to create a rooted quadrangulation by adding a twisted diagonal inside a digon. This is the only difference between the oriented and non-oriented case, and it adds one term to Equation (11) in [21]. Once this difference is taken into account, the proof of [21, Lemma 7] can be copied verbatim. \square

A direct consequence of what precedes is the recurrence formula stated as Theorem 1.1 in the introduction. One can also obtain a version with control on vertices and faces, from which Theorem 1.1 follows immediately.

THEOREM 5.2 (Counting maps by vertices, faces, and genus). *The generating polynomial*

$$H_n^g \equiv H_n^g(u, z) = \sum_{i+j=n+2-2g} H_n^{i,j} u^i z^j$$

of rooted maps of genus g with n edges, orientable or not, with weight u per vertex and z per face, can be computed from the following recurrence formula:

(42)

$$\begin{aligned} H_n^g &= \frac{2}{(n+1)(n-2)} \times \left(n(2n-1)((u+z)H_{n-1}^g + H_{n-1}^{g-1/2}) + \frac{(2n-3)(2n-2)(2n-1)(2n)}{2} H_{n-2}^{g-1} \right. \\ &\quad + 12 \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n \\ n_1+n_2=n}} \frac{(2n_2-1)(2n_1-1)n_1}{2} H_{n_2-1}^{g_2} H_{n_1-1}^{g_1} \\ &\quad - \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n-1 \\ n_1+n_2=n}} \sum_{\substack{g_0=0..g_1 \\ g_1-g_0 \in \mathbb{N}}} \sum_{p+q=n_1+2-2g_0} 2^{2(1+g_1-g_0)} \phi_{p,q,n_1-2g_1}(u, z) H_{n_1}^{p,q} \\ &\quad \left. \left(\frac{(2n_2-1)(2n_2-2)(2n_2-3)}{2} H_{n_2-2}^{g_2-1} - \delta_{(n_2, g_2) \neq (n, g)} \frac{n_2+1}{4} H_{n_2}^{g_2} + \frac{2n_2-1}{2} ((u+z)H_{n_2-1}^{g_2} + H_{n_2-1}^{g_2-1/2}) \right. \right. \\ &\quad \left. \left. + 6 \sum_{\substack{g_3=0..g_2 \\ g_3+g_4=g_2}} \sum_{\substack{n_3=0..n_2 \\ n_3+n_4=n_2}} \frac{(2n_3-1)(2n_4-1)}{4} H_{n_3-1}^{g_3} H_{n_4-1}^{g_4} \right) \right) \end{aligned}$$

for $n > 2$, with the initial conditions $H_0^0 = uz$, $H_1^0 = uz(u+z)$, $H_2^0 = uz(2u^2 + 5uz + 2z^2)$, $H_1^{1/2} = uz$, $H_2^{1/2} = 5uz(u+z)$, $H_2^1 = 5uz$, and $H_n^g = 0$ if $n < 2g$, and where

$$\phi_{p,q,m}(u, z) := \sum_{\substack{i+j=m \\ p \geq i, q \geq j}} \binom{p}{i} \binom{q}{j} u^i z^j.$$

Proof. We substitute $u \rightarrow ur, z \rightarrow zr$ in (41) and we extract the coefficient of $[t^{2n+1}r^{n+2-2g}]$ after applying θ_2 . We obtain from Lemma 5.1

$$\begin{aligned} &- \frac{n(n+1)}{2} H_n^g + n(2n-1)((u+z)H_{n-1}^g + H_{n-1}^{g-1/2}) + \frac{(2n-3)(2n-2)(2n-1)(2n)}{2} H_{n-2}^{g-1} \\ &\quad + 12 \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n \\ n_1+n_2=n}} \frac{(2n_2-1)(2n_1-1)n_1}{2} H_{n_2-1}^{g_2} H_{n_1-1}^{g_1} \\ &= \sum_{\substack{g_1=0..g \\ g_1+g_2=g}} \sum_{\substack{n_1=0..n-1 \\ n_1+n_2=n}} \left(\sum_{\substack{g_0=0..g_1 \\ g_1-g_0 \in \mathbb{N}}} \sum_{p+q=n_1+2-2g_0} 2^{2(1+g_1-g_0)} \phi_{p,q,n_1-2g_1}(u, z) H_{n_1}^{p,q} \right) \\ &\quad \left(\frac{(2n_2-1)(2n_2-2)(2n_2-3)}{2} H_{n_2-2}^{g_2-1} + \frac{-n_2-1}{4} H_{n_2}^{g_2} + \frac{2n_2-1}{2} ((u+z)H_{n_2-1}^{g_2} + H_{n_2-1}^{g_2-1/2}) \right. \\ &\quad \left. + 6 \sum_{\substack{g_3=0..g_2 \\ g_3+g_4=g_2}} \sum_{\substack{n_3=0..n_2 \\ n_3+n_4=n_2}} \frac{(2n_3-1)(2n_4-1)}{4} H_{n_3-1}^{g_3} H_{n_4-1}^{g_4} \right). \end{aligned}$$

Here we have extracted, respectively, in (41) (after applying θ_2 , and substitution $u \rightarrow ur, z \rightarrow zr$) the coefficient of $[t^{2n+1}r^{n+2-2g}]$, of $[t^{2n_1+1}r^{n_1-2g_1}]$, and of $[t^{2n_2}r^{n_2+2-2g_2}]$, in the LHS, in the first factor of the RHS, and in the second factor of the RHS. The summation over n_1 in the RHS stops at $(n-1)$ since from the last sentence of Lemma 5.1, the term $n_2 = 0$

does not contribute. Moreover we have used

$$\begin{aligned}
 & [r^m](f(ru + 2, rz + 2) + f(ru - 2, rz - 2) - 2f(ru, rz)) \\
 &= \sum_{i+j=m} u^i z^j \sum_{\substack{p \geq i, q \geq j \\ p+q \neq m}} \binom{p}{i} \binom{q}{j} (2^{p+q-m} + (-2)^{p+q-m}) [u^p z^q] f(u, z) \\
 &= \sum_{i+j=m} u^i z^j \sum_{\substack{p \geq i, q \geq j \\ p+q \geq m+2, p+q-m \in 2\mathbb{N}}} \binom{p}{i} \binom{q}{j} 2^{1+p+q-m} [u^p z^q] f(u, z) \\
 &= \sum_{\substack{k \geq m+2 \\ k-m \in 2\mathbb{N}}} 2^{1+k-m} \sum_{p+q=k} \phi_{p,q,m}(u, z) [u^p z^q] f(u, z),
 \end{aligned}$$

which we use with the parametrization $m = n_1 - 2g_1$, $k = n_1 + 2 - 2g_0$ (the condition $k \geq m + 2$ translates into $g_0 \leq g_1$, and the summand is null when $g_0 < 0$).

It now only remains to group the two terms of the form H_n^g (namely: the first term of the LHS and the term $H_{n_2}^{g_2}$ in the RHS when $n_1 = 0, n_2 = n, g_1 = 0, g_2 = g$). They appear in the difference LHS–RHS with coefficient $-\frac{n(n+1)}{2} + 2^2 \times \frac{n+1}{4} = -\frac{(n+1)(n-2)}{2}$, which leads to the main equation of the theorem after dividing by this factor. The identification of the initial conditions for $n \geq 2$ can be done in many ways including: by hand drawing, or from the OEIS, or from explicit expansions in small genera using the Virasoro constraints, or from the expansion in Zonal polynomials up to order $n = 2$. □

Proof of Theorem 1.1 stated in the introduction. Note that $\mathfrak{h}_n^g = H_n^g(1, 1)$ and

$$\phi_{p,q,n_1-2g_1}(1, 1) = \sum_{\substack{i+j=m \\ p \geq i, q \geq j}} \binom{p}{i} \binom{q}{j} = \binom{p+q}{n_1-2g_1},$$

therefore (4) is a specialization of (42) at $u = z = 1$. □

APPENDIX A. SOME TABLES

We provide here some tables computed with our recurrences, see also [13].

A.1. ROOTED MAPS OF GENUS g WITH n EDGES (ORIENTABLE OR NOT).

$n \setminus g$	0	1/2	1	3/2	2
1	2	1	0	0	0
2	9	10	5	0	0
3	54	98	104	41	0
4	378	983	1647	1380	509
5	2916	10062	23560	31225	24286
6	24057	105024	320198	592824	724866
7	208494	1112757	4222792	10185056	17312568
8	1876446	11934910	54617267	164037704	361811054
9	17399772	129307100	696972524	2525186319	6912864180
10	165297834	1412855500	8807574390	37596421940	123814835628
11	1602117468	15548498902	110483092984	545585129474	2111880200672
12	15792300756	172168201088	1377998069826	7758174844664	34669329147582
13	157923007560	1916619748084	17108920039328	108518545261360	551879941676492
14	1598970451545	21436209373224	211636362018548	1497384373878512	8565305839025180
15	16365932856990	240741065193282	2609949110616064	20426386710028260	130146976774282440
16	169114639522230	2713584138389838	32104324480419131	275940187259609296	1942255149093281772

$n \setminus g$	5/2	3	7/2	4
5	8229	0	0	0
6	516958	166377	0	0
7	19381145	13093972	4016613	0
8	562395292	595145086	382630152	113044185
9	13929564070	20431929240	20549348578	12704958810
10	309411522140	587509756150	818177659640	790343495467
11	6344707786945	14923379377192	26881028060634	35918779737610
12	122357481545872	345651571125768	770725841809552	1330964564940140
13	2247532739398856	7452363840633244	19946409152977346	42611002435124552
14	39681114425793904	151717486205709730	476412224477845444	1220973091185233106
15	677939355268197412	2946794762696249280	10665684328125155376	32054128913697072040
16	11265765391845733784	55029552840385680100	226357454725004343024	783804517126931272890

A.2. ROOTED BIPARTITE MAPS OF GENUS g WITH n EDGES (ORIENTABLE OR NOT).

$n \setminus g$	0	1/2	1	3/2	2
1	1	0	0	0	0
2	3	1	0	0	0
3	12	9	4	0	0
4	56	69	63	20	0
5	288	510	720	480	148
6	1584	3738	7254	7584	4860
7	9152	27405	68460	99372	99036
8	54912	201569	621315	1169640	1607432
9	339456	1488762	5496208	12841632	22759560
10	2149888	11043318	47759130	134278720	293971176
11	13891584	82257890	409620156	1354371348	3553592152
12	91287552	615092178	3478672642	13287239184	40855164228
13	608583680	4615882908	29315742924	127526774024	451592018748
14	4107939840	34752865332	245539064736	1202371430148	4836001359644
15	28030648320	262437282621	2046309441924	11170818315900	50454786158100
16	193100021760	1987229885913	16983591315267	102508926612240	515031678182160

$n \setminus g$	5/2	3	7/2	4
6	1348	0	0	0
7	57204	15104	0	0
8	1445760	793260	198144	0
9	28251720	24092916	12500640	2998656
10	470885712	553335140	446044020	222034464
11	7034561160	10652501508	11827897444	9139492032
12	96964428080	181251943620	259263273912	275741173612
13	1256403317832	2812951666460	4965451637328	6799083573828
14	15499423803780	40643437847436	85911625991020	145094953853052
15	183709516250796	554529301430940	1372607347932900	2774708761422460
16	2106284848285632	7218066635434760	20563312515574176	48658560979911312

A.3. ROOTED TRIANGULATIONS OF GENUS g WITH $2n$ FACES (ORIENTABLE OR NOT).

$n \setminus g$	0	1/2	1
1	4	9	7
2	32	118	202
3	336	1773	4900
4	4096	28650	112046
5	54912	484578	2490132
6	786432	8457708	54442636
7	11824384	151054173	1177912344
8	184549376	2745685954	25302706734
9	2966845440	50606020854	540709469284
10	48855252992	943283037684	11509659737732
11	820675092480	17746990547634	244254583041960
12	14018773254144	336517405188900	5170993925895980
13	242919827374080	6423775409047716	109258058984867592
14	4261707069259776	123332141503711704	2304778527410416728
15	75576645116559360	2379824766494404317	48552885599587471920

$n \setminus g$	3/2	2	5/2
1	0	0	0
2	128	0	0
3	6786	3885	0
4	249416	309792	163840
5	7820190	15536592	17742726
6	224154528	626073960	1140086560
7	6064485588	22147258392	56574101430
8	157592065776	718135826112	2394618429216
9	3975252852294	21875815507824	90903502798380
10	98013064376240	635740513124184	3186926652389376
11	2373323509105164	17808561973715832	105134232237568182
12	56632532943141168	484348105828421472	3305475583204245376
13	1335091307453227116	12857728996745420112	99951709676667034212
14	31155184166556067968	334487003255090327376	2926388895694344300864
15	720738499764872647080	8553392225715199201200	83383518174303020028732

$n \setminus g$	3	7/2	4
4	0	0	0
5	8878870	0	0
6	1227058016	587202560	0
7	96836144376	99359372628	45877917085
8	5738654714432	9344829276160	9227542480640
9	283959455776728	646430229699516	1011244742721480
10	12391917590699520	36729466978572288	80222081136864896
11	492702239182522512	1816211696054002632	5159782135287908304
12	18229054925434379424	80930038930104447744	285723552389864612352
13	636795227835309684024	3324906134317505727756	14128927461188199914592
14	21225085309259820837824	127965696661596592413184	639196545524077326637824
15	680321493460375920656880	4667484955217376877322616	26909217174327495052218480

APPENDIX B. A FIXED-CHARGE BKP EQUATION

THEOREM B.1. Let $\tau(N)$ be a BKP tau function. Then for $k \in \mathbb{N}, k \geq 1$ the following identity holds in $\mathbb{C}(N)[\mathbf{p}, \mathbf{q}][[t]]$:

$$(43) \quad 2F_{1^3} \text{KP1}^3 = (\text{KP3}_1 - 2 \text{KP2}_2) \text{KP1}^2 - (\text{KP3} - 3 \text{KP1}_{1^2}) \text{KP1} \text{KP1}_1 + 2(\text{KP1}_2 - \text{KP2}_1) \text{KP1} \text{KP2} + 2 \text{KP2}^2 \text{KP1}_1 - 2 \text{KP1}^3_1 - \text{KP1}^2 \text{KP1}_{1^3},$$

where

$$\begin{aligned} \text{KP1} &= -F_{3,1} + F_{2^2} + \frac{1}{2}F_{1^2}^2 + \frac{1}{12}F_{1^4}, \\ \text{KP2} &= -2F_{4,1} + 2F_{3,2} + 2F_{2,1}F_{1^2} + \frac{1}{3}F_{2,1^3}, \\ \text{KP3} &= -6F_{5,1} + 4F_{4,2} + 2F_{3^2} + 4F_{3,1}F_{1^2} + \frac{2}{3}F_{3,1^3} + 4F_{2,1}^2 + 2F_{2^2}F_{1^2} + F_{2^2,1^2} \\ &\quad + \frac{1}{3}F_{1^2}^3 + \frac{1}{6}F_{1^4}F_{1^2} + \frac{1}{180}F_{1^6}, \end{aligned}$$

with $F \equiv F(N) = \log \tau(N)$.

Proof. Denote $\Delta f(N) = f(N + 2) - f(N - 2)$ and $\nabla f = f(N + 2) + f(N - 2)$, and

$$(44) \quad E = S_2(N)e^{\nabla F(N) - 2F(N)}.$$

Using (3), (11), (12) we obtain the following equations

$$(45) \quad E = \text{KP1}, \quad E\Delta(F_1) = \text{KP2}, \quad E(\nabla(F_{1^2}) + 2\Delta(F_2) + (\Delta F_1)^2) = \text{KP3},$$

where KP1, KP2, KP3 are given in the statement and they are the LHS in (3), (11), (12). We need to differentiate the 3rd equation w.r.t. p_1 ,

$$(46) \quad \Delta 2F_{2,1} + \nabla F_{1^3} + 2\Delta F_1 \Delta F_{1^2} = \left(\frac{\text{KP3}}{\text{KP1}}\right)_1$$

and rewrite the LHS using derivatives of the first 2 BKP equations,

$$(47) \quad \begin{aligned} \Delta 2F_{2,1} + \nabla F_{1^3} + 2\Delta F_1 \Delta F_{1^2} &= 2\left(\frac{\text{KP2}}{\text{KP1}}\right)_2 + 2F_{1^3} + \left(\frac{\text{KP1}_1}{\text{KP1}}\right)_{1^2} + 2\left(\frac{\text{KP2}}{\text{KP1}}\right)\left(\frac{\text{KP2}}{\text{KP1}}\right)_1 \\ &= 2F_{1^3} + \frac{1}{\text{KP1}^3} \left(-2 \text{KP1} \text{KP2} \text{KP1}_2 + 2 \text{KP1}^2 \text{KP2}_2 - 2 \text{KP2}^2 \text{KP1}_1 \right. \\ &\quad \left. + 2 \text{KP1}^3_1 + 2 \text{KP1} \text{KP2} \text{KP2}_1 - 3 \text{KP1} \text{KP1}_1 \text{KP1}_{1^2} + \text{KP1}^2 \text{KP1}_{1^3} \right). \end{aligned}$$

We finally equate the RHS of the above two equations and multiply by KP1^3 to obtain (43). □

REFERENCES

- [1] M. Adler and P. van Moerbeke, *Hermitian, symmetric and symplectic random ensembles: PDEs for the distribution of the spectrum*, Ann. of Math. (2) **153** (2001), no. 1, 149–189.
- [2] N. M. Adrianov, *An analog of the Harer–Zagier formula for unicellular bicolored maps*, Funktsional. Anal. i Prilozhen. **31** (1997), no. 3, 1–9.
- [3] M. Albenque and M. Lepoutre, *Blossoming bijection for higher-genus maps, and bivariate rationality*, Preprint arXiv:2007.07692, 2019.
- [4] A. Alexandrov, G. Chapuy, B. Eynard, and J. Harnad, *Weighted Hurwitz numbers and topological recursion*, Comm. Math. Phys. **375** (2020), no. 1, 237–305.
- [5] R. Belliard, S. Charbonnier, B. Eynard, and E. Garcia-Failde, *Topological recursion for generalised Kontsevich graphs and r -spin intersection numbers*, Preprint arXiv:2105.08035, 2021, <https://arxiv.org/abs/2105.08035>.
- [6] E. A. Bender and E. R. Canfield, *The asymptotic number of rooted maps on a surface*, J. Combin. Theory Ser. A **43** (1986), no. 2, 244–257.
- [7] ———, *The number of rooted maps on an orientable surface*, J. Combin. Theory Ser. B **53** (1991), no. 2, 293–299.
- [8] ———, *The number of degree-restricted rooted maps on the sphere*, SIAM J. Discrete Math. **7** (1994), no. 1, 9–15.
- [9] E. A. Bender, Z. Gao, and L. B. Richmond, *The map asymptotics constant t_g* , Electron. J. Combin. **15** (2008), no. 1, Research paper 51, 8 pp.
- [10] O. Bernardi and M. Bousquet-Mélou, *Counting colored planar maps: Algebraicity results*, J. Combin. Theory Ser. B **101** (2011), no. 5, 315–377.
- [11] ———, *Counting coloured planar maps: Differential equations*, Comm. Math. Phys. **354** (2017), no. 1, 31–84.
- [12] O. Bernardi and G. Chapuy, *Counting unicellular maps on non-orientable surfaces*, Adv. in Appl. Math. **47** (2011), no. 2, 259–275.
- [13] V. Bonzom, G. Chapuy, and M. Dołęga, *Worksheet accompanying the present paper, available at <http://www.irif.fr/~chapuy/worksheets/countingmaps-nonoriented.html> or <http://www.irif.fr/~chapuy/worksheets/countingmaps-nonoriented.mw>*, 2021.
- [14] Valentin Bonzom, Guillaume Chapuy, Séverin Charbonnier, and Elba Garcia-Failde, *Topological recursion for Orlov-Scherbin tau functions, and constellations with internal faces*, 2022, <https://arxiv.org/abs/2206.14768>.
- [15] Valentin Bonzom, Guillaume Chapuy, and Maciej Dołęga, *b -Monotone Hurwitz Numbers: Virasoro Constraints, BKP Hierarchy, and $O(N)$ -BGW Integral*, Int. Math. Res. Not. IMRN (2022), article no. rnac177.
- [16] J. Bouttier and E. Guitter, *On irreducible maps and slices*, Combin. Probab. Comput. **23** (2014), no. 6, 914–972.
- [17] J. Bouttier, E. Guitter, and G. Miermont, *Bijjective enumeration of planar bipartite maps with three tight boundaries, or how to slice pairs of pants*, Preprint arXiv:2104.10084, 2021, <https://arxiv.org/abs/2104.10084>.
- [18] T. Budzinski and B. Louf, *Local limits of uniform triangulations in high genus*, Invent. Math. **223** (2020), no. 1, 1–47.
- [19] B. Bychkov, P. Dunin-Barkowski, M. Kazarian, and S. Shadrin, *Topological recursion for Kadomtsev-Petviashvili tau functions of hypergeometric type*, Preprint arXiv:2012.14723, 2020, <https://arxiv.org/abs/2012.14723>.
- [20] S. R. Carrell, *The Non-Orientable Map Asymptotics Constant p_g* , Preprint arXiv:1406.1760, June 2014, <https://arxiv.org/abs/1406.1760>.
- [21] S. R. Carrell and G. Chapuy, *Simple recurrence formulas to count maps on orientable surfaces*, J. Combin. Theory Ser. A **133** (2015), 58–75.
- [22] G. Chapuy and M. Dołęga, *A bijection for rooted maps on general surfaces*, J. Combin. Theory Ser. A **145** (2017), 252–307.
- [23] G. Chapuy, M. Marcus, and G. Schaeffer, *A bijection for rooted maps on orientable surfaces*, SIAM J. Discrete Math. **23** (2009), no. 3, 1587–1611.
- [24] L. Chekhov, B. Eynard, and N. Orantin, *Free energy topological expansion for the 2-matrix model*, J. High Energy Phys. (2006), no. 12, 053, 31.
- [25] M. Dołęga and M. Lepoutre, *Blossoming bijection for bipartite pointed maps and parametric rationality of general maps of any surface*, Adv. in Appl. Math. **141** (2022), Paper No. 102408.
- [26] B. Eynard, *Counting surfaces*, Progress in Mathematical Physics, vol. 70, Birkhäuser/Springer, [Cham], 2016, CRM Aisenstadt chair lectures.
- [27] B. Eynard and N. Orantin, *Invariants of algebraic curves and topological expansion*, Commun. Number Theory Phys. **1** (2007), no. 2, 347–452.
- [28] Bertrand Eynard, *An overview of the topological recursion*, in Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. III, Kyung Moon Sa, Seoul, 2014, pp. 1063–1085.

- [29] I. P. Goulden and D. M. Jackson, *The KP hierarchy, branched covers, and triangulations*, Adv. Math. **219** (2008), no. 3, 932–951.
- [30] V. Kac and J. van de Leur, *The geometry of spinors and the multicomponent BKP and DKP hierarchies*, in CRM Proceedings and Lecture Notes, vol. 14, 1998, pp. 159–202.
- [31] V. G. Kac, A. K. Raina, and N. Rozhkovskaya, *Bombay lectures on highest weight representations of infinite dimensional Lie algebras*, vol. 29, World Scientific, 2013.
- [32] V. A. Kazakov, I. K. Kostov, and N. Nekrasov, *D-particles, matrix integrals and KP hierarchy*, Nuclear Physics B **557** (1999), no. 3, 413–442.
- [33] M. Kazarian and P. Zograf, *Virasoro constraints and topological recursion for Grothendieck’s dessin counting*, Lett. Math. Phys. **105** (2015), no. 8, 1057–1084.
- [34] M. Ledoux, *A recursion formula for the moments of the Gaussian orthogonal ensemble*, Ann. Inst. Henri Poincaré Probab. Stat. **45** (2009), no. 3, 754–769.
- [35] M. Lepoutre, *Blossoming bijection for higher-genus maps*, J. Combin. Theory Ser. A **165** (2019), 187–224.
- [36] B. Louf, *Simple formulas for constellations and bipartite maps with prescribed degrees*, Canad. J. Math. **73** (2019), no. 1, 160–176.
- [37] T. Miwa, M. Jimbo, and E. Date, *Solitons*, Cambridge Tracts in Mathematics, vol. 135, Cambridge University Press, Cambridge, 2000. Differential equations, symmetries and infinite-dimensional algebras, Translated from the 1993 Japanese original by Miles Reid.
- [38] A. Okounkov, *Toda equations for Hurwitz numbers*, Math. Res. Lett. **7** (2000), no. 4, 447–453.
- [39] W. T. Tutte, *A census of planar triangulations*, Canad. J. Math **14** (1962), no. 1, 21–38.
- [40] ———, *A census of slicings*, Canad. J. Math **14** (1962), no. 4, 708–722.
- [41] ———, *A census of planar maps*, Canad. J. Math. **15** (1963), 249–271.
- [42] J. Van de Leur, *Matrix integrals and the geometry of spinors*, J. Nonlinear Math. Phys. **8** (2001), no. 2, 288–310.
- [43] T. R. S. Walsh and A. B. Lehman, *Counting rooted maps by genus. I*, J. Combinatorial Theory Ser. B **13** (1972), 192–218.

VALENTIN BONZOM, Université Sorbonne Paris Nord, LIPN, CNRS, UMR 7030, F-93430 Villetaneuse, France
E-mail: bonzom@lipn.univ-paris13.fr

GUILLAUME CHAPUY, CNRS, IRIF UMR 8243, Université Paris Cité.
E-mail: guillaume.chapuy@irif.fr

MACIEJ DOŁĘGA, Institute of Mathematics, Polish Academy of Sciences, ul. Śniadeckich 8, 00-956 Warszawa, Poland.
E-mail: mdolega@impan.pl