



Citation for published version:

Kinsella, AG, Tibbetts, JD, Stead, D & Cresswell, AJ 2022, 'N-tosylhydrazones as acceptors for nucleophilic alkyl radicals in photoredox catalysis: A short case study on possible side reactions', *Synthetic Communications*, vol. 52, no. 3, pp. 413-423. <https://doi.org/10.1080/00397911.2022.2028844>

DOI:

[10.1080/00397911.2022.2028844](https://doi.org/10.1080/00397911.2022.2028844)

Publication date:

2022

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

CC BY

This is an Accepted Manuscript of an article published by Taylor & Francis in *Synthetic Communications* on 21 January 2022, available online: <http://www.tandfonline.com/10.1080/00397911.2022.2028844>

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

***N*-Tosylhydrazones as acceptors for nucleophilic alkyl radicals in photoredox catalysis: a short case study on possible side reactions**

Anna G. Kinsella,¹ Joshua D. Tibbetts,¹ Darren Stead,² and Alexander J. Cresswell^{1,*}

¹ Department of Chemistry, University of Bath, Claverton Down, Bath, U.K., BA2 7AY

² AstraZeneca, Oncology R&D, Research & Early Development, Darwin Building, 310, Cambridge Science Park, Milton Road, Cambridge, CB4 0WG, U.K.


* Correspondence: a.j.cresswell@bath.ac.uk

Abstract: We report the attempted development of a photoredox-catalysed α -C–H alkylation reaction of aliphatic amine derivatives, using *N*-tosylhydrazones as radical alkylating partners. The original intention was to intercept α -aminoalkyl radical intermediates with *N*-tosylhydrazones, followed by expulsion of a sulfonyl radical by β -scission to generate *N*-H diazene species. Facile denitrogenation of these intermediates would remove all trace of the hydrazone moiety and provide a net C–H alkylation process. However, our plans were derailed by issues with the low reactivity of *N*-tosylhydrazones towards intermolecular capture by nucleophilic radicals, and a number of unexpected side reactions. Our findings, though unsuccessful, do serve to identify challenges for future researchers attempting to develop similar transformations.

Keywords: hydrazones; amines; radicals; photoredox; side reactions


Introduction

The centrality of aliphatic amines in drug discovery[1] and the ever increasing drive to expand chemical space in this area has spurred the design of new and creative approaches to these important building blocks.[2] As part of this effort, chemists have made significant strides in the area of C–H functionalisation of aliphatic amines, addressing C–H bonds α - and β - to nitrogen, as well as more remote C–H bonds.[3] For example, we recently reported a powerful new approach for the catalytic α -C–H alkylation of primary amines **1**, [4] using photoredox catalysis[5] in conjunction with hydrogen atom transfer (HAT) catalysis[6] (e.g., **1** + **2** \rightarrow **3**, Scheme 1a). Given the profound effect that introduction of simple alkyl groups – including methyl groups[7] – can have upon the physicochemical properties of drugs, we were motivated to develop an α -C–H alkylation reaction for alkylamines that could introduce such motifs without the need for transition metal co-catalysis.[3a,b] We were inspired by reports utilising *N*-tosylhydrazones as alkylating agents for either alkyllithium reagents[8] or alkyl radicals,[9] in each case relying upon denitrogenation of transient diazenes[10] to enable a ‘traceless’ C–C bond-forming process (e.g., **4** \rightarrow **8** from ref. 9c, Scheme 1b). Our intended reaction design was to harness established photoredox protocols for the generation of α -amino radicals **11**[11] from *N*-protected aliphatic amines **9** and intercept these radicals with *N*-tosylhydrazones **10**. A subsequent β -scission[12] event from hydrazinyl radical **12** would expel a sulfonyl radical (to oxidatively turnover the photoredox catalyst) and furnish *N*-H diazenes **13**. Subsequent denitrogenative decomposition of these species would give the desired α -alkylated amines **14** (Scheme 1c). However, a number of reactivity problems and unexpected side reactions conspired to derail our ambitions, including poor radicalophilicity of *N*-tosylhydrazones towards intermolecular capture by highly nucleophilic α -amino radicals, competitive SET reduction of the putative hydrazinyl radical intermediate **12**, and a suspected intramolecular HAT process possible with cyclic amine substrates (Scheme 1d).



Scheme 1. (a) Prior art: Catalytic γ -lactam synthesis from unprotected 1° alkylamines; (b) Inspiration: traceless alkyl-alkyl cross-coupling using *N*-tosylhydrazones; (c) Original aim of this work: metal-free α -C–H alkylation of amines; (d) Problems and side reactions.

As the studies described herein were nearing completion, König *et al.* reported the successful execution of a very similar concept to our intended ‘traceless’ alkylation, albeit with electrophilic thiyl (RS•) or trifluoromethyl (F₃C•) radicals *in lieu* of nucleophilic α -amino radicals (Scheme 2).[9a] In this instance, the *N*-tosylhydrazones are displaying *nucleophilic* reactivity[10] at carbon, in contrast to the *electrophilic* behaviour[8,9] we have envisioned. The successful β -scission step to expel a sulfonyl radical, leading to diazenes **18** and **24** as key intermediates, is also noteworthy, and contrasts with our own findings (*vide infra*).




Scheme 2. König’s merger of photoredox catalysis with Wolff-Kishner-type reactivity of *N*-tosylhydrazones.

Results and Discussion

Our investigations commenced with the intention of adding photogenerated simple alkyl radicals to *N*-tosylhydrazone **28** (derived from 4-fluorobenzaldehyde), using the *para*-fluoro group as a spectroscopic handle for ¹⁹F NMR (Scheme 3). Initial attempts with acridinium photocatalysts employing carboxylic acids (e.g., **27**) or trifluoroborates (i.e., **30**) were unsuccessful, with only a complex mixture generated in the former case and no reaction in the latter. Control experiments


verified that hydrazone **28** is sensitive to extensive decomposition on exposure to Cs_2CO_3 , even in the absence of light, likely *via* generation of the corresponding diazo species. We next tested the easily oxidised bis(catecholato)silicate **32** ($E_{\text{p}/2}^{\text{red}} = +0.69 \text{ V vs SCE}$ in DMF)[13] as an alkyl radical precursor, and 2,4,5,6-tetra(9*H*-carbazol-9-yl)isophthalonitrile (4CzIPN) **33** [$E_{1/2}(\text{PC}^*/\text{PC}^-) = +1.43 \text{ V vs SCE}$ in MeCN][14] as the photoredox catalyst. Again, with *N*-tosylhydrazone **28**, no reactivity was observed, but with the much more activated hydrazone **34** (derived from trifluoroacetaldehyde), a successful reaction ensued to give $\alpha\text{-CF}_3$ hydrazide **35** in 41% isolated yield (albeit alongside a number of other, unidentified minor products). This result suggests that simple aromatic *N*-tosylhydrazones like **28** are not sufficiently electrophilic to efficiently capture nucleophilic radicals [9a], but that strong electron-withdrawing groups on the hydrazone carbon (as in **34**) can remedy this situation. Unfortunately, this result also indicated that our originally conceived β -scission[12] to expel a sulfonyl radical and generate a transient *N*-H diazene is not operative in this case, and that the hydrazinyl radical generated after hydrazone addition is rapidly quenched *via* SET reduction by the reduced photocatalyst [$E_{1/2}(\text{PC}/\text{PC}^-) = -1.24 \text{ V vs SCE}$ in MeCN]. We also attempted to use an $\alpha\text{-CF}_3$ hydrazone substituted with a 4-(trifluoromethyl)phenylsulfonyl group instead of a Ts group, but the photoredox reaction with **32** was significantly messier and lower-yielding than that with hydrazone **34**, and the product could not be isolated by chromatography without extensive decomposition.



Scheme 3. Attempts at photocatalytic alkyl radical addition to *N*-tosylhydrazones.

We next investigated some other alkyl radical precursors[15] with $\alpha\text{-CF}_3$ hydrazone **34** as the acceptor, in the hope that we might identify an instance in which the β -scission step might be competitive with reduction of the *N*-centred radical (Scheme 4). With redox-active ester **36** as the radical precursor, a set of reaction conditions previously reported for photocatalytic conjugate additions[15a] gave $\alpha\text{-CF}_3$ hydrazide **35** in 38% NMR yield, but no sign of the desired alkyl-alkyl cross-coupling product (*via* a β -scission and denitrogenation pathway analogous to that in Scheme 1c). The outcome was similar for benzylic trifluoroborate **38**, from which the $\alpha\text{-CF}_3$ hydrazide **40** was isolated in 17% yield. In both cases, the crude product mixtures were complex and difficult to deconvolute *via* ^{19}F NMR, with several unidentified, minor fluorinated species also evident. We also attempted the reaction of


tetrahydrofuran **41** (as the solvent and radical precursor) with *N*-tosylhydrazone **34**, using eosin Y **42** as a photo-HAT catalyst.[15b] In this case also, an α -CF₃ hydrazide **43** was isolated in 59% yield (70:30 dr), but again with no sign of products resulting from a putative *N*-H diazene. Unfortunately this process did not prove to be general with other substrates (**44–48**), all of which gave no reaction, despite being used in a 5-fold excess, relative to **34** [15b].



Scheme 4. Further studies using other alkyl radical precursors.


Given that there is precedent for the controlled decomposition of *N*-tosyl hydrazides by a Wolff-Kishner-type pathway,[9a,b] we reasoned that our original goal of a 'traceless' α -C–H alkylation process for aliphatic amines may still be within reach, albeit as a two-step process (i.e., *via* an *N*-tosyl hydrazide intermediate). Thus, we next investigated the reactivity of *N*-tosylhydrazone **34** with *N,N*-dimethylaniline **49** – a known precursor to α -amino radicals *via* SET oxidation-deprotonation.[11] Using an iridium photocatalyst (**50**) previously employed with *N*-Ph amines,[16] the coupling of *N,N*-dimethylaniline **49** ($E_{1/2} = +0.74 \text{ V vs SCE}$)[17] with *N*-tosylhydrazone **34** generated hydrazide **51** in 37% isolated yield (Scheme 5a). Minor unidentified by-products were observed by ¹⁹F NMR, alongside remaining hydrazone starting material **34**. However, allowing the reaction to proceed until all of the starting hydrazone **34** had been consumed (41 h) led to a decrease in product **51** yield, suggesting that **51** may be decomposing under the conditions. Moreover, on repetition of this reaction, the yield of **51** and the extent of decomposition varied considerably. Attempts to stem the decomposition by varying both the photocatalyst and the reaction solvent (i.e., MeOH, MeCN, CH₂Cl₂) were fruitless. Moreover, employing a large excess of hydrazone **34** (8.0 equiv) conferred no benefit, providing an NMR yield of 30% of **51** after 20 h. Concurrently, the reaction of *N*-Ph pyrrolidine **52** ($E_{1/2} = +0.70 \text{ V vs SCE}$)[16] with *N*-tosylhydrazone **34** was also studied, and a hydrazide product **53** was isolated in 12% yield and >99:1 dr (though it is possible that a minor diastereomer was removed during chromatographic purification) (Scheme 5b). In this case, a new hydrazide species **54** (a reduction

product of **34**) was detected, and mass spectroscopic evidence consistent with enamine **55** was also found (i.e., m/z peaks of 146 for $[M+H]^+$ and 289 for $[2M+H]^+$).[18]



Scheme 5. (a) Photoredox coupling of *N,N*-dimethylaniline **49** with *N*-Ts hydrazone **34**. (b) Photoredox coupling of *N*-Ph pyrrolidine **52** with *N*-Ts hydrazone **34**.


Given that none of the reduced hydrazone by-product **54** was detected in the reaction of *N,N*-dimethylaniline **49**, this suggests that cyclic amine substrate **52** is prone to a side-reaction to afford **54** that is not possible with compound **49**. On the basis that *N*-Ph enamine **55** was also detected by MS analysis from **52** (see Scheme 5b), a speculative mechanistic rationalisation to account for these findings is presented in Scheme 6. Assuming that the reaction proceeds *via* α -amino radical attack onto the C=N bond of the CF_3 hydrazone **34**, the resultant hydrazinyl radical **57** may be able to engage in a 1,5-HAT with a C–H bond on the ring of the cyclic amine. β -Scission of the alkyl radical would then generate enamine by-product **55**, alongside the α - CF_3 radical species **59**. Proton transfer (to give **60**) could potentially be followed by an intermolecular HAT from the amine substrate, delivering the reduced hydrazone species **54**. Concomitant generation of an α -amino radical from **52** could then feasibly lead to a chain process.



Scheme 6. Proposed mechanistic rationale for the generation of reduced hydrazone by-product **54** when using cyclic amine **52**.


In parallel to investigations with *N*-Ph amines as radical precursors, *N*-Boc amines were also tested as substrates for a photoredox-catalysed α -C–H functionalisation with α - CF_3 hydrazone **34**. Given that carbamates possess much higher oxidation potentials than *N*-Ph amines (i.e., $E_{1/2} = +1.96$ V vs SCE for *N*-Boc piperidine)[19], we opted to use the strongly oxidising acridinium photocatalyst **62** [$E_{1/2}$ ($\text{PC}^*/\text{PC}^{\bullet-}$) = +2.15 V vs SCE in MeCN] recently developed by Nicewicz and co-workers.[19] Thus, *N*-Boc dimethylamine **61** in MeCN was reacted with α - CF_3 hydrazone **34** in the presence of acridinium

catalyst **62** (5 mol%) with irradiation by Kessil blue LEDs for 20 h (Scheme 7a). Under these conditions, the α -CF₃ hydrazone **63** was isolated in 63% yield. Unfortunately, attempts to reduce the stoichiometry of *N*-Boc dimethylamine **61** to 1.0 equivalent led to extensive by-product formation and a complex mixture. A similar reaction using *N*-Boc pyrrolidine **64** gave the corresponding α -CF₃ hydrazone **65** in only 39% isolated yield (67:33 dr). The reduced hydrazone by-product **54** was only observed at a trace level (<5% by NMR), suggesting that the side reaction operative for *N*-Ph pyrrolidine **52** does not make a significant contribution in this case (Scheme 7b).



Scheme 7. (a) Photoredox coupling of *N*-Boc dimethylamine **61** with *N*-Ts hydrazone **34**. (b) Photoredox coupling of *N*-Boc pyrrolidine **64** with *N*-Ts hydrazone **34**.

At this stage, we attempted a base-induced denitrogenative decomposition of *N*-Ts hydrazone products **51** and **63**, to gauge the possibility of developing a one-pot protocol for ‘traceless’ α -C–H alkylation (or, more specifically, trifluoroethylation if using **34**). Using literature conditions for *N*-sulfonyl hydrazone decomposition with NaOAc (*via* a radical pathway from an intermediate *N*-H diazene),^[9b,20] no reaction was observed. The complete lack of reactivity is surprising, and we postulate that the strongly inductively-withdrawing CF₃ group may render the transient *N*-centered anion too thermodynamically stable to eliminate the β -sulfinyl group. Further attempts to induce denitrogenative decomposition of **51** and **63** using the stronger base *t*-BuOK^[21] were also unsuccessful, with both compounds suffering extensive degradation to afford complex mixtures of unidentifiable products (Scheme 8). A β -elimination of fluoride ion from a putative α -CF₃ carbanion in the latter case may be to blame.^[22]



Scheme 8. Attempted denitrogenative decomposition of *N*-Ts hydrazides **51** and **63**.

Conclusions

The reactivity of *N*-tosylhydrazones as acceptors for nucleophilic radicals generated by photoredox catalysis has been briefly explored, as part of studies directed towards a ‘traceless’, metal-free α -C–H alkylation of aliphatic amine derivatives. The introduction of an electron-withdrawing trifluoromethyl group on the hydrazone carbon proved necessary to switch on reactivity for intermolecular capture with nucleophilic radicals, presumably by increasing the electrophilicity of what is otherwise a relatively nucleophilic site. The intended β -scission step from the hydrazinyl radical intermediates –

to expel a sulfonyl radical and generate *N*-H diazenes – was outcompeted by SET reduction by the reduced form of the photocatalyst, leading to *N*-Ts hydrazide products. A side reaction involving a suspected 1,5-HAT process and subsequent fragmentation appears to operate with *N*-Ph pyrrolidine as the substrate, eroding the yield of *N*-Ts hydrazide product. Attempted denitrogenative decomposition of *N*-Ts hydrazide products was not fruitful, and gave either no reaction or complex mixtures of unidentified species. Although unsuccessful, these studies do serve to identify challenges for future researchers attempting to develop similar radical-based transformations with *N*-tosylhydrazones.

Experimental

Synthesis of N-(1-cyclohexyl-2,2,2-trifluoroethyl)-4-methylbenzenesulfonylhydrazide (35)

An oven-dried, 20-mL scintillation vial equipped with a stirrer bar was charged with silicate **32** (96 mg, 0.22 mmol, 1.0 equiv), hydrazone **34** (238 mg, 0.89 mmol, 4.0 equiv) and 4CzIPN **33** (1.8 mg, 2.3 μ mol, 1 mol%). The vial was sealed with an upturned septa and evacuated and backfilled with nitrogen three times. DMF (anhydrous, 2.5 mL) was added and the mixture was sparged with nitrogen for 15 min. The resultant solution was stirred under irradiation (Kessil H150-Blue LED lamp) for 18 h. The reaction mixture was diluted with EtOAc (15 mL) and washed with H₂O (5 \times 15 mL). The organic layer was dried with a phase separating cartridge and then concentrated *in vacuo*. Purification *via* preparative HPLC (Waters CSH C₁₈ OBD column, 5 μ silica, 30 mm diameter, 100 mm length) using decreasingly polar mixtures of H₂O [containing 1% by volume of NH₄OH (28–30% in H₂O)] and MeCN as eluents gave **35** as a white, crystalline solid (32 mg, 41%). mp 161 °C. ¹H NMR (500 MHz, chloroform-*d*) 7.71 – 7.91 (m, 2H, ArH), 7.34 (d, *J* = 7.9 Hz, 2H, ArH), 6.10 (s, 1H, NH), 3.71 (d, *J* = 8.3 Hz, 1H, NH), 2.94–3.11 (m, 1H, C(1'')H), 2.45 (s, 3H, C(9)H₃), 1.57–1.78 (m, 5H, Cy), 1.49 (s, 1H, Cy), 1.04–1.25 (m, 4H, Cy), 0.83–1.01 (m, 1H, Cy). ¹³C NMR (126 MHz, DMSO-*d*₆) δ 143.7 (Ar), 137.0 (Ar), 130.2 (Ar), 128.7 (Ar), 127.4 (q, *J* = 287 Hz, C(2'')), 67.2 (q, *J* = 24.4 Hz, C(1'')), 37.3 (C(1)), 29.9 (Cy), 29.2 (Cy), 26.7 (Cy), 26.7 (Cy), 26.6 (Cy), 21.9 (C(9)). ¹⁹F NMR (376 MHz, chloroform-*d*) δ –73.3 (d, *J* = 8.0 Hz). IR (neat, cm⁻¹) 3297 (w), 3249 (w), 2923 (w), 2853 (w), 1601 (w), 1509 (w), 1497 (w), 1455 (w), 1354 (w), 1320 (w), 1258 (w), 1228 (w), 1189 (w), 1156 (m), 1145 (m), 1086 (w), 971 (w), 842 (w), 817 (w), 801 (w), 748 (w), 707 (w), 689 (w), 666 (w). HRMS (ESI⁺) calculated for C₁₅H₂₁F₃N₂O₂S: [M+H]⁺ 351.1354; found: [M+H]⁺ 351.1363.

Supplementary Materials: Supporting Information: Full experimental detail, ¹H and ¹³C NMR spectra. This material can be found *via* the “Supplementary Content” section of this article’s webpage.

Acknowledgments: The authors gratefully acknowledge the technical staff within Chemistry at the University of Bath for technical support and assistance in this work, including the Material and Chemical Characterisation Facility (MC²) (<https://doi.org/10.15125/mx6j-3r54>).

Funding: This work was funded by the Engineering and Physical Sciences Research Council (EP/S028595/1). A.J.C. thanks the Royal Society for a University Research Fellowship (UF150533), the University of Bath for a PhD studentship (A.G.K.), and AstraZeneca for generous financial support.

References

1. Blakemore, D. C.; Castro, L.; Churcher, I.; Rees, D. C.; Thomas, A. W.; Wilson, D. M.; Wood, A. [Organic Synthesis Provides Opportunities to Transform Drug Discovery](#). *Nat. Chem.* **2018**, *10*, 383–394.

2. Nugent, T. C. *Chiral Amine Synthesis: Methods, Developments and Applications*; Wiley-VCH, 2010.
3. Selected reviews: (a) Trowbridge, A.; Walton, S. M.; Gaunt, M. J. [New Strategies for the Transition-Metal Catalyzed Synthesis of Aliphatic Amines](#). *Chem. Rev.* **2020**, *120*, 2613–2692. (b) Gonnard, L.; Guérinot, A.; Cossy, J. [Transition metal catalyzed \$\alpha\$ -alkylation of amines by \$Csp^3\$ -H bond activation](#). *Tetrahedron* **2019**, *75*, 145–163. (c) Mitchell, E. A.; Peschiulli, A.; Lefevre, N.; Meerpoel, L.; Maes, B. U. W. [Direct \$\alpha\$ -Functionalization of Saturated Cyclic Amines](#). *Chem. Eur. J.* **2012**, *18*, 10092–10142. (d) Shi, L.; Xia, W. [Photoredox functionalization of C–H bonds adjacent to a nitrogen atom](#). *Chem. Soc. Rev.* **2012**, *41*, 7687–7697. (e) Campos, K. R. [Direct \$sp^3\$ C–H bond activation adjacent to nitrogen in heterocycles](#). *Chem. Soc. Rev.* **2007**, *36*, 1069–1084.
4. (a) Grayson, J. D.; Cresswell, A. J. [\$\gamma\$ -Amino Phosphonates via the Photocatalytic \$\alpha\$ -C–H Alkylation of Primary Amines](#). *Tetrahedron* **2021**, *81*, 131896. (b) Ryder, A. S. H.; Cunningham, W. B.; Ballantyne, G.; Mules, T.; Kinsella, A. G.; Turner-Dore, J.; Alder, C. M.; Edwards, L. J.; McKay, B. S. J.; Grayson, M. N.; Cresswell, A. J. [Photocatalytic \$\alpha\$ -Tertiary Amine Synthesis via C–H Alkylation of Unmasked Primary Amines](#). *Angew. Chem. Int. Ed.* **2020**, *59*, 14986–14991.
5. Reviews of visible-light photoredox catalysis: (a) Zilate, B.; Fischer, C.; Sparr, C. [Design and application of aminoacridinium organophotoredox catalysts](#). *Chem. Commun.* **2020**, *56*, 1767–1775. (b) McAtee, R. C.; McClain, E. J.; Stephenson, C. R. J. [Illuminating Photoredox Catalysis](#). *Trends Chem.* **2019**, *1*, 111–125. (c) Shang, T.-Y.; Lu, L.-H.; Cao, Z.; Liu, Y.; He, W.-M.; Yu, B. [Recent advances of 1,2,3,5-tetrakis\(carbazol-9-yl\)-4,6-dicyanobenzene \(4CzIPN\) in photocatalytic transformation](#). *Chem. Commun.* **2019**, *55*, 5408–5419. (d) *Visible light photocatalysis in organic chemistry*, Stephenson, C.R.J.; Yoon, T.; MacMillan, D.W.C., Eds.; Wiley-VCH, Berlin, 2018. (e) Marzo, L.; Pagire, S. K.; Reiser, O.; König, B. [Visible-Light Photocatalysis: Does It Make a Difference in Organic Synthesis?](#) *Angew. Chem. Int. Ed.* **2018**, *57*, 2–41. (f) Srivastava, V.; Singh, P. P. [Eosin Y catalysed photoredox synthesis: a review](#). *RSC Adv.* **2017**, *7*, 31377–31392. (g) Romero, N. A.; Nicewicz, D. A. [Organic Photoredox Catalysis](#). *Chem. Rev.* **2016**, *116*, 10075–10166. (h) Ravelli, D.; Protti, S.; Fagnoni, M. [Carbon-Carbon Bond Forming Reactions via Photogenerated Intermediates](#). *Chem. Rev.* **2016**, *116*, 9850–9913. (i) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C. [Photoredox Catalysis in Organic Chemistry](#). *J. Org. Chem.* **2016**, *81*, 6898–6926. (j) Schultz, D. M.; Yoon, T. P. [Solar Synthesis: Prospects in Visible Light Photocatalysis](#). *Science* **2014**, *343*, 1239176. (k) Prier, C. K.; Rankic, D. A.; Macmillan, D. W. C. [Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis](#). *Chem. Rev.* **2013**, *113*, 5322–5363.
6. Selected reviews on HAT catalysis, see: (a) Chu, J. C. K.; Rovis, T. [Complementary Strategies for Directed \$C\(sp^3\)\$ -H Functionalization: A Comparison of Transition-Metal-Catalyzed Activation, Hydrogen Atom Transfer, and Carbene/Nitrene Transfer](#). *Angew. Chem. Int. Ed.* **2018**, *57*, 62–101. (b) Capaldo, L.; Ravelli, D. [Hydrogen Atom Transfer \(HAT\): A Versatile Strategy for Substrate Activation in Photocatalyzed Organic Synthesis](#). *Eur. J. Org. Chem.* **2017**, 2056–2071.
7. (a) Aynedtinova, D.; Callens, M. C.; Hicks, H. B.; Poh, C. Y. X.; Shennan, B. D. A.; Boyd, A. M.; Lim, Z. H.; Leitch, J. A.; Dixon, D. J. [Installing the "magic methyl"- C–H methylation in synthesis](#). *Chem. Soc. Rev.* **2021**, *50*, 5517–5563. (b) Schonherr, H.; Cernak, T. [Profound Methyl Effects in Drug Discovery and a Call for New C–H Methylation Reactions](#). *Angew. Chem. Int. Ed.* **2013**, *52*, 12256–12267.

8. For reactions of *N*-silylated/metalated *N*-Ts hydrazones with organometallic nucleophiles, followed by denitrogenative decomposition see: (a) Myers, A. G.; Movassaghi, M. [Highly Efficient Methodology for the Reductive Coupling of Aldehyde Tosylhydrazones with Alkylolithium Reagents](#). *J. Am. Chem. Soc.* **1998**, *120*, 8891–8892. (b) Myers, A. G.; Kukkola, P. J. [Stereoselective synthesis of olefins from silylated sulfonylhydrazones](#). *J. Am. Chem. Soc.* **1990**, *112*, 8208–8210. (c) Vedejs, E.; Dolphin, J.; Stolle, W. [A new olefin synthesis: condensation of aldehyde tosylhydrazones with stabilized carbanions](#). *J. Am. Chem. Soc.* **1979**, *101*, 249–251. (d) Vedejs, E.; Stolle, W. T. [Reductive alkylation of aldehyde tosylhydrazones with organolithium reagents](#). *Tetrahedron Lett.* **1977**, *18*, 135–138.
9. (a) Wang, S.; Cheng, B.-Y.; Sršen, M.; König, B. [Umpolung Difunctionalization of Carbonyls via Visible-Light Photoredox Catalytic Radical-Carbanion Relay](#). *J. Am. Chem. Soc.* **2020**, *142*, 7524–7531. (b) Dao, H. T.; Li, C.; Michaudel, Q.; Maxwell, B. D.; Baran, P. S. [Hydromethylation of Unactivated Olefins](#). *J. Am. Chem. Soc.* **2015**, *137*, 8046–8049. (c) Campbell, N. E.; Sammis, G. M. [Single-Electron/Pericyclic Cascade for the Synthesis of Dienes](#). *Angew. Chem. Int. Ed.* **2014**, *53*, 6228–6231. (d) Kim, S.; Cho, J. R. [Radical Cyclization of Mesitylsulfonylhydrazones](#). *Synlett* **1992**, 629–630.
10. For umpolung reactions of *N*-deprotonated *N*-Ts hydrazones with electrophiles, followed by denitrogenative decomposition see: (a) Reyes, J. R.; Rawal, V. H. [Reductive Chlorination and Bromination of Ketones via Trityl Hydrazones](#). *Angew. Chem. Int. Ed.* **2016**, *128*, 3129–3132. (b) Baldwin, J. E.; Adlington, R. M.; Bottaro, J. C.; Kolhe, J. N.; Newington, I. M.; Perry, M. W. D. [Azo Anions in Synthesis: Use of Trityl- and Diphenyl-4-pyridylmethylhydrazones for Reductive C–C Bond Formation](#). *Tetrahedron* **1986**, *42*, 4235–4246. (c) Baldwin, J. E.; Bottaro, J. C.; Kolhe, J. N.; Adlington, R. M. [Azo Anions in Synthesis. Use of Trityl- and Diphenyl-4-Pyridylmethyl-Hydrazones for Reductive C–C Bond Formation from Aldehydes and Ketones](#). *J. Chem. Soc., Chem. Commun.* **1984**, 22–23.
11. For reviews, see: (a) Ref. 3a. (b) Leitch, J. A.; Rossolini, T.; Rogova, T.; Maitland, J. A. P.; Dixon, D. J. [\$\alpha\$ -Amino Radicals via Photocatalytic Single-Electron Reduction of Imine Derivatives](#). *ACS Catal.* **2020**, *10*, 2009–2025. (c) Y.-Q. Zou, W.-J. Xiao in *Visible light photocatalysis in organic chemistry* (Eds.: C. R. J. Stephenson, T. Yoon, D. W. C. MacMillan), Wiley-VCH, Berlin, **2018**, Chap. 4, pp 93–128. (d) K. Nakajima, Y. Miyake, Y. Nishibayashi, *Acc. Chem. Res.* **2016**, *49*, 1946–1956.
12. Zhang, H.; Hay, E. B.; Geib, S. J.; Curran, D. P. [Fates of imine intermediates in radical cyclizations of *N*-sulfonylindoles and ene-sulfonamides](#). *Beilstein J. Org. Chem.* **2015**, *11*, 1649–1655.
13. Corcé, V.; Chamoreau, L. M.; Derat, E.; Goddard, J. P.; Ollivier, C.; Fensterbank, L. [Silicates as Latent Alkyl Radical Precursors: Visible-Light Photocatalytic Oxidation of Hypervalent Bis-Catecholato Silicon Compounds](#). *Angew. Chem. Int. Ed.*, **2015**, *54*, 11414–11418.
14. Luo, J.; Zhang, J. [Donor–Acceptor Fluorophores for Visible-Light-Promoted Organic Synthesis: Photoredox/Ni Dual Catalytic C\(sp³\)-C\(sp²\) Cross Coupling](#). *ACS Catal.* **2016**, *6*, 873–877.
15. (a) Pratsch, G.; Lackner, G. L.; Overman, L. E. *J. Org. Chem.* **2015**, *80*, 6025–6036. (b) Fan, X.-Z.; Rong, J.-W.; Wu, H.-L.; Zhou, Q.; Deng, H.-P.; Da Tan, J.; Xue, C.-W.; Wu, L.-Z.; Tao, H.-R.; Wu, J. [Eosin Y as a Direct Hydrogen-Atom Transfer Photocatalyst for the Functionalization of C–H Bonds](#). *Angew. Chem. Int. Ed.* **2018**, *57*, 8514–8518.

16. Ahneman, D. T.; Doyle, A. G. [C–H functionalization of amines with aryl halides by nickel-photoredox catalysis](#). *Chem. Sci.* **2016**, *7*, 7002–7006.
17. Das, P.; Begam, H. M.; Bhunia, S.; Jana, R. [Photoredox-Catalyzed Tandem Demethylation of *N,N*-Dimethyl Anilines Followed by Amidation with \$\alpha\$ -Keto or Alkynyl Carboxylic Acids](#). *Adv. Synth. Catal.* **2019**, *361*, 4048–4054.
18. Analysis of the ^1H NMR spectrum was not fruitful, given the complexity of the mixture and signal overlap.
19. McManus, J. B.; Onuska, N. P. R.; Nicewicz, D. A. [Generation and Alkylation of \$\alpha\$ -Carbamyl Radicals via Organic Photoredox Catalysis](#). *J. Am. Chem. Soc.* **2018**, *140*, 9056–9060.
20. Zhang, J.; Eisink, N. N. H. M.; Witte, M. D.; Minnaard, A. J. [Regioselective Manipulation of GlcNAc Provides Allosamine, Lividosamine, and Related Compounds](#). *J. Org. Chem.* **2019**, *84*, 516–525.
21. Kedrowski, S. M. A.; Dougherty, D. A. [Room-Temperature Alternative to the Arbuzov Reaction: The Reductive Deoxygenation of Acyl Phosphonates](#). *Org. Lett.* **2010**, *12*, 3990–3993.
22. Ramanathan, S.; Lemal, D. M. [Synthesis, Mechanism of Formation, and Dynamics of a Highly Fluorinated Methylenecyclobutene](#). *J. Org. Chem.* **2007**, *72*, 1570–1574.