

Association between XRCC3 p.Thr241Met polymorphism and risk of glioma

Tan, Shing Cheng; Low, Teck Yew; Hussain, Hafiz Muhammad Jafar; Sharzehan, Mohamad Ayub Khan; Sito, Hilary; Kord-Varkaneh, Hamed; Islam, Md Asiful

DOI:

[10.1371/journal.pone.0276313](https://doi.org/10.1371/journal.pone.0276313)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Tan, SC, Low, TY, Hussain, HMJ, Sharzehan, MAK, Sito, H, Kord-Varkaneh, H & Islam, MA 2022, 'Association between XRCC3 p.Thr241Met polymorphism and risk of glioma: a systematic review and meta-analysis', *PLoS ONE*, vol. 17, no. 10, e0276313. <https://doi.org/10.1371/journal.pone.0276313>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

RESEARCH ARTICLE

Association between *XRCC3* p.Thr241Met polymorphism and risk of glioma: A systematic review and meta-analysis

Shing Cheng Tan¹*, Teck Yew Low¹, Hafiz Muhammad Jafar Hussain², Mohamad Ayub Khan Sharzehan¹, Hilary Sito¹, Hamed Kord-Varkaneh³, Md Asiful Islam⁴*

1 UKM Medical Molecular Biology Institute, Universiti Kebangsaan Malaysia, Kuala Lumpur, Malaysia, **2** Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, United States of America, **3** Department of Clinical Nutrition and Dietetics, Student Research Committee, Faculty of Nutrition and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran, **4** Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, United Kingdom

* These authors contributed equally to this work.

* sctan@ukm.edu.my, shingchengtan@gmail.com (SCT); ayoncx70@yahoo.com (MAI)



OPEN ACCESS

Citation: Tan SC, Low TY, Hussain HMJ, Sharzehan MAK, Sito H, Kord-Varkaneh H, et al. (2022) Association between *XRCC3* p.Thr241Met polymorphism and risk of glioma: A systematic review and meta-analysis. PLoS ONE 17(10): e0276313. <https://doi.org/10.1371/journal.pone.0276313>

Editor: Yanhong Liu, Baylor College of Medicine, UNITED STATES

Received: October 16, 2021

Accepted: October 4, 2022

Published: October 20, 2022

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0276313>

Copyright: © 2022 Tan et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Abstract

Background

The *XRCC3* p.Thr241Met (rs861539) polymorphism has been extensively studied for its association with glioma risk, but results remain conflicting. Therefore, we performed a systematic review and meta-analysis to resolve this inconsistency.

Methods

Studies published up to June 10, 2022, were searched in PubMed, Web of Science, Scopus, VIP, Wanfang, and China National Knowledge Infrastructure databases and screened for eligibility. Then, the combined odds ratio (OR) of the included studies was estimated based on five genetic models, i.e., homozygous (Met/Met vs. Thr/Thr), heterozygous (Thr/Met vs. Thr/Thr), dominant (Thr/Met + Met/Met vs. Thr/Thr), recessive (Met/Met vs. Thr/Thr + Thr/Met) and allele (Met vs. Thr). The study protocol was preregistered at PROSPERO (registration number: CRD42021235704).

Results

Overall, our meta-analysis of 14 eligible studies involving 12,905 subjects showed that the p.Thr241Met polymorphism was significantly associated with increased glioma risk in both homozygous and recessive models (homozygous, OR = 1.381, 95% CI = 1.081–1.764, P = 0.010; recessive, OR = 1.305, 95% CI = 1.140–1.493, P < 0.001). Subgroup analyses by ethnicity also revealed a statistically significant association under the two aforementioned genetic models, but only in the Asian population and not in Caucasians (P > 0.05).

Funding: Research in SCT's laboratory is supported by the Research University Grant of Universiti Kebangsaan Malaysia (No. GUP-2020-076), the Fundamental Research Grant Scheme of the Ministry of Higher Education, Malaysia (No. FRGS/1/2019/SKK08/UKM/02/9), and the Higher Institution Center of Excellence (HiCoE) grant of the Ministry of Higher Education, Malaysia (No. AKU49). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of this manuscript.

Competing interests: The authors declare no conflicts of interest.

Conclusion

We demonstrated that the *XRCC3* p.Thr241Met polymorphism is associated with an increased risk of glioma only in the homozygous and recessive models.

Introduction

Gliomas refer to a type of tumor that originates from the glial cells located in the brain or spine. This type of cancer accounts for ~30% of tumors of the brain and central nervous system and ~80% of total cases of malignant brain tumors [1, 2]. The tumor has a five-year survival rate of approximately 60%. However, this rate can be improved to as high as 74% if adequate surveillance by biopsy is performed in conjunction with early resection for low-grade gliomas [3, 4]. Depending on their aggressiveness, gliomas can be classified into four different grades, namely grade I, II, III, and IV. While grade I and II gliomas (low-grade gliomas) are comparatively benign and grow slowly, grade III (malignant glioma) and IV (glioblastoma multiforme or glioblastoma) gliomas are very aggressive and grow rapidly. Notably, glioblastoma is the most common and deadly form of primary malignant glioma in adults. It accounts for ~70% of gliomas and has a median survival of only 12–14 months [2, 5].

Although gliomas are a heterogeneous disease with comparatively unclear etiology, risk factors for gliomas are thought to consist of genetic predisposition and environmental influences [2, 3]. Genetic disorders such as type 1 and type 2 neurofibromatosis and tuberous sclerosis complex [6], as well as obesity and body height [7], have been shown to predispose to gliomas. In addition, several genome-wide association studies (GWAS) have reported the involvement of multiple genetic polymorphisms in mediating glioma risk [8, 9]. On the other hand, environmental factors such as ionizing radiation, ultraviolet radiation, cytomegalovirus infection, environmental carcinogens, and diet have been associated with an increased risk of developing glioma, although only the first two risk factors have been clearly demonstrated [3, 10–13].

Among a variety of genetic factors, germline polymorphisms have been widely associated with cancer risk [14, 15]. In particular, polymorphisms of DNA repair genes such as *ERCC1* [16], *ERCC2 (XPD)* [17], *XRCC1* [18], *XRCC3* [18], *XRCC4* [19], *XRCC5*, *XRCC6* [20], *XRCC7* [21], *MGMT* [22], *CHAF1* [23], *LIG4* [19], and *GLTSCR1* [17] have been associated with glioma risk [24, 25]. As genetic material, DNA is constantly exposed to endogenous and exogenous attacks that can manifest themselves in cellular metabolic processes and genotoxic or clastogenic stresses, including ionizing radiation and ultraviolet radiation mentioned earlier. These attacks can cause cross-links between DNA and proteins, oxidative damage to DNA, and single- and double-strand breaks in DNA chains [26]. However, this damage, which can affect the integrity of the genome, is continuously and effectively repaired by various inherent DNA repair pathways. These cellular DNA repair pathways can function in several ways and rely on the mechanisms of homologous recombination repair (HRR), nonhomologous end joining, base excision repair, and nucleotide excision repair. However, common polymorphisms in these DNA repair genes can impair DNA repair, leading to a higher risk of developing gliomas and other forms of malignancy [18–22].

This meta-analysis focuses on the *XRCC3* (X-ray repair cross complementing 3) gene, which encodes a RecA/Rad51-related protein involved in HRR. Of the polymorphisms in *XRCC3*, only p.Thr241Met (rs861539) has been frequently studied with respect to its association with glioma risk. This polymorphism is located in exon 7 of the gene and involves a C-to-T transition at codon 241 [27]. Unlike other polymorphisms in *XRCC3*, the nonsynonymous

p.Thr241Met polymorphism causes the replacement of a polar amino acid (threonine) with a nonpolar one (methionine), which can significantly affect the functionality of the protein product. For this reason, this polymorphism has been studied in numerous malignancies, including bladder cancer, head and neck cancer, ovarian cancer, and colorectal cancer, and significant associations have been demonstrated in some cases [28–32]. However, in gliomas, the association between this polymorphism and susceptibility to this cancer remains unclear [18, 21, 24, 33–44]. To clarify this, we performed a meta-analysis to further investigate the association between the XRCC3 p.Thr241Met polymorphism and glioma risk.

Materials and methods

Literature search

We comprehensively searched PubMed, Web of Science, Scopus, VIP, Wanfang, and China National Knowledge Infrastructure databases for relevant studies, including grey literature, published through June 10, 2022. The terms used in these searches were: (brain tumor OR glioma OR glioblastoma OR astrocytoma OR oligodendroglioma OR GBM OR glioblastoma multiforme) AND (XRCC3 OR X-Ray repair cross complementing 3 OR DNA repair gene) AND (polymorphism OR mutation OR variant OR variation). No language restriction was applied. We then included studies that met our prespecified inclusion criteria, i.e., (i) examined the association between the XRCC3 p.Thr241Met polymorphism and glioma risk, (ii) were observational studies, such as case-control studies, and (iii) reported genotype and/or allele frequencies or adequate information to infer these frequencies. We focused only on observational studies because other types of studies, such as randomized controlled trials, are not ethically acceptable in the context of genetic studies. In contrast, we excluded a study if (i) it was not an original study, (ii) the research was performed on cell lines, animal models, or other non-human subjects, and (iii) the study was a duplication of other publications. We also pre-registered our study protocol with PROSPERO (registration number: CRD42021235704).

Extraction of data and quality appraisal

Two investigators independently extracted the following information from the eligible studies: (i) first author's name, (ii) publication year, (iii) ethnicity, (iv) source of controls (hospital-based vs. population-based), (v) country where the study was conducted, (vi) number of subjects per group, (vii) frequency of polymorphic genotypes and alleles, and (viii) Hardy-Weinberg equilibrium (HWE) of genotype distribution in controls. If data on HWE were missing in any of the studies, a goodness-of-fit test was performed. If data were missing, the corresponding author of the included studies was contacted by email. The quality of the studies was then assessed using a modified version of the Newcastle-Ottawa Scale [45], which assessed (i) whether the definition of cases was appropriate, (ii) whether cases were representative of populations, (iii) whether controls were recruited from the community and the genotype distribution conformed to HWE, (iv) whether controls were defined as having no history of disease, (v) whether cases and controls were comparable in terms of ethnic homogeneity, (vi) whether there was no evidence of population stratification, (vii) whether quality control procedures and blinding were used in genotyping, (viii) whether cases and controls were genotyped using the same method, and (ix) whether the genotyping call rate exceeded 99% (S1 File). If the answer to each of the above criteria was "yes", the study received one star for each criterion. A study had to receive at least 5 stars to be considered of high methodological quality.

Meta-analysis

STATA (ver. 16.0) was used to pool data from included studies. $P < 0.05$ was considered statistically significant, unless otherwise stated. The association between XRCC3 p.Thr241Met polymorphism and glioma risk was assessed using five genetic models: homozygous (Met/Met vs. Thr/Thr), heterozygous (Thr/Met vs. Thr/Thr), dominant (Thr/Met + Met/Met vs. Thr/Thr), recessive (Met/Met vs. Thr/Thr + Thr/Met), and allele (Met vs. Thr). Examination of multiple genetic models was required because genetic association studies do not adopt a specific model [46]. Subsequently, Cochran's Q and I^2 tests were used to measure the heterogeneity of the studies, which was considered high when $P < 0.1$ or $I^2 > 50\%$ [47]. In this case, a random-effects method was used to derive the odds ratio (OR). On the other hand, when heterogeneity was low, a fixed-effects method was used. The Z test was then used to measure the significance of the pooled OR, and forest plots were created to visually display the results. Subgroup analyses were then conducted by several variables, including ethnicity, source of control, and HWE status, as these variables have long been known to affect genetic associations [48–50]. To examine whether the results were primarily influenced by one of the studies, we also performed a sensitivity analysis using the leave-one-out method. Publication bias was assessed using Begg's test, and Egger's test, and visual inspection of funnel plots.

Results

Study selection

We first identified 2,052 entries from the search databases (PubMed, N = 1,127; Scopus, N = 46; WoS, N = 838; CNKI, N = 9; Wanfang, N = 19; VIP, N = 13). We then removed 488 duplicate entries and subjected the remaining 1,564 articles to title and abstract screening. This process identified only 27 articles as potentially relevant. We then screened these 27 articles for eligibility by reviewing the full texts. This resulted in a further exclusion of 13 articles, so that finally only 14 articles describing 14 relevant studies were considered for quantitative data synthesis. The flow diagram of the study selection is shown in Fig 1.

Characteristics of the studies

The final 14 eligible studies included a total of 12,905 study participants, including 5,852 cases and 7,053 controls (Table 1). Eight (8) of these 14 studies were conducted in China, with subjects from the Asian population [28, 33, 37, 39, 40, 42–44]; the other seven studies were predominantly Caucasian—with three studies from the United States [21, 24, 36] and one each from Spain [41] and Brazil [34], and one study from multiple countries [18]. In addition, 11 studies used hospital-based controls [21, 24, 33, 36, 37, 39–44], whereas the remaining 3 studies opted for population-based controls [18, 34, 36]. In addition, all studies except Liu et al. [36] reported the frequencies of the three genotypes (Thr/Thr, Thr/Met, and Met/Met) separately. Instead, Liu et al. 2009 [36] combined the homozygous wild type (Thr/Thr) and heterozygous genotypes (Thr/Met). Moreover, of all 14 studies, the distribution of control genotypes did not deviate significantly from HWE in only five studies ($P > 0.05$) [21, 33, 37, 39, 41]. Nevertheless, all 14 studies were of high quality (S1 Table).

Quantitative data synthesis

The combined results of the 14 studies are shown in Table 2, and the corresponding forest plots are shown in Fig 2. High statistical heterogeneity was detected under in all genetic models; therefore, a random-effects model was used in the data synthesis. We detected a statistically significant association between the XRCC3 p.Thr241Met polymorphism and increased glioma




Fig 1. Flow diagram of study selection.

<https://doi.org/10.1371/journal.pone.0276313.g001>

Table 1. Characteristics of the included studies.

Study ID ^{reference}	Country	Ethnicity	Source of control*	Cases			Controls			HWE P-value (controls)
				Thr/Thr	Thr/Met	Met/Met	Thr/Thr	Thr/Met	Met/Met	
Huang 2015 [39]	China	Asian	HB	310	72	7	239	111	8	0.237
Gao 2014 [40]	China	Asian	HB	158	146	22	202	159	15	0.016
Rodriguez-Hernandez 2014 [41]	Spain	Caucasian	HB	43	56	16	87	92	21	0.646
Xu 2014 [42]	China	Asian	HB	472	343	71	485	356	45	0.047
Luo 2013 [43]	China	Asian	HB	145	131	21	229	168	17	0.042
Pan 2013 [44]	China	Asian	HB	217	198	28	234	200	9	<0.001
Zhao 2013 [33]	China	Asian	HB	336	47	1	340	41	3	0.165
Custódio 2012 [34]	Brazil	Caucasian	PB	53	18	9	86	9	5	<0.001
Liu 2012 [35]	China	Asian	HB	223	154	66	254	147	42	0.003
Rajaraman 2010 [24]	USA	Caucasian	HB	135	162	53	185	208	86	0.042
Liu 2009 [36]	USA	Caucasian	PB	308		61	315		45	-
Zhou 2009 [37]	China	Asian	HB	677	80	3	629	75	4	0.286
Kiuru 2008 [18]	Multiple	Caucasian	PB	288	319	94	630	761	169	0.006
Wang 2004 [21]	USA	Caucasian	HB	134	138	37	147	147	48	0.254

* HB, hospital-based; PB, population-based

<https://doi.org/10.1371/journal.pone.0276313.t001>

risk when the homozygous model and the recessive model were applied (homozygous, OR = 1.381, 95% CI = 1.081–1.764, P = 0.010; recessive, OR = 1.305, 95% CI = 1.140–1.493, P < 0.001). However, no significant association was found in the heterozygous (OR = 1.040, 95% CI = 0.908–1.191, P = 0.574), dominant (OR = 1.082, 95% CI = 0.939–1.247, P = 0.277), and allele (OR = 1.097, 95% CI = 0.972–1.237, P = 0.133) models. Sensitivity analysis revealed that the pooled OR was not significantly affected by any of the studies (S1 Fig).

Subgroup analysis

In the ethnicity-based subgroup analysis, we again found a statistically significant association between XRCC3 p.Thr241Met and increased glioma risk in the homozygous (OR = 1.704, 95% CI = 1.358–2.317, P < 0.001) and recessive models (OR = 1.305, 95% CI = 1.140–1.493, P < 0.001) in the Asians (Table 2). In contrast, no significant association was observed in Caucasians in all genetic models (P > 0.05). Subgroup analysis by source of controls revealed that the association in hospital-based controls was significant only in the homozygous model (OR = 1.364, 95% CI = 1.015–1.832, P = 0.040), whereas in population-based controls, the polymorphism was associated with increased glioma risk only in the recessive model (OR = 1.338, 95% CI = 1.071–1.670, P = 0.010). Interestingly, subgroup analysis by HWE status revealed no significant association for studies whose genotype distribution conformed to the equilibrium. For studies in which the genotype distribution deviated significantly from HWE, a significant association was found in the homozygous (OR = 1.588, 95% CI = 1.200–2.102, P = 0.001), dominant (OR = 1.124, 95% CI = 1.027–1.229, P = 0.011), recessive (OR = 1.518, 95% CI = 1.164–1.981, P = 0.002), and allele (OR = 1.185, 95% CI = 1.053–1.333, P = 0.005) models.

Publication bias

We did not detect any noticeable asymmetry in any of the funnel plots (Fig 3), indicating that there was no publication bias. This finding was also supported by the results of Begg's and Egger's tests (homozygous, Begg's P = 1.000, Egger's P = 0.811; heterozygous, Begg's P = 0.222,

Table 2. Summary of the association between XRCC3 p.Thr241Met polymorphism and glioma risk.

Comparison model	No. of studies	No. of cases	No. of controls	Effect model	OR (95% CI)	P
Homozygous model						
Overall	13	3,619	4,219	Random	1.381 (1.081–1.764)	0.010
Asian	8	2,757	2,755	Fixed	1.704 (1.358–2.137)	<0.001
Caucasian	5	862	1,464	Fixed	1.094 (0.895–1.338)	0.378
Hospital-based controls	11	3,175	3,329	Random	1.364 (1.015–1.832)	0.040
Population-based controls	2	444	890	Random	1.570 (0.720–3.425)	0.257
Conform to HWE	5	1,564	1,526	Fixed	0.915 (0.636–1.317)	0.634
Deviate from HWE	8	2,055	2,693	Random	1.588 (1.200–2.102)	0.001
Heterozygous model						
Overall	13	5,055	6,221	Random	1.040 (0.908–1.191)	0.574
Asian	8	3,709	3,869	Random	1.009 (0.841–1.210)	0.925
Caucasian	5	1,346	2,352	Random	1.102 (0.872–1.394)	0.415
Hospital-based controls	11	4,377	4,735	Random	1.028 (0.896–1.180)	0.694
Population-based controls	2	678	1,486	Random	1.602 (0.468–5.480)	0.453
Conform to HWE	5	1,893	1,908	Random	0.923 (0.663–1.284)	0.633
Deviate from HWE	8	3,162	4,313	Fixed	1.060 (0.964–1.164)	0.228
Dominant model						
Overall	13	5,483	6,693	Random	1.082 (0.939–1.247)	0.277
Asian	8	3,928	4,012	Random	1.056 (0.872–1.279)	0.578
Caucasian	5	1,555	2,681	Random	1.129 (0.888–1.435)	0.321
Hospital-based controls	11	4,702	5,033	Random	1.058 (0.915–1.224)	0.446
Population-based controls	2	781	1,660	Random	1.649 (0.527–5.162)	0.390
Conform to HWE	5	1,957	1,992	Random	0.914 (0.667–1.253)	0.577
Deviate from HWE	8	3,526	4,701	Fixed	1.124 (1.027–1.229)	0.011
Recessive model						
Overall	14	5,852	7,053	Fixed	1.305 (1.140–1.493)	<0.001
Asian	8	3,928	4,012	Fixed	1.655 (1.327–2.065)	<0.001
Caucasian	6	1,924	3,041	Fixed	1.132 (0.954–1.343)	0.155
Hospital-based controls	11	4,702	5,033	Random	1.316 (0.992–1.748)	0.057
Population-based controls	3	1,150	2,020	Fixed	1.338 (1.071–1.670)	0.010
Conform to HWE	5	1,957	1,992	Fixed	0.911 (0.645–1.286)	0.594
Deviate from HWE	8	3,526	4,701	Random	1.518 (1.164–1.981)	0.002
Allele model						
Overall	13	5,483	6,693	Random	1.097 (0.972–1.237)	0.133
Asian	8	3,928	4,012	Random	1.087 (0.923–1.281)	0.316
Caucasian	5	1,555	2,681	Random	1.106 (0.910–1.344)	0.311
Hospital-based controls	11	4,702	5,033	Random	1.069 (0.942–1.214)	0.302
Population-based controls	2	781	1,660	Random	1.621 (0.625–4.203)	0.320
Conform to HWE	5	1,957	1,992	Random	0.915 (0.714–1.174)	0.486
Deviate from HWE	8	3,526	4,701	Random	1.185 (1.053–1.333)	0.005

<https://doi.org/10.1371/journal.pone.0276313.t002>

Egger's P = 0.185; dominant, Begg's P = 0.542, Egger's P = 0.340; recessive, Begg's P = 0.784, Egger's P = 0.857; allele, Begg's P = 0.542, Egger's P = 0.643).

Discussion

The XRCC3 protein belongs to the RecA/Rad51-related family of proteins involved in HRR. This protein is encoded by a ~18kb gene, XRCC3, located on chromosome 14q32.33 [51]. The




Fig 2. Forest plots of the association between XRCC3 p.Thr241Met polymorphism and glioma risk.

<https://doi.org/10.1371/journal.pone.0276313.g002>




Fig 3.

<https://doi.org/10.1371/journal.pone.0276313.g003>

gene product is a ~38kDa protein that belongs to the five paralogs of RAD51, i.e., XRCC2, XRCC3, RAD51B, RAD51C, and RAD51D, all of which share 20–30% structural similarity with RAD51 [52]. These paralogs typically function by forming complex assemblies that can facilitate RAD51-mediated activities to repair DNA double-strand breaks via HRR. Therefore, depletion of any of these paralogs often leads to a reduction in the DNA damage response [53]. Two important protein complexes formed by these paralogs are (i) the BCDX2 complex, which includes XRCC2, RAD51B, RAD51C, and RAD51D, and (ii) the CX3 complex, which includes XRCC3 as well as RAD51C [54, 55]. These two complexes have been reported to be involved in two distinct steps of HRR: the BCDX2 complex contributes to the stabilization of the nucleoprotein filament of RAD51, whereas the CX3 complex acts after the recruitment of RAD51 to the damage sites [53]. Although the exact biochemical mechanisms of the CX3 complex have remained unclear, the human CX3 complex has been shown to promote RAD51 nucleofilament remodeling and stability, as well as strand invasion, whereas its other proposed role is to help recruit specialized factors to catalyze fork restoration through branch migration or controlled fork processing [56].

Reduced levels of XRCC3 protein have been shown to lead to increased rates of chromosome segregation errors, aneuploidy, and other chromosome aberrations [57]. Notably, cells deficient in XRCC3 have been shown to have lower HRR and hypersensitivity to cross-linking agents, genotoxic alkylating agents, UV radiation, and ionizing radiation [58, 59]. Similarly, common polymorphisms in this gene have been shown to lead to a derailed DNA damage response that may increase the likelihood of tumor development [60]. It is also known that XRCC3 polymorphisms play an important role in the treatment of gliomas. In current practice, temozolomide is the chemotherapeutic agent of choice for high-grade gliomas [61]. Recently, the XRCC3 polymorphism was found to contribute to temozolomide resistance in glioblastoma cells by mediating the repair of DNA double-strand breaks [62]. However, this may not be the whole story about the contribution of XRCC3 to temozolomide resistance. XRCC3 has also been found to contribute to mitochondrial biogenesis by promoting mitochondrial DNA integrity [63]. Mitochondria are known to enhance temozolomide chemoresistance, suggesting a role for XRCC3 in the treatment of gliomas [64, 65].

Given the important role of XRCC3 in HRR, its abundance and 3-D conformation require well-tuned regulation. As both protein structure and expression can be affected by genetic polymorphisms, a number of GWAS have investigated the association between XRCC3 polymorphisms and risk for cancers, including gliomas [33, 37, 39, 66–68]. Among the polymorphisms in XRCC3, we focused on p.Thr241Met, an exonic polymorphism involving a C-to-T transition at codon 241 in exon 7 [27]. This transition replaces threonine (Thr) with methionine (Met) and may affect XRCC3 functions, activities, and interactions [69]. We chose to perform a meta-analysis of this polymorphism because p.Thr241Met has been extensively studied with respect to its association with gliomas, with often conflicting results. For example, whereas Custódio et al. [34] found that the Met allele of p.Thr241Met was significantly associated with a 3-fold increase in glioma risk, Huang et al. [39] found that the same allele was associated with a 1.6-fold decrease in glioma risk. On the other hand, Rodriguez-Hernandez et al. [41] showed that there was no significant association between the polymorphism and glioma risk. These inconsistencies could be due to numerous factors, such as the ethnicity of the study population, and the sample size (and thus statistical power) of the studies [70, 71]. A well-conducted meta-analysis is essential to address these inconsistencies. However, the most recent meta-analyses on this topic were conducted many years ago [72–77] and did not include several eligible studies. The lack of meticulous data extraction in meta-analysis can lead to misleading results [78]. Therefore, in this work, we meticulously performed a systematic review and meta-analysis to investigate the association between the XRCC3 p.Thr241Met polymorphism and glioma risk.

Our results showed that the p.Thr241Met polymorphism was associated with increased glioma risk only in the homozygous and recessive models, but not in the heterozygous, dominant, and allele models. Within each model, we additionally clustered our analysis by different subgroups, i.e., ethnicity, source of controls, and HWE status. Our analysis in the Asian subgroup again showed that only the homozygous and recessive models were associated with increased glioma risk, whereas in the Caucasian subgroup, no significance was found in any of the genetic models. In addition, studies using hospital- and population-based controls appeared to result in different risk profiles, and studies in which the genotypic distribution of controls deviated from HWE showed more significant associations. It should be noted that the number of participants in each subgroup was much smaller than in the overall analysis. Therefore, there is a possibility that the subgroup analyses were not sufficiently powered to detect possible associations [79].

The greatest strength of our meta-analysis lies in the extensive collection of genetic association studies on the XRCC3 p.Thr241Met polymorphism performed on a large population. Moreover, the selected parameter studied, i.e., the p.Thr241Met polymorphism, is a well-established genetic variation that can be accurately genotyped using contemporary genotyping technologies. Therefore, the genotyping results obtained in different studies are reliable and well comparable with each other. Nevertheless, there are some limitations to this study as well. First, the influence of gene-gene and gene-environment interactions was not investigated because information on this aspect was lacking in the included studies. In addition, the study population, especially that of Asian populations, was mainly from China and therefore not very diverse. Furthermore, our meta-analysis examined only one polymorphism, whereas analysis of multiple genetic polymorphisms in the same gene might provide a more complete picture of the role of XRCC3 in mediating glioma risk.

Conclusions

In conclusion, our results show that there is a significant association between the XRCC3 p.Thr241Met polymorphism and increased glioma risk, but only when the homozygous and recessive models were adopted. When divided into subgroups by ethnicity, similar results were observed only in the Asian population. In addition, no obvious trend was observed when data were stratified by source of controls and HWE status, presumably because of low study power. Therefore, further genetic association studies are needed to provide a more accurate assessment of the association between the polymorphism and glioma risk.

Supporting information

S1 Table. Quality assessment of the included studies.
(DOCX)

S1 Fig. Sensitivity analysis of the meta-analysis.
(ZIP)

S1 File. Modified Newcastle-Ottawa scale for meta-analysis of genetic association studies.
(DOCX)

Acknowledgments

The authors thank Academic Proofreading (<https://www.academicproofreading.uk>) for English language editing.

Author Contributions

Conceptualization: Shing Cheng Tan.

Data curation: Shing Cheng Tan, Mohamad Ayub Khan Sharzehan, Hilary Sito, Hamed Kord-Varkaneh.

Formal analysis: Shing Cheng Tan.

Funding acquisition: Shing Cheng Tan.

Investigation: Shing Cheng Tan, Teck Yew Low, Hafiz Muhammad Jafar Hussain, Mohamad Ayub Khan Sharzehan, Hilary Sito.

Methodology: Shing Cheng Tan, Mohamad Ayub Khan Sharzehan, Hilary Sito, Hamed Kord-Varkaneh.

Software: Hamed Kord-Varkaneh.

Supervision: Hafiz Muhammad Jafar Hussain.

Validation: Teck Yew Low, Mohamad Ayub Khan Sharzehan, Hilary Sito.

Visualization: Shing Cheng Tan.

Writing – original draft: Teck Yew Low.

Writing – review & editing: Shing Cheng Tan, Md Asiful Islam.

References

1. Goodenberger ML, Jenkins RB. Genetics of adult glioma. *Cancer Genet.* 2012; 205: 613–621. <https://doi.org/10.1016/j.cancergen.2012.10.009> PMID: 23238284
2. Wen PY, Kesari S. Malignant gliomas in adults. *N Engl J Med.* 2008; 359: 492–507. <https://doi.org/10.1056/NEJMra0708126> PMID: 18669428
3. Ohgaki H, Kleihues P. Epidemiology and etiology of gliomas. *Acta Neuropathol.* 2005; 109: 93–108. <https://doi.org/10.1007/s00401-005-0991-y> PMID: 15685439
4. Jakola AS, Myrmet KS, Kloster R, Torp SH, Lindal S, Unsgård G, et al. Comparison of a strategy favoring early surgical resection vs a strategy favoring watchful waiting in low-grade gliomas. *JAMA.* 2012; 308: 1881–1888. <https://doi.org/10.1001/jama.2012.12807> PMID: 23099483
5. Stupp R, Hegi ME, Mason WP, van den Bent MJ, Taphoorn MJ, Janzer RC, et al. Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol.* 2009; 10: 459–466. [https://doi.org/10.1016/S1470-2045\(09\)70025-7](https://doi.org/10.1016/S1470-2045(09)70025-7) PMID: 19269895
6. Reuss D, von Deimling A. Hereditary tumor syndromes and gliomas. *Recent Results Cancer Res.* 2009; 171: 83–102. https://doi.org/10.1007/978-3-540-31206-2_5 PMID: 19322539
7. Moore SC, Rajaraman P, Dubrow R, Darefsky AS, Koebnick C, Hollenbeck A, et al. Height, body mass index, and physical activity in relation to glioma risk. *Cancer Res.* 2009; 69: 8349–8355. <https://doi.org/10.1158/0008-5472.CAN-09-1669> PMID: 19808953
8. Shete S, Hosking FJ, Robertson LB, Dobbins SE, Sanson M, Malmer B, et al. Genome-wide association study identifies five susceptibility loci for glioma. *Nat Genet.* 2009; 41: 899–904. <https://doi.org/10.1038/ng.407> PMID: 19578367
9. Wrensch M, Jenkins RB, Chang JS, Yeh RF, Xiao Y, Decker PA, et al. Variants in the CDKN2B and RTEL1 regions are associated with high-grade glioma susceptibility. *Nat Genet.* 2009; 41: 905–908. <https://doi.org/10.1038/ng.408> PMID: 19578366
10. Dziurzynski K, Chang SM, Heimberger AB, Kalejta RF, McGregor Dallas SR, Smit M, et al. Consensus on the role of human cytomegalovirus in glioblastoma. *Neuro Oncol.* 2012; 14: 246–255. <https://doi.org/10.1093/neuonc/nor227> PMID: 22319219
11. Strong MJ, Blanchard E 4th, Lin Z, Morris CA, Baddoo M, Taylor CM, et al. A comprehensive next generation sequencing-based virome assessment in brain tissue suggests no major virus—tumor association. *Acta Neuropathol Commun.* 2016; 4: 71. <https://doi.org/10.1186/s40478-016-0338-z> PMID: 27402152

12. Sadetzki S, Chetrit A, Freedman L, Stovall M, Modan B, Novikov I. Long-term follow-up for brain tumor development after childhood exposure to ionizing radiation for tinea capitis. *Radiat Res.* 2005; 163: 424–432. <https://doi.org/10.1667/rr3329> PMID: 15799699
13. Davis F, Il'yasova D, Rankin K, McCarthy B, Bigner DD. Medical diagnostic radiation exposures and risk of gliomas. *Radiat Res.* 2011; 175: 790–796. <https://doi.org/10.1667/RR2186.1> PMID: 21466382
14. Tan SC, Low TY, Mohamad Hanif EA, Sharzehan MAK, Kord-Varkaneh H, Islam MA. The rs9340799 polymorphism of the estrogen receptor alpha (ESR1) gene and its association with breast cancer susceptibility. *Sci Rep.* 2021; 11: 18619. <https://doi.org/10.1038/s41598-021-97935-8> PMID: 34545128
15. Najafi S, Tan SC, Raei P, Rahmati Y, Asemiani Y, Lee EHC, et al. Gene regulation by antisense transcription: A focus on neurological and cancer diseases. *Biomed Pharmacother.* 2022; 145: 112265. <https://doi.org/10.1016/j.biopha.2021.112265> PMID: 34749054
16. Wrensch M, Kelsey KT, Liu M, Miike R, Moghadassi M, Sison JD, et al. ERCC1 and ERCC2 polymorphisms and adult glioma. *Neuro Oncol.* 2005; 7: 495–507. <https://doi.org/10.1215/S1152851705000037> PMID: 16212814
17. Yang P, Kollmeyer TM, Buckner K, Bamlet W, Ballman KV, Jenkins RB. Polymorphisms in GLTSCR1 and ERCC2 are associated with the development of oligodendrogliomas. *Cancer.* 2005; 103: 2363–2372. <https://doi.org/10.1002/cncr.21028> PMID: 15834925
18. Kiuru A, Lindholm C, Heinavaara S, Ilus T, Jokinen P, Haapasalo H, et al. XRCC1 and XRCC3 variants and risk of glioma and meningioma. *J Neurooncol.* 2008; 88: 135–142. <https://doi.org/10.1007/s11060-008-9556-y> PMID: 18330515
19. Liu Y, Zhou K, Zhang H, Shugart YY, Chen L, Xu Z, et al. Polymorphisms of LIG4 and XRCC4 involved in the NHEJ pathway interact to modify risk of glioma. *Hum Mutat.* 2008; 29: 381–389. <https://doi.org/10.1002/humu.20645> PMID: 18165945
20. Liu Y, Zhang H, Zhou K, Chen L, Xu Z, Zhong Y, et al. Tagging SNPs in non-homologous end-joining pathway genes and risk of glioma. *Carcinogenesis.* 2007; 28: 1906–1913. <https://doi.org/10.1093/carcin/bgm073> PMID: 17389609
21. Wang LE, Bondy ML, Shen H, El-Zein R, Aldape K, Cao Y, et al. Polymorphisms of DNA repair genes and risk of glioma. *Cancer Res.* 2004/08/18. 2004; 64: 5560–5563. <https://doi.org/10.1158/0008-5472.CAN-03-2181> PMID: 15313891
22. Felini MJ, Olshan AF, Schroeder JC, North KE, Carozza SE, Kelsey KT, et al. DNA repair polymorphisms XRCC1 and MGMT and risk of adult gliomas. *Neuroepidemiology.* 2007; 29: 55–58. <https://doi.org/10.1159/000108919> PMID: 17898525
23. Bethke L, Webb E, Murray A, Schoemaker M, Johansen C, Christensen HC, et al. Comprehensive analysis of the role of DNA repair gene polymorphisms on risk of glioma. *Hum Mol Genet.* 2008; 17: 800–805. <https://doi.org/10.1093/hmg/ddm351> PMID: 18048407
24. Rajaraman P, Hutchinson A, Wichner S, Black PM, Fine HA, Loeffler JS, et al. DNA repair gene polymorphisms and risk of adult meningioma, glioma, and acoustic neuroma. *Neuro Oncol.* 2010; 12: 37–48. <https://doi.org/10.1093/neuonc/nop012> PMID: 20150366
25. Adel Fahmideh M, Schwartzbaum J, Frumentio P, Feychting M. Association between DNA repair gene polymorphisms and risk of glioma: a systematic review and meta-analysis. *Neuro Oncol.* 2014; 16: 807–814. <https://doi.org/10.1093/neuonc/nou003> PMID: 24500421
26. Vogelstein B, Kinzler KW. Cancer genes and the pathways they control. *Nat Med.* 2004; 10: 789–799. <https://doi.org/10.1038/nm1087> PMID: 15286780
27. Shen MR, Jones IM, Mohrenweiser H. Nonconservative Amino Acid Substitution Variants Exist at Polymorphic Frequency in DNA Repair Genes in Healthy Humans. 1998.
28. Yin QH, Liu C, Li L, Zu XY, Wang YJ. Association between the XRCC3 T241M polymorphism and head and neck cancer susceptibility: a meta-analysis of case-control studies. *Asian Pac J Cancer Prev.* 2012; 13: 5201–5205. <https://doi.org/10.7314/apjcp.2012.13.10.5201> PMID: 23244135
29. Zhao Y, Deng X, Wang Z, Wang Q, Liu Y. Genetic polymorphisms of DNA repair genes XRCC1 and XRCC3 and risk of colorectal cancer in Chinese population. *Asian Pac J Cancer Prev.* 2012; 13: 665–669. <https://doi.org/10.7314/apjcp.2012.13.2.665> PMID: 22524842
30. Cheng CX, Xue M, Li K, Li WS. Predictive value of XRCC1 and XRCC3 gene polymorphisms for risk of ovarian cancer death after chemotherapy. *Asian Pac J Cancer Prev.* 2012; 13: 2541–2545. <https://doi.org/10.7314/apjcp.2012.13.6.2541> PMID: 22938418
31. Zhu X, Zhong Z, Zhang X, Zhao X, Xu R, Ren W, et al. DNA repair gene XRCC3 T241M polymorphism and bladder cancer risk in a Chinese population. *Genet Test Mol Biomarkers.* 2012; 16: 640–643. <https://doi.org/10.1089/gtmb.2011.0334> PMID: 22299591
32. Tan SC. Low penetrance genetic polymorphisms as potential biomarkers for colorectal cancer predisposition. *J Gene Med.* 2018; 20: e3010. <https://doi.org/10.1002/jgm.3010> PMID: 29424105

33. Zhao P, Zou P, Zhao L, Yan W, Kang C, Jiang T, et al. Genetic polymorphisms of DNA double-strand break repair pathway genes and glioma susceptibility. *BMC Cancer*. 2013/05/15. 2013; 13: 234. <https://doi.org/10.1186/1471-2407-13-234> PMID: 23663450
34. Custódio AC, Almeida LO, Pinto GR, Santos MJ, Almeida JRW, Clara CA, et al. Variation in DNA repair gene XRCC3 affects susceptibility to astrocytomas and glioblastomas. *Genet Mol Res*. 2012; 11: 332–339. <https://doi.org/10.4238/2012.February.10.4> PMID: 22370935
35. Liu HB, Peng YP, Dou CW, Su XL, Gao NK, Tian FM, et al. Comprehensive study on associations between nine SNPs and glioma risk. *Asian Pac J Cancer Prev*. 2012/12/19. 2012; 13: 4905–4908. <https://doi.org/10.7314/apjcp.2012.13.10.4905> PMID: 23244079
36. Liu Y, Scheurer ME, El-Zein R, Cao Y, Do KA, Gilbert M, et al. Association and interactions between DNA repair gene polymorphisms and adult glioma. *Cancer Epidemiol Biomarkers Prev*. 2009/01/07. 2009; 18: 204–214. <https://doi.org/10.1158/1055-9965.EPI-08-0632> PMID: 19124499
37. Zhou K, Liu Y, Zhang H, Liu H, Fan W, Zhong Y, et al. XRCC3 haplotypes and risk of gliomas in a Chinese population: A hospital-based case-control study. *Int J Cancer*. 2009; 124: 2948–2953. <https://doi.org/10.1002/ijc.24307> PMID: 19330829
38. Yang P, Liu H, Lin J, Yang H. The Association of rs4753426 Polymorphism in the Melatonin Receptor 1B (MTNR1B) Gene and Susceptibility to Adolescent Idiopathic Scoliosis: A Systematic Review and Meta-analysis. *Pain Physician*. 2015/10/03. 2015; 18: 419–431. PMID: 26431121
39. Huang JY, Yang JF, Qu Q, Qu J, Liu F, Liu FE, et al. DNA repair gene XRCC3 variants are associated with susceptibility to glioma in a Chinese population. *Genet Mol Res*. 2015; 14: 10569–10575. <https://doi.org/10.4238/2015.September.8.18> PMID: 26400288
40. Gao K, Mu SQ, Wu ZX. Investigation of the effects of single-nucleotide polymorphisms in DNA repair genes on the risk of glioma. *Genet Mol Res*. 2014; 13: 1203–1211. <https://doi.org/10.4238/2014.February.27.5> PMID: 24634177
41. Rodríguez-Hernández I, Perdomo S, Santos-Briz A, García JL, Gómez-Moreta JA, Cruz JJ, et al. Analysis of DNA repair gene polymorphisms in glioblastoma. *Gene*. 2014; 536: 79–83. <https://doi.org/10.1016/j.gene.2013.11.077> PMID: 24325908
42. Xu G, Wang M, Xie W, Bai X. DNA repair gene XRCC3 Thr241Met polymorphism and susceptibility to glioma: A case-control study. *Oncol Lett*. 2014/07/12. 2014; 8: 864–868. <https://doi.org/10.3892/ol.2014.2192> PMID: 25013509
43. Luo KQ, Mu SQ, Wu ZX, Shi YN, Peng JC. Polymorphisms in DNA repair genes and risk of glioma and meningioma. *Asian Pac J Cancer Prev*. 2013/03/29. 2013; 14: 449–452. <https://doi.org/10.7314/apjcp.2013.14.1.449> PMID: 23534771
44. Pan WR, Li G, Guan JH. Polymorphisms in DNA repair genes and susceptibility to glioma in a Chinese population. *Int J Mol Sci*. 2013/02/07. 2013; 14: 3314–3324. <https://doi.org/10.3390/ijms14023314> PMID: 23385236
45. Tan SC, Ismail MP, Duski DR, Othman NH, Ankathil R. FAS c.-671A>G polymorphism and cervical cancer risk: a case-control study and meta-analysis. *Cancer Genet*. 2017; 211. <https://doi.org/10.1016/j.cancergen.2017.01.004> PMID: 28279307
46. Lee YH. Meta-analysis of genetic association studies. *Ann Lab Med*. 2015; 35: 283–287. <https://doi.org/10.3343/alm.2015.35.3.283> PMID: 25932435
47. Deeks JJ, Higgins JPT, Altman DG, on behalf of the Cochrane Statistical Methods Group. Analysing data and undertaking meta-analyses. *Cochrane Handbook for Systematic Reviews of Interventions*. John Wiley & Sons, Ltd; 2019. pp. 241–284. <https://doi.org/10.1002/9781119536604.ch10>
48. Hattersley AT, McCarthy MI. What makes a good genetic association study? *Lancet* (London, England). 2005; 366: 1315–1323. [https://doi.org/10.1016/S0140-6736\(05\)67531-9](https://doi.org/10.1016/S0140-6736(05)67531-9) PMID: 16214603
49. Salanti G, Sanderson S, Higgins JPT. Obstacles and opportunities in meta-analysis of genetic association studies. *Genet Med Off J Am Coll Med Genet*. 2005; 7: 13–20. <https://doi.org/10.1097/01.gim.0000151839.12032.1a> PMID: 15654223
50. Trikalinos TA, Salanti G, Khoury MJ, Ioannidis JPA. Impact of violations and deviations in Hardy-Weinberg equilibrium on postulated gene-disease associations. *Am J Epidemiol*. 2006; 163: 300–309. <https://doi.org/10.1093/aje/kwj046> PMID: 16410351
51. Tebbs RS, Zhao Y, Tucker JD, Scheerer JB, Siciliano MJ, Hwang M, et al. Correction of chromosomal instability and sensitivity to diverse mutagens by a cloned cDNA of the XRCC3 DNA repair gene. *Proc Natl Acad Sci U S A*. 1995; 92: 6354–6358. <https://doi.org/10.1073/pnas.92.14.6354> PMID: 7603995
52. Sullivan MR, Bernstein KA. RAD-ical new insights into RAD51 regulation. *Genes* (Basel). 2018;9. <https://doi.org/10.3390/genes9120629> PMID: 30551670

53. Chun J, Buechelmaier ES, Powell SN. Rad51 paralog complexes BCDX2 and CX3 act at different stages in the BRCA1-BRCA2-dependent homologous recombination pathway. *Mol Cell Biol.* 2013; 33: 387–395. <https://doi.org/10.1128/MCB.00465-12> PMID: 23149936
54. Masson JY, Stasiak AZ, Stasiak A, Benson FE, West SC. Complex formation by the human RAD51C and XRCC3 recombination repair proteins. *Proc Natl Acad Sci U S A.* 2001; 98: 8440–8446. <https://doi.org/10.1073/pnas.111005698> PMID: 11459987
55. Masson JY, Tarsounas MC, Stasiak AZ, Stasiak A, Shah R, McIlwraith MJ, et al. Identification and purification of two distinct complexes containing the five RAD51 paralogs. *Genes Dev.* 2001; 15: 3296–3307. <https://doi.org/10.1101/gad.947001> PMID: 11751635
56. Berti M, Teloni F, Mijic S, Ursich S, Fuchs J, Palumbieri MD, et al. Sequential role of RAD51 paralog complexes in replication fork remodeling and restart. *Nat Commun.* 2020; 11: 3531. <https://doi.org/10.1038/s41467-020-17324-z> PMID: 32669601
57. Thacker J. The RAD51 gene family, genetic instability and cancer. *Cancer Lett.* 2005; 219: 125–135. <https://doi.org/10.1016/j.canlet.2004.08.018> PMID: 15723711
58. Caldecott K, Jeggo P. Cross-sensitivity of gamma-ray-sensitive hamster mutants to cross-linking agents. *Mutat Res.* 1991; 255: 111–121. [https://doi.org/10.1016/0921-8777\(91\)90046-r](https://doi.org/10.1016/0921-8777(91)90046-r) PMID: 1922147
59. Pierce AJ, Johnson RD, Thompson LH, Jasin M. XRCC3 promotes homology-directed repair of DNA damage in mammalian cells. *Genes Dev.* 1999; 13: 2633–2638. <https://doi.org/10.1101/gad.13.20.2633> PMID: 10541549
60. Jiao L, Hassan MM, Bondy ML, Wolff RA, Evans DB, Abbruzzese JL, et al. XRCC2 and XRCC3 gene polymorphism and risk of pancreatic cancer. *Am J Gastroenterol.* 2008; 103: 360–367. <https://doi.org/10.1111/j.1572-0241.2007.01615.x> PMID: 17986315
61. Attarian F, Taghizadeh-Hesary F, Fanipakdel A, Javadinia SA, Porouhan P, PeyroShabany B, et al. A Systematic Review and Meta-Analysis on the Number of Adjuvant Temozolomide Cycles in Newly Diagnosed Glioblastoma. *Frontiers in oncology.* 2021. p. 779491. <https://doi.org/10.3389/fonc.2021.779491> PMID: 34900732
62. Roos WP, Frohnapfel L, Quiros S, Ringel F, Kaina B. XRCC3 contributes to temozolomide resistance of glioblastoma cells by promoting DNA double-strand break repair. *Cancer Lett.* 2018; 424: 119–126. <https://doi.org/10.1016/j.canlet.2018.03.025> PMID: 29574277
63. Mishra A, Saxena S, Kaushal A, Nagaraju G. RAD51C/XRCC3 Facilitates Mitochondrial DNA Replication and Maintains Integrity of the Mitochondrial Genome. *Mol Cell Biol.* 2018;38. <https://doi.org/10.1128/MCB.00489-17> PMID: 29158291
64. Taghizadeh-Hesary F, Akbari H, Bahadori M. Anti-mitochondrial Therapy: A Potential Therapeutic Approach in Oncology. *Preprints.* 2022. p. 2022010171. <https://doi.org/10.20944/preprints202201.0171.v2>
65. Oliva CR, Nozell SE, Diers A, McClugage SG 3rd, Sarkaria JN, Markert JM, et al. Acquisition of temozolomide chemoresistance in gliomas leads to remodeling of mitochondrial electron transport chain. *J Biol Chem.* 2010; 285: 39759–39767. <https://doi.org/10.1074/jbc.M110.147504> PMID: 20870728
66. Han S, Zhang H-T, Wang Z, Xie Y, Tang R, Mao Y, et al. DNA repair gene XRCC3 polymorphisms and cancer risk: a meta-analysis of 48 case-control studies. *Eur J Hum Genet* 2006 1410. 2006; 14: 1136–1144. <https://doi.org/10.1038/sj.ejhg.5201681> PMID: 16791138
67. Ali AM, Abdulkareem H, Anazi M Al, Parine NR, Shaik JP, Alamri A, et al. Polymorphisms in DNA Repair Gene XRCC3 and Susceptibility to Breast Cancer in Saudi Females. *Biomed Res Int.* 2016;2016. <https://doi.org/10.1155/2016/8721052> PMID: 26881229
68. Tan SC, Ankathil R. Genetic susceptibility to cervical cancer: role of common polymorphisms in apoptosis-related genes. *Tumor Biol.* 2015;36. <https://doi.org/10.1007/s13277-015-3868-2> PMID: 26242271
69. Williams K, Sobol RW. Mutation research/fundamental and molecular mechanisms of mutagenesis: special issue: DNA repair and genetic instability. *Mutat Res.* 2013; 743–744: 1–3. <https://doi.org/10.1016/j.mrfmmm.2013.04.009> PMID: 23688353
70. Tan SC, Lim PY, Fang J, Mokhtar MFM, Hanif EAM, Jamal R. Association between MIR499A rs3746444 polymorphism and breast cancer susceptibility: a meta-analysis. *Sci Rep.* 2020; 10: 1–10. <https://doi.org/10.1038/s41598-020-60442-3> PMID: 32103099
71. Huang T, Shu Y, Cai Y-D. Genetic differences among ethnic groups. *BMC Genomics.* 2015; 16: 1093. <https://doi.org/10.1186/s12864-015-2328-0> PMID: 26690364
72. Liang H-J, Yan Y-L, Liu Z-M, Chen X, Peng Q-L, Wang J, et al. Association of XRCC3 Thr241Met polymorphisms and gliomas risk: evidence from a meta-analysis. *Asian Pac J Cancer Prev.* 2013; 14: 4243–4247. <https://doi.org/10.7314/apjcp.2013.14.7.4243> PMID: 23991984
73. Zhao B, Ye J, Li B, Ma Q, Su G, Han R. DNA repair gene XRCC3 Thr241Met polymorphism and glioma risk: a meta-analysis. *Int J Clin Exp Med.* 2013; 6: 438–443. PMID: 23844267

74. Lin J, Kou Y. Association between the Thr241Met polymorphism of X-ray repair cross-complementing group 3 gene and glioma risk: evidence from a meta-analysis based on 4,136 cases and 5,233 controls. *Tumour Biol J Int Soc Oncodevelopmental Biol Med.* 2014; 35: 425–432. <https://doi.org/10.1007/s13277-013-1059-6> PMID: 23918307
75. Jiang J, Quan X-F, Zhang L, Wang Y-C. The XRCC3 Thr241Met polymorphism influences glioma risk—a meta-analysis. *Asian Pac J Cancer Prev.* 2013; 14: 3169–3173. <https://doi.org/10.7314/apjcp.2013.14.5.3169> PMID: 23803098
76. Feng Y, Zeng M, Xu Q. Association between XRCC3 T241M polymorphism and glioma risk: A meta-analysis. *Tumor Biol.* 2014; 35: 5589–5592. <https://doi.org/10.1007/s13277-014-1738-y> PMID: 24633885
77. Qi L, Yu HH-Q, Zhang Y, Ding L, Zhao D-HD, Lv P, et al. A Comprehensive Meta-analysis of Genetic Associations Between Key Polymorphic Loci in DNA Repair Genes and Glioma Risk. *Mol Neurobiol.* 2017; 54: 1314–1325. <https://doi.org/10.1007/s12035-016-9725-5> PMID: 26843108
78. Low TY, Chin K-Y, Tan SC. On the critical importance of meticulous data extraction for meta-analysis of genetic association study. *Arch Physiol Biochem.* 2021; 1–2. <https://doi.org/10.1080/13813455.2021.1890132> PMID: 33689514
79. Button KS, Ioannidis JPA, Mokrysz C, Nosek BA, Flint J, Robinson ESJ, et al. Power failure: why small sample size undermines the reliability of neuroscience. *Nat Rev Neurosci.* 2013; 14: 365–376. <https://doi.org/10.1038/nrn3475> PMID: 23571845