

# The formation and function of the neutrophil phagosome

Emily Naish<sup>1</sup> | Alexander JT Wood<sup>2,3</sup>  | Andrew P Stewart<sup>4</sup> | Matthew Routledge<sup>4,5</sup> | Andrew Conway Morris<sup>4,5</sup> | Edwin R Chilvers<sup>1</sup>  | Katharine M Lodge<sup>1</sup> 

<sup>1</sup>National Heart and Lung Institute, Imperial College London, London, UK

<sup>2</sup>Medical School, University of Western Australia, Perth, Australia

<sup>3</sup>Department of Critical Care, University of Melbourne, Melbourne, Australia

<sup>4</sup>Department of Medicine, University of Cambridge, Cambridge, UK

<sup>5</sup>Division of Immunology, Department of Pathology, University of Cambridge, Cambridge, UK

## Correspondence

Dr Katharine M Lodge and Professor Edwin R Chilvers, National Heart and Lung Institute, Imperial College London, Level 5 ICTEM Building, Du Cane Road, London, W12 0NN UK.

Email: [k.lodge@imperial.ac.uk](mailto:k.lodge@imperial.ac.uk) and [e.chilvers@imperial.ac.uk](mailto:e.chilvers@imperial.ac.uk)

## Funding information

Academy of Medical Sciences, Grant/Award Number: SGL024\1086; Avont Foundation, Grant/Award Number: 2021/000023; British Infection Association; Evelyn Trust; Medical Research Council, Grant/Award Number: MR/V006118/1; NIHR Imperial Biomedical Research Centre; Academy of Medical Sciences, Grant/Award Number: SGL023\1108

## Summary

Neutrophils are the most abundant circulating leukocyte and are crucial to the initial innate immune response to infection. One of their key pathogen-eliminating mechanisms is phagocytosis, the process of particle engulfment into a vacuole-like structure called the phagosome. The antimicrobial activity of the phagocytic process results from a collaboration of multiple systems and mechanisms within this organelle, where a complex interplay of ion fluxes, pH, reactive oxygen species, and antimicrobial proteins creates a dynamic antimicrobial environment. This complexity, combined with the difficulties of studying neutrophils *ex vivo*, has led to gaps in our knowledge of how the neutrophil phagosome optimizes pathogen killing. In particular, controversy has arisen regarding the relative contribution and integration of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase-derived antimicrobial agents and granule-delivered antimicrobial proteins. Clinical syndromes arising from dysfunction in these systems in humans allow useful insight into these mechanisms, but their redundancy and synergy add to the complexity. In this article, we review the current knowledge regarding the formation and function of the neutrophil phagosome, examine new insights into the phagosomal environment that have been permitted by technological advances in recent years, and discuss aspects of the phagocytic process that are still under debate.

## KEYWORDS

neutrophil, phagocytosis, phagosome

## 1 | INTRODUCTION

Neutrophils are among the first immune cells to respond following infection or injury; this makes the possession of proficient pathogen-killing mechanisms essential. Phagocytosis, the process of detecting and engulfing particles into an organelle called the phagosome, is key to the ability of neutrophils to kill pathogens. In neutrophils, phagocytosis is a highly specialized and efficient event. Thus, neutrophils are the archetypal “professional” phagocyte, although monocytes, macrophages, eosinophils, and dendritic cells also display phagocytic

ability to a somewhat lesser extent.<sup>1</sup> As well as killing ingested pathogens, innate immune phagocytes may present antigens to adaptive immune cells, highlighting the importance of phagocytosis for both arms of the immune system.<sup>2</sup> An intriguing neutrophil-dendritic cell hybrid phenotype has been identified in mice, exhibiting retained phagocytic and microbial-killing capacity as well as typical dendritic cell properties, such as antigen presentation.<sup>3</sup>

The neutrophil phagosome is a distinctive organelle, formed from an invagination of the plasma membrane to completely enclose an engulfed particle. A host of complementary processes

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. *Immunological Reviews* published by John Wiley & Sons Ltd.

and pathways then transforms the phagosome environment from a largely inert cellular inclusion into one optimized for the degradation of ingested particles.<sup>4</sup> Within the phagosome, two major cytotoxic events take place: the production of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase-derived reactive oxygen species (ROS) and the delivery of microbicidal proteins from pre-formed granules. The potential of these mechanisms to wreak havoc intracellularly and extracellularly necessitates exhaustive and complex regulatory systems, as excessive or aberrant neutrophil activation has been implicated in tissue damage in multiple inflammatory and autoimmune diseases.<sup>5</sup>

The complex mechanisms governing the phagocytic process in neutrophils have been a particular area of contention. Neutrophils are challenging to study, in part due to their short life span, abundance of degradative enzymes, and significant challenges in genetic manipulation.<sup>6</sup> As a result, there is a relative paucity of neutrophil-specific phagocytosis research. Many aspects of phagosome maturation have been studied in macrophages and then assumed to apply to neutrophils, despite disparities in phagocytic function (reviewed in [7]). One notable difference is that in most phagocytes (including macrophages), phagosome maturation follows an endocytic maturation pathway, whereby the phagosome fuses with lysosomes to form a phagolysosome,<sup>7</sup> whereas in the neutrophil, the process of maturation is more rapid due to the presence of pre-formed granules situated within the cytoplasm. This, combined with the neutrophil's streamlined killing mechanisms, leads to a very different phagosome environment. Despite these differences, study of the phagosome in other cell types has been informative to a degree as certain parallels can be drawn.

In this article, we will review the processes and mechanisms which shape the environment of the neutrophil phagosome and how these aid its pathogen-killing abilities, including the formation of ROS by NADPH oxidase, the movement of ions across the phagosomal membrane, the pH of the formed phagosome, and the delivery of granule contents. We will highlight controversies in the complex field of neutrophil phagocytosis, where further investigation is paramount to understand their role in health and disease, which is often a fine balance between harm and help (Figure 1).

## 2 | STUDYING THE PHAGOSOME THROUGHOUT HISTORY


Elie Metchnikoff (1845–1916), termed the father of cellular innate immunity, was the first to identify phagocytosis as a central host defense mechanism. He initially observed motile cells encircling an inserted thorn in starfish larvae, following which he realized the broader significance of phagocyte recruitment in host defense.<sup>8</sup> At around the same time, Paul Ehrlich identified complementary humoral adaptive immunity. Ehrlich's precise staining techniques led to the modern era of leukocyte biology; his neutral dyes were able

to identify "epsilon granules" in neutrophils, which he called "cells with polymorphous nuclei".<sup>8</sup> In 1960, it was shown that, following engulfment of bacteria, neutrophils underwent granule fusion (degranulation) with the phagocytic vacuole, and that the contents of the granules were "consumed," indicating their usage against pathogens.<sup>9</sup> A substance named "phagocytin" was the first microbicidal granule element to be identified,<sup>10</sup> which was later shown to comprise a number of different cationic antimicrobial proteins.<sup>11</sup> Subsequently, the advancement of staining and cell fractionation techniques, and cell imaging modalities including electron microscopy established the granule subset classification we know today, of which azurophil, specific, and gelatinase are the predominant subtypes.

Research into neutrophil phagocytosis has been full of contradictions and controversies. Shortcomings have been in part due to the inherent limitations of all *in vitro* and *ex vivo* studies. As neutrophils do not proliferate *in vitro*, much research has been undertaken using freshly isolated human donor cells, meaning donor-dependent variation is unavoidable. The neutrophil-like cells differentiated from HL-60 and PLB-985 cell lines have been used as an alternative; however, these cells are imperfect models for phagocytosis as intracellular killing is much less efficient, potentially due to the lack of specific granules in HL-60 cells.<sup>12</sup> The recent use of swimming zebrafish larvae, which are small, transparent, and only utilize innate immunity, in combination with high-resolution live imaging has been revolutionary in following neutrophil responses to microorganisms *in vivo* and in real time.<sup>14</sup> Additionally, *intra-vital* two-photon microscopy has enabled real-time live imaging of neutrophils phagocytosing bacteria or viral prey in murine lymph nodes, with important implications for antigen presentation.<sup>13,14</sup>

Other difficulties in phagocytosis research have arisen due to the transience of the phagosome, and an array of confounding factors present within this organelle; these include phagosomal membrane potential, osmotic strength, pH of the phagosome and cytosol, and the complex interplay of many different ion channels, enzymes, and chemical reactions.<sup>15</sup> However, recent technological advances have increased our knowledge of neutrophil phagocytosis: In particular, the use of automated fluorescence microscopy to observe phagosome formation and maturation<sup>16,17</sup> and gene-targeting technology in murine neutrophils<sup>18</sup> have permitted more focused experiments.

Despite information from these new experimental techniques, several controversies remain, including the role of ROS in establishing the *intra-phagosomal* environment. The increased oxygen consumption seen during phagocytosis in neutrophils was originally thought to be due to mitochondrial respiration, until the production of ROS within the phagosome was discovered. However, the extent to which ROS are directly antimicrobial is still actively debated. Another area of continued uncertainty is the *intra-phagosomal* pH; due to the difficulties of measuring pH in such a small volume, purported phagosomal pH values have varied hugely over the years, ranging from acidification to neutral to alkalinization.



**FIGURE 1** Phagosome formation and maturation. (A) Overview of the major steps in neutrophil phagosome formation following pathogen detection. Events 1–4 are depicted in further detail in panel B. (B) 1: An opsonized pathogen engages Fc receptors (Fc $\gamma$ R) or complement receptors (e.g., CR3) to initiate phagocytosis. Both Fc $\gamma$ R and CR3 can employ immunoreceptor signaling pathways: SH2-domain-bearing proteins (e.g., Syk) associate with phosphorylated ITAM, signaling downstream through phosphatidylinositol 3-kinase (PI3K) and/or phospholipase C $\gamma$  (PLC $\gamma$ ). CR3 also employs independent inside-out and outside-in integrin signaling pathways. 2: Phagocytic receptor signaling induces regulation of the actin cytoskeleton via Rac and/or Rho. Myosin motor control of actin rearrangement drives extending pseudopod protrusions from the plasma membrane to form the phagocytic cup around the pathogen. 3: Cytosolic specific/gelatinase granules deliver proteins to the membrane of the forming phagosome, for example, the membrane-bound subunits of NADPH oxidase, gp91<sup>phox</sup> (NOX2), and p22<sup>phox</sup>. Actin polymerization at the pseudopod tips facilitates membrane sealing to complete the phagocytic vacuole around the pathogen. 4: The formed pathogen-containing phagosome translocates toward the granule-rich centriole within the neutrophil cytosol. NADPH oxidase generates antimicrobial reactive oxygen species inside the phagosome. The negative charge generated by this process is compensated by an influx of protons. Cytosolic azurophil granules, containing cytotoxic proteins, for example, elastase, fuse with the phagosome membrane to deliver their contents to the lumen of the phagosome

### 3 | PHAGOCYTOSIS SIGNALING

During phagocytosis, multiple signaling cascades are activated, which result in the rearrangement of the actin cytoskeleton into a

nascent phagosome (i.e., the stage of the phagosome immediately following initial sealing), after which phagosome translocation toward the center of the cell occurs, alongside phagosomal maturation and pathogen killing.<sup>19–21</sup>

The magnitude of the neutrophil phagocytic response to pathogens is substantial, which is unsurprising given that a primary function of neutrophils is pathogen destruction. Recent work from our group demonstrated that although there were minimal changes in protein expression between neutrophils exposed to *Staphylococcus aureus* bioparticles for 15 minutes compared to untreated cells, approximately one third of the phospho-proteome was altered.<sup>22</sup> These datasets are publicly available in the PRIDE database (<https://www.ebi.ac.uk/pride/> reference PXD017092). This dramatic change in protein phosphorylation indicates that *S. aureus* encounter and phagocytosis result in a significant signaling stimulus for the neutrophil. Reactome database<sup>23</sup> pathways enriched in our dataset of phagocytosing neutrophils include Rho GTPase signaling, neutrophil degranulation, membrane trafficking and nuclear membrane breakdown, vesicle-mediated transport and phosphatidylinositol signaling, which are discussed in detail below. The use of unbiased techniques to understand dynamic signaling in neutrophils has been hampered by the release of degradative enzymes during cell lysis, though our group<sup>22</sup> and others<sup>24–26</sup> have demonstrated these techniques are becoming feasible with modern technologies. It is anticipated that further insights into complex signaling networks will be gained through the ongoing application of these methods.

Phagocytosis is initiated by the engagement of various receptors on the surface of neutrophils, often by endogenous opsonic ligands such as immunoglobulins or iC3b, a product of the complement system. Opsonins coating the pathogen are recognized by the neutrophil, stimulating phagocytosis, and they play an additional role by helping to overcome the repellent forces between the neutrophil and the negatively charged cell wall of many bacteria.<sup>27</sup> Microbial pathogen-associated molecular patterns (PAMPs) also bind to a variety of neutrophil receptors, such as dectin-1.<sup>28,29</sup> Thus, neutrophils can internalize both opsonized and non-opsonized particles. Here, we focus on opsonin-mediated phagocytosis signaling pathways activated by Fc gamma receptors (FcγR) and complement receptors, which exhibit distinct mechanisms.

FcγRs bind immunoglobulin G (IgG)-coated targets whereas complement receptors bind activated complement-coated targets. Following receptor activation, pseudopods form from the plasma membrane to produce a cup-shaped enclosure of the target particle that is enabled by the rearrangement of the actin cytoskeleton.<sup>30</sup> Actin polymerizes at the leading edge and around the phagosomal cup, and this polymerization persists until the constriction and closure of the phagosome<sup>31</sup>. As the phagosome matures, the actin network disassembles to complete closure of the phagosome cup and allow the fusion of granules.<sup>32,33</sup>

Originally, pseudopod extension to surround a micro-organism prior to engulfment was thought to be exclusive to FcγR-mediated phagocytosis as macrophage studies described “sinking” of complement-opsonized targets.<sup>34,35</sup> More recent live-cell imaging of phagocytosis in macrophages has contested this, with evidence of membrane ruffles and protrusions encapsulating targets in complement receptor 3 (CR3)-mediated phagocytosis.<sup>36–38</sup> However, slower “sinking” phagocytosis also occurs as an alternative mechanism

requiring less involvement of membrane extensions. Moreover, in macrophages, the sinking phenomenon was shown to be CR3-dependent but FcγR-independent, whereas both CR3 and FcγR appear necessary for the formation and closure of FcγR-mediated phagocytic cups.<sup>39</sup> This is likely to be the case in neutrophils also, as neutrophils with CR3 mutations displayed decreased ability to ingest IgG-opsonized targets.<sup>40</sup>

### 3.1 | FcγR receptors

Neutrophils express FcγRI (CD64), FcγRIIA (CD32), and FcγRIIIB (CD16),<sup>41</sup> where the predominant Fc receptor subtypes are FcγRIIA and FcγRIIIB.<sup>42</sup> The class I and III receptors form multimeric complexes, while class II receptors exist as monomers with a unique phosphorylation motif and inhibitory action when engaged.<sup>43</sup> As circulating unbound IgGs are ubiquitous, phagocytes must be able to distinguish between these and IgG-associated with particles or immune complexes. Immunoglobulin receptors are therefore activated by clustering, mediated by the simultaneous engagement of multiple ligands, as opposed to ligand-induced conformational change.<sup>44,45</sup> Clustering induces the activation and recruitment of Src family kinases, which results in the phosphorylation of tyrosine residues in the immunoreceptor tyrosine activation motif (ITAM) within the FcγR signaling subunit.

Phosphorylation of ITAM generates docking sites for proteins bearing SH2 domains, including the tyrosine kinase, Syk. Subsequent phosphorylation of Syk leads to the recruitment of various signaling proteins to the activated FcγR complex. Highlighting the importance of Syk in FcγR-mediated phagocytosis, Jaumouillé et al found that in macrophages, Syk regulated FcγR responsiveness by increasing lateral receptor mobility and clustering through a reduction in actin polymerization.<sup>46</sup> Although this mechanism was not demonstrated directly, neutrophils from Syk-deficient mice displayed a similar reduced ability to ingest IgG-opsonized particles.<sup>47,48</sup>

Phosphorylation of Syk leads to the recruitment of adaptor proteins to the activated FcγR complex, leading to the activation of lipid-modification enzymes, including phosphatidylinositol 3-kinase (PI3K) and phospholipase Cγ (PLCγ). PI3K is responsible for the accumulation of phosphatidylinositol 3,4,5-trisphosphate (PtdIns(3,4,5)P<sub>3</sub>), derived from phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P<sub>2</sub>). PtdIns(4,5)P<sub>2</sub> is also the substrate of PLCγ, and hydrolyzes into inositol1,4,5-trisphosphate (Ins(1,4,5)P<sub>3</sub>) and diacylglycerol (DAG), which also act as messengers in the phagocytic signaling cascade.<sup>29</sup>

Despite the actions of PI3K and PLCγ, levels of PtdIns(4,5)P<sub>2</sub> increase in the early stages of macrophage phagocytosis, accumulating at the site of particle engagement and at the pseudopod tips. Levels abruptly decrease upon internalization, as actin disassembles to allow phagosome detachment, suggesting that accumulation of PtdIns(4,5)P<sub>2</sub> is associated with initial actin recruitment, and its hydrolysis is associated with subsequent actin degradation and remodeling.<sup>32,33</sup> This is evidenced by the reduced actin disassembly during phagocytosis of IgG-opsonized latex beads when PtdIns(4,5)P<sub>2</sub>

hydrolysis was inhibited in macrophages.<sup>33</sup> The role of PtdIns(4,5)P<sub>2</sub> in neutrophil phagosomes is less certain. Similar phosphoinositide dynamics in the phagosomal cup have been demonstrated between macrophages and the neutrophil-like HL-60 cell line, although this study did not differentiate between PtdIns(4,5)P<sub>2</sub> and PtdIns(3,4,5)P<sub>3</sub> accumulation.<sup>49</sup> Conversely, a study of human and murine neutrophils showed that although PtdIns(4,5)P<sub>2</sub> was abundant in the plasma membrane, unlike macrophages it decreased rapidly during phagosome formation and was undetectable after sealing.<sup>21</sup>

The synthesis of PtdIns(3)P on phagosomal membranes appears to be universal across phagocytosis,<sup>50</sup> where similar dynamics have been demonstrated in neutrophils and macrophages. The generation of PtdIns(3)P by Class III PI3K in the phagosome membrane aids in the assembly of NADPH oxidase, making it essential for phagosome maturation. PtdIns(3)P binds the PX domain of p40<sup>phox</sup> (a component of NADPH oxidase) with high affinity and selectivity<sup>51</sup> and is critical for intracellular ROS production: Neutrophils from a patient carrying a mutation in the p40<sup>phox</sup> PX domain (compromising PtdIns(3)P binding) had markedly decreased production of intracellular ROS during phagocytosis with subsequent impaired bacterial killing capacity.<sup>52</sup>

### 3.2 | Complement receptors

Complement receptors are categorized into CR1 and CR2, formed by short consensus repeat (SCR) elements; CR3 and CR4, which belong to the  $\beta$ 2 integrin family; and CR1g, which belongs to the immunoglobulin Ig-superfamily.<sup>53</sup> CR3 (also called CD11b/CD18 or Mac-1) is the most efficient phagocytic complement receptor.<sup>54,55</sup> CR3 receptors are  $\alpha$ M $\beta$ 2 integrins that are activated by outside-in (binding to extracellular ligands) and inside-out (intracellular protein binding that changes integrin conformation and thus affinity state) signaling (reviewed in<sup>56,57</sup>). With regard to phagocytosis, the complement fragment C3bi, a potent serum opsonin, is the most important ligand; however, CR3 is promiscuous and can bind a range of other ligands, including extracellular matrix proteins, surface receptors, blood coagulation proteins, and microbial surface molecules.

Patients with leukocyte adhesion deficiency (LAD), a condition caused by genetic mutations in  $\beta$ 2 integrins (and thus CR3), experience severe recurrent bacterial infections.<sup>58</sup> Neutrophils from patients with LAD type 1 are CR3-deficient and defective in phagocytosis when stimulated *in vitro* but appear to have normal unstimulated IgG-dependent phagocytosis.<sup>40</sup> This suggests that there are CR3-dependent and CR3-independent mechanisms of phagocytosis and that the recurrent infections in LAD patients may be associated with failure to increase phagocytosis in response to inflammatory stimuli. However, neutrophils from these patients also display significant migration defects so it is difficult to tease out the relative contribution of this impairment, compared with phagocytosis defects, to infection susceptibility *in vivo*.

Inside-out activation appears critical for CR3-mediated phagocytosis. Inside-out activation involves the transduction of an intracellular signal to the cytoplasmic and then the extracellular domain of

an integrin, with signaling often initiated by G protein coupled receptor (GPCR) or Toll-like receptor (TLR) activation, or cytokine stimulation.<sup>53</sup> Once inside-out signaling has facilitated conformational changes, from low to high affinity, outside-in signaling occurs.<sup>38</sup> This always involves linkage of integrins to the actin cytoskeleton but, depending on the effector response required, recruits and activates different adaptor proteins (reviewed in<sup>59</sup>).

As well as the distinct signaling pathways described above, CR3 also employs an immunoreceptor-like signaling mechanism through phosphorylation of ITAMs on receptor-associated transmembrane adaptors: Neutrophils lacking ITAM-containing adaptor proteins are defective in integrin-mediated phagocytosis.<sup>60</sup> Similar to Fc $\gamma$ R signaling, protein adaptors provide docking sites for the SH2 domains of Syk kinases which are then activated, followed by downstream signaling events. This study also found that Src family kinase phosphorylation of the adaptor proteins was essential for the association of the SH2 Syk domains. These signaling events are important for rapid target ingestion: In macrophages, deletion of Syk or ITAM adaptors accentuated the inefficient and slow sinking method of phagocytosis.<sup>39</sup> An alternative mechanism, where inefficient phagocytic cups were formed via the extension of membrane ruffles, was also inhibited by the deletion of Syk or ITAM adaptors, suggesting that Syk signaling induces ruffling, a phenomenon which can also be observed during Fc $\gamma$ R-mediated phagocytosis.<sup>39</sup>

The majority of early work on integrin signaling was performed in macrophages, initially describing Fc $\gamma$ R- or CR3-mediated phagocytosis as two discrete mechanisms.<sup>61</sup> More recently, it has been suggested that CR3 and its downstream effectors are essential for phagocytosis of both opsonized and non-opsonized targets.<sup>62</sup>

### 3.3 | GTPase and cytoskeleton dynamics

Following protein and lipid kinase activity, signaling cascades induce actin polymerization and localized membrane remodeling. Rho family small GTPases play a central role in actin dynamic regulation and can switch between an active (GTP-bound) and inactive (GDP-bound) state. Fc $\gamma$ R-mediated phagocytosis is thought to involve predominantly Rac1, Rac2, and Cdc42, whereas complement-mediated phagocytosis utilizes Rho. Rac and Cdc42 direct lamellipodial and filopodial membrane protrusions, respectively, whereas Rho induces the assembly of contractile actomyosin filaments.<sup>63,64</sup> However, in macrophages, RhoG involvement has also been identified in Fc $\gamma$ R-mediated phagocytosis and iC3b-opsonized particle uptake is greatly reduced in Rac1- and Rac2-deficient cells,<sup>36</sup> suggesting some crossover of these pathways. It is likely that this is also true for neutrophils but has yet to be established.

In neutrophils, Fc $\gamma$ R-mediated phagocytosis was markedly impaired in Rac2- but not Rac1-deficient cells,<sup>65</sup> indicating a non-redundant role for the Rac2 isoform. A rare inhibitory Rac2 mutation has been described in a patient who presented with recurrent and severe bacterial infection; however, phagocytic capacity was not assessed directly and Rac2 is also implicated in granule

translocation (section 7.4) and NADPH oxidase function.<sup>66</sup> Due to its polybasic domain, Rac can interact with anionic phospholipids. Thus, as well as actin modulation, Rac has an important role as a targeting signal to localize and correctly position proteins at the phagosomal membrane. Faure et al. showed that the transient recruitment of Rac2 and the NADPH oxidase component, p47<sup>phox</sup>, to the phagosomal cup membrane of neutrophil-like PLB-985 cells at the beginning of phagocytosis was phosphatidylserine-dependent, and that inhibition of this recruitment resulted in reduced ROS production.<sup>67</sup> In murine neutrophils, assembly and activation of NADPH oxidase following FcγR-mediated phagocytosis was completely dependent on Rac2 whereas following complement-mediated phagocytosis, there was redundancy between Rac1 and Rac2. Further differences were identified, where the oxidase rapidly (less than 6 seconds) accumulated on sealed phagosomes formed in response to complement-opsonized prey, but during slower (more than 10 seconds) phagocytosis of IgG-opsonized targets the oxidase could assemble at the base of a forming phagosome.<sup>68</sup>

In FcγR-mediated phagocytosis, fluorescence resonance energy transfer (FRET) stoichiometry of macrophages revealed that Cdc42 was activated early and localized to actin in the extending pseudopod, Rac1 was active across the phagocytic cup and during closure, and Rac2 was active predominantly during contractile activities and closure of the phagosome.<sup>31</sup> Similarly, in neutrophils, Rac1 was preferentially recruited to actin-rich pseudopods due to the negative charge generated by phospholipids whereas Rac2 localized to the intermediately-charged phagosomal membrane.<sup>69</sup> However, neutrophil and macrophage GTPase dynamics are not identical as genetic deletion or pharmacological inhibition of murine or human neutrophil Cdc42 does not impair phagocytosis, although bacterial killing was compromised.<sup>70</sup> These spatio-temporal phagocytic control mechanisms for the different GTPases suggest that differential regulation of these enzymes coordinates the localized membrane remodeling.

Intriguingly, the use of pharmacological depolymerization agents showed that an intact actin cytoskeleton is required to target uptake and Rac2 translocation to the site of particle attachment in neutrophil-like PLB-985 cells, but disruption of the actin cytoskeleton once phagocytosis had been initiated does not prevent translocation of Rac2 toward the phagosome.<sup>71</sup> Indeed, actin depolymerization following FcγR-crosslinking was shown to enhance the oxidative burst,<sup>65</sup> indicating that the timing of actin assembly and disassembly is important.

The formation of membrane protrusions and closure of the phagocytic cup are facilitated by myosin “motors,” which can control actin assembly, crosslinking and rearrangement, and are thus important regulators of membrane deformation, protein localization and phagosome translocation during phagocytosis (reviewed in detail in<sup>72</sup>). Myosin contractility is thought to be particularly important in FcγR-mediated phagocytosis in neutrophils, where inhibitors of myosin ATPase prevented particle uptake,<sup>73</sup> and mechanical models suggest there is a requirement for protrusive force.<sup>30</sup>

Rho and Rac activation are thought to be downstream events that follow the phosphorylation of the guanine exchange factor (GEF) Vav by Syk.<sup>36,74</sup> Neutrophils from Vav knockout mice are significantly deficient in FcγR- and integrin-mediated phagocytosis.<sup>65,75</sup> Downstream of Rho and Rac, the Arp2/3 complex is responsible for actin polymerization. In murine neutrophils, Rac1 and Rac2 isoforms were shown to differentially regulate actin assembly using different pathways, where Rac2 mediated the majority of its effect on actin via the Arp2/3 complex.<sup>76</sup>

During CR3-mediated phagocytosis, actin polymerization is propagated by two distinct mechanisms via RhoA. In one pathway, RhoA can activate Rho kinase, which phosphorylates and activates myosin II, leading to recruitment of Arp2/3; inhibition of Rho kinase or myosin II activity results in reduced Arp2/3 recruitment and actin cup assembly in murine macrophages.<sup>77</sup> A non-redundant role for Arp2/3 in neutrophils is less certain, however, as neutrophils isolated from a patient with a rarely described *ARPC1B* deficiency (lacking Arp2/3) did not demonstrate any defect in phagocytosis.<sup>78</sup> In a separate pathway, RhoA can recruit the actin nucleator mDia1, which is recruited to the phagocytic cup via the microtubule-associated protein CLIP-170.<sup>79</sup> Inhibition of mDia1 reduces actin polymerization and particle uptake.<sup>80</sup> Highlighting the importance of RhoA signaling, the complement product, C5a (an anaphylatoxin implicated in sepsis pathogenesis), inhibits RhoA activation in a PI3Kδ-dependent manner, which prevents actin polymerization and reduces neutrophil phagocytosis.<sup>81</sup> Importantly, neutrophils from critically ill patients exhibit a similar phenotype, that is, reduced phagocytosis and a failure to activate RhoA or polymerise actin, which may contribute to the increased risk of infection in this patient group.

Various actin-binding proteins have been shown to be important during neutrophil phagocytosis. The mammalian actin-binding protein 1 (mAbp1), an adaptor protein phosphorylated by Syk, has been implicated in complement-mediated phagocytosis.<sup>82</sup> Syk was found to be necessary for the translocation of mAbp1 to the site of engulfment during phagocytosis and down-regulation of mAbp1 led to a depletion of clustered β2 integrins in high-affinity conformation. This high-affinity conformation is essential to generate the tensile strength required for phagocytosis and the absence of mAbp1 led to severe defects in β2 integrin-mediated phagocytosis.<sup>82</sup> Inhibition of the actin-binding protein, coronin-1, has also been found to arrest phagocytosis. Coronin-1 is recruited to the phagosomal cup early, alongside actin. This suggestion that it may regulate actin is supported by the presence of coronin-1 at the leading edge of migrating cells.<sup>83</sup> Coronin-1 co-localizes with filamins, the most potent actin crosslinkers, at the phagocytic cup, where it appears to be involved in non-opsonic phagocytosis in response to PAMPs.<sup>84</sup> In macrophage phagosomes, phosphorylated Syk and paxillin colocalize and recruit vinculin, which facilitates target internalization by anchoring F-actin. These exact phosphorylation sites have not been detected in neutrophils, but it is likely that a similar process occurs.<sup>38</sup>

Overall, distinct signaling mechanisms characterize FcγR and CR3-initiated phagocytosis, which results in different, though overlapping, processes for particle uptake and phagosome formation. In

vitro, it is often necessary to dissect the roles of various signaling molecules individually. These experiments have highlighted important redundant and non-redundant roles for selected proteins, which have also been identified clinically through mutations, albeit rarely. In practice, however, there is likely substantial crossover and synergy between the immunoglobulin and complement pathways, which optimizes phagocytosis *in vivo*.

## 4 | FORMATION OF THE PHAGOSOME

### 4.1 | NADPH oxidase

The formation of NADPH oxidase gives rise to some of the key changes in the intra-phagosomal environment, starting with ROS production. In most cell types, ROS are produced as a by-product from a variety of processes, such as mitochondrial respiration,<sup>85</sup> but in neutrophils, the majority of ROS are actively generated by NADPH oxidase (NOX enzyme complexes). NOX enzymes are “professional” generators of ROS, as reducing molecular oxygen to form ROS is their sole enzymatic function. In leukocytes, NOX2 is the main catalytic subunit, which is highly expressed in neutrophils.<sup>86</sup> The enzyme NOX2 is dormant in the circulating quiescent neutrophil: Activation requires an initial priming step followed by full assembly of the NADPH oxidase complex from membrane-bound and regulatory cytosolic subunits.<sup>86</sup>

At rest, the two membrane-bound subunits, gp91<sup>phox</sup> (which is the main redox center, NOX2) and p22<sup>phox</sup>, are embedded in specific granules. Upon phagocytosis, and before the phagocytic cup is even sealed, these subunits are delivered to the phagosomal membrane by granule exocytosis.<sup>87</sup> Translocation of additional regulatory cytosolic subunits to the membrane-bound gp91<sup>phox</sup>/p22<sup>phox</sup> heterodimer to form the active NADPH oxidase complex is tightly controlled via the activation of a series of kinases, including protein kinase A (PKA), phosphoinositide 3-kinase (PI3K), and mitogen-activated protein kinases (MAPKs).<sup>50,88</sup> The small GTPase, Rac, is also important for NADPH oxidase activation. Guanine nucleotide exchange factors (GEFs) convert Rac-GDP to the active Rac-GTP, which translocates to the phagosomal membrane and there binds gp91<sup>phox</sup> and p67<sup>phox</sup>.<sup>87-89</sup> The obligate binding partner, p22<sup>phox</sup>, ensures stability of the heterodimer and accommodates docking of the regulatory subunits.<sup>89</sup>

NADPH oxidase activation is a dynamic process whereby a cascade of kinase signaling, calcium release, and small GTPase (predominantly Rac2) activation lead to subsequent NOX2 activation.<sup>90,91</sup> Activation of voltage-gated proton channels, chloride channels, and subsequent calcium fluxes is also utilized to regulate the enzyme complex.<sup>87</sup> The activation process is differentially regulated by various receptors, depending on the stimulus.<sup>92</sup> Robust activation of NOX2 is seen following FcγR and integrin receptor ligation. Other receptors, such as formyl peptide receptors, which detect bacterial cell wall products, or GPCRs, can activate the enzyme complex directly but to a lesser extent. Neutrophil priming is an important

control mechanism, which prevents inappropriate triggering of ROS generation. Here, an initial priming stimulus, for example, ligand binding to TLRs or TNF exposure, generates a pre-activated neutrophil phenotype, which results in a substantially enhanced activation of the NADPH oxidase when the cell encounters a second stimulus.<sup>93</sup> The range of upstream signaling pathways indicates that the threshold for NADPH oxidase activation is high, and that the ROS response is heterogenous, dictated by which signal is received. This complexity ensures tight regulation, with the aim of avoiding prolonged inflammation and oxidative damage.

### 4.2 | The production of ROS in the phagosome

ROS are chemically reactive radical and non-radical derivatives of oxygen, the former containing at least one unpaired electron. NOX2 (gp91<sup>phox</sup>) is the main redox center, transferring two electrons provided by cytosolic NADPH via internal heme moieties to the phagosome, whereupon oxygen is reduced to superoxide (O<sub>2</sub><sup>·-</sup>). ROS react with many biomolecules, including DNA, proteins, lipids, and carbohydrates, to cause damage. The study of ROS has historically been focused on their capacity to cause cellular toxicity. In the 1930s, the rapid release of ROS was found to be associated with the formation of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>),<sup>94</sup> which can generate potentially cytotoxic derivatives, such as the highly reactive hydroxyl radical. These findings were reinforced by the early discovery of bacterial ROS-eliminating enzymes, such as superoxide dismutase.<sup>95</sup>

### 4.3 | Reactive oxygen intermediates

The long-held consensus has been that the ROS produced by NOX2, derived from superoxide anions, worked directly to kill bacteria,<sup>88,96,97</sup> although this has now been contested (section 8). The initial question was which product was key to the microbial killing? The direct product of NOX2, superoxide anion (O<sub>2</sub><sup>·-</sup>), is thought not to be highly bactericidal as it is weakly reactive and unable to migrate far from the site of production. However, a few of its derivatives are more reactive and able to diffuse through pathogen membranes.<sup>88,98</sup>

O<sub>2</sub><sup>·-</sup> readily forms several reactive products, including hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). This either happens spontaneously, or through superoxide dismutase catalysis from two O<sub>2</sub><sup>·-</sup>.<sup>88</sup> H<sub>2</sub>O<sub>2</sub> is converted to other forms of ROS by the myeloperoxidase-hydrogen peroxide-halide system: Myeloperoxidase (MPO) catalyzes the oxidation of different substrates by H<sub>2</sub>O<sub>2</sub>, with the most common substrates being halides. The products are hypohalous acids, including hypochlorous acid (HOCl). H<sub>2</sub>O<sub>2</sub> also interacts with transition metals, such as iron, to produce hydroxyl radicals (OH·). However, iron is bound to lactoferrin within the phagosome and so is unlikely to contribute to bacterial killing.<sup>98</sup> In addition to catalyzing H<sub>2</sub>O<sub>2</sub> oxidation, MPO has the potential to act as a superoxide dismutase, contributing to H<sub>2</sub>O<sub>2</sub> production. Furthermore, superoxide anion can react with redox intermediates of MPO that could impact its chlorination

activity.<sup>99</sup> Whichever reaction kinetics are favorable within the phagosome likely influence the degree of MPO-dependent bacterial killing.

H<sub>2</sub>O<sub>2</sub> can also interact with O<sub>2</sub><sup>•−</sup> or HOCl to generate singlet oxygen (<sup>1</sup>O<sub>2</sub>), an electronically excited state of molecular oxygen that is highly reactive but short-lived.<sup>100</sup> Some studies have suggested its presence in neutrophil phagosomes,<sup>101,102</sup> and there is some evidence acquired using <sup>1</sup>O<sub>2</sub> and antibodies to support the formation of ozone in neutrophils.<sup>103–105</sup> However, the abundance and microbicidal effects of <sup>1</sup>O<sub>2</sub> remain ambiguous due to the imperfect methods of detection.<sup>106</sup>

An alternative reaction occurs with nitric oxide and superoxide anions, which generates various downstream reactive nitrogen species (RNS). Evidence of RNS-mediated microbial killing in neutrophils is variable, with some studies showing little effect of RNS deficiency on neutrophil phagocytosis or killing ability,<sup>107</sup> while others demonstrated increased infection susceptibility.<sup>108,109</sup> Discrepancies are likely explained by the varying susceptibility of different pathogens to RNS, which is thought to be a more important mechanism in tuberculosis and *Salmonella* species infection.

#### 4.4 | The role of ROS in the phagosome

The complex nature of ROS production illustrates the dynamic environment of the phagosome. Early findings by Klebanoff suggested that the bactericidal mechanism of neutrophils was mediated through MPO-catalyzed iodination and chlorination.<sup>110,111</sup> He hypothesized that MPO, rather than having a direct bactericidal effect itself, might act indirectly in the phagosome to catalyze the production of substances with potent antibacterial properties. Initial investigations using iodine, which, when oxidized with H<sub>2</sub>O<sub>2</sub>, become the potent germicide iodide, demonstrated that MPO was only bactericidal when combined with iodide ions and H<sub>2</sub>O<sub>2</sub>. MPO-dependent iodination was traced within close proximity of phagocytosed bacteria using electron-microscopic autoradiographs on silver grains and demonstrated that the reaction did not happen in resting neutrophils, or in those treated with an MPO inhibitor.<sup>110</sup> Similar results were obtained with chloride, which is converted into HOCl<sup>111</sup>; further studies have shown evidence of chlorination inside the phagosome and concluded that the amount of HOCl produced in the phagosome is sufficient to kill engulfed bacteria.<sup>112–115</sup> These findings elucidated roles for the respiratory burst, H<sub>2</sub>O<sub>2</sub>, and MPO in the neutrophil. However, these results have been contested, with subsequent experiments suggesting that ROS are not directly microbicidal; this aspect is further explored in section 8.

### 5 | METABOLISM DURING PHAGOCYTOSIS

Phagocytosis is an active process: Protein phosphorylation and the generation of ROS require an energy source. Unlike monocytes,

whose oxygen consumption, and thus phagocytic capacity, is substantially reduced by mitochondrial respiratory chain inhibitors, neutrophils derive most of their energy from glycolysis, even under aerobic conditions.<sup>116,117</sup> Early studies of metabolism in phagocytosing neutrophils have shown varied results: Some studies indicated increased glucose utilization and lactate production while others did not.<sup>118–121</sup> Variations in host species, phagocytic prey, or the availability of extracellular glucose may explain these experimental discrepancies.

The accepted dogma is that neutrophils rely on glucose-fueled glycolysis to provide the energy for phagocytosis. Recently, transcriptomic analysis of neutrophils from patients with sepsis compared with healthy controls revealed upregulation of glycolytic gene expression, as well as a reduction in phagocytosis when healthy neutrophils were treated with glycolytic pathway inhibitors.<sup>122</sup> However, in vivo the availability of glucose may be limited, for example in a purulent exudate. Detailed studies using genetically modified mice have shown that intracellular glucose and glycogen shuttling, and glycogen storage capacity are all important for neutrophils to maintain their ability to generate ROS and kill bacteria in situations where nutrients are lacking.<sup>123,124</sup>

### 6 | ION CHANNELS

The movement of NADPH oxidase-generated electrons into the phagosome requires an ionic current to balance the negative charge produced, without which there would be huge depolarization of the membrane. The enzyme complex is inhibited from 0 mV to +190 mV, at which point electron translocation is abolished, highlighting the necessity of an outward cation or inward anion flux.<sup>125</sup> If the neutrophil phagosome pH remains close to neutral, as some studies suggest,<sup>112,126,127</sup> most of the compensated charge must be due to the movement of protons (H<sup>+</sup>), which are osmotically neutral<sup>128</sup>; only up to 5% of the charge could be compensated by any other ion.

In the 1980s, Henderson et al. found NADPH oxidase to be electrogenic and suggested that the negative charge was wholly compensated by H<sup>+</sup> movement into the phagosome.<sup>129</sup> There was speculation that the passage of H<sup>+</sup> could be through gp91<sup>phox</sup> itself<sup>130,131</sup>; however, bioinformatic analysis of known voltage-gated cation channels, followed by cloning of the H<sup>+</sup> voltage-gated channel 1 (Hv1) gene and examination of Hv1 knockout mouse neutrophils, confirmed the existence of a specific voltage-gated H<sup>+</sup> channel.<sup>18</sup>

Hv1, activated at depolarizing voltages, opens or closes a voltage-gated H<sup>+</sup> channel. Usually voltage-gated ion channels comprise both voltage-sensing and pore domains, but as Hv1 lacks the pore domain it is highly selective and tightly regulated by the transmembrane pH gradient, as well as voltage.<sup>132,133</sup> In neutrophils, Hv1 is highly expressed and delivered to the phagosome by granules with, but remaining independent of, NOX2.<sup>134,135</sup> Vacuolar-type ATPase (V-ATPase) is another H<sup>+</sup> transport



mechanism that actively pumps  $H^+$  into the phagosome using ATP hydrolysis.

The electrogenicity of NADPH oxidase and the consequent need for charge compensation was classically thought to be a nuisance. However, it is possible that charge compensation mechanisms serve a purpose for the cell<sup>136</sup>  $H^+$  flux from the cytoplasm into the phagosome not only compensates the electrogenicity of the NADPH oxidase, but helps prevent osmotic disruptions, provides substrates for  $H_2O_2$  and HOCl generation, and minimizes pH disturbances, both in the phagosome and cytoplasm. The  $H_2O$  and HOCl formed from  $H^+$  and electrons are membrane permeable and thus have no osmotic effect. Strong acidification of the neutrophil cytoplasm is avoided through consumption of  $H^+$  during the production of ROS, and significant alkalization of the phagosome is avoided through the dismutation of  $O_2^{\cdot-}$ .

Surprisingly, deletion of Hv1 does not prevent ROS production completely, suggesting that other charge compensation mechanisms either exist in parallel or can compensate for lack of  $H^+$  transport.<sup>24</sup> Chloride ( $Cl^-$ ) has been suggested as a compensatory ion<sup>137</sup>; however, it is estimated that approximately 90% of the oxygen consumed by NADPH oxidase is utilized for MPO-catalyzed HOCl production and, given this requirement for  $Cl^-$  supply, it is unlikely that  $Cl^-$  movement out of the phagosome is responsible for all of the charge compensation.<sup>138</sup> In stimulated neutrophils, the cytosolic level of  $Cl^-$  is lower than the estimated concentration of 70 mM in the phagosome, suggesting active transport mechanisms enable  $Cl^-$  accumulation in the phagosome.<sup>139</sup>

Two  $Cl^-$  channels (CIC), the cystic fibrosis transmembrane conductance regulator (CFTR) and CIC-3, have been identified on neutrophil phagosomes.<sup>140,141</sup> Neutrophils from *Clcn3(-/-)* mice display reduced phagocytosis and ROS generation, demonstrating its importance in neutrophil phagocytic function.<sup>141</sup> It has been suggested that CIC-3 is a  $Cl^-/H^+$  antiporter rather than an ion channel, meaning it could extrude  $H^+$  against the electrochemical gradient.<sup>142,143</sup> This appears counter-productive to the accumulation of  $H^+$  in the phagosome; however, CIC-3 may be partly responsible for  $H^+$  leak out of the phagosome (section 6).

CFTR is a cAMP-activated chloride channel, which contributes approximately 50% of the total halide transport activity in neutrophils and is responsible for maintaining phagosomal HOCl levels.<sup>144</sup> CFTR is defective in patients with cystic fibrosis (CF)<sup>145</sup>; neutrophils from CF patients with the typical  $\Delta F508$  homozygous mutation exhibit significantly reduced CFTR content in the phagosomal membrane<sup>146</sup> and have impaired HOCl production.<sup>140</sup> Although there are additional reasons why CF patients suffer recurrent infections, an impaired antimicrobial environment in the neutrophil phagosome due to defective  $Cl^-$  transport may be an important contributing factor, and killing of *Pseudomonas aeruginosa* (a common pathogen in CF) by neutrophils has been shown to be chloride-dependent (reviewed in<sup>147</sup>) (Table 1).

The  $K^+/Cl^-$  cotransporter KCC3 is also postulated to be present on the neutrophil phagosome: KCC3-deficient murine neutrophils exhibited reduced NADPH oxidase activity due to disturbances in

the recruitment and phosphorylation of its subunits.<sup>162</sup> Furthermore, KCC activity was increased synchronously with neutrophil activation and was required for ROS generation, supporting the theory that KCC3 contributes to microbial killing and potentially to charge compensation.

Calcium ion ( $Ca^{2+}$ ) transport through the phagosomal membrane is more contentious; although it has been thought to regulate phagocytosis,<sup>163</sup> the source of  $Ca^{2+}$  is still debated and the requirement of  $Ca^{2+}$  release for phagosomal oxidase activity remains controversial. This ion does, however, play an important role in regulating signaling and cytoskeletal dynamics during particle ingestion and granule fusion.

Ionic fluxes are dictated by dynamic interactions of channels, transporters, and pumps on the phagosome, which change as the phagosome matures, and have not been fully elucidated. Changes in pH and vacuolar volume provide clues as to the movement of ions in and out of the phagosome, but research into the pH of the neutrophil phagosome has posed challenges.

## 7 | PH OF PHAGOSOME AFTER INGESTION

The idea of phagosomal acidification has been in existence for as long as the concept of phagocytosis itself: Metchnikoff originally hypothesized that acidification was the cause of death for ingested organisms.<sup>164</sup> However, the phagosomal pH of neutrophils is inherently different from other phagocytes as it acidifies more weakly, at least at early timepoints.<sup>126,165</sup> Over the years, there have been many contradictions regarding neutrophil phagosomal pH. Investigators initially reported an acidification,<sup>164</sup> but further studies suggested a biphasic pH change, with initial alkalization followed by a modest acidification to pH ~6.5.<sup>165,166</sup> Subsequently, the intra-phagosomal pH was suggested to remain neutral.<sup>126,127</sup>

Proponents of the biphasic pH change proposed that initial alkalization was due to the consumption of  $H^+$  by the formation of  $H_2O_2$ , and that this alkalinity was beneficial as it optimized bacterial killing by granule proteases.<sup>165</sup> This theory was supported by the sustained acidification seen when NADPH oxidase is inhibited. Similarly, neutrophils from patients with chronic granulomatous disease (CGD) (which lack a functional NOX2) undergo a rapid and extreme acidification during phagocytosis, which is associated with impaired bactericidal function.<sup>165</sup> In contrast to healthy neutrophils, CGD neutrophils did not display the usual phagosome swelling, which was found to be independent of hydrolytic enzyme content and degranulation. It was suggested that normal phagocytic vacuoles enlarge due to an increase in the osmotically active products of bacterial digestion but that this does not occur in CGD neutrophils as the extreme acidification impairs digestion and thus swelling.

The second step in the proposed biphasic pH change is that of modest acidification, occurring once the respiratory burst subsides, and suggested to enhance activity of proteins with acidic pH optima,

TABLE 1 Genetic defects compromising phagocytic machinery and their clinical implications

Clinical condition	Genetic mutations	Clinical consequence	Neutrophil phagosome phenotype
Chronic granulomatous disease (CGD)	Genes encoding components of NADPH oxidase. Commonest affected subunit: gp91 <sup>phox</sup> – X-linked mutations in <i>CYBB</i> gene. Disease severity correlates with NOX2 function.	Recurrent severe bacterial and fungal infections, commonly lung, skin, and liver with abscesses. Persistent inflammation with granulomata.	Absent/diminished production of all phagosomal ROS. Reduced bacterial killing, for example, <i>S. aureus</i> . <sup>148-151</sup>
Myeloperoxidase (MPO) deficiency	<i>MPO</i> gene encoding myeloperoxidase. Commonest mutation: R569W.	Predominantly asymptomatic unless concurrent diabetes. Increased susceptibility to <i>Candida</i> species infection.	Increased phagosomal H <sub>2</sub> O <sub>2</sub> due to i) absent/diminished MPO-catalyzed production of HOCl and ii) prolonged activity of NADPH oxidase. Possible increased bioactivity of granule proteins, for example, elastase due to reduced HOCl-dependent inactivation. Slower <i>S. aureus</i> killing markedly reduced <i>C. albicans</i> killing. <sup>152-155</sup>
Cystic fibrosis (CF)	Cystic fibrosis transmembrane conductance regulator <i>CFTR</i> gene encoding CFTR chloride channel. Commonest mutation: ΔF508. Disease severity correlates with CFTR function.	Abnormally viscous mucus affecting multiple organs, for example, lungs and pancreas. Recurrent bacterial, fungal, and mycobacterial lung infections.	Defective targeting of mutant CFTR to phagosomes. Impaired chloride transport into phagosomes. Reduced phagosomal HOCl production. Reduced <i>P. aeruginosa</i> killing. <sup>144,146,147,156,157</sup>
Elastase deficiency	<i>ELANE</i> gene encoding neutrophil elastase. Missense mutations cause severe congenital neutropenia (SCN), probably due to mis-trafficked or unfolded elastase. Whole <i>ELANE</i> deletion occurs with chromosome 19p terminal deletions.	Unclear clinical effects of elastase deficiency versus neutropenia. Recurrent severe bacterial infection in SCN but not described with <i>ELANE</i> deletion. G-CSF recovery of neutrophil count in SCN does not prevent lethal infection.	Abnormal granule structure and additional antimicrobial peptide deficiency, for example, α-defensins in SCN neutrophils, may also affect MPO activity. Reduced <i>Escherichia coli</i> and <i>C. albicans</i> killing in SCN neutrophils despite G-CSF treatment. Reduced particle ingestion in some SCN neutrophils. <sup>158-161</sup>

for example, hydrolases. The H<sup>+</sup> pump, V-ATPase, was hypothesized to be coupled to NADPH oxidase activity and to be responsible for the moderate acidification. However, in opposition of this theory, the rate of superoxide anion production was unrelated to V-ATPase activity and inhibition of the pump did not provoke alkalization of oxidase-active phagosomes.<sup>126</sup>

Potential limitations of the studies suggesting biphasic pH change<sup>165</sup> include use of the pH sensor fluorescein, non-sealed phagosomes, and dye leakage.<sup>127,166</sup> In addition, pH changes to optimize granule proteins may not be necessary; there are examples of prominent neutrophil enzymes that potentiate MPO-mediated killing even when heat inactivated, for example, elastase and cathepsin G.<sup>167,168</sup> This could be because their cationic nature enables strong, disruptive electrostatic interactions with microbial surfaces, that consequently cause the microbe to be more susceptible to attack.

The current consensus is that neutrophil phagosomes acidify more weakly (certainly compared to macrophages) or remain neutral, due to robust NADPH oxidase activity.<sup>169</sup> This would certainly


make biological sense as NADPH oxidase functions optimally at neutral pH, so maintenance of this would allow maximal ROS production.<sup>170</sup> Indeed, studies of single cells<sup>126</sup> and direct measurement of intra-phagosomal pH<sup>126,127,166</sup> support this. More recently, genetic deletion of Hv1 has confirmed its role in sustaining NADPH oxidase activity (as a key charge compensatory channel), controlling both the homeostasis of ROS production and pH.<sup>169</sup> H<sup>+</sup> consumption, which should alkalinize the phagosome, is counteracted by a combination of active pumping of H<sup>+</sup> into the phagosome by V-ATPase (in of itself not enough to cause acidification as previously thought) and passive movement of H<sup>+</sup> through Hv1.

Consistent with the theory of maintaining neutral pH, Jankowski et al. found that activity of NADPH oxidase itself hindered acidification through several complementary mechanisms: impairing V-ATPase recruitment to the neutrophil phagosome, thus reducing H<sup>+</sup> delivery; increasing phagosomal membrane permeability, therefore increasing passive H<sup>+</sup> back-flux into the cytosol; and increasing luminal H<sup>+</sup> consumption.<sup>127</sup> The mechanisms of the impaired V-ATPase

recruitment and increased  $H^+$  leak are uncertain. Theories for the former include V-ATPase inhibition by ROS and oxidase-induced depolarization of the phagosomal membrane, causing a depressed flux of  $Ca^{2+}$ .<sup>127</sup> In support of these findings, Chemaly et al. confirmed an inverse correlation of intra-phagosomal ROS production and V-ATPase accumulation on phagosomes.<sup>170</sup>

Our group has generated recent evidence for the development of a more acidic pH as phagocytosis progresses (Figure 2). Using bacterial targets conjugated to a pH-sensitive dye, we demonstrated progressive acidification of the maturing neutrophil phagosome at later time points (up to 120 minutes) that was class 3 PI3K- and V-ATPase-dependent.<sup>22</sup> However, calibrated probes were not used at early time points to measure the exact pH. Of

clinical relevance, exposure to the pro-inflammatory complement product, C5a, reduced the proportion of neutrophils phagocytosing (as expected given its effects on RhoA signaling) but also impaired phagosomal acidification in neutrophils that had ingested prey, indicating a distinct mechanism of failed phagosomal acidification. The C5a-mediated reduction in acidification was associated with a reduced phosphorylation of the V-ATPase G subunit. Defective phagosomal acidification was also demonstrated in neutrophils from critically ill patients vulnerable to secondary infection, suggesting that acquired phagosome dysfunction in the context of systemic inflammation may increase susceptibility to infection.<sup>24</sup> Another interesting observation is that when neutrophils phagocytose other apoptotic neutrophils, for example



**FIGURE 2** Maturation of phagosomes formed by human peripheral blood neutrophils ingesting *Staphylococcus aureus*. (A) Neutrophil that has not encountered bacteria (Blue: Nucleus labeled with Hoescht, White: Actin labeled with Alexa Fluor(AF)647-phalloidin). (B) Early ingestion of *S. aureus* (Green: AF488-labeled *S. aureus* bioparticles), with limited change in phagosomal pH as assessed by co-localized lysotracker signal (Red: Lyso Tracker Red DND-99, Yellow: co-localization with AF488 *S. aureus*). (C): Mid-maturation with mixture of low pH (mature) and high pH (immature) phagosomes. (D): Late maturation with the majority of phagosomes demonstrating low pH. (E) Early endosomal antigen 1 (EEA1) staining on the surface of *S. aureus* containing phagosomes. Note the penumbral rather than co-localized signature indicating membrane distribution (Red: EEA1, mouse anti-human EEA1 with secondary anti-mouse AF568 phalloidin staining omitted for clarity)

at sites of acute inflammation where neutrophils significantly outnumber macrophages, their oxidative burst is reduced with a resultant acidification of the phagosome, although pH was again measured at later timepoints (90 minutes).<sup>171</sup> This demonstrates that the content of the phagosome may dictate its pH and consequently its maturation.

The disparity of phagosome pH in different phagocytes highlights the importance of pH regulation as a functional mechanism itself. The neutrophil's counter mechanisms against alkalization and acidification suggest a likely neutral pH, certainly at earlier timepoints, with evidence of a later acidification as the phagosome matures. Further research is needed to examine the impact of systemic inflammation and extreme environments, such as the anaerobic hypoxic environment of infected or inflamed tissue, on pH regulation as the conditions may vary dramatically from *in vitro* studies. For example, the pH of the phagosome may be affected by the extracellular pH, due to knock-on effects on the cytosolic and granule pH.

## 8 | GRANULES

During the process of phagosome formation and maturation, granule fusion with the phagosome plays two fundamental roles: the delivery of antimicrobial granule proteins to the lumen and the insertion of functionally important membrane proteins into the phagosomal membrane, including receptors, ion channels, and gp91<sup>phox</sup>(NOX2)/p22<sup>phox</sup>. This fusion of pre-formed granules is key to the neutrophil's rapid speed of engagement, engulfment, and killing of prey.<sup>7</sup> Granules are estimated to make up about 40% of the phagosome volume<sup>172</sup> and are commonly classified as peroxidase-positive (azurophil) or peroxidase-negative based on whether they contain MPO. However, they are heterogenous and can also be characterized by size, morphology, electron density, or protein content.<sup>92</sup> The generally accepted subtypes are designated azurophil (primary), specific (secondary), and gelatinase (tertiary) granules, and secretory vesicles; this description is predominantly based on the timing of formation and subsequent differential contents, although there is a degree of overlap.<sup>93</sup>

Similar to signaling during phagosome formation, receptor ligation initiates intracellular signaling cascades during neutrophil degranulation (the process of granule delivery to plasma/phagosomal membranes). Although the signaling pathways that regulate degranulation are not completely delineated, activation of integrins,<sup>173</sup> Fc receptors,<sup>174</sup> and GPCRs<sup>175</sup> has been implicated; the precise signaling pattern likely depends on the range, concentration, and context of external stimuli, and downstream signaling events may occur in parallel. As with activation of the NADPH oxidase, priming is also required for maximal degranulation response. However, these pathways converge on common degranulation effector mechanisms, principally cytoskeletal rearrangement, and granule/vesicle trafficking and membrane fusion.<sup>176</sup>

### 8.1 | Granule contents

Neutrophil granules are produced from the Golgi complex by aggregation of immature transport vesicles; distinct stages of secretory activity produce granules with different contents (reviewed in<sup>177</sup>). However, they can also be classified by protein expression more loosely as a continuum. The prevailing hypothesis of granule heterogeneity is that of protein targeting by timing of biosynthesis, proposed by Borregaard et al. whereby concomitantly formed proteins are directed into granules together, and different granule subtypes are formed as protein expression changes during neutrophil maturation.<sup>178</sup> However, it has been shown that active sorting of proteins to granules also occurs.<sup>179</sup>

Granules contain a range of different proteins: Enzymes and defense proteins are present in the lumen, whereas receptors, signaling proteins, and adhesion molecules exist in the membrane. Neutrophil granules mobilize to the plasma membrane for extracellular release and phagocytic receptor delivery (gelatinase and secretory vesicles) or to the phagosomal membrane for intracellular release (specific and azurophil).<sup>180</sup> Specific granules are rich in antimicrobial substances, for example, lactoferrin and lipocalin-2, and also contain membrane proteins, including NOX2.<sup>10</sup> Azurophil granules are highly microbicidal, containing MPO, lysozyme, bactericidal/permeability-increasing protein, defensins, and the serine proteinases elastase, cathepsin G, and proteinase 3.<sup>7</sup> These proteins contribute to microbial killing in various ways, including inducing membrane permeabilization and lysis (e.g., cathelicidins and defensins) and scavenging metals needed for growth and enzyme function (e.g., lactoferrin).<sup>181,182</sup>

### 8.2 | Role of calcium in granule-phagosome fusion

Sengeløv et al.<sup>173</sup> showed an *in vivo* hierarchy of granule subtype mobilization, with complete mobilization of secretory vesicles, and a stepped reduction in the degranulation of gelatinase, specific and azurophil granules, respectively, that was dependent on intracellular calcium concentration.<sup>175</sup> This hierarchical mobilization allows tight functional control, which is essential due to the large number of tissue-destructive proteases contained within azurophil granules<sup>183</sup>; thus, their mobilization requires the most strongest of stimuli.

Granule fusion with the phagosome is inherently different from fusion with the plasma membrane due to the reversed curvature of the membrane<sup>184</sup> and phagosome maturation. Defining the role of Ca<sup>2+</sup> in phagocytosis has proven controversial; transient increases in cytosolic Ca<sup>2+</sup> have been widely, but inconsistently observed during phagocytosis and the functional relevance and source of the Ca<sup>2+</sup> remain a subject of debate (reviewed in<sup>185</sup>). Fusion of specific granules with phagosomes has been shown to be calcium-dependent,<sup>186</sup> but the role of calcium in azurophil granule fusion is uncertain.<sup>187</sup> This distinction could be due to function and timing of fusion: Specific granules can fuse with the plasma membrane anywhere and deliver plasma membrane proteins to the phagosome, such as the subunits of NADPH, whereas azurophil granules mainly fuse with the

phagosomal membrane and contain an array of enzymes and proteins that can damage surrounding tissue.<sup>187,188</sup> Nordenfelt et al. found that azurophil granule-phagosome fusion and late-stage killing in FcγR-mediated phagocytosis was Ca<sup>2+</sup>-independent but that a separate mechanism of early azurophil delivery in the forming phagosome was Ca<sup>2+</sup>-dependent.<sup>189</sup> This suggests that it is only a transient early phase of azurophil granule, and potentially other granule, fusion and killing that is Ca<sup>2+</sup>-dependent and that later phagosome maturation mechanisms, such as ROS production and ion fluxes, make it redundant.

Actin polymerization is also independent of Ca<sup>2+</sup> in neutrophils.<sup>190</sup> Conversely, actin severing and depolymerization by gelsolin, which is crucial to dissolve the thick polymerized actin ring that surrounds the forming phagosome, is Ca<sup>2+</sup>-dependent.<sup>191</sup> Other possible targets for Ca<sup>2+</sup> transients during phagosome formation include the Ca<sup>2+</sup>-dependent protease calpain, which has been implicated in clustering of β2 integrins<sup>192</sup>; the Ca<sup>2+</sup>-binding protein synaptotagmin, which translocates to CR3-initiated phagosomes and is involved in particle uptake in macrophages<sup>193,194</sup>; or the various Ca<sup>2+</sup>-binding annexins, which promotes membrane fusion and may be involved in actin dynamics during phagocytosis.<sup>195,196</sup>

### 8.3 | Cytoskeletal reorganization

Actin polymerization, initiated by signaling cascades after receptor ligation, is essential for the formation of phagosomal cups: Inhibition of actin polymerization with cytochalasin B inhibits phagocytosis, although azurophil granule fusion with the phagosome persists.<sup>197</sup> This suggests that the actin cytoskeleton may not be involved in granule-phagosome fusion.

Microtubule polymerization is also promoted by phosphorylation signaling cascades in activated neutrophils (distinct from actin); late phagosomes are found in close proximity to the centriole, part of the microtubule organizing center (MTOC) where granules are rich.<sup>198</sup> The use of colchicine, which binds to tubulin and prevents its polymerization into microtubules, led to phagosome disorganization and no preferential association with centrioles within the neutrophil, suggesting that microtubules are required to translocate the phagosome toward the granule-rich centriole;<sup>198</sup> in macrophages, this was observed directly.<sup>199</sup> In neutrophils, colchicine also inhibited the association of MTOC and azurophil granules with the phagosome, implicating microtubule involvement in the movement of granules and their delivery to the phagosome, as well as movement of the phagosome itself.<sup>197</sup> The association of kinesin, a motor protein involved in microtubule mediated transport, with neutrophil granules and microtubules also supports this theory.<sup>200</sup>

### 8.4 | GTPases

The cytoskeleton's involvement in neutrophil trafficking is complex and the different granules are variously associated with actin and

microtubules, which may explain how granules are differentially directed and regulated.<sup>183,197</sup> Actin and microtubule dynamics are predominantly controlled by Rac and Rab GTPases, which facilitate granule migration to the phagosome by orchestrating cytoskeletal rearrangement. Vesicle-membrane tethering, docking, and fusion are then regulated by Munc family proteins and SNAREs (soluble NSF-attachment protein receptors).

The precise secretory machinery associated with a particular granule is linked to its classification and function. Neutrophils display a non-redundant role for Rac2 in the secretion of azurophil, but not specific or gelatinase granules: Neutrophils from mice deficient in Rac2 showed absent azurophil granule exocytosis whereas specific and gelatinase granule release was unaffected.<sup>201</sup> Rac2 deficiency appears to mainly affect actin function in phagocytosis.<sup>202</sup> Rac1 and Rac2 are also required for the formation of NADPH oxidase, depending on the microorganism phagocytosed.<sup>203</sup>

An example of directional granule trafficking control is demonstrated by Rab27a. Subcellular fractionation and immunoelectron microscopy has shown that Rab27a is located predominantly on gelatinase and specific granules, with lesser localization to azurophil granules.<sup>204</sup> Rab27a directs all granules for plasma membrane fusion but granules lacking Rab27a are still able to fuse with the phagosome: Neutrophils from Rab27a-deficient mice exhibited normal phagosome maturation and azurophil granule recruitment.<sup>205,206</sup> Conversely, Munc13-4, an effector of Rab27a present on all three granule subtypes, was found to be essential for phagosomal maturation and delivery of azurophilic granules to the phagosome, with its absence leading to impaired intracellular bacterial killing.<sup>205</sup> Therefore, Munc13-4 is a protein of key significance in phagosomal killing due to the importance of MPO in formation of oxidized halides, and the delivery of toxic proteases and bactericidal peptides. Munc13-4 may regulate granule tethering to the phagosome, which correlates with the finding that tethering of secretory lysosomes to the plasma membrane in cytotoxic T lymphocytes requires the Munc13-4-Rab27 complex.<sup>207</sup> Alternatively, Munc13-4 may form complexes with SNARE proteins to regulate azurophil granule fusion.

Regulation of differential granule fusion and mobilization is also mediated by Src tyrosine kinases and granule associated-SNARE complexes, whereby association with certain family members dictates whether a granule is directed to the phagosome or plasma membrane.<sup>208,209</sup> For example, the Src family member, Hck, was found to be localized to azurophil granules, with translocation directed toward the phagosome<sup>210</sup> whereas Fgr was associated with specific granules, and increased their fusion with the plasma membrane.<sup>211</sup> Similarly, differential granule fusion is directed by various members of SNARE complexes.<sup>208,212</sup>

Overall, a number of secretory control mechanisms ensure that the majority of azurophil granules (unlike specific and gelatinase granules) are preferentially targeted to the phagosome rather than the cell surface membrane, thus establishing a highly toxic phagosomal environment while limiting the capacity for host tissue damage.

## 9 | OXIDATIVE VERSUS NON-OXIDATIVE KILLING

Conventionally, the two arms of oxidative (ROS) and non-oxidative (granule protein) killing in neutrophil phagosomes have been seen as opposing mechanisms, with evidence both for and against a directly microbicidal role for ROS in the phagosome. However, current research suggests a more synergistic approach.

As outlined in section 3, it was elucidated early on by Klebanoff that the respiratory burst in neutrophils produces a large amount of hydrogen peroxide ( $H_2O_2$ ).<sup>94</sup> As  $H_2O_2$  is a potential substrate of MPO, the primary hypothesis was that MPO had an indirect microbicidal effect in the phagosome via MPO-catalyzed oxidation of antimicrobial substances.<sup>59</sup> In support of this theory, there is evidence of chlorination of ingested material,<sup>112-115</sup> and iodination is reduced in neutrophils from patients with MPO deficiency.<sup>213,214</sup> However, no loss of bacterial viability was observed in the pH range of 7.0–8.0, unless the  $H_2O_2$  and iodide concentrations were markedly increased, suggesting that these results may not accurately reflect the phagosome environment. Similar findings were demonstrated using chloride but to a lesser extent.

Evidence for the importance of NOX2, and thus ROS generation, is provided by patients with CGD. CGD occurs as the result of a genetic mutation in any of the five subunits of NADPH oxidase and manifests as a primary immunodeficiency characterized by severe, prolonged, and often fatal infections. Neutrophils from CGD patients are unable to generate ROS, manifesting clinically as a profound susceptibility to bacterial and fungal infections. However, neutrophil microbial killing in these patients is not completely absent, which is best explained by the action of cytotoxic granule proteins (Table 1). This is demonstrated by the preserved ability of CGD neutrophils to kill *Escherichia coli*, where bacterial clearance is thought to be mediated by Bactericidal/permeability-increasing protein,<sup>215,216</sup> indicating that non-oxidative mechanisms can produce sufficient microbicidal effect against certain bacteria.

The original theory<sup>217</sup> of a direct microbicidal role for ROS was subsequently contested (although the importance of MPO was not disputed). Segal et al. utilized stoichiometry to calculate that only a very small proportion of the oxygen consumed during the respiratory burst was utilized for iodination.<sup>216,217</sup> Furthermore, the pH used in previous experiments demonstrating chlorination, around 4.6–5.0, was thought to be too acidic, and it was also suggested that the initial in vitro experiments on the MPO/ $H_2O_2$ /Cl<sup>-</sup> system had used unrealistically low granule protein concentrations. A later reassessment, using higher concentrations of superoxide radicals and granule proteins and a less acidic pH, reported that the directly microbicidal effects of  $H_2O_2$  and HOCl were abolished.<sup>218</sup> Segal's group proposed that as the majority of iodinated proteins were of host rather than bacterial origin, ROS may be ineffective against bacteria within the confines of the phagosome due to consumption of HOCl by the large amount of host-derived protein. Despite this finding, there is evidence of HOCl-mediated oxidation of bacterial methionine residues.<sup>219</sup> Moreover, *E. coli* ingested by healthy

neutrophils upregulated oxidant-sensing genes (e.g., OxyR), and disruption of the bacterial methionine sulfoxide repair system ( $\Delta oxyRS$  mutant) rendered *E. coli* more susceptible to neutrophil-mediated killing.<sup>220</sup> Intriguingly, neither upregulation of the oxidant-sensing system or enhanced susceptibility of the mutant was observed when *E. coli* were phagocytosed by neutrophils from CGD patients.

An opposing theory of non-oxidative killing was thus proposed whereby the microbicidal capacity of neutrophils was due to the abundant granule proteins instead of ROS. It was suggested that the purpose of the electrogenic current of NADPH oxidase was to drive compensatory ion fluxes to optimize the conditions in the phagosome for granule enzymes (rather than primarily to generate ROS), which could also explain why pathogen killing is impaired in CGD.<sup>136</sup> However, whether any cations other than H<sup>+</sup> significantly influx into the phagosome is debatable and depends largely on the pH of the phagosome: If the pH remains close to neutral (as currently thought), then the influx will be predominantly H<sup>+</sup> whereas if the pH rises (initial alkalinization) then other cations must be involved (section 6). Calculations based on the small amount of osmotic swelling of the phagosome estimate that most of the charge compensation must be osmotically neutral.<sup>128</sup> Thus, it is likely that the only cation entering the phagosome is H<sup>+</sup>.

Evidence for the importance of neutrophil granule enzyme-mediated killing is provided by mice deficient in the proteases, cathepsin G, and elastase. Neutrophils from these mice exhibited normal respiratory burst, ROS production, and iodination, but the mice were unable to resist infection with *S. aureus* (a prominent cause of infection in CGD) or *Candida albicans* (which also causes severe infection in MPO-deficient mice).<sup>136</sup> Reeves et al. suggested a role for MPO in protecting granule enzymes against oxidative damage, for example cathepsin G, which is very sensitive to oxidation by  $H_2O_2$ . In this scenario, the impaired killing observed in neutrophils from MPO-deficient mice<sup>221</sup> was suggested to be due to increased accumulation of  $H_2O_2$ ,<sup>222</sup> potentially leading to the oxidative inactivation of antimicrobial proteins. Conversely, as HOCl can also inactivate antimicrobial proteins, such as elastase, a lack of MPO may actually enhance their activity. Assessing the role of single proteases in humans is even more complex; for example, mutations in elastase may cause disease through protein misfolding and mis-trafficking, resulting in neutropenia, as well as from deficiency of the protease (Table 1).

The disparity in the clinical manifestations of CGD, where patients suffer significant infection-related morbidity and mortality, and MPO deficiency, which somewhat surprisingly goes largely unnoticed, questions the predominance of a single killing mechanism. CGD patients experience severity of disease correlated to the degree of impairment of NADPH oxidase; however, the precise mechanism of antimicrobial defense due to NADPH oxidase is ambiguous.<sup>148</sup> Although MPO deficiency in humans does not commonly present with critical illness,<sup>223</sup> MPO-deficient neutrophils exhibit less efficient bacterial killing<sup>224</sup> (Table 1). The reduced killing speed observed in MPO-deficient neutrophils *in vitro*<sup>152,153</sup> suggests that MPO may be involved early, and that proteases, potentially acting later as the

pH is optimized, can compensate for the deficiency in vivo. An important early report highlighted the potential compensatory contributions of peroxidase-dependent and peroxidase-independent microbicidal systems by comparing neutrophils from patients with CGD or MPO deficiency.<sup>225</sup> Furthermore, when both systems are operational in healthy neutrophils, oxidative and non-oxidative microbial killing may be synergistic, as demonstrated by the ability of elastase to potentiate MPO-dependent killing of *S. aureus* or *E. coli* in vitro in an acidic environment.<sup>167</sup>

The mechanism of oxidative killing is still controversial. Evolutionarily, the theory that NADPH oxidase only produces ROS for the purpose of membrane potential changes is unlikely due to the toxicity of its products. However, as the lifespan of neutrophils is so short, it is questionable whether neutrophils need to be protected against ROS. Presently, the conclusion is that there is not a single predominant mechanism, but a complex synergistic relationship between oxidative and non-oxidative killing that provides alternative compensatory systems, as is often the case with the immune system (Figure 3).


## 10 | WHEN THE PHAGOSOME CANNOT FORM

Neutrophils are powerful and efficient killers, but bacteria are formidable opponents and with their rapid evolutionary speed they have developed a range of evasion tactics. A common evasion tactic of bacteria is the formation of aggregates that circumvent phagocytosis. Large fungal hyphae also obviate phagocytosis. However, ligation of the phagocytic dectin-1 receptor, which detects fungal elements, can divert degradative proteins from the phagosome to the nucleus, thereby enabling the formation of neutrophil extracellular traps (NETs) in situations where a pathogen is too large to engulf.<sup>226</sup> NETs, composed of decondensed chromatin acting as a scaffold for cytosolic and granule proteins, act to trap pathogens which cannot


be phagocytosed, and can exert antimicrobial effects via exposed antimicrobial molecules.<sup>227</sup>

Metzler et al. found that in a subset of resting neutrophils there is a protein complex, termed the azurosome, which is localized on the azurophil membrane.<sup>228</sup> This complex acts as a scaffold for proteins, including elastase and MPO, and is regulated by oxidation, specifically by  $H_2O_2$ . When activated by  $H_2O_2$ , some proteins, including elastase, dissociate from intact granules into the cytosol by an unknown mechanism. Elastase is then activated in an MPO-dependent manner and is free to enter the nucleus and cleave histones, producing NETs.<sup>228</sup> Thus, similar to pathogen killing during

### (A) Granule delivery of phagocytic machinery



### (B) Antimicrobial environment of the phagosome



**FIGURE 3** Role of granules and reactive oxygen species in phagosome formation and function. (A): Granules deliver phagocytic machinery to the plasma and phagosomal membranes during phagocytosis. Secretory vesicles supply phagocytic receptors to the plasma membrane. Specific and gelatinase granules deliver NADPH oxidase components and ion channels to the phagosomal membrane. Azurophil granules supply myeloperoxidase (MPO) and cytotoxic proteins and proteases, including elastase. (B) Reactive oxygen species (ROS) and granules contribute synergistically to microbial killing. NADPH oxidase translocates electrons into the phagosome which react with molecular oxygen to form superoxide ( $O_2^-$ ). Electrogenic charge is compensated by proton ( $H^+$ ) influx, predominantly through Hv1 channels. MPO catalyzes the reaction of  $O_2^-$  and  $H^+$  to form hydrogen peroxide ( $H_2O_2$ ). Chloride ( $Cl^-$ ) enters the phagosome, for example, through CFTR channels. MPO oxidizes  $Cl^-$  to form HOCl which is likely to be directly microbicidal. Non-oxidative proteins and proteases, for example, elastase and cathepsin G, are also directly microbicidal. Numbers in blue indicate clinical syndromes associated with mutations in various steps of the pathway (Table 1)

phagocytosis, NET-generation may be dependent on both ROS and granule proteins.

The ability of neutrophils to “sense” pathogen size during attempted phagocytosis is intriguing. When neutrophils were exposed to *Candida albicans* hyphae, over 70% of total elastase was seen in the nucleus, accompanied by NET release, compared to less than 20% when *Candida albicans* was in the easily phagocytosable yeast form.<sup>226</sup> The exact mechanism is unclear, but it is possible that the “decision” for NETosis versus phagocytosis is dependent on whether elastase is delivered to the phagosome: phagocytosis, which acts more rapidly than NETosis (seconds to minutes versus minutes to hours), sequesters elastase to the phagosome and inhibits the production of NETs. An additional response to fungal hyphae is mediated via CR3 ligation by the fungal PAMP,  $\beta$  glucan, and components of the extracellular matrix. Here, temporally regulated CR3 cross-talk with different  $\beta_1$  integrins can instigate neutrophil swarming, whereby neutrophil aggregates can form a so-called attack complex, prior to NETosis.<sup>229</sup> This size-dependent regulation ensures that the destructive collateral damage from NETosis is only permitted when phagocytosis is not a viable option. Intriguingly, NETosis does not appear to preclude the ability of neutrophils to also perform phagocytosis, and anuclear neutrophils containing bacteria have been observed, implying that NET generation and phagosome maturation can be compartmentalized within the same cell.<sup>230</sup>

This “decision making” during neutrophil phagocytosis highlights the complexity and plasticity of the neutrophil and shows that even when neutrophils are unable to form a phagosome, there are other microbicidal mechanisms they can employ to protect the host.

## 11 | PHAGOSOME RESOLUTION

Phagosome resolution is the concluding stage in the phagocytic process, comprising disposal or recycling of ingested contents and membrane resources, as well as providing the opportunity for antigen presentation.<sup>231,232</sup> Resolution is an important step that allows phagocytes to return to homeostasis but, in comparison with phagosome formation and maturation, resolution remains largely unexplored. Consequently, the molecular basis of phagosome resolution is rather speculative, predominantly based upon related processes such as lysosome turnover and autophagy. Limited experiments in phagosomes have almost exclusively been performed using macrophages (reviewed in [4]), whose phagosomes mature through fusion with endosomes and lysosomes in a manner which is very different from neutrophil phagosome-granule fusion. Therefore, although knowledge acquired from these experiments is often extrapolated, findings in macrophages during this phagocytic stage may not be truly representative of the situation in neutrophils.

The objectives of phagosome resolution are manifold: safe disposal of destroyed prey; waste management (including recycling) of ingested contents, including nucleic acids, proteins, and lipids; antigen presentation to lymphoid cells; and resorption of the phagosomal membrane. Resolution was originally thought to proceed in

a similar fashion to unicellular eukaryotes, which egest indigestible content and plasma membrane.<sup>233</sup> However, recent work in macrophages has shown evidence of shrinkage and fission,<sup>234,235</sup> with resolution of phagolysosomes through fragmentation, by vesicle budding, tubulation, and constriction.<sup>236</sup> It is unclear how this reformation of lysosome-like organelles during fragmentation would apply to neutrophils as their granules are pre-formed. In addition to mechanical contraction, exportation of organic osmolytes produced by target degradation is essential for volume loss during phagosome resolution to prevent osmotically-induced hydrostatic pressure (reviewed in<sup>237</sup>).

One conundrum of the phagosome maturation and resolution process is that phagocytes must be able to target the intraphagosomal lipids of ingested prey for degradation while maintaining an intact phagosomal membrane that is protected from lipolytic enzymes, but also be able to resorb this membrane once internal degradation is completed. The mechanisms controlling phagosomal membrane recycling are largely unknown but possibilities for the handling of various lipid species are discussed in.<sup>4</sup>

There is a particular paucity of information regarding signaling events for phagosome resolution. In macrophages, the fragmentation process is thought to involve phosphoinositides, particularly PtdIns4P, and Rab GTPases,<sup>237</sup> which direct tethering to the endoplasmic reticulum.<sup>235</sup> Whether this process occurs in neutrophils is uncertain.

Given that phagosome resolution is a pivotal step in the successful completion of phagocytosis, this stage has garnered surprisingly little attention. In neutrophils, the processes required for phagosome resolution are almost completely unknown. Macrophages must be equipped to perform multiple rounds of phagocytosis while maintaining phagocytic and degradative capacity but it is not clear how vital this step is in the relatively short-lived neutrophil, which is itself efferocytosed by macrophages during inflammation resolution. Given the unique nature of neutrophil granules, and the involvement of proteins required for phagosome fragmentation throughout phagosome maturation, there will be considerable challenges in elucidating this process.

## 12 | CONCLUSION

For the past few decades, progress in neutrophil biology has been extremely slow; many researchers have opted to study cell types that are more easily cultured and genetically manipulated. Over time, however, the neutrophil has been revealed as unique, and its plasticity and complexity underestimated. The view that neutrophils are single function suicide killers has been overtaken by evidence of integration of complex signals to make “decisions” and instigate a range of different signaling pathways in the face of invading pathogens. This is particularly apparent in the neutrophil phagosome, which has phenotypic and functional plasticity. We now know that the neutrophil phagosome environment changes to optimize killing, but an interesting debate still surrounds the complex interplay between various systems.



One key area of contention regards the antimicrobial role of ROS, which are small, transient, and ubiquitous, juxtaposed with that of granule proteins, which are macromolecules with specific killing mechanisms. For years, researchers have believed a single mechanism to be predominant; however, more recent research suggests synergism. This is highlighted by the different but overlapping consequences of CGD, MPO deficiency, and cathepsin G and/or elastase knockout. As may be expected, the ability of the compromised host to prevent bacterial or fungal infection also varies depending on the target organism.<sup>136,224,238,239</sup> It is likely that some pathogens are more susceptible to a particular mode of killing. For example, neutrophil killing of *S. aureus* appears to require ROS as cells experiencing a lack of molecular oxygen under hypoxia displayed impaired bacterial killing (but not ingestion).<sup>240</sup> This makes sense in the context of the wider immune system, which employs a combination of different effector systems to tackle infection.

Another area of phagosome research which has not yet provided a conclusive answer is that of intra-phagosomal pH. Variation in experimental techniques and theories has led to a range of different persuading arguments; however, there is consensus that neutrophils are unique phagocytes, with a weaker acidification than macrophages due to their increased ROS production, and it is likely that the phagosome acidifies more as it matures. The lack of decisive data on pH has also led to ambiguity about the phagosomal ion fluxes. H<sup>+</sup> and some Cl<sup>-</sup> transportation mechanisms are well established, but the involvement of other ions, such as K<sup>+</sup> and Ca<sup>2+</sup>, is still unclear. Ion transport mechanisms are also intrinsic to other systems, such as MPO-mediated chlorination and NADPH oxidase activity. This interconnected web of components and factors, which in turn react together to create many derivatives, are the crux of the dynamic phagosome. Consequently, trying to model the changing biochemistry of the phagosome in vitro is challenging and extrapolating from studies of the systems in isolation, for example, inherited defects and murine knockouts, can undermine the complexity of neutrophil phagocytosis.

Some aspects of the neutrophil phagosome are well established, for example, the formation of NADPH oxidase, production of ROS, and granule fusion mechanisms. However, due to the functional plasticity of neutrophil phagocytosis, we must continue to investigate the unique and distinct mechanisms to understand the complex interaction of neutrophils with each other, with various pathogens, and with other cells. Advances in proteomics, live imaging techniques, and genome sequencing have advanced our understanding in this fascinating fusion of cell biology and microbiology. To further our knowledge of neutrophil phagocytosis, we must employ careful experimental design and analyze potentially limited technological approaches, while acknowledging and appreciating the complexity of the immune system.

## ACKNOWLEDGMENTS

The authors thank Philip Ball for providing illustrations for Figures 1 and 3. Supported by Avant Doctor in Training Research Scholarship 2021/000023 (Alexander JT Wood); Evelyn Trust Clinical Research

Fellowship (Matthew Routledge); British Infection Association (Matthew Routledge); Academy of Medical Sciences Starter Grant for Clinical Lecturers SGL023\1108 (Andrew P Stewart); Medical Research Council Clinician Scientist Fellowship MR/V006118/1 (Andrew Conway Morris); Academy of Medical Sciences Starter Grant for Clinical Lecturers SGL024\1086 (Katharine M Lodge); and NIHR Imperial Biomedical Research Centre (Katharine M Lodge, Edwin R Chilvers)

## CONFLICT OF INTEREST

Andrew Conway Morris is a member of the scientific advisory board of Cambridge Infection Diagnostics Ltd and reports speaking fees from Boston Scientific.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in PRIDE PRoteomics IDentifications Database at <https://www.ebi.ac.uk/pride/>, reference number PXD017092.<sup>241</sup>

## ORCID

Alexander JT Wood  <https://orcid.org/0000-0001-7819-0447>

Edwin R Chilvers  <https://orcid.org/0000-0002-4230-9677>

Katharine M Lodge  <https://orcid.org/0000-0002-3203-9941>

## REFERENCES

- Rabinovitch M. Professional and non-professional phagocytes: an introduction. *Trends Cell Biol.* 1995;5(3):85-87.
- Uribe-Querol E, Rosales C. Phagocytosis: our current understanding of a universal biological process. *Front Immunol.* 2020;11:1066.
- Matsushima H, Geng S, Lu R, et al. Neutrophil differentiation into a unique hybrid population exhibiting dual phenotype and functionality of neutrophils and dendritic cells. *Blood.* 2013;121(10):1677-1689.
- Levin R, Grinstein S, Canton J. The life cycle of phagosomes: formation, maturation, and resolution. *Immunol Rev.* 2016;273(1):156-179.
- Mócsai A. Diverse novel functions of neutrophils in immunity, inflammation, and beyond. *J Exp Med.* 2013;210(7):1283-1299.
- Luerman GC, Uriarte SM, Rane MJ, McLeish KR. Application of proteomics to neutrophil biology. *J Proteomics.* 2010;73(3):552-561.
- Nordenfelt P, Tapper H. Phagosome dynamics during phagocytosis by neutrophils. *J Leukoc Biol.* 2011;90(2):271-284.
- Kay AB. Paul Ehrlich and the early history of granulocytes. *Microbiol Spectr.* 2016;4(4). Available from: <https://doi.org/10.1128/microbiolspec.MCHD-0032-2016>
- Cohn ZA, Hirsch JG. The influence of phagocytosis on the intracellular distribution of granule-associated components of polymorphonuclear leucocytes. *J Exp Med.* 1960;112:1015-1022.
- Hirsch JG. Phagocytin: a bactericidal substance from polymorphonuclear leucocytes. *J Exp Med.* 1956;103(5):589-611.
- Zeya HI, Spitznagel JK. Arginine-rich proteins of polymorphonuclear leukocyte lysosomes. Antimicrobial specificity and biochemical heterogeneity. *J Exp Med.* 1968;127(5):927-941.
- Blanter M, Gouwy M, Struyf S. Studying neutrophil function in vitro: Cell models and environmental factors. *J Inflamm Res.* 2021;14:141-162.
- Kamenyeva O, Boularan C, Kabat J, et al. Neutrophil recruitment to lymph nodes limits local humoral response to *Staphylococcus aureus*. *PLoS Pathog.* 2015;11(4):e1004827.

14. Pizzagalli DU, Latino I, Pulfer A, et al. Characterization of the dynamic behavior of neutrophils following influenza vaccination. *Front Immunol*. 2019;10:2621.
15. Winterbourn CC, van Dalen CJ, Hampton MB, Kettle AJ. Reactions of myeloperoxidase and production of Hypochlorous acid in neutrophil phagosomes. In: Petrides PE, Nauseef WM, eds. *The Peroxidase Multigene Family of Enzymes*. Springer, Berlin; 2000. doi:10.1007/978-3-642-58314-8\_8
16. Stuart LM, Boulais J, Charriere GM, et al. A systems biology analysis of the drosophila phagosome. *Nature*. 2007;445(7123):95-101.
17. Steinberg BE, Grinstein S. Analysis of macrophage phagocytosis: quantitative assays of phagosome formation and maturation using high-throughput fluorescence microscopy. *Methods Mol Biol*. 2009;531:45-56.
18. Ramsey IS, Ruchti E, Kaczmarek JS, Clapham DE. Hv1 proton channels are required for high-level NADPH oxidase-dependent superoxide production during the phagocyte respiratory burst. *Proc Natl Acad Sci USA*. 2009;106(18):7642-7647.
19. Chow C, Downey GP, Grinstein S. Measurements of phagocytosis and phagosomal maturation. *Curr Protoc Cell Biol*. 2004;cb1507s22. Available from: <https://doi.org/10.1002/0471143030.cb1507s22>
20. Nauseef WM. How human neutrophils kill and degrade microbes: an integrated view. *Immunol Rev*. 2007;219(1):88-102.
21. Minakami R, Maehara Y, Kamakura S, Kumano O, Miyano K, Sumimoto H. Membrane phospholipid metabolism during phagocytosis in human neutrophils. *Genes Cells*. 2010;15(5):409-424.
22. Wood AJT, Vassallo AM, Ruchaud-Sparagano MH, et al. C5a impairs phagosomal maturation in the neutrophil through phosphoproteomic remodeling. *JCI Insight*. 2020;5(15):e137029.
23. David C, Fabregat MA, Robin H, et al. The Reactome pathway knowledgebase. *Nucleic Acids Res*. 2014;42:D472-D477.
24. Muschter S, Berthold T, Bhardwaj G, et al. Mass spectrometric phosphoproteome analysis of small-sized samples of human neutrophils. *Clin Chim Acta*. 2015;451:199-207.
25. Tak T, Wijten P, Heeres M, et al. Human CD62Ldim neutrophils identified as a separate subset by proteome profiling and in vivo pulse-chase labeling. *Blood*. 2017;129(26):3476-3485.
26. Amara N, Cooper M, Voronkova M, Cell BW. 2021 U. selective activation of PFKL suppresses the phagocytic oxidative burst. *Cell*. 2021;184(17):4480-4494.
27. Thau L, Asuka E, Mahajan K. Physiology, Opsonization. *StatPearls [Internet]*. Statpearls Publishing. updated 2022. Available from; 2019. <http://www.ncbi.nlm.nih.gov/pubmed/30480954>
28. Hajishengallis G, Reis ES, Mastellos DC, Ricklin D, Lambris JD. Novel mechanisms and functions of complement. *Nat Immunol*. 2017;18(12):1288-1298.
29. Flannagan RS, Jaumouillé V, Grinstein S. The cell biology of phagocytosis. *Annu Rev Pathol*. 2012;7:61-98.
30. Herant M, Heinrich V, Dembo M. Mechanics of neutrophil phagocytosis: experiments and quantitative models. *J Cell Sci*. 2006;119(9):1903-1913.
31. Nobes CD, Hall A. Rho, Rac, and Cdc42 GTPases regulate the assembly of multimolecular focal complexes associated with Actin stress fibers, lamellipodia, and filopodia. *Cell*. 1995;81(1):53-62.
32. Botelho RJ, Teruel M, Dierckman R, et al. Localized biphasic changes in Phosphatidylinositol-4,5-bisphosphate at sites of phagocytosis. *J Cell Biol*. 2000;151(7):1353-1368.
33. Scott CC, Dobson W, Botelho RJ, et al. Phosphatidylinositol-4,5-bisphosphate hydrolysis directs Actin remodeling during phagocytosis. *J Cell Biol*. 2005;169(1):139-149.
34. Kaplan G. Differences in the mode of phagocytosis with fc and C3 receptors in macrophages. *Scand J Immunol*. 1977;6(8):797-807.
35. Nunes P, Demaurex N. The role of calcium signaling in phagocytosis. *J Leukoc Biol*. 2010;88(1):57-68.
36. Hall AB, Gakidis MAM, Glogauer M, et al. Requirements for Vav guanine nucleotide exchange factors and rho GTPases in FcγR- and complement-mediated phagocytosis. *Immunity*. 2006;24(3):305-316.
37. Patel PC, Harrison RE. Membrane ruffles capture C3bi-opsonized particles in activated macrophages. *Mol Biol Cell*. 2008;19(11):4628-4639.
38. Jaumouillé V, Cartagena-Rivera AX, Waterman CM. Coupling of β2 integrins to Actin by a mechanosensitive molecular clutch drives complement receptor-mediated phagocytosis. *Nat Cell Biol*. 2019;21(11):1357-1369.
39. Walbaum S, Ambrosy B, Schütz P, et al. Complement receptor 3 mediates both sinking phagocytosis and phagocytic cup formation via distinct mechanisms. *J Biol Chem*. 2021;296:100256.
40. Gresham HD, Graham IL, Anderson DC, Brown EJ. Leukocyte adhesion-deficient neutrophils fail to amplify phagocytic function in response to stimulation: evidence for CD11b/CD18-dependent and -independent mechanisms of phagocytosis. *J Clin Invest*. 1991;88(2):588-597.
41. McKenzie SE, Schreiber AD. Fc gamma receptors in phagocytes. *Curr Opin Hematol*. 1998;5(1):16-21.
42. John Looney R, Ryan DH, Takahashi K, et al. Identification of a second class of IgG Fc receptors on human neutrophils. A 40 kilodalton molecule also found on eosinophils. *J Exp Med*. 1986;163(4):826-836.
43. Indik ZK, Park JG, Hunter S, Schreiber AD. The molecular dissection of Fcγ receptor mediated phagocytosis. *Blood*. 1995;86(12):4389-4399.
44. Odin JA, Edberg JC, Painter CJ, Kimberly RP, Unkeless JC. Regulation of phagocytosis and [Ca<sup>2+</sup>]<sub>i</sub> flux by distinct regions of an fc receptor. *Science*. 1991;254(5039):1785-1788.
45. Jones DH, Nusbacher J, Anderson CL. Fc receptor-mediated binding and endocytosis by human mononuclear phagocytes: monomeric IgG is not Endocytosed by U937 cells and monocytes. *J Cell Biol*. 1985;100(2):558-564.
46. Jaumouillé V, Farkash Y, Jaqaman K, Das R, Lowell CA, Grinstein S. Actin cytoskeleton reorganization by syk regulates fcγ receptor responsiveness by increasing its lateral mobility and clustering. *Dev Cell*. 2014;29(5):534-546.
47. Kiefer F, Brumell J, Al-Alawi N, et al. The Syk protein tyrosine kinase is essential for Fcγ receptor signaling in macrophages and neutrophils. *Mol Cell Biol*. 1998;18(7):4209-4220.
48. Ozaki N, Suzuki S, Ishida M, et al. Syk-dependent signaling pathways in neutrophils and macrophages are indispensable in the pathogenesis of anti-collagen antibody-induced arthritis. *Int Immunol*. 2012;24(9):539-550.
49. Dewitt S, Tian W, Hallett MB. Localised PtdIns(3,4,5)P3 or PtdIns(3,4)P2 at the phagocytic cup is required for both phagosome closure and Ca<sup>2+</sup> signalling in HL60 neutrophils. *J Cell Sci*. 2006;119(3):443-451.
50. Hawkins PT, Stephens LR, Suire S, Wilson M. PI3K Signaling in Neutrophils. *Curr Top Microbiol Immunol*. 2010;346:183-202.
51. Ellson CD, Gobert-Gosse S, Anderson KE, et al. PtdIns(3)P regulates the neutrophil oxidase complex by binding to the PX domain of p40phox. *Nat Cell Biol*. 2001;3(7):679-682.
52. Matute JD, Arias AA, Wright NAM, et al. A new genetic subgroup of chronic granulomatous disease with autosomal recessive mutations in p40phox and selective defects in neutrophil NADPH oxidase activity. *Blood*. 2009;114(15):3309-3315.
53. Freeman SA, Grinstein S. Phagocytosis: receptors, signal integration, and the cytoskeleton. *Immunol Rev*. 2014;262(1):193-215.
54. Tohyama Y, Life HYI. 2006 U. complement-mediated phagocytosis: the role of Syk. *IUBMB Life*. 2006;58(5-6):304-308.
55. Rosales C. Fc receptor and integrin signaling in phagocytes. *Signal Transduct*. 2007;7(5-6):386-401.

56. Zhang Y, Wang H. Integrin signalling and function in immune cells. *Immunology*. 2012;135(4):268-275.
57. Torres-Gomez A, Cabañas C, Lafuente EM. Phagocytic Integrins: activation and signaling. *Front Immunol*. 2020;11:738.
58. Anderson DC, Schmalsteig FC, Finegold MJ, et al. The severe and moderate phenotypes of heritable mac-1, LFA-1 deficiency: their quantitative definition and relation to leukocyte dysfunction and clinical features. *J Infect Dis*. 1985;152(4):668-689.
59. Bouti P, Webb SDS, Fagerholm SC, et al.  $\beta 2$  integrin signaling Cascade in neutrophils: more than a single function. *Front Immunol*. 2021;11:3930.
60. Mócsai A, Abram CL, Jakus Z, Hu Y, Lanier LL, Lowell CA. Integrin signaling in neutrophils and macrophages uses adaptors containing immunoreceptor tyrosine-based activation motifs. *Nat Immunol* 2006 7:12. 2006;7(12):1326-1333.
61. Freeman SA, Goyette J, Furuya W, et al. Integrins form an expanding diffusional barrier that coordinates phagocytosis. *Cell*. 2016;164(1-2):128-140.
62. le Cabec V, Carréno S, Moisan A, Bordier C, Maridonneau-Parini I. Complement receptor 3 (CD11b/CD18) mediates type I and type II phagocytosis during nonopsonic and opsonic phagocytosis, respectively. *J Immunol*. 2002;169(4):2003-2009.
63. Ridley AJ, Paterson HF, Johnston CL, Diekmann D, Hall A. The small GTP-binding protein rac regulates growth factor-induced membrane ruffling. *Cell*. 1992;70(3):401-410.
64. Ridley A, Hall A. The small GTP-binding protein rho regulates the assembly of focal adhesions and Actin stress fibers in response to growth factors. *Cell*. 1992;70:389-399.
65. Utomo A, Cullere X, Glogauer M, Swat W, Mayadas TN. Vav proteins in neutrophils are required for Fc $\gamma$ R-mediated signaling to Rac GTPases and nicotinamide adenine dinucleotide phosphate oxidase component p40(phox). *J Immunol*. 2006;177(9):6388-6397.
66. Ambruso DR, Knall C, Abell AN, et al. Human neutrophil immunodeficiency syndrome is associated with an inhibitory Rac2 mutation. *Proc Natl Acad Sci USA*. 2000;97(9):4654-4659.
67. Faure MC, Sulpice JC, Delattre M, et al. The recruitment of p47phox and Rac2G12V at the phagosome is transient and phosphatidylserine dependent. *Biol Cell*. 2013;105(11):501-518.
68. Anderson KE, Chessa TAM, Davidson K, et al. PtdIns3P and Rac direct the assembly of the NADPH oxidase on a novel, pre-phagosomal compartment during FcR-mediated phagocytosis in primary mouse neutrophils. *Blood*. 2010;116(23):4978-4989.
69. Magalhaes MAO, Glogauer M. Pivotal advance: phospholipids determine net membrane surface charge resulting in differential localization of active Rac1 and Rac2. *J Leukoc Biol*. 2010;87(4):545-555.
70. Tackenberg H, Möller S, Filippi MD, Laskay T. The small GTPase Cdc42 is a major regulator of neutrophil effector functions. *Front Immunol*. 2020;11:1197.
71. van Bruggen R, Anthony E, Fernandez-Borja M, Roos D. Continuous translocation of Rac2 and the NADPH oxidase component p67 phox during phagocytosis. *J Biol Chem*. 2004;279(10):9097-9102.
72. Barger SR, Gauthier NC, Krendel M. Squeezing in a meal: myosin functions in phagocytosis. *Trends Cell Biol*. 2020;30(2):157-167.
73. Mansfield PJ, Shayman JA, Boxer LA. Regulation of polymorphonuclear leukocyte phagocytosis by myosin light chain kinase after activation of mitogen-activated protein kinase. *Blood*. 2000;95(7):2407-2412.
74. Patel JC, Hall A, Caron E. Vav regulates activation of Rac but not Cdc42 during Fc $\gamma$ R-mediated phagocytosis. *Mol Biol Cell*. 2002;13(4):1215-1226.
75. Martinez Gakidis MA, Cullere X, Olson T, et al. Vav GEFs are required for  $\beta 2$  integrin-dependent functions of neutrophils. *J Cell Biol*. 2004;166(2):273-282.
76. Chun XS, Magalhães MAO, Glogauer M. Rac1 And Rac2 differentially regulate Actin free barbed end formation downstream of the fMLP receptor. *J Cell Biol*. 2007;179(2):239-245.
77. Olazabal IM, Caron E, May RC, Schilling K, Knecht DA, Machesky LM. Rho-kinase and myosin-II control phagocytic cup formation during CR, but not Fc $\gamma$ R. *Phagocytosis Curr Biol*. 2002;12(16):1413-1418.
78. Kuijpers TW, Tool ATJ, van der Bijl I, et al. Combined immunodeficiency with severe inflammation and allergy caused by ARPC1B deficiency. *J Allergy Clin Immunol*. 2017;140(1):273-277.e10.
79. Lewkowicz E, Herit F, le Clainche C, Bourdoncle P, Perez F, Niedergang F. The microtubule-binding protein CLIP-170 coordinates mDia1 and Actin reorganization during CR3-mediated phagocytosis. *J Cell Biol*. 2008;183(7):1287-1298.
80. Colucci-Guyon E, Niedergang F, Wallar BJ, Peng J, Alberts AS, Chavrier P. A role for mammalian diaphanous-related Formins in complement receptor (CR3)-mediated phagocytosis in macrophages. *Curr Biol*. 2005;15(22):2007-2012.
81. Morris AC, Brittan M, Wilkinson TS, et al. C5a-mediated neutrophil dysfunction is RhoA-dependent and predicts infection in critically ill patients. *Blood*. 2011;117(19):5178-5188.
82. Schymeinsky J, Gerstl R, Mannigel I, et al. A fundamental role of mAbp1 in neutrophils: impact on  $\beta 2$  integrin-mediated phagocytosis and adhesion in vivo. *Blood*. 2009;114(19):4209-4220.
83. Yan M, di Ciano-Oliveira C, Grinstein S, Trimble WS. Coronin function is required for chemotaxis and phagocytosis in human neutrophils. *J Immunol*. 2007;178(9):5769-5778.
84. Roth H, Samereier M, Begandt D, et al. Filamin a promotes efficient migration and phagocytosis of neutrophil-like HL-60 cells. *Eur J Cell Biol*. 2017;96(6):553-566.
85. Buriez O, Labbé E. Disclosing the redox metabolism of drugs: the essential role of electrochemistry. *Curr Opin Electrochem*. 2020;24:63-68.
86. DeLeo FR, Allen LAA, Apicella M, Nauseef WM. NADPH oxidase activation and assembly during phagocytosis. *J Immunol*. 1999;163(12):6732-6740.
87. Zeng MY, Miralda I, Armstrong CL, Uriarte SM, Bagaitkar J. The roles of NADPH oxidase in modulating neutrophil effector responses. *Mol Oral Microbiol*. 2019;34(2):27-38.
88. Lam GY, Huang J, Brumell JH. The many roles of NOX2 NADPH oxidase-derived ROS in immunity. *Semin Immunopathol*. 2010;32(4):415-430.
89. Groemping Y, Rittinger K. Activation and assembly of the NADPH oxidase: a structural perspective. *Biochem J*. 2005;386:401-416.
90. Bokoch GM, Gilman AG. Inhibition of receptor-mediated release of arachidonic acid by pertussis toxin. *Cell*. 1984;39:301-308.
91. Sklar LA, Hyslop PA, Oades ZG, et al. Signal transduction and ligand-receptor dynamics in the human neutrophil. Transient responses and occupancy-response relations at the formyl peptide receptor. *J Biol Chem*. 1985;260(21):11461-11467.
92. Nguyen GT, Green ER, Meccas J. Neutrophils to the ROScues: mechanisms of NADPH oxidase activation and bacterial resistance. *Front Cell Infect Microbiol*. 2017;7:373.
93. Guthrie LA, McPhail LC, Henson PM, Johnston RB. Priming of neutrophils for enhanced release of oxygen metabolites by bacterial lipopolysaccharide. Evidence for increased activity of the superoxide-producing enzyme. *J Exp Med*. 1984;160(6):1656-1671.
94. Steinberg SE, BD GYNI, Islam MF, Prj HQ. Biochemical aspects of phagocytosis. *Nature*. 1961;192:535-541.
95. McCord J, Fridovich I. The utility of superoxide dismutase in studying free radical reactions. *J Biol Chem*. 1970;245(6):1374-1377.
96. Babior BM, Curnutte JT, Kipnes RS. Biological defense mechanisms. Evidence for the participation of superoxide in bacterial killing by xanthine oxidase. *J Lab Clin Med*. 1975;85(2):235-244.
97. Rosen H, Klebanoff SJ. Bactericidal activity of a superoxide anion-generating system. A model for the polymorphonuclear leukocyte. *J Exp Med*. 1979;149(1):27-39.
98. Rada B, Hably C, Meczner A, et al. Role of Nox2 in elimination of microorganisms. *Semin Immunopathol*. 2008;30(3):237-253.

99. Kettle AJ, Anderson RF, Hampton MB, Winterbourn CC. Reactions of superoxide with myeloperoxidase. *Biochemistry*. 2007;46(16):4888-4897.
100. Rosen H, Klebanoff SJ. Formation of singlet oxygen by the myeloperoxidase-mediated antimicrobial system. *J Biol Chem*. 1977;252(14):4803-4810.
101. Allen R, Stjernholm R, Steele R. Evidence for the generation of an electronic excitation state (s) in human polymorphonuclear leukocytes and its participation in bactericidal activity. *Biochem Biophys Res Commun*. 1972;47(4):679-684.
102. Steinbeck MJ, Khan AU, Karnovsky MJ. Intracellular singlet oxygen generation by phagocytosing neutrophils in response to particles coated with a chemical trap. *J Biol Chem*. 1992;267(19):13425-13433.
103. Yamashita K, Miyoshi T, Arai T, et al. Ozone production by amino acids contributes to killing of bacteria. *Proc Natl Acad Sci USA*. 2008;105(44):16912-16917.
104. Babior BM, Takeuchi C, Ruedi J, Gutierrez A, Wentworth P. Investigating antibody-catalyzed ozone generation by human neutrophils. *Proc Natl Acad Sci USA*. 2003;100(6):3031-3034.
105. Kettle AJ, Winterbourn CC. Do neutrophils produce ozone? An appraisal of current evidence. *Biofactors*. 2005;24(1-4):41-45.
106. Clifford DP, Repine JE. Hydrogen peroxide mediated killing of bacteria. *Mol Cell Biochem*. 1982;49(3):143-149.
107. Segal AW. How neutrophils kill microbes. *Annu Rev Immunol*. 2005;23:197-223.
108. Wang L, Taneja R, Razavi HM, Law C, Gillis C, Mehta S. Specific role of neutrophil inducible nitric oxide synthase in murine sepsis-induced lung injury in vivo. *Shock*. 2012;37(5):539-547.
109. Elks PM, Brizee S, van der Vaart M, et al. Hypoxia inducible factor signaling modulates susceptibility to mycobacterial infection via a nitric oxide dependent mechanism. *PLoS Pathog*. 2013;9(12):1-16.
110. Klebanoff SJ. Iodination of bacteria: a bactericidal mechanism. *J Exp Med*. 1967;126(6):1063-1078.
111. Klebanoff SJ. Myeloperoxidase-Halide-Hydrogen Peroxide Antibacterial System. *J Bacteriol*. 1968;95(6):2131-2138.
112. Jiang Q, Griffin DA, Barofsky DF, Hurst JK. Intraphagosomal chlorination dynamics and yields determined using unique fluorescent bacterial mimics. *Chem Res Toxicol*. 1997;10(10):1080-1089.
113. Hampton MB, Vissers MCM, Keenan JI, Winterbourn CC. Oxidant-mediated phosphatidylserine exposure and macrophage uptake of activated neutrophils: possible impairment in chronic granulomatous disease. *J Leukoc Biol*. 2002;71(5):775-781.
114. Klebanoff SJ. Oxygen metabolites from phagocytes. In: Gallin JI, Syndermann R, eds. *Inflammation: Basic Principles and Clinical Correlates*. Lippincott Williams and Wilkins; 1992:541-588.
115. Rosen H, Crowley JR, Heinecke JW. Human neutrophils use the myeloperoxidase-hydrogen peroxide-chloride system to chlorinate but not nitrate bacterial proteins during phagocytosis. *J Biol Chem*. 2002;277(34):30463-30468.
116. Reiss M, Roos D. Differences in oxygen metabolism of phagocytosing monocytes and neutrophils. *J Clin Invest*. 1978;61(2):480-488.
117. Fossati G, Moulding DA, Spiller DG, Moots RJ, White MRH, Edwards SW. The mitochondrial network of human neutrophils: role in chemotaxis, phagocytosis, respiratory burst activation, and commitment to apoptosis. *J Immunol*. 2003;170(4):1964-1972.
118. Sbarra AJ, Karnovsky ML. The biochemical basis of phagocytosis. *J Biol Chem*. 1959;234(6):1355-1362.
119. Reed PW, Tepperman J. Phagocytosis-associated metabolism and enzymes in the rat polymorphonuclear leukocyte. *Am J Physiol*. 1969;216(2):223-230.
120. Borregaard N, Herlin T. Energy metabolism of human neutrophils during phagocytosis. *J Clin Invest*. 1982;70(3):550-557.
121. Boxer LA, Baehner RL, Davis J. The effect of 2-deoxyglucose on Guinea pig polymorphonuclear leukocyte phagocytosis. *J Cell Physiol*. 1977;91(1):89-102.
122. Pan T, Sun S, Chen Y, et al. Immune effects of PI3K/Akt/HIF-1 $\alpha$ -regulated glycolysis in polymorphonuclear neutrophils during sepsis. *Crit Care*. 2022;26(1):1-17.
123. Yuk YC, So YK, Wai HY, et al. Impaired neutrophil activity and increased susceptibility to bacterial infection in mice lacking glucose-6-phosphatase- $\beta$ . *J Clin Invest*. 2007;117(3):784-793.
124. Sadiku P, Willson JA, Ryan EM, et al. Neutrophils fuel effective immune responses through gluconeogenesis and glycogenesis. *Cell Metab*. 2021;33(2):411-423.e4.
125. DeCoursey TE, Morgan D, Chervy V v. The voltage dependence of NADPH oxidase reveals why phagocytes need proton channels. *Nature*. 2003;422(6931):531-534.
126. Jankowski A, Scott CC, Grinstein S. Determinants of the phagosomal pH in neutrophils. *J Biol Chem*. 2002;277(8):6059-6066.
127. Hurst JK, Albrich JM, Green TR, Rosen H, Klebanoff S. Myeloperoxidase-dependent fluorescein chlorination by stimulated neutrophils. *J Biol Chem*. 1984;259(8):4812-4821.
128. Murphy R, DeCoursey TE. Charge compensation during the phagocyte respiratory burst. *Biochim Biophys Acta*. 2006;1757(8):996-1011.
129. Henderson LM, Chappell JB, Jones OTG. The superoxide-generating NADPH oxidase of human neutrophils is electrogenic and associated with an H<sup>+</sup> channel. *Biochem J*. 1987;246(2):325-329.
130. Henderson LM, Meech RW. Evidence that the product of the human X-linked Cgd gene, Gp91-phox, is a voltage-gated H<sup>+</sup> pathway. *J Gen Physiol*. 1999;114(6):771-786.
131. Bánfi B, Schrenzel J, Nüsse O, et al. A novel H<sup>+</sup> conductance in eosinophils: unique characteristics and absence in chronic granulomatous disease. *J Exp Med*. 1999;190(2):183-194.
132. Ramsey IS, Moran MM, Chong JA, Clapham DE. A voltage-gated proton-selective channel lacking the pore domain. *Nature*. 2006;440(7088):1213-1216.
133. Sasaki M, Takagi M, Okamura Y. A voltage sensor-domain protein is a voltage-gated proton channel. *Science*. 2006;312(5773):589-592.
134. Petheo GL, Orient A, Baráth M, et al. Molecular and functional characterization of Hv1 proton channel in human granulocytes. *PLoS One*. 2010;5(11):e14081.
135. Okochi Y, Sasaki M, Iwasaki H, Okamura Y. Voltage-gated proton channel is expressed on phagosomes. *Biochem Biophys Res Commun*. 2009;382(2):274-279.
136. Reeves EP, Lu H, Jacobs HL, et al. Killing activity of neutrophils is mediated through activation of proteases by K<sup>+</sup> flux. *Nature*. 2002;416(6878):291-297.
137. Schwingshackl A, Moqbel R, Duszyk M. Involvement of ion channels in human eosinophil respiratory burst. *J Allergy Clin Immunol*. 2000;106(2):272-279.
138. Winterbourn CC, Hampton MB, Livesey JH, Kettle AJ. Modeling the reactions of superoxide and myeloperoxidase in the neutrophil phagosome: implications for microbial killing. *J Biol Chem*. 2006;281(52):39860-39869.
139. Painter RG, Wang G. Direct measurement of free chloride concentrations in the phagolysosomes of human neutrophils. *Anal Chem*. 2006;78(9):3133-3137.
140. Painter RG, Valentine VG, Lanson NA, et al. CFTR expression in human neutrophils and the phagolysosomal chlorination defect in cystic fibrosis. *Biochemistry*. 2006;45(34):10260-10269.
141. Moreland JG, Davis AP, Bailey G, Nauseef WM, Lamb FS. Anion channels, including CIC-3, are required for normal neutrophil oxidative function, phagocytosis, and transendothelial migration. *J Biol Chem*. 2006;281(18):12277-12288.

142. Scheel O, Zdebek AA, Lourdel S, Jentsch TJ. Voltage-dependent electrogenic chloride/proton exchange by endosomal CLC proteins. *Nature*. 2005;436(7049):424-427.
143. Picollo A, Pusch M. Chloride/proton antiporter activity of mammalian CLC proteins CIC-4 and CIC-5. *Nature*. 2005;436(7049):420-423.
144. Dickerhof N, Isles V, Pattermore P, Hampton MB, Kettle AJ. Exposure of *Pseudomonas aeruginosa* to bactericidal hypochlorous acid during neutrophil phagocytosis is compromised in cystic fibrosis. *J Biol Chem*. 2019;294(36):13502-13514.
145. Riordan JR, Rommens JM, Kerem B s, et al. Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. *Science* (1979). 1989;245(4922):1066-1073.
146. Zhou Y, Song K, Painter RG, et al. Cystic fibrosis transmembrane conductance regulator recruitment to phagosomes in neutrophils. *J Innate Immun*. 2013;5(3):219-230.
147. Wang G, Nauseef WM. Neutrophil dysfunction in the pathogenesis of cystic fibrosis. *Blood*. 2022;139(17):2622-2631.
148. Kuhns DB, Alvord WG, Heller T, et al. Residual NADPH oxidase and survival in chronic granulomatous disease. *N Engl J Med*. 2010;363(27):2600-2610.
149. Baehner RL, Nathan DG. Leukocyte oxidase: defective activity in chronic granulomatous disease. *Science*. 1967;155(3764):835-836.
150. Quie PG, White JG, Holmes B, Good RA. In vitro bactericidal capacity of human polymorphonuclear leukocytes: diminished activity in chronic granulomatous disease of childhood. *J Clin Invest*. 1967;46(4):668-679.
151. Holland SM. Chronic granulomatous disease. *Hematol Oncol Clin North Am*. 2013;27(1):89-99.
152. Hampton MB, Kettle AJ, Winterbourn CC. Involvement of superoxide and myeloperoxidase in oxygen-dependent killing of *Staphylococcus aureus* by neutrophils. *Infect Immun*. 1996;64(9):3512-3517.
153. Lehrer RI, Hanifin J, Cline MJ. Defective bactericidal activity in myeloperoxidase-deficient human neutrophils. *Nature*. 1969;223(5201):78-79.
154. Nauseef WM, Metcalf JA, Root RK. Role of myeloperoxidase in the respiratory burst of human neutrophils. *Blood*. 1983;61(3):483-492.
155. Lehrer R, Cline MJ. Leukocyte myeloperoxidase deficiency and disseminated candidiasis: the role of myeloperoxidase in resistance to *Candida* infection. *J Clin Invest*. 1969;48(8):1478-1488.
156. Ng HP, Zhou Y, Song K, Hodges CA, Drumm ML, Wang G. Neutrophil-mediated phagocytic host defense defect in myeloid Cftr-inactivated mice. *PLoS One*. 2014;9(9):e106813.
157. Painter RG, Marrero L, Lombard GA, Valentine VG, Nauseef WM, Wang G. CFTR-mediated halide transport in phagosomes of human neutrophils. *J Leukoc Biol*. 2010;87(5):933-942.
158. Donini M, Fontana S, Savoldi G, et al. G-CSF treatment of severe congenital neutropenia reverses neutropenia but does not correct the underlying functional deficiency of the neutrophil in defending against microorganisms. *Blood*. 2007;109(11):4716-4723.
159. Horwitz MS, Laurino MY, Keel SB. Normal peripheral blood neutrophil numbers accompanying ELANE whole gene deletion mutation. *Blood Adv*. 2019;3(16):2470-2473.
160. Sabo P, Makaryan V, Dicken Y, et al. Mutant allele knockout with novel CRISPR nuclease promotes myelopoiesis in ELANE neutropenia. *Mol Ther Methods Clin Dev*. 2022;26:119-131.
161. Horwitz MS, Corey SJ, Grimes HL, Tidwell T. ELANE mutations in cyclic and severe congenital neutropenia. *Hematol Oncol Clin North Am*. 2013;27:19-41.
162. Sun YT, Shieh CC, Delpire E, Shen R, Shen R. K<sup>+</sup>-Cl<sup>-</sup> cotransport mediates the bactericidal activity of neutrophils by regulating NADPH oxidase activation. *J Physiol*. 2012;590(14):3231-3243.
163. Stossel TP. Quantitative studies of phagocytosis: kinetic effects of cations and heat-labile opsonin. *J Cell Biol*. 1973;58(2):346-356.
164. Jensen MS, Bainton DF. Temporal changes in pH within the phagocytic vacuole of the polymorphonuclear neutrophilic leukocyte. *J Cell Biol*. 1973;56(2):379-388.
165. Segal AW, Geisow M, Garcia R, Harper A, Miller R. The respiratory burst of phagocytic cells is associated with a rise in vacuolar pH. *Nature*. 1981;290(5805):406-409.
166. Cech P, Lehrer RI. Phagolysosomal pH of human neutrophils. *Blood*. 1984;63(1):88-95.
167. Odeberg H, Olsson I. Microbicidal mechanisms of human granulocytes: synergistic effects of granulocyte elastase and myeloperoxidase or chymotrypsin like cationic protein. *Infect Immun*. 1976;14(6):1276-1283.
168. Odeberg H, Olsson I. Antibacterial activity of cationic proteins from human granulocytes. *J Clin Invest*. 1975;56(5):1118-1124.
169. el Chemaly A, Nunes P, Jimaja W, Castelbou C, Demaurex N. Hv1 proton channels differentially regulate the pH of neutrophil and macrophage phagosomes by sustaining the production of phagosomal ROS that inhibit the delivery of vacuolar ATPases. *J Leukoc Biol*. 2014;95(5):827-839.
170. Morgan D, Capasso M, Musset B, Cherny V. v., Ríos E, dyer MJS, et al. voltage-gated proton channels maintain pH in human neutrophils during phagocytosis. *Proc Natl Acad Sci USA*. 2009;106(42):18022-18027.
171. Esmann L, Idel C, Sarkar A, et al. Phagocytosis of apoptotic cells by neutrophil granulocytes: diminished Proinflammatory neutrophil functions in the presence of apoptotic cells. *J Immunol*. 2010;184(1):391-400.
172. Hampton MB, Kettle AJ, Winterbourn CC. Inside the neutrophil phagosome: oxidants, myeloperoxidase, and bacterial killing. *Blood*. 1998;92:3007-3017.
173. Sengelov H, Follin P, Kjeldsen L, Lollike K, Dahlgren C, Borregaard N. Mobilization of granules and secretory vesicles during in vivo exudation of human neutrophils. *J Immunol*. 1995;154:4157-4165.
174. Huizinga TW, Dolman KM, van der Linden NJ, et al. Phosphatidylinositol-linked FcR $\text{III}$  mediates exocytosis of neutrophil granule proteins, but does not mediate initiation of the respiratory burst. *J Immunol*. 1990;144:1432-1437.
175. Ito N, Yokomizo T, Sasaki T, et al. Requirement of phosphatidylinositol 3-kinase activation and calcium influx for leukotriene B $_4$ -induced enzyme release. *J Biol Chem*. 2002;277:44898-44904.
176. Borregaard N, Cowland JB. Granules of the human neutrophilic Polymorphonuclear leukocyte. *Blood*. 1997;89(10):3503-3521.
177. Bainton DF, Farquhar MG. Origin of granules in polymorphonuclear leukocytes. Two types derived from opposite faces of the Golgi complex in developing granulocytes. *J Cell Biol*. 1966;28(2):277-301.
178. Borregaard N, Sehested M, Nielsen BS, Sengeløv H, Kjeldsen L. Biosynthesis of granule proteins in normal human bone marrow cells. Gelatinase is a marker of terminal neutrophil differentiation. *Blood*. 1995;85:812-817.
179. Fontana S, Parolini S, Vermi W, et al. Innate immunity defects in Hermansky-Pudlak type 2 syndrome. *Blood*. 2006;107:4857-4864.
180. Faurschou M, Borregaard N. Neutrophil granules and secretory vesicles in inflammation. *Microbes Infect*. 2003;5(14):1317-1327.
181. Lehrer RI, Lichtenstein AK, Ganz T. Defensins: antimicrobial and cytotoxic peptides of mammalian cells. *Annu Rev Immunol*. 1993;11:105-128.
182. Flannagan RS, Cosío G, Grinstein S. Antimicrobial mechanisms of phagocytes and bacterial evasion strategies. *Nat Rev Microbiol*. 2009;7:355-366.
183. Jog NR, Rane MJ, Lominadze G, Luerman GC, Ward RA, McLeish KR. The Actin cytoskeleton regulates exocytosis of all neutrophil granule subsets. *Am J Physiol Cell Physiol*. 2007;292(5):C1690-C1700.
184. Epand RM. Membrane fusion. *Biosci Rep*. 2000;20(6):435-441.

185. Westman J, Grinstein S, Maxson ME. Revisiting the role of calcium in phagosome formation and maturation. *J Leukoc Biol.* 2019;106(4):837-851.
186. Jaconi MEE, Lew DP, Carpentier JL, Magnusson KE, Sjogren M, Stendahl O. Cytosolic free calcium elevation mediates the phagosome-lysosome fusion during phagocytosis in human neutrophils. *J Cell Biol.* 1990;110(5):1555-1564.
187. Nordenfelt P, Tapper H. The role of calcium in neutrophil granule-phagosome fusion. *Commun Integr Biol.* 2010;3(3):224-226.
188. Cowland JB, Borregaard N. Granulopoiesis and granules of human neutrophils. *Immunol Rev.* 2016;273(1):11-28.
189. Nordenfelt P, Winberg ME, Lönnbro P, Rasmusson B, Tapper H. Different requirements for early and late phases of Azurophilic granule-phagosome fusion. *Traffic.* 2009;10(12):1881-1893.
190. Downey GP, Chan CK, Trudel S, Grinstein S. Actin assembly in electroporated neutrophils: role of intracellular calcium. *J Cell Biol.* 1990;110(6):1975-1982.
191. Bengtsson T, Jaconi MEE, Gustafson M, et al. Actin dynamics in human neutrophils during adhesion and phagocytosis is controlled by changes in intracellular free calcium. *Eur J Cell Biol.* 1993;62(1):49-58.
192. Dewitt S, Hallett MB. Cytosolic free Ca<sup>2+</sup> changes and calpain activation are required for  $\beta$  integrin-accelerated phagocytosis by human neutrophils. *J Cell Biol.* 2002;159(1):181-189.
193. Vinet AF, Fukuda M, Descoteaux A. The exocytosis regulator Synaptotagmin V controls phagocytosis in macrophages. *J Immunol.* 2008;181(8):5289-5295.
194. Arango Duque G, Fukuda M, Descoteaux A. Synaptotagmin XI regulates phagocytosis and cytokine secretion in macrophages. *J Immunol.* 2013;190(4):1737-1745.
195. Ernst JD. Annexin III translocates to the periphagosomal region when neutrophils ingest opsonized yeast. *J Immunol.* 1991;146(9):3110-3114.
196. Majeed M, Perskvist N, Ernst JD, Orselius K, Stendahl O. Roles of calcium and annexins in phagocytosis and elimination of an attenuated strain of mycobacterium tuberculosis in human neutrophils. *Microb Pathog.* 1998;24(5):309-320.
197. Tapper H, Furuya W, Grinstein S. Localized exocytosis of primary (lysosomal) granules during phagocytosis: role of Ca<sup>2+</sup>-dependent tyrosine phosphorylation and microtubules. *J Immunol.* 2002;168(10):5287-5296.
198. Hoffstein S, Goldstein IM, Weissmann G. Role of microtubule assembly in lysosomal enzyme secretion from human polymorphonuclear leukocytes. A reevaluation. *J Cell Biol.* 1977;73(1):242-256.
199. Blocker A, Griffiths G, Olivo JC, Hyman AA, Severin FF. A role for microtubule dynamics in phagosome movement. *J Cell Sci.* 1998;111(3):303-312.
200. Rothwell SW, Deal CC, Pinto J, Wrigth DG. Affinity purification and subcellular localization of kinesin in human neutrophils. *J Leukoc Biol.* 1993;53(4):372-380.
201. Abdel-Latif D, Steward M, Macdonald DL, Francis GA, Dinauer MC, Lacy P. Rac2 is critical for neutrophil primary granule exocytosis. *Blood.* 2004;104:832-839.
202. Williams DA, Tao W, Yang F, et al. Dominant negative mutation of the hematopoietic-specific Rho GTPase, Rac2, is associated with a human phagocyte immunodeficiency. *Blood.* 2000;96(5):1646-1654.
203. Nauseef WM. Assembly of the phagocyte NADPH oxidase. *Histochem Cell Biol.* 2004;122(4):277-291.
204. Herrero-Turrión MJ, Calafat J, Janssen H, Fukuda M, Mollinedo F. Rab27a regulates exocytosis of tertiary and specific granules in human neutrophils. *J Immunol.* 2008;181:3793-3803.
205. Monfregola J, Johnson JL, Meijler MM, Napolitano G, Catz SD. MUNC13-4 protein regulates the oxidative response and is essential for phagosomal maturation and bacterial killing in neutrophils. *J Biol Chem.* 2012;287(53):44603-44618.
206. Munafó DB, Johnson JL, Ellis BA, Rutschmann S, Beutler B, Catz SD. Rab27a is a key component of the secretory machinery of azurophilic granules in granulocytes. *Biochem J.* 2007;402(2):229-239.
207. Elstak ED, Neeft M, Nehme NT, et al. The munc13-4-rab27 complex is specifically required for tethering secretory lysosomes at the plasma membrane. *Blood.* 2011;118(6):1570-1578.
208. Mollinedo F, Calafat J, Janssen H, et al. Combinatorial SNARE complexes modulate the secretion of cytoplasmic granules in human neutrophils. *J Immunol.* 2006;177(5):2831-2841.
209. Hirahashi J, Mekala D, van Ziffle J, et al. Mac-1 signaling via Src-family and Syk kinases results in elastase-dependent Thrombohemorrhagic vasculopathy. *Immunity.* 2006;25(2):271-283.
210. Möhn H, Le Cabec V, Fischer S, Maridonneau-Parini I. The src-family protein-tyrosine kinase p59hck is located on the secretory granules in human neutrophils and translocates towards the phagosome during cell activation. *Biochem J.* 1995;309:657-665.
211. Gutkind JS, Robbins KC. Translocation of the FGR protein-tyrosine kinase as a consequence of neutrophil activation. *Proc Natl Acad Sci USA.* 1989;86:8783-8787.
212. Catz SD, McLeish KR. Therapeutic targeting of neutrophil exocytosis. *J Leukoc Biol.* 2020;107(3):393-408.
213. Pincus SH, Klebanoff SJ. Quantitative leukocyte iodination. *N Engl J Med.* 1971;284(14):744-750.
214. Lehrer RI. Functional aspects of a second mechanism of candidacidal activity by human neutrophils. *J Clin Invest.* 1972;51(10):2566-2572.
215. Rosen H, Michel BR. Redundant contribution of myeloperoxidase-dependent systems to neutrophil-mediated killing of *Escherichia coli*. *Infect Immun.* 1997;65(10):543-3947.
216. Weiss J, Kao L, Victor M, Elsbach P. Oxygen-independent intracellular and oxygen-dependent extracellular killing of *Escherichia coli* S15 by human polymorphonuclear leukocytes. *J Clin Invest.* 1985;76(1):206-212.
217. Segal AW, Garcia RC, Harper AM, Banga JP. Iodination by stimulated human neutrophils. Studies on its stoichiometry, subcellular localization and relevance to microbial killing. *Biochem J.* 1983;210(1):215-225.
218. Reeves EP, Nagl M, Godovac-Zimmermann J, Segal AW. Reassessment of the microbicidal activity of reactive oxygen species and hypochlorous acid with reference to the phagocytic vacuole of the neutrophil granulocyte. *J Med Microbiol.* 2003;52(8):643-651.
219. Rosen H, Klebanoff SJ, Wang Y, Brot N, Heinecke JW, Fu X. Methionine oxidation contributes to bacterial killing by the myeloperoxidase system of neutrophils. *Proc Natl Acad Sci U S A.* 2009;106(4):18686-18691.
220. Staudinger BJ, Oberdoerster MA, Lewis PJ, Rosen H. mRNA expression profiles for *Escherichia coli* ingested by normal and phagocyte oxidase-deficient human neutrophils. *J Clin Invest.* 2002;110(8):1151-1163.
221. Aratani Y, Kura F, Watanabe H, et al. Critical role of myeloperoxidase and nicotinamide adenine dinucleotide phosphate-oxidase in high-burden systemic infection of mice with *Candida albicans*. *J Infect Dis.* 2002;185(12):1833-1837.
222. Locksley R, Wilson C, Klebanoff S. Increased respiratory burst in myeloperoxidase-deficient monocytes. *Blood.* 1983;62(4):902-909.
223. Nauseef WM. Myeloperoxidase Deficiency. *Hematol Oncol Clin North Am.* 1988;2(1):135-158.
224. Klebanoff SJ, Kettle AJ, Rosen H, Winterbourn CC, Nauseef WM. Myeloperoxidase: a front-line defender against phagocytosed microorganisms. *J Leukoc Biol.* 2013;93(2):185-198.
225. Klebanoff SJ. Myeloperoxidase: contribution to the microbicidal activity of intact leukocytes. *Science.* 1970;169(3950):1095-1097.

226. Branzk N, Lubojemska A, Hardison SE, et al. Neutrophils sense microbial size and selectively release neutrophil extracellular traps in response to large pathogens. *Nat Immunol*. 2014;15(11):1017-1025.
227. Brinkmann V, Reichard U, Goosmann C, et al. Neutrophil extracellular traps kill bacteria. *Science (1979)*. 2004;303(5663):1532-1535.
228. Metzler KD, Goosmann C, Lubojemska A, Zychlinsky A, Papayannopoulos V. A myeloperoxidase-containing complex regulates neutrophil elastase release and Actin dynamics during NETosis. *Cell Rep*. 2014;8(3):883-896.
229. Johnson CM, O'Brien XM, Byrd AS, et al. Integrin cross-talk regulates the human neutrophil response to fungal  $\beta$ -Glucan in the context of the extracellular matrix: a prominent role for VLA3 in the antifungal response. *J Immunol*. 2016;198(1):318-334.
230. Yipp BG, Petri B, Salina D, et al. Dynamic NETosis is carried out by live neutrophils in human and mouse bacterial abscesses and during severe gram-positive infection. *Nat Med*. 2012;18(9):1386-1393.
231. Zent C, Elliott MR. Maxed out macs: physiologic cell clearance as a function of macrophage phagocytic capacity. *FEBS J*. 2017;284(7):1021-1039.
232. Mantegazza AR, Zajac AL, Twelvetrees A, Holzbaur ELF, Amigorena S, Marks MS. TLR-dependent phagosome tubulation in dendritic cells promotes phagosome cross-talk to optimize MHC-II antigen presentation. *Proc Natl Acad Sci USA*. 2014;111(43):15508-15513.
233. Stewart JR, Weisman RA. Exocytosis of latex beads during the encystment of *Acanthamoeba*. *J Cell Biol*. 1972;52(1):117-130.
234. Krajcovic M, Krishna S, Akkari L, Joyce JA, Overholtzer M. MTOR regulates phagosome and entotic vacuole fission. *Mol Biol Cell*. 2013;24(23):3736-3745.
235. Levin-Konigsberg R, Montaño-Rendón F, Keren-Kaplan T, et al. Phagolysosome resolution requires contacts with the ER and phosphatidylinositol 4-phosphate signaling. *Nat Cell Biol*. 2019;21(10):1234-1247.
236. Lancaster CE, Fountain A, Dayam RM, et al. Phagosome resolution regenerates lysosomes and maintains the degradative capacity in phagocytes. *J Cell Biol*. 2021;220(9):e202005072.
237. Freeman SA, Grinstein S. Resolution of macropinosomes, phagosomes and autolysosomes: osmotically driven shrinkage enables tubulation and vesiculation. *Traffic*. 2018;19(12):965-974.
238. Vethanayagam RR, Almyroudis NG, Grimm MJ, et al. Role of NADPH oxidase versus neutrophil proteases in antimicrobial host defense. *PLoS One*. 2011;6(12):28149.
239. MacIvor DM, Shapiro SD, Pham CTN, Belaouaj A, Abraham SN, Ley TJ. Normal neutrophil function in Cathepsin G-deficient mice. *Blood*. 1999;94(12):4282-4293.
240. McGovern NN, Cowburn AS, Porter L, et al. Hypoxia selectively inhibits respiratory burst activity and killing of *Staphylococcus aureus* in human neutrophils. *J Immunol*. 2010;186(1):453-463.
241. Tejedo CG. (2020) Complement C5a impairs phagosomal maturation in the neutrophil through phosphoproteomic remodelling. PRIDE Archive. Available from <https://www.ebi.ac.uk/pride/archive?keyword=PXDD017092>

**How to cite this article:** Naish E, Wood AJ, Stewart AP, et al. The formation and function of the neutrophil phagosome. *Immunol Rev*. 2022;00:1-23. doi:[10.1111/imr.13173](https://doi.org/10.1111/imr.13173)