

Northumbria Research Link

Citation: Raza, Zahid, Akhter, Shehnaz and Shang, Yilun (2023) Expected value of first Zagreb connection index in random cyclooctatetraene chain, random polyphenyls chain, and random chain network. *Frontiers in Chemistry*, 10. p. 1067874. ISSN 2296-2646

Published by: Frontiers

URL: <https://doi.org/10.3389/fchem.2022.1067874>
<<https://doi.org/10.3389/fchem.2022.1067874>>

This version was downloaded from Northumbria Research Link:
<https://nrl.northumbria.ac.uk/id/eprint/51205/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



OPEN ACCESS

EDITED BY

Anakuthil Anoop,
Indian Institute of Technology
Kharagpur, India

REVIEWED BY

Parvez Ali,
Qassim University, Saudi Arabia
Rogers Mathew,
Indian Institute of Technology
Hyderabad, India

*CORRESPONDENCE

Zahid Raza,
✉ zraza@sharjah.ac.ae
Yilun Shang,
✉ yilun.shang@northumbria.ac.uk

SPECIALTY SECTION

This article was submitted to Theoretical
and Computational Chemistry,
a section of the journal
Frontiers in Chemistry

RECEIVED 12 October 2022

ACCEPTED 07 December 2022

PUBLISHED 04 January 2023

CITATION

Raza Z, Akhter S and Shang Y (2023),
Expected value of first Zagreb
connection index in random
cyclooctatetraene chain, random
polyphenyls chain, and random
chain network.
Front. Chem. 10:1067874.
doi: 10.3389/fchem.2022.1067874

COPYRIGHT

© 2023 Raza, Akhter and Shang. This is
an open-access article distributed
under the terms of the [Creative
Commons Attribution License \(CC BY\)](#).
The use, distribution or reproduction in
other forums is permitted, provided the
original author(s) and the copyright
owner(s) are credited and that the
original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution
or reproduction is permitted which does
not comply with these terms.

Expected value of first Zagreb connection index in random cyclooctatetraene chain, random polyphenyls chain, and random chain network

Zahid Raza^{1*}, Shehnaz Akhter² and Yilun Shang^{3*}

¹Department of Mathematics, College of Sciences, University of Sharjah, Sharjah, United Arab Emirates,

²School of Natural Sciences, National University of Sciences and Technology, Islamabad, Pakistan,

³Department of Computer and Information Sciences, Northumbria University, Newcastle upon Tyne, United Kingdom

The Zagreb connection indices are the known topological descriptors of the graphs that are constructed from the connection cardinality (degree of given nodes lying at a distance 2) presented in 1972 to determine the total electron energy of the alternate hydrocarbons. For a long time, these connection indices did not receive much research attention. Ali and Trinajstić [Mol. Inform. 37, Art. No. 1800008, 2018] examined the Zagreb connection indices and found that they compared to basic Zagreb indices and that they provide a finer value for the correlation coefficient for the 13 physico-chemical characteristics of the octane isomers. This article acquires the formulae of expected values of the first Zagreb connection index of a random cyclooctatetraene chain, a random polyphenyls chain, and a random chain network with l number of octagons, hexagons, and pentagons, respectively. The article presents extreme and average values of all the above random chains concerning a set of special chains, including the meta-chain, the ortho-chain, and the para-chain.

KEYWORDS

average value, expected value, random cyclooctatetraene chain, random polyphenyls chain, zagreb connection indices

1 Introduction

Graph theory is vital to various disciplines, including the chemical and biological sciences. One of the objectives of chemical graph theory is its primary and significant role in studying physico-chemical reactions and biological activities and pointing out the structural properties of molecular graphs, *etc.*, Topological descriptors have played a significant role in achieving the desired properties of molecular graphs. Topological descriptors are molecular structural invariants that theoretically and mathematically explain the connectivity characteristics of nano-materials and chemical compounds. Therefore, topological indices produce sharper approaches to measuring their behavior and characteristics.

For 20 years, hydrocarbons and their derivatives have received attention from researchers because these compounds only have two members, carbon and hydrogen. We can acquire various types of hydrocarbon derivatives by replacing their molecular hydrogen atoms with different types of other atomic groups. A large number of valuable hydrocarbons are available in plants and some valuable characteristics of hydrocarbons are important to chemical raw materials and fuel.

Throughout this article, the vertex and edge sets of a graph \mathcal{H} are represented as $V(\mathcal{H})$ and $E(\mathcal{H})$, respectively. We denote the degree of a vertex $v \in V(\mathcal{H})$ by $d_{\mathcal{H}}(v)$, which is defined as the cardinality of edges joined with v . Let the order and size of \mathcal{H} be n and m , respectively. The l -degree of a given vertex $v \in V(\mathcal{H})$, presented by $d_l(v)$, is the cardinality of set of vertices of $V(\mathcal{H})$ whose distance from v is l , where $d_1(v) = d_{\mathcal{H}}(v)$ and $d_2(v) = \tau_v$ [this is known as the connection number of v (Todeschini and Consonni, 2000)].

Suppose that \mathbb{Z} is a collection of all connected simple graphs. There is a function $\mathcal{P}: \mathbb{Z} \rightarrow \mathbb{R}^+$ that describes a topological invariant if for any two isomorphic members \mathcal{M}_1 and \mathcal{M}_2 of \mathbb{Z} , we have $\mathcal{P}(\mathcal{M}_1) = \mathcal{P}(\mathcal{M}_2)$. Thousands of degree and distance-related topological invariants have been proposed, but some are better known because of their high predictive power for many characteristics like density, boiling point, molecular weight, refractive index, etc., Topological invariants have so many implementations in numerous areas of sciences such as drug discovery, physico-chemical research, toxicology, biology, and chemistry. To date, topological indices are the most notable field of graphical research. For more discussion on numerous invariants, we refer readers to studies by (Gutman, 2013; Akhter et al., 2016; Akhter and Imran, 2016; Akhter et al., 2017; Akhter et al., 2018; Akhter et al., 2020).

The Zagreb indices are the most notable invariants, and they have many valuable applications in chemistry. In 1972 Gutman and Trinajstić (Gutman and Trinajstić, 1972) established the first vertex degree dependent Zagreb index of a graph \mathcal{H} . Two renowned Zagreb indices of a graph \mathcal{H} can be described in the following manner:

$$\begin{aligned} \mathcal{M}_1(\mathcal{H}) &= \sum_{x \in V(\mathcal{H})} d_{\mathcal{H}}^2(x), \\ \mathcal{M}_2(\mathcal{H}) &= \sum_{xy \in E(\mathcal{H})} d_{\mathcal{H}}(x)d_{\mathcal{H}}(y). \end{aligned}$$

Motivated by how influential they have become and the many important applications of primary Zagreb indices, Naji et al. (Soner and Naji, 2016; Gutman et al., 2017) presented the concept of Zagreb connection indices (leap Zagreb indices), constructed from the second degrees of the vertices of a graph \mathcal{H} . The first, second, and modified Zagreb connection indices of \mathcal{H} can be defined as:

$$\begin{aligned} ZC_1(\mathcal{H}) &= \sum_{y \in V(\mathcal{H})} \tau_y^2, \\ ZC_2(\mathcal{H}) &= \sum_{xy \in E(\mathcal{H})} \tau_x \tau_y, \\ ZC_1^*(\mathcal{H}) &= \sum_{x \in V(\mathcal{H})} d_{\mathcal{H}}(x) \tau_x. \end{aligned}$$

The chemical applications of ZC_1 were presented in (8), indicating that the given index has a wide co-relation with the physical characteristics of chemical compounds, for instance, boiling point, enthalpy of evaporation, entropy, acentric factor, and standard enthalpy of vaporization. Let f_l present the cardinality of the subset of vertices of \mathcal{H} with connection number l . The next formula for the first Zagreb connection index is equal to the above definition.

$$ZC_1(\mathcal{H}) = \sum_{0 \leq l \leq n-2} f_l(G)l^2. \quad (1.1)$$


Naji and Soner (2018), (Gutman et al., 2017) determined the leap Zagreb descriptors of some graph operations and families. Leap Zagreb indices are presented in a recently published survey (Gutman et al., 2020). In (39), the authors establish sharp bounds for the leap Zagreb indices of trees and unicyclic graphs and also determined the corresponding extremal graphs. For more studies on Zagreb connection indices, we refer the readers to (Ducoffe et al., 2018a; Ali and Trinajstić, 2018; Shao et al., 2018a; Basavanagoud and Chitra, 2018; Ducoffe et al., 2018b; Khalid et al., 2018; Manzoor et al., 2018; Du et al., 2019; Fatima et al., 2019; Tang et al., 2019; Ye et al., 2019; Raza, 2020a; Bao et al., 2020; Raza, 2020b; Cao et al., 2020; Naji et al., 2020; Raza, 2022).

Huang et al. (2014) determined the expected values for Kirchhoff indices of random polyphenyl and spiro chains. Ma et al. (2017), Yang and Zhang. (2012), and Qi et al. (2022) independently acquired the expected value of Wiener indices of random polyphenyl chain and random spiro chain. Zhang et al. (2020) have provided expected values of the Schultz, Gutman, multiplicative degree-Kirchhoff, and additive degree-Kirchhoff indices of random polyphenylene chains. Raza and Imran. (2021) obtained expected values of modified second Zagreb, symmetric difference, inverse symmetric, and augmented Zagreb indices in random cyclooctane chains. Zhang et al. (2021) established the formulae for expected values of Sombor indices of a general random chain. Recently, many studies have explored the expected values of different topological indices. For further information, we refer readers to the following studies (Raza, 2020b; Fang et al., 2021; Raza, 2021; Jahanbanni, 2022; Raza et al., 2022).

Motivated by the above research, the present study determined the explicit formulae for expected values of the first Zagreb connection index of the random cyclooctatetraene chain, random polyphenyls chain, and random chain network with l octagons, hexagons, and pentagons, respectively. Moreover, we examined the average and extreme values of the Zagreb connection index among all the above-mentioned random chains corresponding to their set.

2 The first Zagreb connection index of random cyclooctatetraene chain

Cyclooctatetraene, having chemical formula C_8H_8 , is an organic compound whose full name is '1, 3, 5, 7 - cyclooctene.



Its structure is a cyclic polyolefin-like benzene, but it is not aromatic, see (Willis et al., 1952; Mathews and Lipscomb, 1959; Traetteberg et al., 1970). It has the same chemical characteristics as unsaturated hydrocarbons and is easy to construct explosive organic peroxides, (Milas and Nolan Petrus, 1958; Donald and Whitehead, 1969; Garavelli et al., 2002; Schwamm et al., 2019).

Spiro compounds are valuable types of cycloalkanes in organic chemistry. A spiro union is a join of two rings that have a common atom between both rings, and a join of a direct union among the rings is known as a free spiro union in spiro compounds. In a cyclooctatylene chain, octagons are

joined by cut vertices or cut edges. A random cyclooctatetraene chain COC_l , has l octagons, and can be constructed by a cyclooctatetraene chain COC_{l-1} with $l-1$ octagons attached to a new octagon G_l by a bridge (see Figure 1).

The COC_l is a cyclooctatetraene chain with $l \geq 2$ having G_1, G_2, \dots, G_l octagons. The new octagon can be joined by four different schemes, which give the local orderings. We use these as $COC_l^1, COC_l^2, COC_l^3, COC_l^4$ (see Figure 2).


A random cyclooctatetraene chain $COC_l(k_1, k_2, k_3)$ is a cyclooctatetraene chain constructed by step-by-step attachment of new octagons. At every step $p = 2, 3, \dots, l$ a random choice is constructed from one of the four possible chains:

- 1 $COC_{p-1} \rightarrow COC_p^1$ with probability k_1 ,
- 2 $COC_{p-1} \rightarrow COC_p^2$ with probability k_2 ,
- 3 $COC_{p-1} \rightarrow COC_p^3$ with probability k_3 ,
- 4 $COC_{p-1} \rightarrow COC_p^4$ with probability $k_4 = 1 - k_1 - k_2 - k_3$,

Where all the given probabilities are constant. In this section, we will discuss the expected value for the first Zagreb connection index among random cyclooctatetraene chains with l octagons.

Theorem 2.1. For $l \geq 2$, the expected value for the first Zagreb connection index of random cyclooctatetraene chain COC_l is

$$E(ZC_1(COC_l)) = (8k_1 + 2k_2 + 76)l - 16k_1 - 4k_2 - 44.$$



Proof. **Case-I:** When $l = 2$, we get the result by direct calculations as:

$$E(ZC_1(COC_l)) = 10 \times (2)^2 + 4 \times (3)^2 + 2 \times (4)^2 = 108.$$

Case-II: When $l \geq 3$, it is obvious that $f_2(COC_l)$, $f_3(COC_l)$, $f_4(COC_l)$ and $f_5(COC_l)$ depends on the four possible cases as following:

1 If $COC_{l-1} \rightarrow COC_l^1$ with probability k_1 , we acquire

$$\begin{aligned} f_2(COC_l^1) &= f_2(COC_{l-1}) + 4, \\ f_3(COC_l^1) &= f_3(COC_{l-1}) + 2, \\ f_4(COC_l^1) &= f_4(COC_{l-1}) + 2, \\ f_5(COC_l^1) &= f_5(COC_{l-1}) + 2. \end{aligned}$$

By using the above values in Eq. 1.1, we get

$$\begin{aligned} ZC_1(COC_l^1) &= ZC_1(COC_{l-1}) + 4 \times 2^2 \\ &\quad + 2 \times 3^2 + 2 \times 5^2 \\ &= ZC_1(COC_{l-1}) + 84. \end{aligned}$$

2 If $COC_{l-1} \rightarrow COC_l^2$ with probability k_2 , we acquire

$$\begin{aligned} f_2(COC_l^2) &= f_2(COC_{l-1}) + 3, \\ f_3(COC_l^2) &= f_3(COC_{l-1}) + 2, \\ f_4(COC_l^2) &= f_4(COC_{l-1}) + 3, \end{aligned}$$

By using the above values in Eq. 1.1, we get

$$\begin{aligned} ZC_1(COC_l^2) &= ZC_1(COC_{l-1}) + 3 \times 2^2 + 2 \times 3^2 \\ &\quad + 3 \times 4^2 \\ &= ZC_1(COC_{l-1}) + 78. \end{aligned}$$

3 If $COC_{l-1} \rightarrow COC_l^3$ with probability k_3 , we acquire

$$\begin{aligned} f_2(COC_l^3) &= f_2(COC_{l-1}) + 2, \\ f_3(COC_l^3) &= f_3(COC_{l-1}) + 4, \\ f_4(COC_l^3) &= f_4(COC_{l-1}) + 2, \end{aligned}$$

By using the above values in Eq. 1.1, we get

$$\begin{aligned} ZC_1(COC_l^3) &= ZC_1(COC_{l-1}) + 2 \times 2^2 + 4 \times 3^2 \\ &\quad + 2 \times 4^2 \\ &= ZC_1(COC_{l-1}) + 76. \end{aligned}$$

4 If $COC_{l-1} \rightarrow COC_l^4$ with probability $1 - k_1 - k_2 - k_3$, we acquire

$$\begin{aligned} f_2(COC_l^4) &= f_2(COC_{l-1}) + 2, \\ f_3(COC_l^4) &= f_3(COC_{l-1}) + 4, \\ f_4(COC_l^4) &= f_4(COC_{l-1}) + 2, \end{aligned}$$

By using above the values in Eq. 1.1, we get

$$\begin{aligned} ZC_1(COC_l^4) &= ZC_1(COC_{l-1}) + 2 \times 2^2 + 4 \times 3^2 \\ &\quad + 2 \times 4^2 \\ &= ZC_1(COC_{l-1}) + 76. \end{aligned}$$

Now

$$\begin{aligned} E_l^i &= E(ZC_1(COC_l)) = k_1 ZC_1(COC_l^1) + k_2 ZC_1(COC_l^2) + k_3 ZC_1(COC_l^3) \\ &\quad + (1 - k_1 - k_2 - k_3) ZC_1(COC_l^4) \\ &= ZC_1(COC_{l-1}) + 8k_1 + 2k_2 + 76. \end{aligned} \quad (2.1)$$

Note that $E[E_l^i] = E_l^i$. By applying the expression operator to Eq. 2.1 and also $l \geq 3$, we get

$$E_l^i = E_{l-1}^i + 8k_1 + 2k_2 + 76. \quad (2.2)$$

The Eq. 2.2 is a first-order non-homogeneous linear difference result with constant coefficients. We easily see that the general solution of the homogeneous equation of Eq. 2.2 is $E^i = C$. Suppose $E^i = bl$ is a particular solution of Eq. 2.2, using E^i into Eq. 2.2, we acquire

$$b = 8k_1 + 2k_2 + 76.$$

Finally the general solution of Eq. 2.2 is

$$\begin{aligned} E_l^i &= E^i + E^{i'} \\ &= E(ZC_1(COC_l)) = (8k_1 + 2k_2 + 76)l + C. \end{aligned}$$

Applying the initial condition $l = 3$, we get the following

$$C = -16k_1 - 4k_2 - 44.$$

Therefore

$$\begin{aligned} E_l^i &= E(ZC_1(COC_l)) \\ &= (8k_1 + 2k_2 + 76)l - 16k_1 - 4k_2 - 44. \end{aligned}$$

If $k_1 = 1$ (respectively, $k_2 = 1$) and $k_2 = k_3 = k_4 = 0$ (respectively, $k_1 = k_3 = k_4 = 0$), then $COC_l = M_l$ (respectively, $COC_l = O_l^1$). Similarly, if $k_3 = 1$ (respectively, $k_4 = 1$) and $k_1 = k_2 = k_4 = 0$ (respectively, $k_1 = k_2 = k_3 = 0$), then $COC_l = Q_l^2$ (respectively $COC_l = L_l$). By Theorem 2.1, we can acquire the first Zagreb connection index of the cyclooctatetraene meta-chain M_l , ortho-chains O_l^1 , O_l^2 and para-chain L_l as:

$$\begin{aligned} ZC_1(M_l) &= 84l - 60, \quad ZC_1(O_l^1) = 78l - 48, \\ ZC_1(O_l^2) &= 76l - 44, \quad ZC_1(L_l) = 76l - 44. \end{aligned}$$

Corollary 2.2. For a random cyclooctatetraene chain $COC_l(l \geq 3)$, the para-chain L_l and ortho chain O_l^1 , and the meta-chain M_l achieves the minimum and the maximum of $E(ZC_1(COC_l))$, respectively.

Proof. Using Theorem 2.1, we acquire

$$E_l^i = E(ZC_1(COC_l)) = (8l - 16)k_1 + (2l - 4)k_2 + 76l - 44.$$

By taking partial derivatives, we acquire $\frac{\partial E}{\partial k_1} = 8l - 16 > 0$, $\frac{\partial E}{\partial k_2} = 2l - 4 > 0$. When $k_1 = k_2 = k_3 = 0$ (i.e. $k_4 = 1$), the para-chain L_l and ortho chain O_l^1 achieve the minimum of $E(ZC_1(COC_l))$, that is $COC_l \cong L_l$ or $COC_l \cong O_l^1$. If $k_3 = 1 - k_1 - k_2$ ($0 \leq k_1, k_2 \leq 1$), we have

$$E_l^i = E(ZC_1(COC_l)) \\ = (8l - 16)k_1 + (2l - 4)k_2 + 76l - 44.$$

But $k_1 = k_2 = 0$ (when $k_3 = 1$), $E(ZC_1(COC_l))$ can not attain the maximum value. If $k_1 = 1 - k_2$ ($0 \leq k_2 \leq 1$), we acquire

$$E_l^i = E(ZC_1(COC_l)) \\ = (8l - 16)(1 - k_2) + (2l - 4)k_2 + 76l - 44.$$

Therefore $\frac{\partial E}{\partial k_2} = -6l + 12 < 0$. Thus $E(ZC_1(COC_l))$ achieves the maximum value, if $k_2 = 0$ ($k_1 = 1$), that is $COC_l \cong M_l$.

3 The first Zagreb connection index of a random polyphenyl chain

Polyphenyls showed a molecular graph corresponding to a type of macrocyclic aromatic hydrocarbons, and these molecular graphs of polyphenyls construct a polyphenyl structure. Polyphenyls and their derivatives have applications in drug synthesis, organic synthesis, heat exchangers, etc., and have received attention from chemists. A random polyphenyl chain PPC_l with l hexagons can be constructed by a polyphenyl chain PPC_{l-1} using $l-1$ hexagons attached to a new hexagon G_l by a bridge (see Figure 3).

The PPC_l will be a polyphenyl chain with $l \geq 2$ having G_1, G_2, \dots, G_l hexagons. PPC_l is the meta-chain M_l , the ortho-chain O_l^1 and the para-chain L_l . The new hexagon can be

joined in three arrangements, which construct the local orderings. We use these as $PPC_l^1, PPC_l^2, PPC_l^3$ (see Figure 4).

A random polyphenyl chain $PPC_l(k_1, k_2)$ is a polyphenyl chain constructed by step-by-step attachment of new hexagons. At every step $p = 2, 3, \dots, l$, a random choice construct one of the three possible chains:

- 1 $PPC_{p-1} \rightarrow PPC_p^1$ with probability k_1 ,
- 2 $PPC_{p-1} \rightarrow PPC_p^2$ with probability k_2 ,
- 3 $PPC_{p-1} \rightarrow PPC_p^3$ with probability $k_3 = 1 - k_1 - k_2$,

Where all the given probabilities are constant. In this section, we discuss the expected value for the first Zagreb connection index of the random polyphenyl chain with l hexagons.

Theorem 3.1. For $l \geq 2$, the expected value for the first Zagreb connection index of the random polyphenyl chain PPC_l is

$$E(ZC_1(PPC_l)) = (8k_1 + 2k_2 + 68)l - 16k_1 - 4k_2 - 44.$$

Proof. Case-I: When $l = 2$, one can get

$$E(ZC_1(PPC_2)) = 6 \times (2)^2 + 4 \times (3)^2 + 2 \times (4)^2 = 92.$$

Case-II: When $l \geq 3$, it is obvious that $f_2(PPC_l), f_3(PPC_l), f_4(PPC_l)$ and $f_5(PPC_l)$ depends on the four possible cases, as follows:

- 1 If $PPC_{l-1} \rightarrow PPC_l^1$ having probability k_1 , we acquire





FIGURE 3
The three types of local arrangements of hexagons PPC_l^1, PPC_l^2 , and PPC_l^3 .



$$\begin{aligned} f_2(PPC_l^1) &= f_2(PPC_{l-1}) + 2, \\ f_3(PPC_l^1) &= f_3(l-1) + 2, \\ f_4(PPC_l^1) &= f_4(PPC_{l-1}) = 2, \\ f_5(PPC_l^1) &= f_5(PPC_{l-1}) + 2. \end{aligned}$$

By using the above values in Eq. 1.1, we get

$$\begin{aligned} ZC_1(PPC_l^1) &= ZC_1(PPC_{l-1}) + 2 \times 2^2 + 2 \times 3^2 \\ &\quad + 2 \times 5^2 \\ &= ZC_1(PPC_{l-1}) + 76. \end{aligned}$$

2 If $PPC_{l-1} \rightarrow PPC_l^2$ having probability k_2 , we acquire

$$\begin{aligned} f_2(PPC_l^2) &= f_2(PPC_{l-1}) + 1, \\ f_3(PPC_l^2) &= f_3(PPC_{l-1}) + 2, \\ f_4(PPC_l^2) &= f_4(PPC_{l-1}) + 3, \end{aligned}$$

By using the above values in Eq. 1.1, we get

$$\begin{aligned} ZC_1(PPC_l^2) &= ZC_1(PPC_{l-1}) + 1 \times 2^2 + 2 \times 3^2 \\ &\quad + 3 \times 4^2 \\ &= ZC_1(PPC_{l-1}) + 70. \end{aligned}$$

3 If $PPC_{l-1} \rightarrow PPC_l^3$ having probability k_3 , we acquire

$$\begin{aligned} f_2(PPC_l^3) &= f_2(PPC_{l-1}) = 6, \\ f_3(PPC_l^3) &= f_3(PPC_{l-1}) + 4, \\ f_4(PPC_l^3) &= f_4(PPC_{l-1}) + 2, \end{aligned}$$

By using the above values in Eq. 1.1, we get

$$\begin{aligned} ZC_1(PPC_l^3) &= ZC_1(PPC_{l-1}) + 4 \times 3^2 + 2 \times 4^2 \\ &= ZC_1(PPC_{l-1}) + 68. \end{aligned}$$

Now

$$\begin{aligned} E_l^i &= E(ZC_1(PPC_l)) \\ &= k_1 ZC_1(PPC_l^1) + k_2 ZC_1(PPC_l^2) \\ &\quad + (1 - k_1 - k_2) ZC_1(PPC_l^3) \\ &= ZC_1(PPC_{l-1}) + 8k_1 + 2k_2 + 68. \end{aligned} \quad (3.1)$$

Note that $E[E_l^i] = E_l^i$. By applying the expression operator to Eq. 3.1 and also $l \geq 3$, we get

$$E_l^i = E_{l-1}^i + 8k_1 + 2k_2 + 68. \quad (3.2)$$

The result Eq. 3.2 is a first-order non-homogeneous linear difference equation with constant coefficients. The general solution of the homogeneous side is Eq. 3.2 is $E^i = C$. Suppose $E^{i'} = bl$ is a particular result of Eq. 3.2, using $E^{i'}$ into Eq. 3.2, we acquire

$$b = 8k_1 + 2k_2 + 68.$$

Finally the general solution of Eq. 3.2 is given by

$$\begin{aligned} E_l^i &= E^i + E^{i'} \\ &= E(ZC_1(PPC_l)) = (8k_1 + 2k_2 + 68)l + C. \end{aligned}$$

Applying the initial condition $l = 3$, we get following

$$C = -16k_1 - 4k_2 - 44.$$

Therefore

$$E_l^i = E(ZC_1(PPC_l)) = (8k_1 + 2k_2 + 68)l - 16k_1 - 4k_2 - 44.$$

If $k_1 = 1$ (respectively, $k_2 = 1$) and $k_2 = k_3 = 0$ (respectively, $k_1 = k_3 = 0$), then $PPC_l = M_l$ (respectively, $PPC_l = O_l$). Similarly, if $k_3 = 1$ and $k_1 = k_2 = 0$, then $PPC_l = L_l$. By Theorem 3.1, we can acquire the first Zagreb connection index of polyphenyl chains like meta M_b , ortho O_b , and para L_b as

$$\begin{aligned} ZC_1(M_l) &= 76l - 60, & ZC_1(O_l) &= 70l - 48, \\ ZC_1(L_l) &= 68l - 44. \end{aligned}$$

Corollary 3.2. For a random polyphenyl chain $PPC_l (l \geq 3)$, the para-chain L_l and the meta-chain M_l achieves the minimum and the maximum $E(ZC_1(PPC_l))$, respectively.

Proof. From Theorem 3.1, we obtain

$$\begin{aligned} E_l^i &= E(ZC_1(PPC_l)) \\ &= (8l - 16)k_1 + (2l - 4)k_2 + 68l - 44. \end{aligned}$$

By taking partial derivatives, we acquire $\frac{\partial E}{\partial k_1} = 8l - 16 > 0$, $\frac{\partial E}{\partial k_2} = 2l - 4 > 0$. When $k_1 = k_2 = 0$ (i.e. $k_3 = 1$), the para-chain L_l has the minimum of $E(ZC_1(COC_l))$, that is $PPC_l \cong L_l$. If $k_1 = 1 - k_2$ ($0 \leq k_2 \leq 1$), we acquire


$$\begin{aligned} E_l^i &= E(ZC_1(PPC_l)) \\ &= (8l - 16)(1 - k_2) + (2l - 4)k_2 + 68l - 44. \end{aligned}$$

Therefore $\frac{\partial E}{\partial k_2} = -6l + 12 < 0$. Thus $E(ZC_1(PPC_l))$ achieves the maximum value, if $k_2 = 0$ ($k_1 = 1$), that is $PPC_l \cong M_l$.

4 The first Zagreb connection index of random chain network PG_l

The random chain networks PG_l with l pentagons can be constructed by PG_{l-1} having $l-1$ pentagons attached to a new pentagon H_l by a bridge (see Figure 5).

The PG_l will be a random chain network with $l \geq 2$, and H_1, H_2, \dots, H_l pentagons. For $l \geq 3$, there are two ways to attach pentagons at the end and get PG_l^1 and PG_l^2 , (see Figure 6). For



such a random chain network, any step for $q = 2, 3, 4, \dots, l$ can be constructed by two possible chains with given probabilities k_1 and k_2 , respectively:

- 1 $PG_{q-1} \rightarrow PG_q^1$ with probability k_1 ,
- 2 $PG_{q-1} \rightarrow PG_q^2$ with probability $k_2 = 1 - k_1$,

Where all the given probabilities are constant.

This section discusses the expected value for the first Zagreb connection index of the random chain network with l pentagons. The proof of [Theorem 4.1](#) is the same as the proofs of [Theorem 2.1](#) and [Theorem 3.1](#); therefore, we omit it here.

Theorem 4.1. For $l \geq 2$, the expected value for the first Zagreb connection index of random chain network PG_l is $E(ZC_1(PG_l)) = (6k_1 + 66)L - 12k_1 - 48$.

If $k_1 = 1$ (respectively, $k_2 = 1$) and $k_2 = 0$ (respectively, $k_1 = 0$), then $PG_l = PG_l^1$ (respectively, $PG_l = PG_l^2$). By [Theorem 4.1](#), we can acquire the first Zagreb connection index of the meta-chain PG_l^1 and para-chain PG_l^2 , as

$$ZC_1(PG_l^1) = 72l - 60, \quad ZC_1(PG_l^2) = 66l - 48.$$

Corollary 4.2. For a random chain network $PG_l (l \geq 3)$, the para-chain PG_l^2 and the meta-chain PG_l^1 achieves the minimum and the maximum of $E(ZC_1(PG_l))$, respectively.

5 The average values for the first Zagreb connection index

This section finds the average values for the first Zagreb connection index concerning the sets of all cyclooctatetraene chains with l octagons, polyphenyl chains with l hexagons, and chain networks with l pentagons. Let \mathbb{G}_l , \mathbb{R}_l and \mathbb{Q}_l be the sets of all cyclooctatetraene chains, polyphenyl chains, and random chain network, respectively. The average values for the first Zagreb connection index for the sets \mathbb{G}_l , \mathbb{R}_l and \mathbb{Q}_l are given below:

$$ZC_1^{avg}(\mathbb{G}_l) = \frac{1}{|\mathbb{G}_l|} \sum_{\mathcal{H} \in \mathbb{G}_l} ZC_1(\mathcal{H}),$$

$$ZC_1^{avg}(\mathbb{R}_l) = \frac{1}{|\mathbb{R}_l|} \sum_{\mathcal{H} \in \mathbb{R}_l} ZC_1(\mathcal{H}),$$

$$ZC_1^{avg}(\mathbb{Q}_l) = \frac{1}{|\mathbb{Q}_l|} \sum_{\mathcal{H} \in \mathbb{Q}_l} ZC_1(\mathcal{H}).$$


The average value concerning sets \mathbb{G}_l , \mathbb{R}_l , and \mathbb{Q}_l are expected values for the first Zagreb connection index of the random chains. From [Theorem 2.1](#), [Theorem 3.1](#) and [Theorem 4.1](#), we have.

Theorem 5.1. The average value for the first Zagreb connection index concerning the set \mathbb{G}_l is given as:

$$ZC_1^{avg}(\mathbb{G}_l) = \frac{157}{2}l - 49.$$

After calculation, we acquire

$$ZC_1^{avg}(\mathbb{G}_l) = \frac{1}{4} (ZC_1(M_l) + ZC_1(O_l^1) + ZC_1(O_l^2) + ZC_1(L_l)).$$



Theorem 5.2. The average value for the first Zagreb connection index concerning \mathbb{R}_l is

$$ZC_1^{avg}(\mathbb{R}_l) = \frac{214}{3}l - \frac{152}{3}.$$

After calculation, we acquire

$$ZC_1^{avg}(\mathbb{R}_l) = \frac{1}{3}(ZC_1(M_l) + ZC_1(O_l) + ZC_1(L_l)).$$

Theorem 5.3. The average value for the first Zagreb connection index concerning \mathbb{Q}_l is $ZC_1^{avg}(\mathbb{Q}_l) = 69l - 54$. It is also:

$$ZC_1^{avg}(\mathbb{Q}_l) = \frac{1}{2}(ZC_1(PG_l^1) + ZC_1(PG_l^2)).$$

6 Conclusion

This study computed the expected values of the first Zagreb connection index in a random cyclooctatetraene chain, random polyphenyls chain, and random chain network with l , octagons, hexagons, and pentagons, respectively. It has discussed the maximum chain and the minimum chain of the COC_b , PPC_b , and PG_b , respectively, concerning the expected values of these chains. The average values discussed in all of the above are considered random chains for unique chains.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material, further inquiries can be directed to the corresponding authors.

References

- Akhter, S., Imran, M., Farahan, M. R., and Javaid, I. (2017). On topological properties of hexagonal and silicate networks. *Hac. J. Math. Stat.* 48 (3), 1–13. doi:10.15672/hjms.2017.541
- Akhter, S., Imran, M., Gao, W., and Farahani, M. R. (2018). On topological indices of honeycomb networks and graphene networks. *Hac. J. Math. Stat.* 47 (1), 19–35. doi:10.15672/hjms.2017.464
- Akhter, S., Imran, M., and Iqbal, Z. (2020). Mostar indices of SiO_2 nanostructures and melem chain nanostructures. *Int. J. Quantum Chem.* 121 (5). doi:10.1002/qua.26520
- Akhter, S., and Imran, M. (2016). On molecular topological properties of benzenoid structures. *Can. J. Chem.* 94 (8), 687–698. doi:10.1139/cjc-2016-0032
- Akhter, S., Imran, M., and Raza, Z. (2016). On the general sum-connectivity index and general Randić index of cacti. *J. Inequal. Appl.* 2016 (1), 300–309. doi:10.1186/s13660-016-1250-6
- Ali, A., and Trinajstić, N. (2018). A novel/old modification of the first Zagreb index. *Mol. Inf.* 37 (6–7), 1800008. doi:10.1002/minf.201800008
- Bao, L. J., Raza, Z., and Javaid, M. (2020). Zagreb connection numbers for cellular neural networks. *Discrete Dyn. Nat. Soc.* 2020, 1–8. doi:10.1155/2020/8038304
- Basavanagoud, B., and Chitra, E. (2018). On the leap Zagreb indices of generalized xyz -point-line transformation graphs $T_{xyz}(G)$, when $z = 1$. *Int. J. Math. Comb.* 2 (44–66).
- Basavanagoud, B., and Jakkannavar, P. (2018). Computing the first leap Zagreb index of some nanostructures. *Inter. J. Appl. Math.* 6 (2–B), 141–150.
- Cao, J., Ali, U., Javaid, M., and Huang, C. (2020). Zagreb connection indices of molecular graphs based on operations. *Complexity* 2020, 1–15. doi:10.1155/2020/7385682
- Donald, H. L., and Whitehead, M. A. (1969). Molecular geometry and bond energy. III. cyclooctatetraene and related compounds. *J. Am. Chem. Soc.* 91 (2), 238–242. doi:10.1021/ja01030a003
- Du, Z., Ali, A., and Trinajstić, N. (2019). Alkanes with the first three maximal/minimal modified first Zagreb connection indices. *Mol. Inf.* 38 (4), 1800116. doi:10.1002/minf.201800116
- Ducoffe, G., Marinescu-Ghemeci, R., and Obreja, C. (2018). International symposium on symbolic and numeric algorithms for scientific computing (SYNASC). *Timisoara, Rom.* 141–148.
- Ducoffe, G., Marinescu-Ghemeci, R., and Obreja, C. (2018). Proceedings of the 16th cologne-twente workshop on graphs and combinatorial optimization. *Paris, Fr. CNAM* 65–68.
- Fang, X., You, L., and Liu, H. (2021). The expected values of Sombor indices in random hexagonal chains, phenylene chains, and Sombor indices of some chemical graphs. *Int. J. Quantum. Chem.* 121 (17). doi:10.1002/qua.26740

Author contributions

Investigation: ZR, SA, and YS; Writing: ZR, SA, and YS; Review: ZR, SA, and YS.

Funding

This research was funded by the University of Sharjah.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fchem.2022.1067874/full#supplementary-material>

- Fatima, N., Bhatti, A. A., Ali, A., and Gao, W. (2019). Zagreb connection indices of two dendrimer nanostars. *Acta Chem. Iasi* 27 (1), 1–14. doi:10.2478/achi-2019-0001
- Garavelli, M., Bernardi, F., Cembran, A., Castaño, O., Frutos, L. M., Merchán, M., et al. (2002). Cyclooctatetraene computational photo- and thermal chemistry: A reactivity model for conjugated hydrocarbons. *J. Am. Chem. Soc.* 124 (46), 13770–13789. doi:10.1021/ja020741v
- Gutman, I. (2013). Degree-based topological indices. *Croat. Chem. Acta.* 86, 351–361. doi:10.5562/cca2294
- Gutman, I., Milovanović, E., and Milovanović, I. (2020). Beyond the Zagreb indices. *AKCE Int. J. Graphs Comb.* 17 (1), 74–85. doi:10.1016/j.akcej.2018.05.002
- Gutman, I., Naji, A. M., and Soner, N. D. (2017). The first leap Zagreb index of some graph operations. *Commun. Comb. Optim.* 2 (2), 99–117.
- Gutman, I., and Trinajstić, N. (1972). Graph theory and molecular orbitals, Total π electron energy of alternant hydrocarbons. *Chem. Phys. Lett.* 17 (4), 535–538. doi:10.1016/0009-2614(72)85099-1
- Huang, G. H., Kuang, M. J., and Deng, H. Y. (2014). The expected values of Kirchhoff indices in the random polyphenyl and spiro chains. *ARS Math. Contem.* 9 (2), 197–207. doi:10.26493/1855-3974.458.7b0
- Jahanbanni, A. (2022). The expected values of the first Zagreb and Randić indices in random polyphenyl chains. *Polycycl. Aromat. Compd.* 42 (4), 1851–1860. doi:10.1080/10406638.2020.1809472
- Khalid, S., Kok, J., and Ali, A. (2018). Zagreb connection indices of TiO_2 nanotubes. *Chem. Bulg. J. Sci. Edu.* 27 (1), 86–92.
- Ma, L., Bian, H., Liu, B. J., and Yu, H. Z. (2017). The expected values of the Wiener index in the random phenylene and spiro chains. *ARS Comb.* 130, 267–274.
- Manzoor, S., Fatima, N., Bhatti, A. A., and Ali, A. (2018). Zagreb connection indices of some nanostructures. *Acta Chem. Iasi* 26 (2), 169–180. doi:10.2478/achi-2018-0011
- Mathews, F. S., and Lipscomb, W. N. (1959). The structure of silver cyclooctatetraene nitrate. *J. Phy. Chem.* 63 (6), 845–850. doi:10.1021/j150576a017
- Milas, N., and NolanPetrus, J., Jr. (1958). Notes-ozonization of cyclooctatetraene. *J. Org. Chem.* 23 (4), 624–625. doi:10.1021/jo01098a611
- Naji, A. M., Davvaz, B., and Mahde, S. S. (2020). A study on some properties of leap graphs. *Commun. Comb. Optim.* 5 (1), 9–17.
- Naji, A. M., and Soner, N. D. (2018). The first leap Zagreb index of some graph operations. *Int. J. Appl. Graph Theor.* 2 (1), 7–18.
- Qi, J., Fang, M., and Geng, X. (2022). The expected value for the Wiener index in the random spiro chains. *Polycycl. Aromat. Compd.* 1–11.
- Raza, Z., and Imran, M. (2021). Expected values of some molecular descriptors in random cyclooctane chains. *Symmetry* 13 (11), 2197. doi:10.3390/sym13112197
- Raza, Z. (2020). Leap Zagreb connection numbers for some networks models. *J. Chem.* 20 (6), 1407–1413. doi:10.22146/jc.53393
- Raza, Z., Naz, K., and Ahmad, S. (2022). Expected values of molecular descriptors in random polyphenyl chains. *Emerg. Sci. J.* 6 (1), 151–165. doi:10.28991/esj-2022-06-01-012
- Raza, Z. (2020). The expected values of arithmetic bond connectivity and geometric indices in random phenylene chains. *Heliyon* 6 (7), e04479. doi:10.1016/j.heliyon.2020.e04479
- Raza, Z. (2021). The expected values of some indices in random phenylene chains. *Eur. Phys. J. Plus* 136 (11–15), 91. doi:10.1140/epjp/s13360-021-01082-y
- Raza, Z. (2022). Zagreb connection indices for some Benzenoid systems. *Polycycl. Aromat. Compd.* 42 (4), 1814–1827. doi:10.1080/10406638.2020.1809469
- Schwamm, R. J., Anker, M. D., Lein, M., and Coles, M. P. (2019). Reduction vs. Addition: The reaction of an aluminyl anion with 1, 3, 5, 7 - cyclooctatetraene. *Angew. Chem. Int. Ed.* 58 (5), 1489–1493. doi:10.1002/anie.201811675
- Shao, Z., Gutman, I., and Li, Z. (2018). Leap Zagreb indices of trees and unicyclic graphs. *Commun. Comb. Optim.* 3 (2), 179–194.
- Shao, Z., Gutman, I., Li, Z., Wang, S., and Wu, P. (2018). Leap Zagreb indices of trees and unicyclic graphs. *Commun. Comb. Optim.* 3 (2), 179–194.
- Soner, N. D., and Naji, A. M. (2016). The k -distance neighborhood polynomial of a graph. *Int. J. Math. Comput. Sci.* 3 (9), 2359–2364.
- Tang, J. H., Ali, U., Javaid, M., and Shabbir, K. (2019). Zagreb connection indices of subdivision and semi-total point operations on graphs. *J. Chem.*
- Todeschini, R., and Consonni, V. (2000). *Handbook of molecular descriptors*. Weinheim: Wiley VCH.
- Traetteberg, M., Hagen, G., and Cyvin, S. J. (1970). IV. 1, 3, 5, 7-Cyclooctatetraene. *Z. Für Naturforsch. B* 25 (2), 134–138. doi:10.1515/znb-1970-0201
- Willis, B. P., George, C. P., and Kenneth, S. P. (1952). The structure of cyclooctatetraene. *J. Am. Chem. Soc.* 74 (13), 3437–3438. doi:10.1021/ja01133a524
- Yang, W., and Zhang, F. (2012). Wiener index in random polyphenyl chains. *MATCH Commun. Math. Comp. Chem.* 68, 371–376.
- Ye, A., Qureshi, M. I., Fahad, A., Aslam, A., Jamil, M. K., Zafar, A., et al. (2019). Zagreb connection number index of nanotubes and regular hexagonal lattice. *Open Chem.* 17 (1), 75–80. doi:10.1515/chem-2019-0007
- Zhang, L., Li, Q., Li, S., and Zhang, M. (2020). The expected values for the Schultz index, Gutman index, multiplicative degree Kirchhoff index, and additive degree Kirchhoff index of a random polyphenylene chain. *Discrete Appl. Math.* 282, 243–256. doi:10.1016/j.dam.2019.11.007
- Zhang, W., You, L., Liu, H., and Huang, Y. (2021). The expected values and variances for Sombor indices in a general random chain. *App. Math. Comp.* 411, 126521. doi:10.1016/j.amc.2021.126521