

Quantum systems at the brink: existence of bound states, critical potentials and dimensionality

Dirk Hundertmark, Michal Jex, Markus Lange

CRC Preprint 2022/79, December 2022

KARLSRUHE INSTITUTE OF TECHNOLOGY

CRC 1173



Participating universities



Universität Stuttgart

EBERHARD KARLS
UNIVERSITÄT
TÜBINGEN



Funded by

DFG

QUANTUM SYSTEMS AT THE BRINK: EXISTENCE OF BOUND STATES, CRITICAL POTENTIALS AND DIMENSIONALITY

DIRK HUNDERTMARK, MICHAL JEX, AND MARKUS LANGE

ABSTRACT. One of the crucial properties of a quantum system is the existence of bound states. While the existence of eigenvalues below zero, i.e., below the essential spectrum, is well understood, the situation of zero energy bound states at the edge of the essential spectrum is far less understood. We present necessary and sufficient conditions for Schrödinger operators to have a zero energy bound state. Our sharp criteria show that the existence and non-existence of zero energy ground states depends strongly on the dimension and the asymptotic behavior of the potential. There is a spectral phase transition with dimension four being critical.

CONTENTS

1. Introduction	1
2. Definitions and preparations	7
3. Proof of the non-existence result	13
4. Proof of the existence result	16
Appendix A. An example in search of a theorem	22
References	27

1. INTRODUCTION

The existence of bound states plays a crucial role for the properties of quantum systems. Of special importance is the ground state, i.e., the eigenfunction corresponding to the lowest eigenvalue of the Hamiltonian describing the system. In this paper we consider a Schrödinger operator of the form

$$H = -\Delta + V \tag{1.1}$$

on $L^2(\mathbb{R}^d)$ where $V \in L^1_{loc}(\mathbb{R}^d)$ is a real-valued potential such that the operator H is a well-defined self-adjoint realization of the formal differential operator $-\Delta + V$ which is bounded from below. Moreover, we need that eigenfunctions of H are continuous. The precise conditions are given in Assumption 1.1 below.

We are particularly interested in the special case when the ground state energy of the Schrödinger operator H is at the threshold of the essential spectrum. By shifting the potential by a constant, one can assume that the essential spectrum of H starts at zero. One also

2010 *Mathematics Subject Classification.* 81Q05 (primary); 35Q40, 81V45 (secondary).

Key words and phrases. Resonances, virtual levels, threshold eigenvalues.

©2022 by the authors. Faithful reproduction of this article, in its entirety, by any means is permitted for non-commercial purposes.

often assumes that the potential V decays to zero at infinity such that the essential spectrum $\sigma_{\text{ess}}(H) = [0, \infty)$, see Remark 1.10. Under these conditions the zero–energy level is at the edge of two regions with very distinct behavior: the point and the continuous spectrum. It is well-known that positive eigenvalues embedded in the continuum appear only due to a special combination of oscillations and slow decay of the potential. This goes back to [54], see also [13, 17, 24, 49] and the references therein. Also, with the help of the min-max theorem, the existence and non-existence of eigenvalues below zero is well-understood, see, e.g., [47].

Whether zero is actually a threshold eigenvalue, i.e., an eigenvalue at *the edge of the continuum* is a very difficult problem, in general. Early results on existence or non-existence of zero-energy eigenvalues go back to [1, 25, 29, 33, 32, 38, 42, 45, 46, 51]. In [25] the authors studied the behavior of resonances and eigenstates at the zero–energy threshold in $d = 3$. Furthermore, based on the remark of a referee they note that resonances cannot exist in dimensions $d > 4$ based on properties of Riesz potential. However their approach is not applicable for $d = 4$. For slowly decaying *negative* potentials which, amongst other conditions, obey $V(x) \sim -c|x|^{-\gamma}$ for some $c > 0$ and $0 < \gamma < 2$ in the limit $|x| \rightarrow \infty$, the non-existence of zero energy eigenstates was shown in [14, 16], while it was noted in [10] that a long range Coulomb part can create zero energy eigenstates, see also [41, 55]. An analysis of eigenstates and resonances at the threshold for the case of certain nonlocal operators appeared in [27].

In [9] it has been shown that, for Schrödinger operators on $L^2(\mathbb{R}^3)$ with spherical symmetric potentials $V \in L^p(\mathbb{R}^3)$ with $p > 3/2$ whose positive part satisfies $V_+(x) \leq 3/(4|x|^2)$ for $|x|$ large enough, zero is not an eigenvalue corresponding to a positive square integrable ground state eigenfunction. This extends to potentials with $V_+ \leq |x|^{-2} (3/4 + \ln^{-1}(|x|))$ near infinity in \mathbb{R}^3 , the constants $3/4$ and 1 are optimal. For similar results see [19], which reproved a slightly weaker non-existence result compared to [9] and additionally showed that if $V(x) \geq C|x|^{-2}$ for some constant $C > 3/4$ and $|x|$ large then zero is an eigenvalue for critical potential, see Definition 1.5. Thus a repulsive part can stabilize zero energy bound states of quantum systems.

Strictly speaking, the paper [9] deals with continuous potentials on \mathbb{R}^3 but they note that the condition $V \in L^p(\mathbb{R}^3)$ with $p > 3/2$ is enough to guarantee continuity of ground states, due to a Harnack inequality for positive eigenfunctions. We also note that compactly supported zero–energy eigenfunctions were constructed in [30, 34] for potential $V \in L^p(\mathbb{R}^d)$ with $p < d/2$ and compact support. For these potentials, a Harnack inequality for the ground state cannot hold.

In this paper we significantly extend all previous results, in particular the ones of [9] and [19], by proving a *family of sharp criteria* for the existence and non–existence of zero energy ground states at *the edge of the essential spectrum* for Schrödinger operators in *arbitrary* dimensions. In particular, our results apply to Schrödinger operators with a so–called virtual level at zero energy and they explain when such a virtual level is a true ground state or when it is a resonance.

Our results clearly explain why increasing the dimension makes it easier for a virtual level to be a true ground state. In particular, our work explains why dimension $d = 4$ is *critical*. Dimension four shares some similarity with the case of lower dimensions but *higher order corrections* from our criteria are needed to settle this case.

Our main assumption on the potential V are given by

Assumption 1.1. The potential V is in the local Kato–class $K_{d,loc}(\mathbb{R}^d)$ and the negative part $V_- = \sup(-V, 0)$ is relatively form small w.r.t. $-\Delta + V_+$, i.e., there exist $0 \leq a < 1$ and $b \geq 0$ such that

$$\langle \psi, V_- \psi \rangle = \|V_-^{1/2} \psi\|^2 \leq a(\|\nabla \psi\|^2 + \|\sqrt{V_+} \psi\|^2) + b\|\psi\|^2 \quad (1.2)$$

for all $\psi \in H^1(\mathbb{R}^d) \cap \mathcal{D}(\sqrt{V_+})$. Here $\mathcal{D}(\sqrt{V_+})$ is the domain of the multiplication operator $\sqrt{V_+}$ on $L^2(\mathbb{R}^d)$, also called the form domain of V_+ and often written as $\mathcal{Q}(V_+)$.

Note that what we call *relatively form small* is usually called *relatively form-bounded with relative bound $a < 1$* . We will call a potential W infinitesimally form bounded (w.r.t. $-\Delta + V_+$) if for all $a > 0$ there exist $b \geq 0$ such that the positive and negative parts W_{\pm} satisfy (1.2) (with V_- replaced by W_{\pm}).

Remark 1.2. The (local) Kato–class $K_{d,loc}(\mathbb{R}^d) \subset L^1_{loc}(\mathbb{R}^d)$, whose definition is recalled below, see (1.14), contains most, if not all physically relevant potentials. This assumption is only made to guarantee that all weak local eigenfunctions of H are continuous, see [3, 48, 52].

One could relax the assumption that $V \in K^d_{loc}(\mathbb{R}^d)$ to $V \in L^1_{loc}(\mathbb{R}^d)$, if some other condition guaranteed that weak local eigenfunctions of H are continuous. In fact, it would be sufficient to have that eigenfunctions are locally bounded and that a ground state of H is bounded away from zero on compact sets. As will become clear from the proofs, we can allow for severe local singularities. E.g., it is enough to assume that V is in the local Kato–class outside some compact set $K \subset \mathbb{R}^d$.

If $V_{\pm} \in L^1_{loc}$ and (1.2) holds, the KLMN theorem shows that there exists a unique self-adjoint operator H , informally given by the differential operator $-\Delta + V$, such that its quadratic form, which with a slight abuse of notation we write as

$$\langle \psi, H \psi \rangle := \langle \nabla \psi, \nabla \psi \rangle + \langle \sqrt{V_+} \psi, \sqrt{V_+} \psi \rangle - \langle \sqrt{V_-} \psi, \sqrt{V_-} \psi \rangle \quad (1.3)$$

is well-defined for $\psi \in \mathcal{Q}(H) := H^1(\mathbb{R}^d) \cap \mathcal{Q}(V_+)$. Moreover, it is closed and bounded from below on the quadratic form domain $\mathcal{Q}(H)$. See also the discussion at the beginning of the next section.

To formulate our main results we recall the definition of the iterated logarithms \ln_n defined, for natural numbers $n \in \mathbb{N}$, by $\ln_1(r) := \ln(r)$ for $r > 0$ and inductively for $r > e_n$ by $\ln_{n+1}(r) := \ln(\ln_n(r))$. Here $e_0 = 0$ and $e_{n+1} = e^{e_n}$. Our first main result can be summarized as follows

Theorem 1.3 (Absence of a zero energy ground state). *Assume that the potential V satisfies Assumption 1.1 and $\sigma(H) = [0, \infty)$. If for some $m \in \mathbb{N}_0$ and $R > e_m$*

$$V(x) \leq \frac{d(4-d)}{4|x|^2} + \frac{1}{|x|^2} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) \quad (1.4)$$

for all $|x| \geq R$, then zero is not an eigenvalue of the Schrödinger operator H .

As usual the empty product is 1 and the empty sum equals 0.

Remark 1.4. In particular, if $\inf \sigma_{ess}(H) = 0$ then Theorem 1.3 shows that zero is not an eigenvalue at the edge of the essential spectrum. Theorem 1.7 below shows the sharpness of condition (1.4) on the potential V for the absence of an embedded ground state at the edge of the essential spectrum.

Our second main result shows that critical potentials create zero energy ground states if they are not too small at infinity. We call a potential $W \geq 0$ nontrivial, if it is strictly positive on a set of positive Lebesgue measure.

Definition 1.5 (Critical potential). The potential V is *critical* if the Schrödinger operator H has spectrum $\sigma(H) = \sigma_{ess}(H) = [0, \infty)$ and for all nontrivial compactly supported potentials $W \geq 0$ which are infinitesimally form bounded with respect to $-\Delta + V_+$ the family of operators $H_\lambda = H - \lambda W$ has essential spectrum $\sigma_{ess}(H_\lambda) = [0, \infty)$ and a negative energy bound state for all $\lambda > 0$.

Remark 1.6. The potential V is called *subcritical*, if the Schrödinger operator H has spectrum $\sigma(H) = \sigma_{ess}(H) = [0, \infty)$ and there exist a nontrivial potential $W \geq 0$, which is infinitesimally form bounded with respect to $-\Delta + V_+$, such that $H - \lambda W \geq 0$ for some $\lambda > 0$.

Theorem 1.7 (Existence of a zero energy ground state for critical potentials). *Assume that the potential V satisfies Assumption 1.1 and that it is critical. If for some $m \in \mathbb{N}_0$, $\epsilon > 0$, and $R > e_m$*

$$V(x) \geq \frac{d(4-d)}{4|x|^2} + \frac{1}{|x|^2} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) + \frac{\epsilon}{|x|^2} \prod_{k=1}^m \ln_k^{-1}(|x|) \quad (1.5)$$

for all $|x| \geq R$, then zero is an eigenvalue of H .

Remark 1.8. Clearly, the right hand sides of (1.4) and (1.5) are, for each fixed $n \in \mathbb{N}$ complementary. Thus our criteria for existence and non-existence of zero energy ground states at the edge of the essential spectrum are sharp! Considering the simplest case $m = 0$ we have

$$V(x) \leq \frac{d(4-d)}{4|x|^2} \quad (1.6)$$

for the absence and

$$V(x) \geq \frac{d(4-d) + \epsilon}{4|x|^2} \quad (1.7)$$

for the existence with $\epsilon > 0$ and all $|x|$ large enough. For $d = 3$ this recovers the results proved in [19] for the special case of three dimensions.

Using the higher order corrections from Eqs. (1.4) and (1.5) we obtain a sharp distinction between existence and non-existence in the case of a critical potential. For example, the cases $m = 1, 2$ show that if

$$V(x) \leq \frac{d(4-d)}{4|x|^2} + \frac{1}{|x|^2 \ln |x|} \quad \text{or} \quad (1.8)$$

$$V(x) \leq \frac{d(4-d)}{4|x|^2} + \frac{1}{|x|^2 \ln |x|} + \frac{1}{|x|^2 \ln |x| \ln_2 |x|} \quad (1.9)$$

for all large enough $|x|$, then zero will not be a ground state eigenvalue. Conversely, for critical potentials the bound

$$V(x) \geq \frac{d(4-d)}{4|x|^2} + \frac{1+\epsilon}{|x|^2 \ln|x|} \quad \text{or} \quad (1.10)$$

$$V(x) \geq \frac{d(4-d)}{4|x|^2} + \frac{1}{|x|^2 \ln|x|} + \frac{1+\epsilon}{|x|^2 \ln|x| \ln_2|x|} \quad (1.11)$$

for all large enough $|x|$ and some $\epsilon > 0$ implies that zero is a ground state eigenvalue. Using $d = 3$ in (1.8) recovers the non-existence result of [9]. The $d = 3$ case in (1.10) provides a complementary existence result which was missing in [9].

More importantly, our results provide, to arbitrary order, a whole family of complementary sharp criteria which are not restricted to three dimensions and our proofs are considerably simpler than the approaches based on delicate estimates for Green's functions.

One often says that a Schrödinger operator with a critical potential has a *virtual level* (at zero energy), see, e.g. [5, 6, 7]. Theorem 1.3 shows that such a virtual level is *not a bound state* of H if V obeys the bound (1.4), that is, it is a so-called zero energy resonance. Conversely, Theorem 1.7 shows that a virtual level is an eigenvalue *at the edge of the essential spectrum* when the potential V satisfies the *complementary bound* (1.5).

Remark 1.9. In Appendix A we construct a family of potentials $V_{\alpha,d}$ on \mathbb{R}^d for $\alpha \in \mathbb{R}$ and $d \in \mathbb{N}$ such that the Schrödinger operator $H_{\alpha,d} = -\Delta + V_{\alpha,d}$ has spectrum $\sigma(H_{\alpha,d}) = [0, \infty)$. Moreover, $V_{\alpha,d}$ is subcritical for $\alpha < 0$ and critical for $\alpha \geq 0$. The Schrödinger operator $H_{\alpha,d}$ has a zero energy resonance for $0 \leq \alpha \leq 1$ and a zero energy bound state for $\alpha > 1$ in any dimension.

Remark 1.10. The operator H_λ is well-defined with quadratic form methods for all λ , see Remark 2.1. In order to guarantee that $\sigma_{ess}(H_\lambda) = [0, \infty)$ in Definition 1.5, some decay of the potential V is required. A well-known sufficient criteria for this is that V is relatively form compact with respect to the kinetic energy $P^2 = -\Delta$, see [53]. This also implies that V is infinitesimally form bounded, i.e., relatively form small with relative bound zero, w.r.t. $P^2 = -\Delta$, which excludes Hardy type potentials.

A *much less restrictive criterium* for $\sigma_{ess}(H) = [0, \infty)$ only assumes that V *vanishes asymptotically with respect to the kinetic energy*. More precisely, if

$$|\langle \varphi, V\varphi \rangle| \leq a_n \|\nabla \varphi\|^2 + b_n \|\varphi\|^2 \quad (1.12)$$

for all $\varphi \in H^1(\mathbb{R}^d)$ with support $\text{supp}(\varphi) \subset \{|x| \geq R_n\}$ for some sequences $0 \leq a_n, b_n \rightarrow 0$, and $R_n \rightarrow \infty$ as $n \rightarrow \infty$, then $\sigma_{ess}(H) = [0, \infty)$, see [4, 26].

This criterion is clearly in line with the physical heuristic that *only the asymptotic behavior of the potential near infinity* determines the essential spectrum and it allows for strongly singular potentials which are not infinitesimally form bounded. It also shows that $\sigma_{ess}(H_\lambda) = \sigma_{ess}(H) = [0, \infty)$ for all $\lambda > 0$ when W has compact support and is infinitesimally form bounded w.r.t. $-\Delta$ and V is form small w.r.t. $-\Delta$ and satisfies (1.12).

Remark 1.11. The bounds on the potential in Theorems 1.3 and 1.7 are similar in spirit to logarithmic corrections to the Hardy inequality. For $\psi \in C_0^\infty(\mathbb{R}^d \setminus \{|x| < e_m\})$ and one has

$$\langle \nabla \psi, \nabla \psi \rangle \geq \langle \psi, \left(\frac{(d-2)^2}{4|x|^2} + \frac{1}{4|x|^2} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) \right) \psi \rangle, \quad (1.13)$$

see [43], which also discusses conditions on the potential such that $-\Delta + V$ has infinitely many, respectively finitely many negative eigenvalues. Bounds on the number of negative eigenvalues are given in [39, 40]. Certain logarithmic refinements of Hardy's inequality have been used to study the existence of resonances of Schrödinger operators and the Efimov effect in low dimensions, see [8].

Remark 1.12. Theorems 1.3 and 1.7 show a *spectral phase transition* concerning the existence of zero energy ground states for Schrödinger operators with critical dimension $d = 4$:

The sign of the leading order term in (1.4) and (1.5) strongly depends on the dimension d , being positive if $d \leq 3$, zero in dimension $d = 4$, and negative if $d \geq 5$.

Moreover, in dimension $d = 4$, the leading order term vanishes and the next leading order term with $m = 1$ becomes dominant. Thus the four dimensional case is *critical*. Nevertheless, since the new leading order term for $d = 4$ is also positive, the four dimensional case is similar to the case of lower dimensions. In particular, non-positive potentials V cannot support zero energy ground states in dimensions $d \leq 4$ while in dimension $d \geq 5$ non-positive critical potentials have zero energy ground states.

Hence non-positive critical potentials will always create resonances in dimension $d \leq 4$, while in dimension $d \geq 5$ they have zero energy ground states unless their negative part is so long range such that the bound (1.5) does not hold anymore. Nevertheless, in dimension $d \leq 4$ a 'long-range' positive tail of the potential can create zero energy ground states. See also the discussion in Section 2 of [22].

In particular, assume that the potential V is infinitesimally form bounded w.r.t. $-\Delta$ and has compact support. Then $\sigma_{ess}(-\Delta + \beta V) = [0, \infty)$ for all $\beta \geq 0$, see [4, 53], and a simple application of the min-max principle shows that as soon as negative eigenvalues of $-\Delta + \beta V$ exist, they are decreasing in $\beta > 0$, see [47, Proposition after Theorem XIII.2, page 79]. Let $\beta_0 > 0$ be the value of the coupling constant when the ground energy of $-\Delta + \beta V$ hits zero. Theorem 1.3 shows that $-\Delta + \beta_0 V$ has a zero energy resonance when $d \leq 4$ and Theorem 1.7 shows that it has a zero energy ground state in dimension $d \geq 5$. The asymptotic of the eigenvalues of the perturbed operators $-\Delta + \beta V$ in $\beta - \beta_0$ was studied in [31] for all dimensions.

The structure of our paper is as follows: In Section 2 we present all the necessary technical tools to precisely formulate our main results. Theorem 1.3 is proven in Section 3. The proof is by contradiction, assuming that a zero energy ground state exists and then deriving a lower bound which shows that it cannot be square integrable. To construct such a lower bound one only needs to know that a ground state, if it exists, can be chosen to be positive and that it is locally bounded away from zero. It is well-known that ground states of a Schrödinger operator H in $L^2(\mathbb{R}^d)$ are unique, up to global phase, and can be chosen to be strictly positive as soon as they exist, see [15, 18] or [47, Section XIII.12]. Thus, if one knows that the ground state is bounded away from zero, one can relax the condition on V to $V \in L_{loc}^1(\mathbb{R}^d)$

and V_- satisfies (1.2), The assumption that V is in the local Kato–class is only needed to guarantee that eigenfunctions of H are continuous, see [3, 52] and also [48]. This continuity then guarantees that the positive ground state is bounded away from zero on compact sets.

The proof of Theorem 1.7 is given in Section 4. The main tool is an upper bound for the spacial decay of ground states of the approximating Schrödinger operators H_λ , see Definition 1.5, which is *uniform* in $\lambda > 0$.

Since it will be necessary to have a positive ground state for the non-existence proof, we cannot prove the absence of ground state under symmetry constraints which destroy the positivity of ground states, such as fermionic particle statistics. However, the existence proof still works under symmetry restrictions, see Remark 4.10.

In Appendix A we construct an explicit example of a family of potentials which exhibits all possible different scenarios.

Lastly, recall that the Kato–class K_d is given by all real–valued potentials V such that in dimension $d \geq 2$

$$\lim_{\alpha \downarrow 0} \sup_{|x| \in \mathbb{R}^d} \int_{|x-y| \leq \alpha} g_d(x-y) |V(y)| dy = 0, \quad (1.14)$$

where

$$g_d(x) := \begin{cases} |x|^{2-d} & \text{if } d \geq 3 \\ |\ln|x|| & \text{if } d = 2 \end{cases}. \quad (1.15)$$

The Kato class in one dimension is given by $K_1 := L_{loc,unif}^1(\mathbb{R})$, the space of uniformly locally integrable functions on \mathbb{R} . We say that the potential V is in the local Kato–class $K_{d,loc}$ if $V\mathbb{1}_K \in K_d$ for all compact sets $K \subset \mathbb{R}^d$. It is clear that $K_d \subset L_{loc,unif}^1(\mathbb{R}^d)$ and $K_{d,loc} \subset L_{loc}^1(\mathbb{R}^d)$. Moreover, it is well–known that any potential $V \in K_d$ is infinitesimally form small with respect to $-\Delta$, see [11].

Thus if $V = V_+ - V_-$ with $V_\pm \geq 0$, $V_+ \in K_{d,loc}$, and $V_- \in K_d$ then all of the claims of Assumption 1.1 hold. This class of potentials is large enough to include most, if not all, physically relevant potentials, except maybe for some highly oscillatory potentials.

2. DEFINITIONS AND PREPARATIONS

Assume that $V_\pm \in L_{loc}^1(\mathbb{R}^d)$ and (1.2) holds for V_- . The KLMN theorem [47, 53] then shows that there exists a unique self-adjoint operator H corresponding to a quadratic form

$$\langle \psi, H\psi \rangle := \langle \nabla \psi, \nabla \psi \rangle + \langle \sqrt{V_+} \psi, \sqrt{V_+} \psi \rangle - \langle \sqrt{V_-} \psi, \sqrt{V_-} \psi \rangle \quad (2.1)$$

with the usual slight abuse of notation. Here $\psi \in \mathcal{Q}(H) := H^1(\mathbb{R}^d) \cap \mathcal{Q}(V_+)$, the form domain of H , where $H^1(\mathbb{R}^d)$ is the usual L^2 based Sobolev space of functions $\psi \in L^2(\mathbb{R}^d)$ whose weak (distributional) gradient $\nabla \psi \in L^2(\mathbb{R}^d)$, and

$$\mathcal{Q}(V_+) := \mathcal{D}(\sqrt{V_+}) = \{ \psi \in L^2(\mathbb{R}^d) : \sqrt{V_+} \psi \in L^2(\mathbb{R}^d) \} \quad (2.2)$$

is the quadratic form domain of the multiplication operator V_+ .

Since $\sqrt{V_+} \in L_{loc}^2$ we clearly have $\mathcal{C}_0^\infty(\mathbb{R}^d) \subset \mathcal{Q}(H)$. Note that $\mathcal{C}_0^\infty(\mathbb{R}^d)$ is a form core, i.e., dense in $H^1(\mathbb{R}^d) \cap \mathcal{D}(\sqrt{V_+})$ with respect to the norm

$$\|\psi\|_1 := (\|\psi\|_{H^1}^2 + \|\sqrt{V_+} \psi\|^2)^{1/2}, \quad (2.3)$$

see [11, 36]. In addition, Friedrich's extension theorem, see for example [53, Theorem 2.13], implies that the operator H and its domain $\mathcal{D}(H)$ are given by

$$\begin{aligned} D(H) &= \{\psi \in H^1(\mathbb{R}^d) \cap \mathcal{Q}(V_+) : (-\Delta + V)_{\text{distr}} \psi \in L^2(\mathbb{R}^d)\} \\ H\psi &= (-\Delta + V)_{\text{distr}} \psi \end{aligned} \quad (2.4)$$

where $(-\Delta + V)_{\text{distr}} \psi$ is in the sense of distributions when acting on $\psi \in L^2(\mathbb{R}^d)$.

Remark 2.1. If $V, W \in L^1_{\text{loc}}$ and V_- is form small and $|W|$ is form bounded with respect to $-\Delta + V_+$, i.e., (1.2) holds for V_- for some $0 \leq a_1 < 1, b \geq 0$ and it also holds with V_- replaced by $|W|$ for some $a_2, b_2 \geq 0$, then

$$\|\sqrt{V_-} \psi\|^2 + \lambda \|\sqrt{|W|} \psi\|^2 \leq (a_1 + \lambda a_2) (\|\nabla \psi\|^2 + \|\sqrt{V_+} \psi\|^2) + (b_1 + \lambda b_2) \|\psi\|^2 \quad (2.5)$$

for all $\psi \in H^1 \cap \mathcal{Q}(V_+)$. So for any $0 < \lambda_0 < (1 - a_1)/a_2$, we can construct the family of Schrödinger operators H_λ as the unique self-adjoint operator given by the quadratic forms

$$\langle \psi, H_\lambda \psi \rangle := \langle \nabla \psi, \nabla \psi \rangle + \langle \psi, V_+ \psi \rangle - \langle \psi, V_- \psi \rangle - \lambda \langle \psi, W \psi \rangle. \quad (2.6)$$

with quadratic form domain $\mathcal{Q}(H_\lambda) = H^1(\mathbb{R}^d) \cap \mathcal{Q}(V_+) = \mathcal{Q}(H)$ for all $0 \leq \lambda \leq \lambda_0$. For $\lambda = 0$ one recovers H . If W is infinitesimally form bounded w.r.t. $-\Delta + V_+$, then $\lambda_0 = \infty$.

One can relax the conditions on V to hold only on a connected, open set $U \subset \mathbb{R}^d$, which contains infinity. In this case one assumes $V_+ \in L^1_{\text{loc}}(U)$, and (1.2) holds for all $\psi \in H^1_0(U) \cap \mathcal{Q}^U(V_+)$, where $H^1_0(U)$ is the usual Sobolev space with Dirichlet boundary conditions on the boundary ∂U and $\mathcal{Q}^U(V_+) = \{\psi \in L^2(U) : \sqrt{V_+} \psi \in L^2(U)\}$. In this case H is the Schrödinger operator (with Dirichlet boundary conditions) defined by the quadratic form (2.1) which is restricted to $\psi \in \mathcal{Q}^U(H) = H^1_0(U) \cap \mathcal{Q}^U(V_+)$. Again it is well known that $\mathcal{C}^\infty_0(U)$ is dense in $\mathcal{Q}^U(H)$ w.r.t. the norm given in (2.3). The same holds for H_δ and any $\delta > 0$ small enough.

Now assume that the real-valued potential $V \in L^1_{\text{loc}}(\mathbb{R}^d)$, that its negative part V_- is form small w.r.t. $-\Delta + V_+$, i.e., (1.2) holds, and let H be the associated Schrödinger operator defined by quadratic form methods as above. For an open set $U \subset \mathbb{R}^d$ we consider weak (local) eigenfunctions of H at energy E , i.e., (weak local) solutions of the Schrödinger equation

$$H\psi = E\psi \quad \text{in } U. \quad (2.7)$$

We are mainly interested in the case that $E = 0$.

With a slight abuse of notation, we denote by $\langle \varphi, H\psi \rangle$ the sesquilinear form given by

$$\langle \varphi, H\psi \rangle := \langle \nabla \varphi, \nabla \psi \rangle + \langle \varphi, V\psi \rangle = \int (\overline{\nabla \varphi} \cdot \nabla \psi + \overline{\varphi} V \psi) dx \quad (2.8)$$

whenever the right hand sides makes sense. This is the case if $\varphi, \psi \in \mathcal{Q}(H) = H^1(\mathbb{R}^d) \cap \mathcal{Q}(V_+)$ but also if, $\varphi \in \mathcal{Q}^U_c(H)$ and $\psi \in \mathcal{Q}^U_{\text{loc}}(H)$, where for some open set $U \subset \mathbb{R}^d$ the local quadratic form domain

$$\mathcal{Q}^U_{\text{loc}}(H) := \{\psi \in L^2_{\text{loc}}(U) : \chi \psi \in \mathcal{Q}(H) \text{ for all } \chi \in \mathcal{C}^\infty_0(U)\} \quad (2.9)$$

is the vector space of functions which are locally (in U) in the quadratic form domain of H . Moreover,

$$\mathcal{Q}^U_c(H) := \{\psi \in \mathcal{Q}(H) : \text{supp}(\psi) \subset U \text{ is compact}\} \quad (2.10)$$

is the set of functions in $\mathcal{Q}(H)$ with compact support inside U . If $\varphi \in \mathcal{Q}_c^U(H)$ and $\psi \in \mathcal{Q}_{loc}^U(H)$, then the integral on the right-hand-side of (2.8) can be restricted to the set U . Clearly, $\mathcal{Q}_{loc}^U(H) = \{\psi \in H_{loc}^1(U) : \sqrt{V_+}\psi \in L_{loc}^2(U)\} = H_{loc}^1(U) \cap \mathcal{Q}_{loc}^U(V_+)$.

Similarly, one can define the local domain of H , relative to some open set $U \subset \mathbb{R}^d$, by

$$\mathcal{D}_{loc}^U(H) := \{\psi \in L_{loc}^2(U) : \chi\psi \in \mathcal{D}(H) \text{ for all } \chi \in \mathcal{C}_0^\infty(U)\}. \quad (2.11)$$

Remark 2.2. Note that the definitions of $\mathcal{Q}_{loc}^U(H)$ and $\mathcal{D}_{loc}^U(H)$ are consistent in the sense that for any $\chi \in \mathcal{C}_0^\infty(U)$ one has $\chi\psi \in \mathcal{Q}(H)$ (even $\chi\varphi \in \mathcal{Q}_c^U(H)$) for any $\psi \in \mathcal{Q}(H)$ and $\chi\psi \in \mathcal{D}(H)$ for any $\psi \in \mathcal{D}(H)$. This is clear when $\psi \in \mathcal{Q}_{loc}^U(H) = H_{loc}^1(U) \cap \mathcal{Q}_{loc}^U(V_+)$ since for $\chi \in \mathcal{C}_0^\infty(U)$ we have $\chi\psi \in H^1(\mathbb{R}^d)$ for any $\psi \in H_{loc}^1(U)$ and $\chi\psi \in \mathcal{Q}(V_+)$ for any $\psi \in \mathcal{Q}_{loc}^U(V_+)$. In addition, if $\psi \in \mathcal{D}(H)$ then

$$(-\Delta + V)_{\text{distr}} \chi\psi = \chi(-\Delta + V)_{\text{distr}} \psi - 2\nabla\chi\nabla\psi - (\Delta\chi)\psi \in L^2(\mathbb{R}^d),$$

so $\chi\psi \in \mathcal{D}(H)$. Moreover, with $\mathcal{C}^\infty(U)$ the infinitely differentiable functions on U , it is easy to see that

$$\mathcal{C}^\infty(U) \subset \mathcal{Q}_{loc}^U(H). \quad (2.12)$$

since $\mathcal{C}_0^\infty(U) \subset \mathcal{Q}^U(H)$. However, the inclusion $\mathcal{C}^\infty(U) \subset \mathcal{D}_{loc}^U(H)$ is wrong in general, since the construction of the Schrödinger operator H with the help of quadratic forms allows for rather singular potentials V .

Thus we define weak solutions, supersolutions and subsolutions of (2.7) in the following quadratic form sense.

Definition 2.3. a) u is a (weak) eigenfunction of the Schrödinger operator H with energy E if $u \in \mathcal{Q}(H)$ and

$$\langle \varphi, (H - E)u \rangle = 0 \quad (2.13)$$

for every $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d)$.

b) u is a (weak) local eigenfunction of the Schrödinger operator H with energy E in $U \subset \mathbb{R}^d$ if $u \in \mathcal{Q}_{loc}^U(H)$ and

$$\langle \varphi, (H - E)u \rangle = 0 \quad (2.14)$$

for every $\varphi \in \mathcal{C}_0^\infty(U)$.

c) u is a supersolution of the Schrödinger operator H with energy E in $U \subset \mathbb{R}^d$ if $u \in \mathcal{Q}_{loc}^U(H)$ and

$$\langle \varphi, (H - E)u \rangle \geq 0 \quad (2.15)$$

for every nonnegative $\varphi \in \mathcal{C}_0^\infty(U)$.

d) u is a subsolution of the Schrödinger operator H with energy E in $U \subset \mathbb{R}^d$ if $u \in \mathcal{Q}_{loc}^U(H)$ and

$$\langle \varphi, (H - E)u \rangle \leq 0 \quad (2.16)$$

for every nonnegative $\varphi \in \mathcal{C}_0^\infty(U)$.

Remark 2.4. Using the density of \mathcal{C}_0^∞ in $\mathcal{Q}(H)$ it is easy to see that once (2.13) holds for all $\varphi \in \mathcal{C}_0^\infty$, it holds for all $\varphi \in \mathcal{Q}(H)$. Similarly, (2.14) holds for all $\varphi \in \mathcal{Q}_c^U(H)$, and (2.15), respectively (2.16), hold for all $0 \leq \varphi \in \mathcal{Q}_c^U(H)$.

One should note that one does not have to distinguish between weak eigenfunctions and eigenfunctions and similarly for local eigenfunctions.

Lemma 2.5. *Every weak eigenfunction $u \in \mathcal{Q}(H)$ of H is in $\mathcal{D}(H)$ given by (2.4). Similarly, if $u \in \mathcal{Q}_{loc}^U(H)$ is a weak local eigenfunction of H in an open domain $U \subset \mathbb{R}^d$, then u is locally in the domain of H , i.e., $u \in \mathcal{D}_{loc}^U(H)$ given by (2.11).*

Proof. This is probably a standard argument for weak eigenfunctions, but not standard for weak local eigenfunctions. Let $f \in L^2(\mathbb{R}^d)$ and ψ be a weak solution of the equation $H\psi = f$, i.e.,

$$\langle \varphi, H\psi \rangle = \langle \varphi, f \rangle \quad (2.17)$$

for all $\varphi \in \mathcal{Q}(H)$. Then for any $\lambda \in \mathbb{R}$ we have

$$\langle \varphi, (H + \lambda)\psi \rangle = \langle \varphi, \lambda\psi + f \rangle \quad (2.18)$$

for all $\varphi \in \mathcal{Q}(H)$. Since H is bounded from below, all large enough λ will be in the resolvent set of H . So for all large enough λ we can choose $\varphi = (H + \lambda)^{-1}\xi$, with $\xi \in L^2(\mathbb{R}^d)$ in (2.18) to get

$$\langle \xi, \psi \rangle = \langle (H + \lambda)^{-1}\xi, \lambda\psi + f \rangle = \langle \xi, (H + \lambda)^{-1}(\lambda\psi + f) \rangle. \quad (2.19)$$

This holds for all $\xi \in L^2(\mathbb{R}^d)$, so

$$\psi = (H + \lambda)^{-1}(\lambda\psi + f) \in \mathcal{D}(H), \quad (2.20)$$

since $\psi, f \in L^2(\mathbb{R}^d)$ and the resolvent $(H + \lambda)^{-1}$ maps $L^2(\mathbb{R}^d)$ onto $\mathcal{D}(H)$.

Note that if ψ is a weak eigenfunction of H , at energy E , then we can use $f = E\psi$. Thus weak eigenfunctions are eigenfunctions in the domain of H .

Now assume that $f \in L_{loc}^2(U)$ and $\psi \in \mathcal{Q}_{loc}^U(H)$ is a weak local solution of

$$\langle \varphi, H\psi \rangle = \langle \varphi, f \rangle \quad (2.21)$$

for all $\varphi \in \mathcal{Q}_c^U(H)$. Take any $\chi \in C_0^\infty(U)$. Replacing φ by $\bar{\chi}\varphi$ in (2.21) one sees that

$$\langle \bar{\chi}\varphi, H\psi \rangle = \langle \varphi, \chi f \rangle \quad (2.22)$$

for all $\varphi \in \mathcal{Q}(H)$. Using that $\chi, \nabla\chi$, and $\Delta\chi$ have compact supports, a straightforward calculation shows

$$\langle \nabla(\bar{\chi}\varphi), \nabla\psi \rangle = \langle \nabla\varphi, \nabla(\chi\psi) \rangle + \langle \varphi, (\Delta\chi + 2\nabla\chi\nabla)\psi \rangle.$$

Using this and the definition (2.1) of the quadratic form in (2.22) yields

$$\langle \varphi, H\chi\psi \rangle = \langle \varphi, \chi f - (\Delta\chi + 2\nabla\chi\nabla)\psi \rangle \quad (2.23)$$

for all $\varphi \in \mathcal{Q}(H)$. Adding again $\langle \varphi, \lambda\chi\psi \rangle$ on both sides and choosing $\varphi = (H + \lambda)^{-1}\xi$ with λ large that enough, one sees that

$$\langle \xi, \chi\psi \rangle = \langle \xi, (H + \lambda)^{-1}(\chi(\lambda\psi + f) - (\Delta\chi)\psi + 2\nabla\chi\nabla\psi) \rangle \quad (2.24)$$

for all $\xi \in L^2(\mathbb{R}^d)$. Hence

$$\chi\psi = (H + \lambda)^{-1}(\chi(\lambda\psi + f) - (\Delta\chi)\psi + 2\nabla\chi\nabla\psi) \in \mathcal{D}(H), \quad (2.25)$$

for any $\chi \in C_0^\infty(U)$. Thus $\psi \in \mathcal{D}_{loc}^U(H)$. Again, using $f = E\psi$ shows that any weak local eigenfunctions of H at energy E is locally in the domain of H . \blacksquare

Finally let us note that the definition of critical potential and virtual levels are rather natural. It is easy to see that any potential which creates a zero energy ground state is critical.

Lemma 2.6. *Assume that $V \in L^1_{loc}(\mathbb{R}^d)$ and that V_- is form small and $0 \leq W$ is infinitesimally form small w.r.t. $-\Delta + V_+$. Furthermore let H and H_λ , $0 < \lambda \leq \lambda_0$, be the associated Schrödinger operators, see Remark 2.1. Assume also that $\sigma(H) = \sigma_{ess}(H_\lambda) = [0, \infty)$ and that H has a zero energy ground state. Then the potential V is critical.*

Proof. This is probably well-known. We provide the short proof for the convenience of the reader. Let ψ be a zero energy normalized ground state of H . Then for any small enough $\lambda > 0$,

$$\langle \psi, H_\lambda \psi \rangle = \langle \psi, H \psi \rangle - \lambda \langle \psi, W \psi \rangle = -\lambda \langle \psi, W \psi \rangle < 0$$

since we can choose the ground state $\psi > 0$. Thus as soon as $\sigma_{ess}(H_\delta) = [0, \infty)$, the min-max principle shows that H_δ has eigenvalues below zero. \blacksquare

The converse to Lemma 2.6 does not hold. See the example from Appendix A.

Our proofs of Theorems 1.3 and 1.7 rely on the so-called subharmonic comparison lemma which has already seen wide use in the study of the asymptotic decay of eigenfunctions of Schrödinger operators, see, e.g., [12, 21]. We use the version of [2, Theorem 2.7] since it allows for a quadratic form version which needs only minimal regularity assumptions.

Theorem 2.7 (Agmon's version of the comparison principle). *Let w be a positive supersolution of the Schrödinger operator H at energy E in a neighborhood of infinity $U_R := \{x \in \mathbb{R}^d : |x| > R\}$. Let v be a subsolution of H at energy E in U_R . Suppose that*

$$\liminf_{N \rightarrow \infty} \left(\frac{1}{N^2} \int_{N \leq |x| \leq \alpha N} |v|^2 dx \right) = 0 \quad (2.26)$$

for some $\alpha > 1$. If for some $\delta > 0$ and $0 \leq C < \infty$ one has

$$v(x) \leq Cw(x) \text{ on the annulus } R < |x| \leq R + \delta, \quad (2.27)$$

then

$$v(x) \leq Cw(x) \text{ for all } x \in U_R. \quad (2.28)$$

Remark 2.8. We note that the condition (2.26) is trivially satisfied as soon as $v \in L^2(\mathbb{R}^d)$, but it also allows for subsolutions v which are not square integrable at infinity. This is crucial for the proof of our non-existence result. A slight extension of Agmon's comparison principle, which allows to relax the continuity assumptions and works for domains U which are not necessarily neighborhoods of infinity, is derived in [22].

Remark 2.9. Agmon also assumes that the supersolution w and the subsolution v are continuous in $\overline{U_R} = \{|x| \geq R\}$ in [2]. However, the form of Theorem 2.7 is what is really proven in [2], see also [22]. The additional assumption that the supersolution $w > 0$ and the subsolution v are continuous in $\{|x| \geq R\}$ are only made in [2] to guarantee that (2.27) holds with constant $C = c_2/c_1$ where $c_1 := \inf_{R \leq |x| \leq R+\delta} w(x) > 0$ and $c_2 := \sup_{R \leq |x| \leq R+\delta} |v(x)| < \infty$, by continuity, for arbitrary $\delta > 0$.

Before we give the proofs of Theorems 1.3 and 1.7, we sketch the simple proof of the $m = 0$ version of Theorem 1.3 using the comparison theorem:

For $\gamma > 0$ set $\psi_\gamma(x) = |x|^{-\gamma}$, so $\psi_\gamma \in C^\infty(U_R)$ and all $R > 0$. A short calculation shows $\Delta\psi_\gamma(x) = \gamma(\gamma + 2 - d)|x|^{-\gamma-2}$ for $x \neq 0$. Hence with

$$W_\gamma(x) = \frac{\Delta\psi_\gamma(x)}{\psi_\gamma(x)} = \gamma(\gamma + 2 - d)|x|^{-2} \quad (2.29)$$

one sees that $(-\Delta + W_\gamma(x))\psi_\gamma(x) = 0$ for $x \neq 0$. Since $\psi_\gamma \in C^\infty(\mathbb{R}^d \setminus \{0\})$, integration by parts shows that ψ_γ is a weak local eigenfunction of $-\Delta + W_\gamma$ in the sense of Definition 2.3 in the open sets U_R for any $R > 0$.

Moreover, Remark 2.2 shows that $\psi_\gamma \in \mathcal{Q}_{loc}^{U_R}(H)$ for any $R > 0$ and any Schrödinger operator H with potential $V \in L^1_{loc}(\mathbb{R}^d)$ for which V_- is form small w.r.t. $-\Delta + V_+$. Thus if

$$V(x) \leq W_\gamma(x) \quad (2.30)$$

for some $\gamma > 0$ and all large enough $|x| > R$, then

$$\langle \varphi, H\psi_\gamma \rangle \leq \langle \varphi, (-\Delta + W_\gamma)\psi_\gamma \rangle = 0 \quad (2.31)$$

for all $0 \leq \varphi \in C_0^\infty(U_R)$, i.e., ψ_γ is a zero energy subsolution of H . One easily checks that

- a) $\psi_\gamma \notin L^2(U_R)$ if and only if $0 < \gamma \leq d/2$.
- b) ψ_γ satisfies (2.26) if and only if $\gamma > (d-2)/2$.
- c) $0 < \gamma \mapsto W_\gamma$ is increasing if and only if $\gamma > (d-2)/2$.

The last part shows that one should choose γ as large as possible in order to guarantee that (2.31) holds.

Now assume that $H \geq 0$ has zero as an eigenvalue with corresponding unique ground state ψ which can be chosen to be positive [15, 18]. Since V is locally in the Kato class, one also knows that ψ is continuous, [52]. If

$$V(x) \leq \frac{d(4-d)}{4|x|^2} = W_{d/2}(x) \quad (2.32)$$

then the above discussion shows that $u = \psi_{d/2}$ is a zero energy subsolution of H which is not square integrable at infinity but for which (2.26) holds. Using $c_R^1 = \inf_{R \leq |x| \leq R+1} \psi(x) > 0$, $c_R^2 = \sup_{R \leq |x| \leq R+1} u(x)$, and $C = c_R^2/c_R^1$ ensures $u(x) \leq C\psi(x)$ for all $R \leq |x| \leq R+1$. Then Theorem 2.7 shows that

$$u(x) \leq C\psi(x) \quad (2.33)$$

for all $|x| > R$, in particular, $\psi \neq L^2(\mathbb{R}^d)$, hence ψ is not an eigenfunction. Thus zero is not an eigenvalue of the Schrödinger operator H , which proves the $n = 0$ version of Theorem 1.3.

Remark 2.10. Note that for $\gamma > d/2$ the function ψ_γ is in $L^2(U_R)$ for any $R > 0$, so $\gamma = d/2$ is the largest possible choice in order to get the non-existence result. The higher order condition for non-existence will have to use a suitably modified choices of subsolutions at $\gamma = d/2$.

Remark 2.11. Choosing $\gamma = (d-2)/2$ yields the Hardy potential $W_{(d-2)/2}(x) = -\frac{(d-2)^2}{4|x|^2}$. It is well known that a Schrödinger operator with a Hardy potential is nonnegative. It is curious that for the absence of zero energy eigenfunctions the choice $\gamma = d/2$ becomes relevant.

For the existence result we want to reverse the roles of the eigenfunction ψ and ψ_γ . If

$$V(x) \geq \frac{d(4-d) + \varepsilon}{4|x|^2} \quad (2.34)$$

for all $|x| > R$ and some $\varepsilon > 0$ then ψ_γ is a zero energy subsolution of H in U_R where $\gamma > d/2$ is the unique solution of $\gamma(\gamma + 2 - d) = (d(4-d) + \varepsilon)/4$. Arguing as above, one sees that any positive zero energy ground state ψ of H satisfies the upper bound

$$\psi(x) \leq C\psi_\gamma(x) \quad (2.35)$$

for all $|x| > R$, hence it is square integrable at infinity since $\psi_\gamma \in L^2(U_R)$ as soon as $\gamma > d/2$. Of course, this is a circular reasoning, since we need the existence of a square integrable bound state, or at least the existence of a local zero energy bound state which satisfies (2.26). The rigorous argument uses the fact that $H \geq 0$ is assumed to have a virtual level at zero, so the operators H_δ have a negative energy ground states with negative energy for small $\delta > 0$. These ground states will converge to a zero energy ground state of H in the limit $\delta \rightarrow 0$, see Section 4.

3. PROOF OF THE NON-EXISTENCE RESULT

Recall the iterated logarithms \ln_n defined by $\ln_1(r) := \ln(r)$ for $r > 0$ and, for $r > e_j$, inductively by $\ln_{j+1}(r) := \ln(\ln_j(r))$ when $j \in \mathbb{N}$. Here $e_1 = 1$ and $e_{j+1} = e^{e_j}$.

A convenient sequence of functions at the edge of L^2 -integrability near infinity is given by

$$\psi_{\ell,m}(x) := |x|^{-d/2} \prod_{j=1}^m \ln_j^{-1/2}(|x|) \quad \text{for } |x| > e_m. \quad (3.1)$$

As usual, the empty product is one, so $\psi_{\ell,0}(x) = |x|^{-d/2} = \psi_{d/2}(x)$. We still have $\psi_{\ell,m} \in \mathcal{C}^\infty(\{|x| > e_m\})$, in particular, $\psi_{\ell,m} \in \mathcal{Q}_{loc}^{U_R}(H)$, for $R \geq e_m$ and any Schrödinger operator constructed via quadratic form methods as in Section 2. In order to mimic the proof sketched at the end of Section 2, we need to know the potential W_m for which $(-\Delta + W_m)\psi_{\ell,m} = 0$ in $U_{e_m} = \{|x| > e_m\}$. This is a bit more complicated than the previous calculation for ψ_γ .

Lemma 3.1. *For any $m \in \mathbb{N}_0$ we have $(-\Delta + W_m)\psi_{\ell,m} = 0$ in $U_R = \{x \in \mathbb{R}^d : |x| > R\}$, for all large enough $R \geq e_m$, where*

$$\begin{aligned} W_m(x) := & \frac{d(4-d)}{4|x|^2} + \frac{1}{|x|^2} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) + \frac{1}{4|x|^2} \left(\sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) \right)^2 \\ & + \frac{1}{2|x|^2} \sum_{j=1}^m \sum_{l=1}^j \prod_{s=1}^l \prod_{t=1}^j \ln_s^{-1}(|x|) \ln_t^{-1}(|x|) \end{aligned} \quad (3.2)$$

is well-defined for $|x| > e_m$.

Proof. Clearly, if $W_m = \frac{\Delta\psi_{\ell,m}}{\psi_{\ell,m}}$ then $(-\Delta + W_m)\psi_{\ell,m} = 0$ in U_R , for all large enough $R > 0$. For any radial function depending only on the radius $r = |x|$ we have

$$\Delta\psi(x) = \partial_r^2\psi(x) + \frac{d-1}{|x|}\partial_r\psi(x). \quad (3.3)$$

By a straightforward but slightly tedious calculation one sees that

$$\begin{aligned}\partial_r \psi_{\ell,m}(x) &= -\psi_{\ell,m}(x) \left(\frac{d}{2|x|} + \frac{1}{2|x|} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) \right), \\ \partial_r^2 \psi_{\ell,m}(x) &= \psi_{\ell,m}(x) \left(\frac{d}{2|x|} + \frac{1}{2|x|} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) \right)^2 + \psi_{\ell,m}(x) \frac{d}{2|x|^2} \\ &\quad + \psi_{\ell,m}(x) \left(\frac{1}{2|x|^2} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) + \frac{1}{2|x|^2} \sum_{j=1}^m \sum_{l=1}^j \prod_{s=1}^l \prod_{t=1}^j \ln_s^{-1}(|x|) \ln_t^{-1}(|x|) \right)\end{aligned}$$

where we used

$$\partial_r \ln_1(r) = \frac{1}{r} \quad \text{and} \quad \partial_r \ln_j(r) = \frac{1}{\ln_{j-1}(r)} \frac{1}{\ln_{j-2}(r)} \cdots \frac{1}{\ln_1(r)} \frac{1}{r}.$$

Thus

$$\begin{aligned}W_m(x) &= \frac{\Delta \psi_{\ell,m}(x)}{\psi_{\ell,m}(x)} = \frac{d(4-d)}{4|x|^2} + \frac{1}{|x|^2} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) + \frac{1}{4|x|^2} \left(\sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) \right)^2 \\ &\quad + \frac{1}{2|x|^2} \sum_{j=1}^m \sum_{l=1}^j \prod_{s=1}^l \prod_{t=1}^j \ln_s^{-1}(|x|) \ln_t^{-1}(|x|)\end{aligned}$$

and we have $(-\Delta + W_m)\psi_{\ell,m} = 0$ in U_R as long as $R > 0$ is large enough, so that the iterated logarithms are well-defined. \blacksquare

Remark 3.2. Alternatively, one can compute $W_m = \frac{\Delta \psi_{\ell,m}}{\psi_{\ell,m}}$ inductively. For radial functions f, g , i.e., with the usual abuse of notation $f(x) = f(r)$ and $g(x) = g(r)$ for $r = |x|$ we have

$$\Delta(gf) = f\Delta g + 2\partial_r f \partial_r g + (\Delta f)g. \quad (3.4)$$

Using (3.4) and $\psi_{\ell,m+1}(x) = \psi_{\ell,m}(x) \ln_{m+1}^{-\frac{1}{2}}(|x|)$ we obtain

$$W_{m+1}(x) = \frac{\Delta \psi_{\ell,m+1}(x)}{\psi_{\ell,m+1}(x)} = W_m + \frac{\Delta \ln_{m+1}^{-\frac{1}{2}}(|x|)}{\ln_{m+1}^{-\frac{1}{2}}(|x|)} + 2 \frac{\partial_r \psi_{\ell,m}(x)}{\psi_{\ell,m}(x)} \frac{\partial_r \ln_{m+1}^{-\frac{1}{2}}(|x|)}{\ln_{m+1}^{-\frac{1}{2}}(|x|)}.$$

A straightforward calculation yields

$$\begin{aligned}
 \frac{\Delta \ln_{m+1}^{-\frac{1}{2}}(|x|)}{\ln_{m+1}^{-\frac{1}{2}}(|x|)} &= \frac{1}{4|x|^2} \prod_{k=1}^{m+1} \ln_k^{-2}(|x|) + \frac{2-d}{2|x|^2} \prod_{k=1}^{m+1} \ln_k^{-1}(|x|) \\
 &\quad + \frac{1}{2|x|^2} \sum_{j=1}^{m+1} \prod_{s=1}^{m+1} \prod_{t=1}^j \ln_s^{-1}(|x|) \ln_t^{-1}(|x|), \\
 2 \frac{\partial_r \psi_{\ell,m}(x)}{\psi_{\ell,m}(x)} \frac{\partial_r \ln_{m+1}^{-\frac{1}{2}}(|x|)}{\ln_{m+1}^{-\frac{1}{2}}(|x|)} &= \left(\frac{d}{|x|} + \frac{1}{|x|} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) \right) \left(\frac{1}{2|x|} \prod_{s=1}^{m+1} \ln_s^{-1}(|x|) \right), \\
 W_{m+1}(x) &= W_m(x) + \frac{3}{4|x|^2} \prod_{k=1}^{m+1} \ln_k^{-2}(|x|) + \frac{1}{|x|^2} \prod_{k=1}^{m+1} \ln_k^{-1}(|x|) \\
 &\quad + \frac{1}{|x|^2} \sum_{j=1}^m \prod_{s=1}^{m+1} \prod_{t=1}^j \ln_s^{-1}(|x|) \ln_t^{-1}(|x|).
 \end{aligned}$$

Since $W_0(x) = \frac{d(4-d)}{4|x|^2}$, this yields W_m via induction.

Now we come to the

Proof of Theorem 1.3: Using Lemma 3.1 and $\psi_{\ell,m} \in \mathcal{Q}_{loc}^{U_R}(H)$ one sees that

$$\begin{aligned}
 \langle \varphi, H\psi_{\ell,m} \rangle &= \langle \nabla \varphi, \nabla \psi_{\ell,m} \rangle + \langle \varphi, V\psi_{\ell,m} \rangle \leq \langle \nabla \varphi, \nabla \psi_{\ell,m} \rangle + \langle \varphi, W_m \psi_{\ell,m} \rangle \\
 &= \langle \varphi, (-\Delta + W_m)\psi_{\ell,m} \rangle = 0
 \end{aligned} \tag{3.5}$$

for all $0 \leq \varphi \in C_0^\infty(U_R)$ as soon as $V \leq W_m$ in U_R . So $\psi_{\ell,m}$ is a zero energy subsolution of H in U_R as soon as $V \leq W_m$ in U_R . Since V is in the local Kato class we know from [3, 48, 52] that any eigenfunction of H is continuous. Moreover, it is well-known that the ground state eigenfunction can be chosen to be strictly positive [15, 18, 47].

So if $\psi > 0$ is a zero energy ground state of H then, with $c_R^1 = \inf_{R \leq |x| \leq R+1} \psi(x) > 0$ and $c_R^2 = \sup_{R \leq |x| \leq R+1} \psi_{\ell,m}(x) < \infty$, we can set $C := c_R^2/c_R^1$ to see that $\psi_{\ell,m}(x) \leq C\psi(x)$ for $R \leq |x| \leq R+1$. Moreover, ψ being a zero energy solution is also a zero energy supersolution, so Theorem 2.7 shows that

$$\psi_{\ell,m}(x) \leq C\psi(x) \tag{3.6}$$

for all $|x| > R$. In particular, ψ cannot be square integrable as soon as $V \leq W_m$ on U_R for some large enough $R > 0$, since $\psi_{\ell,m}$ is positive and not in $L^2(U_R)$.

Finally, setting $V_m(x) = \frac{d(4-d)}{4|x|^2} + \frac{1}{|x|^2} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(x)$, we note that $W_m(x) \geq V_m(x)$ for all large enough $|x|$. This proves Theorem 1.3. \blacksquare

Remark 3.3. Of course, the proof of Theorem 1.3 given above shows that if

$$V(x) \leq W_m(x) \quad \text{for all } |x| > R \tag{3.7}$$

for large enough $R > 0$ and some $m \in \mathbb{N}_0$, then the Schrödinger operator H cannot have any zero energy ground state.

4. PROOF OF THE EXISTENCE RESULT

For the existence result we need to modify our comparison functions $\psi_{\ell,m}$ to make them barely square integrable near infinity. Given an arbitrary $\varepsilon > 0$ we set

$$\psi_{u,m,\varepsilon}(x) := \psi_{\ell,m}(x) \ln_m^{-\varepsilon/2}(|x|) \quad (4.1)$$

where $\psi_{\ell,m}$ is defined in (3.1). For each $m \in \mathbb{N}_0$ and $\varepsilon > 0$ we have $\psi_{u,m,\varepsilon} \in \mathcal{C}^\infty(\{|x| > e_m\})$ and it is not hard to see that for any $\varepsilon > 0$ and any large enough $R > 0$ the function $\psi_{u,m,\varepsilon}$ is barely in $L^2(U_R)$. The potential $Y_{m,\varepsilon}$ for which $(-\Delta + Y_{m,\varepsilon})\psi_{u,m,\varepsilon} = 0$ in $U_R = \{x \in \mathbb{R}^d : |x| > R\}$ is given by

Lemma 4.1. *For any $m \in \mathbb{N}_0$ we have $(-\Delta + Y_{m,\varepsilon})\psi_{u,m,\varepsilon} = 0$ in $U_{e_m} = \{x \in \mathbb{R}^d : |x| > e_m\}$, where the potential $Y_{m,\varepsilon}$ is given by*

$$\begin{aligned} Y_{m,\varepsilon}(x) &= W_m(x) + \frac{\varepsilon^2}{4|x|^2} \prod_{k=1}^n \ln_k^{-2}(|x|) + \frac{\varepsilon}{|x|^2} \prod_{k=1}^n \ln_k^{-1}(|x|) \\ &\quad + \frac{\varepsilon}{|x|^2} \sum_{j=1}^n \prod_{k=1}^n \prod_{m=1}^j \ln_k^{-1}(|x|) \ln_m^{-1}(|x|), \end{aligned} \quad (4.2)$$

with W_m given in (3.2).

Proof. As in the proof of Lemma 3.1 we have to calculate $Y_{m,\varepsilon} := \frac{\Delta \psi_{u,m,\varepsilon}}{\psi_{u,m,\varepsilon}}$. We use (3.4) to see that

$$\begin{aligned} Y_{m,\varepsilon}(x) &= \frac{\Delta \psi_{\ell,m}(x)}{\psi_{\ell,m}(x)} + \frac{\Delta \ln_{m+1}^{-\frac{\varepsilon}{2}}(|x|)}{\ln_{m+1}^{-\frac{\varepsilon}{2}}(|x|)} + 2 \frac{\partial_r \ln_{m+1}^{-\frac{\varepsilon}{2}}(|x|)}{\ln_{m+1}^{-\frac{\varepsilon}{2}}(|x|)} \frac{\partial_r \psi_{\ell,m}(|x|)}{\psi_{\ell,m}(|x|)} \\ &= W_m(x) + \frac{\varepsilon^2}{4|x|^2} \prod_{k=1}^m \ln_k^{-2}(|x|) + \frac{\varepsilon}{|x|^2} \prod_{k=1}^m \ln_k^{-1}(|x|) \\ &\quad + \frac{\varepsilon}{|x|^2} \sum_{j=1}^m \prod_{s=1}^m \prod_{t=1}^j \ln_s^{-1}(|x|) \ln_t^{-1}(|x|) \end{aligned}$$

which is (4.2). ■

We want to show that ground states of Schrödinger operators H with critical potentials V exist using suitable eigenfunctions of H_λ . For this the following is convenient.

Lemma 4.2. *Assume that the potential V satisfies Assumption 1.1 and W is a positive potential which is infinitesimally form small w.r.t. $-\Delta + V_+$. Let $(H_\lambda)_{\lambda \geq 0}$ be the family of Schrödinger operators constructed in Remark 2.1. Moreover, assume that there exists a sequence $0 < \lambda_n \rightarrow 0$ as $n \rightarrow \infty$ such that the operators $H_n = H_{\lambda_n}$ have eigenvalues $E_n = E_{\lambda_n}$ with corresponding normalized weak eigenfunctions $\psi_n = \psi_{\lambda_n}$. If*

- a) *the sequence of eigenvalues $(E_n)_n$ of H_n is bounded from above and*
- b) *the sequence ψ_n is a Cauchy sequence in L^2 ,*

then the sequence ψ_n is Cauchy w.r.t. the quadratic form norm $\|\cdot\|_1$ given in (2.3), hence its limit $\psi = \lim_{n \rightarrow \infty} \psi_n \in \mathcal{Q}(H)$. Moreover $E = \lim_{n \rightarrow \infty} E_n$ exists and ψ is a normalized weak eigenfunction of H with eigenvalue E .

Remark 4.3. Lemma 2.5 shows even that $\psi \in \mathcal{D}(H)$. Moreover, we do not need that V is in the local Kato–class, only that V_- is relatively form small w.r.t. $-\Delta + V_+$.

Remark 4.4. We apply Lemma 4.2 when λ_n converges monotonically to zero and E_n is a ground state of H_n , in which case one can simplify the proof. For example, if E_n are ground state energies of H_n , then since as quadratic forms $H_\lambda \leq H_{\lambda'}$ for all $0 < \lambda' \leq \lambda \leq \lambda_0$, the limit $\lim_{n \rightarrow \infty} E_n$ exists, by monotonicity. However the result of Lemma 4.2 is needed when one considers not only ground states, but also excited states which hit the bottom of the essential spectrum.

Remark 4.5. Clearly any eigenvalue of H_n is bounded from below uniformly in $n \in \mathbb{N}$ since $H_n \geq H_{\lambda_{max}}$ with $\lambda_{max} = \max_n \lambda_n$ as quadratic forms for all $n \in \mathbb{N}$. In particular, all the eigenvalues E_n are bounded uniformly in $n \in \mathbb{N}$ once they are bounded from above.

Moreover, if the essential spectrum of H is not empty and E_n is an eigenvalue of H_n below the essential spectrum of H_n , then E_n is bounded from above, since as quadratic forms $H_n \leq H$ and by Persson’s theorem [37, 44]

$$\begin{aligned} \inf \sigma_{ess}(H_n) &= \liminf_{R \rightarrow \infty} \{ \langle \varphi, H_n \varphi \rangle : \varphi \in \mathcal{Q}(H), \|\varphi\| = 1, \text{supp}(\varphi) \subset \{|x| > R\} \} \\ &\leq \liminf_{R \rightarrow \infty} \{ \langle \varphi, H \varphi \rangle : \varphi \in \mathcal{Q}(H), \|\varphi\| = 1, \text{supp}(\varphi) \subset \{|x| > R\} \} = \inf \sigma_{ess}(H). \end{aligned} \quad (4.3)$$

Thus

$$-\infty < \inf \sigma(H_{\lambda_{max}}) \leq \inf \sigma(H_n) \leq E_n \leq \inf \sigma_{ess}(H_n) \leq \inf \sigma_{ess}(H)$$

for all $n \in \mathbb{N}$, which shows that $\sup_n |E_n| < \infty$ as soon as $\sigma_{ess}(H)$ is not empty.

Proof of Lemma 4.2: Let ψ_n be a normalized sequence of eigenfunctions of H_n with eigenvalue E_n which is also a Cauchy sequence in L^2 . In particular,

$$\langle \psi_n, H_n \psi_n \rangle = E_n \langle \psi_n, \psi_n \rangle = E_n. \quad (4.4)$$

Let $0 < a_1 < 1$ and $b_1 \geq 0$, respectively $a_2, b_2 \geq 0$, such that (1.2), respectively (1.2) with V_- replaced by W , holds. Then

$$\begin{aligned} \langle \psi_n, H_n \psi_n \rangle &= \|\nabla \psi_n\|^2 + \|\sqrt{V_+} \psi_n\|^2 - \|\sqrt{V_-} \psi_n\|^2 - \lambda_n \|\sqrt{W} \psi_n\|^2 \\ &\geq \|\nabla \psi_n\|^2 + \|\sqrt{V_+} \psi_n\|^2 - (a_1 + \lambda_n a_2) \|\nabla \psi_n\|^2 - (b_1 + \lambda_n b_2) \|\psi_n\|^2 \\ &\geq (1 - a_1 - \lambda_n a_2) \|\psi_n\|_1^2 - (b_1 + \lambda_n b_2), \end{aligned}$$

since ψ_n is normalized. We also used the quadratic form norm $\|\cdot\|_1$ given by (2.3). Using (4.4) this implies

$$(1 - a_1 - \lambda_n a_2) \|\psi_n\|_1^2 \leq b_1 + \lambda_n b_2 + E_n, \quad (4.5)$$

which shows that we have $\limsup_{n \rightarrow \infty} \|\psi_n\|_1 < \infty$, since $a_1 < 1$, $\lambda_n \rightarrow 0$ for $n \rightarrow \infty$, and E_n is bounded from above uniformly in $n \in \mathbb{N}$. Thus both the Sobolev norm $\|\psi_n\|_{H^1}^2 = \|\psi_n\|^2 + \|\nabla \psi_n\|^2$ and $\|\sqrt{V_+} \psi_n\|^2$ are bounded in n .

Now consider

$$\begin{aligned} \langle \varphi, H(\psi_n - \psi_m) \rangle &= \langle \varphi, H_n \psi_n \rangle + \lambda_n \langle \varphi, W \psi_n \rangle - \langle \varphi, H_m \psi_m \rangle - \lambda_m \langle \varphi, W \psi_m \rangle \\ &= E_n \langle \varphi, \psi_n \rangle + \lambda_n \langle \varphi, W \psi_n \rangle - E_m \langle \varphi, \psi_m \rangle - \lambda_m \langle \varphi, W \psi_m \rangle \end{aligned} \quad (4.6)$$

for $\varphi \in \mathcal{Q}(H)$. The choice $\varphi = \psi_n - \psi_m$ and Cauchy–Schwarz yields

$$\begin{aligned}
& \langle \psi_n - \psi_m, H(\psi_n - \psi_m) \rangle \\
& \leq |E_n| \|\varphi\| \|\psi_n\| + \lambda_n \|\sqrt{W}\varphi\| \|\sqrt{W}\psi_n\| + |E_m| \|\varphi\| \|\psi_m\| + \lambda_m \|\sqrt{W}\varphi\| \|\sqrt{W}\psi_m\| \\
& \leq (|E_n| \|\psi_n\| + |E_m| \|\psi_m\|) \|\varphi\| + \frac{\lambda_n}{2} (\|\sqrt{W}\varphi\|^2 + \|\sqrt{W}\psi_n\|^2) \\
& \quad + \frac{\lambda_m}{2} (\|\sqrt{W}\varphi\|^2 + \|\sqrt{W}\psi_m\|^2) \\
& \leq (|E_n| \|\psi_n\| + |E_m| \|\psi_m\|) \|\varphi\| + \frac{\lambda_n + \lambda_m}{2} (a_2 \|\nabla\varphi\|^2 + b_2 \|\varphi\|^2) \\
& \quad + \frac{\lambda_n}{2} (a_2 \|\nabla\psi_n\|^2 + b_2 \|\psi_n\|^2) + \frac{\lambda_m}{2} (a_2 \|\nabla\psi_m\|^2 + b_2 \|\psi_m\|^2)
\end{aligned} \tag{4.7}$$

On the other hand,

$$\begin{aligned}
\langle \psi_n - \psi_m, H(\psi_n - \psi_m) \rangle &= \|\nabla(\psi_n - \psi_m)\|^2 + \|\sqrt{V_+}(\psi_n - \psi_m)\|^2 - \|\sqrt{V_-}(\psi_n - \psi_m)\|^2 \\
&\geq (1 - a_1) \|\nabla(\psi_n - \psi_m)\|^2 + \|\sqrt{V_+}(\psi_n - \psi_m)\|^2 - b_1 \|\psi_n - \psi_m\|^2
\end{aligned}$$

and plugging this lower bound into (4.7) and using that ψ_n is normalized we arrive at

$$\begin{aligned}
& (1 - a_1 - \frac{\lambda_n + \delta_m}{2} a_2) \|\nabla(\psi_n - \psi_m)\|^2 + \|\sqrt{V_+}(\psi_n - \psi_m)\|^2 \\
& \leq (|E_n| + |E_m| + \frac{\lambda_n + \lambda_m}{2} b_2) \|\psi_n - \psi_m\| + b_1 \|\psi_n - \psi_m\|^2 \\
& \quad + \frac{\lambda_n}{2} (a_2 \|\nabla\psi_n\|^2 + b_2) + \frac{\lambda_m}{2} (a_2 \|\nabla\psi_m\|^2 + b_2).
\end{aligned} \tag{4.8}$$

By assumption and Remark 4.5, the sequence of eigenvalues E_n is bounded and, because of (4.5), we also have that $\|\nabla\psi_n\|$ is bounded uniformly in $n \in \mathbb{N}$. Since $\lambda_n \rightarrow 0$ and $\|\psi_n - \psi_m\| \rightarrow 0$ as $n, m \rightarrow \infty$, (4.8) implies

$$\limsup_{n, m \rightarrow \infty} \left((1 - a_1) \|\nabla(\psi_n - \psi_m)\|^2 + \|\sqrt{V_+}(\psi_n - \psi_m)\|^2 \right) \leq 0.$$

That is, the sequence of normalized weak eigenfunctions ψ_n of H_n is Cauchy in $\mathcal{Q}(H)$ with respect to the form norm $\|\cdot\|_1$ as soon as it is Cauchy in L^2 and the sequence of eigenvalues $(E_n)_{n \in \mathbb{N}}$ is bounded. In particular, the limit $\psi = \lim_{n \rightarrow \infty} \psi_n$ exists in $\mathcal{Q}(H)$. Thus $\|\nabla\psi\| = \lim_{n \rightarrow \infty} \|\nabla\psi_n\|$, $\|\sqrt{V_+}\psi\| = \lim_{n \rightarrow \infty} \|\sqrt{V_+}\psi_n\|$, and, since $W \geq 0$ is form bounded w.r.t. $-\Delta + V_+$ also $\|\sqrt{W}\psi\| = \lim_{n \rightarrow \infty} \|\sqrt{W}\psi_n\|$. Hence $\sup_n \|\sqrt{W}\psi_n\| < \infty$.

Now assume additionally that E_n converges to some E as $n \rightarrow \infty$. In this case, using that ψ_n converges to ψ in $\mathcal{Q}(H)$ we get

$$\begin{aligned}
\langle \varphi, H\psi \rangle &= \lim_{n \rightarrow \infty} \langle \varphi, H\psi_n \rangle = \lim_{n \rightarrow \infty} (\langle \varphi, H_n\psi_n \rangle + \lambda_n \langle \sqrt{W}\varphi, \sqrt{W}\psi_n \rangle) \\
&= \lim_{n \rightarrow \infty} (E_n \langle \varphi, \psi_n \rangle + \lambda_n \langle \sqrt{W}\varphi, \sqrt{W}\psi_n \rangle) = E \langle \varphi, \psi \rangle
\end{aligned} \tag{4.9}$$

for all $\varphi \in \mathcal{Q}(H)$ since $\lambda_n \rightarrow 0$ and $\sup_n |\langle \sqrt{W}\varphi, \sqrt{W}\psi_n \rangle| \leq \|\sqrt{W}\varphi\| \sup_n \|\sqrt{W}\psi_n\| < \infty$. Thus we proved that the limit $\psi = \lim_{n \rightarrow \infty} \psi_n \in \mathcal{Q}(H)$ exists, $\|\psi\| = 1$, and ψ is a weak eigenfunction of H with eigenvalue $E = \lim_{n \rightarrow \infty} E_n$ under the *additional assumption* that the limit $E = \lim_{n \rightarrow \infty} E_n$ exists.

Finally, it is easy to see that the sequence of eigenvalues E_n must converge. Assume that E_n does not converge as $n \rightarrow \infty$. Since E_n is bounded in $n \in \mathbb{N}$, there exist two different limit points $E_1 \neq E_2$ of E_n corresponding to two subsequences $E_{\sigma_1(n)} \rightarrow E_1$ and $E_{\sigma_2(n)} \rightarrow E_2$ where $\sigma_1, \sigma_2 : \mathbb{N} \rightarrow \mathbb{N}$ are strictly increasing functions.

Clearly $\psi = \lim_{n \rightarrow \infty} \psi_{\sigma_1(n)} = \lim_{n \rightarrow \infty} \psi_{\sigma_2(n)}$. So (4.9) shows that ψ is a weak eigenfunction of H corresponding to the two different eigenvalues E_1 and E_2 , which is impossible. Hence the eigenvalues E_n converges. This finishes the proof of Lemma 4.2. \blacksquare

Lemma 4.6. *Assume that the potentials V and W satisfy Assumption 1.1, except that the relative bound of W does not have to be less than one. Let $(H_\lambda)_{0 \leq \lambda \leq \lambda_0}$ be the family of perturbed Schrödinger operators constructed in Remark 2.1 for some small enough $0 < \lambda_0$. Moreover, assume that for some sequence $0 < \lambda_n \leq \lambda_0$ the operators $H_n = H_{\lambda_n}$ have eigenvalues E_n with corresponding weak eigenfunctions ψ_n .*

If $\|\psi_n\| = 1$ for all $n \in \mathbb{N}$ and $\sup_n E_n < \infty$, then the weak eigenfunctions ψ_n are pointwise locally bounded uniformly in $n \in \mathbb{N}$, i.e.,

$$\sup_{n \in \mathbb{N}} \sup_{x \in S} |\psi_n(x)| < \infty \quad (4.10)$$

for any bounded set $S \subset \mathbb{R}^d$.

Remark 4.7. Since eigenfunctions are continuous if the potential is locally in the Kato–class, $\psi_n(x)$ makes sense for all $x \in \mathbb{R}^d$ and $n \in \mathbb{N}$.

Proof. Note that ψ_n is a zero energy weak eigenfunction of the Schrödinger operator \tilde{H}_n with potential \tilde{V}_n given by $\tilde{V}_n = V - \lambda_n W - E_n$. If V and W are in the local Kato class, so is \tilde{V}_n . Hence for any $x \in \mathbb{R}^d$ the subsolution estimate

$$|\psi_n(x)| \leq C_{x,n} \int_{|x-y|<1} |\psi_n(y)| dy \quad (4.11)$$

holds, see [52, Theorem C.1.2] and also [3, 48]. Moreover, the constants $C_{x,n}$ depend only on

$$\|\mathbb{1}_{B_1(x)}(\tilde{V}_n)_-\|_{K^d}$$

with $(\tilde{V}_n)_-$ being the negative part of \tilde{V}_n and the Kato norm $\|\cdot\|_{K^d}$ given by

$$\|V\|_{K^d} := \sup_{x \in \mathbb{R}^d} \int_{|x-y| \leq 1} \tilde{g}_d(x-y) |V(y)| dy \quad (4.12)$$

with $\tilde{g}_d = g_d$ when $d \geq 3$, $\tilde{g}_2 = 1 + g_2$, and $\tilde{g}_1 = 1$, where g_d is defined in (1.15). Adding 1 to g_2 is necessary since $g_2(x) = 0$ when $|x| = 1$.

For any set $S \subset \mathbb{R}^d$ and any potential V we have

$$\sup_{x \in S} \|\mathbb{1}_{B_1(x)} V\|_{K^d} \leq \|\sup_{x \in S} \mathbb{1}_{B_1(x)} V\|_{K^d} = \|\mathbb{1}_{S_1} V\|_{K^d} \quad (4.13)$$

where $S_1 = \{y \in \mathbb{R}^d : \text{dist}(y, S) < 1\}$.

Now let $S \subset \mathbb{R}^d$ be bounded. Then S_1 is bounded and, since the Kato norm of a constant function is finite and $(\tilde{V}_n)_- = (V - \lambda_n W - E_n)_- \leq V_- + \lambda_n W_+ + (E_n)_+$, we have

$$\sup_n \|\mathbb{1}_{S_1}(\tilde{V}_n)_-\|_{K^d} \leq (\|\mathbb{1}_{S_1}V_-\|_{K^d} + \sup_n \lambda_n \|\mathbb{1}_{S_1}W_+\|_{K^d} + \sup_n (E_n)_+ \|1\|_{K^d}) < \infty$$

for any bounded set S , using that $\sup_n E_n < \infty$ and $\sup_n \lambda_n < \infty$, by assumption, and $\|\mathbb{1}_{S_1}V_-\|_{K^d} < \infty$ and $\|\mathbb{1}_{S_1}W_+\|_{K^d} < \infty$, since S_1 is bounded and V and W are locally in the Kato class.

Thus for any bounded set $S \subset \mathbb{R}^d$ there exist a constant $C < \infty$ such that

$$|\psi_n(x)| \leq C \int_{|x-y|<1} |\psi_n(y)| dy \quad (4.14)$$

for all $x \in S$ and $n \in \mathbb{N}$. Using the normalization $\|\psi_n\| = 1$ we have

$$\int_{|x-y|<1} |\psi_n(y)| dy \leq |B_1^d|^{1/2} \|\psi_n\| = |B_1^d|^{1/2} \quad (4.15)$$

for all $x \in S$ and $n \in \mathbb{N}$. Hence (4.10) follows immediately from (4.14). \blacksquare

The last result which we need is

Lemma 4.8. *Assume that $V \in L^1_{loc}(\mathbb{R}^d)$ and V_- is form small w.r.t. $-\Delta + V_+$ and ψ is a real-valued weak eigenfunction of H at energy E . Then $|\psi|$ is a subsolution of H at energy E .*

Proof. If ψ is real-valued eigenfunction of H at energy E then it is also a subsolution, hence [2, Lemma 2.9] shows that its positive part $\psi_+ = \sup(\psi, 0)$ is a subsolution. The same argument applied to $-\psi$, which is also a weak solution, shows that its negative part $\psi_- = \sup(-\psi, 0)$ is a subsolution. Hence $|\psi| = \psi_+ + \psi_-$ is a subsolution of H at energy E . \blacksquare

Remark 4.9. It is well-known that for the type of Schrödinger operator H we consider here the eigenfunctions can be chosen to be real-valued. Since H is self-adjoint all eigenvalues are real. Moreover, H commutes with complex conjugation, so for any complex-valued eigenfunction ψ of H also the real and imaginary parts $\operatorname{Re}(\psi) = \frac{1}{2}(\psi + \bar{\psi})$ and $\operatorname{Im}(\psi) = \frac{1}{2i}(\psi - \bar{\psi})$ are eigenfunction of H at energy E . This is not true anymore if one considers Schrödinger operators with magnetic fields, since they do not commute with complex conjugation, in general.

Now we are ready to give the

Proof of Theorem 1.7: By assumption, the potential V is critical. Thus $\sigma(H) = \sigma_{ess}(H) = [0, \infty)$. Moreover, for any non-trivial potential $W \geq 0$ which is infinitesimally form small w.r.t. $-\Delta + V_+$ and has compact support the Schrödinger operators $H_\lambda = H - \lambda W$, constructed in Remark 2.1, have non-trivial discrete spectrum below zero. That is, $\sigma_{ess}(H_\lambda) = [0, \infty)$ and there exist eigenvalues $E_\lambda < 0$ of H_λ with associated normalized weak eigenfunctions ψ_λ for all $\lambda > 0$. We take any sequence $(\lambda_n)_{n \in \mathbb{N}}$ which is monotonically decreasing to zero and abbreviate $H_n = H_{\lambda_n}$, $E_n = E_{\lambda_n}$, and $\psi_n = \psi_{\lambda_n}$.

Recall that we also assume that the potential V satisfies the lower bound

$$V(x) \geq \frac{d(4-d)}{4|x|^2} + \frac{1}{|x|^2} \sum_{j=1}^m \prod_{k=1}^j \ln_k^{-1}(|x|) + \frac{2\epsilon}{|x|^2} \prod_{k=1}^m \ln_k^{-1}(|x|)$$

for $|x| > R$, some $m \in \mathbb{N}_0$, $\epsilon > 0$, and all large enough $R > 0$. We replaced ϵ by 2ϵ in (1.5). Increasing R , if necessary, it is easy to see that this implies

$$V(x) \geq Y_{m,\epsilon}(x) \text{ for all } |x| \geq R, \quad (4.16)$$

where the family of comparison functions $Y_{m,\epsilon}$ is defined in (4.2).

Since W has compact support, we can also assume that R is so large that its support $\text{supp}(W) \subset B_R(x)$. Thus, with $U_R = \{|x| > R\}$ we have $W\varphi = 0$ for all $\varphi \in C_0^\infty(U_R)$. Lemma 4.1 and (4.16) imply

$$\begin{aligned} \langle \varphi, (H_n - E_n)\psi_{u,m,\epsilon} \rangle &= \langle \varphi, (H - E_n)\psi_{u,m,\epsilon} \rangle \\ &= \langle \varphi, (-\Delta + Y_{m,\epsilon} - E_n)\psi_{u,m,\epsilon} \rangle + \langle \varphi, (V - Y_{m,\epsilon})\psi_{u,m,\epsilon} \rangle \geq -E_n \langle \varphi, \psi_{u,m,\epsilon} \rangle \geq 0 \end{aligned} \quad (4.17)$$

for all $0 \leq \varphi \in C_0^\infty(U_R)$. Here $\psi_{u,m,\epsilon} > 0$ is defined in (4.1) and we used that $E_n \leq 0$.

So for fixed $m \in \mathbb{N}$, large enough $R > 0$, and small enough $\epsilon > 0$ the function $\psi_{u,m,\epsilon}$ is a supersolution of H_n at energy E_n in U_R for all $n \in \mathbb{N}$. Moreover, since $\|\psi_n\| = 1$ we have

$$c_R^1 := \sup_{n \in \mathbb{N}} \sup_{R \leq |x| \leq R+1} |\psi_n(x)| < \infty$$

by Lemma 4.6. Since $\psi_{u,m,\epsilon} > 0$ is continuous away from zero, we also have

$$c_R^2 = \inf_{R \leq |x| \leq R+1} \psi_{u,m,\epsilon}(x) > 0$$

and using $C_R = c_R^1/c_R^2$ one gets $|\psi_n(x)| \leq C_R \psi_{u,m,\epsilon}(x)$, hence also

$$\tilde{\psi}_n(x) := |\text{Re}(\psi_n(x))| + |\text{Im}(\psi_n(x))| \leq \sqrt{2}|\psi_n(x)| \leq \sqrt{2}C_R \psi_{u,m,\epsilon}(x) \quad (4.18)$$

for all $R \leq |x| \leq R+1$ and all $n \in \mathbb{N}$. Clearly, $|\psi_n| \leq \tilde{\psi}_n$. Since $\tilde{\psi}_n$ is a nonnegative subsolution of H_n at energy E_n by Lemma 4.8 and Remark 4.9 we can use $w = \psi_{u,m,\epsilon}$ and $v = \tilde{\psi}_n$ in Theorem 2.7 to see that

$$|\psi_n(x)| \leq \tilde{\psi}_n(x) \leq \sqrt{2}C_R \psi_{u,m,\epsilon}(x) \text{ for all } |x| \geq R \quad (4.19)$$

uniformly in $n \in \mathbb{N}$. Since $\psi_{u,m,\epsilon}$ is square integrable at infinity for any fixed $m \in \mathbb{N}$ and $\epsilon > 0$, the bound (4.19) yields tightness in x -space, i.e.,

$$\lim_{R \rightarrow \infty} \sup_{n \in \mathbb{N}} \int_{|x| > R} |\psi_n(x)|^2 dx = 0. \quad (4.20)$$

From (4.5) one gets $\sup_{n \in \mathbb{N}} \|\psi_n\|_{H^1} < \infty$. In particular, we have

$$\lim_{L \rightarrow \infty} \sup_{n \in \mathbb{N}} \int_{|\eta| > L} |\widehat{\psi}_n(\eta)|^2 d\eta = 0, \quad (4.21)$$

which is tightness in momentum space. Here $\widehat{\psi}_n$ is the Fourier transform of ψ_n .

Moreover, since ψ_n is bounded in $H^1(\mathbb{R}^d)$, there exists a subsequence which converges weakly in H^1 and L^2 . By a slight abuse of notation, we also write ψ_n for this subsequence.

Let $\psi \in L^2(\mathbb{R}^d)$ be the weak limit of ψ_n . Tightness and weak convergence then implies that ψ_n converges to ψ in L^2 , see e.g., [23, Appendix A]. Hence $\|\psi\| = \lim_{n \rightarrow \infty} \|\psi_n\| = 1$.

Lemma 4.2 shows that $E = \lim_{n \rightarrow \infty} E_n \leq 0$ exists and that ψ is a normalized weak eigenfunction of H with eigenvalue E . Clearly, $E = 0$ since $\sigma(H) = [0, \infty)$. So zero is the ground state eigenvalue of H which is at the edge of the essential spectrum of H . This finishes the proof of Theorem 1.7. \blacksquare

Remark 4.10. Note that we could have simplified some parts of the proof by using that ground states can be chosen to be strictly positive. We intentionally avoided the use of strict positivity of ground state eigenfunctions. This allows to use Theorem 1.7 also for systems with symmetry restrictions, or for the existence of higher eigenstates with energies above the ground state energy, provided one suitably modifies the assumption of a virtual level for such systems. These modifications are straightforward.

APPENDIX A. AN EXAMPLE IN SEARCH OF A THEOREM

It is well-known that the zero potential is critical in dimensions one and two, see [50] and also [35, Problems 1 and 2 in Chapter 45]. This phenomenon can be explained by the non-integrability of $\eta \mapsto |\eta|^{-2}$ near $\eta = 0$ in \mathbb{R}^d , see [20]. The Iorio-O'Carroll theorem [47, Theorem XII.27] shows that shallow potential wells cannot create ground states in dimension $d \geq 3$ and that the corresponding Schrödinger operators are even unitarily equivalent to the free Laplacian.

Of course, in order to construct zero energy resonances or zero energy ground states, one can take any Schrödinger operator H which has essential spectrum $[0, \infty)$ and finitely many negative eigenvalues. Adding a suitable local positive perturbation then moves the ground state energy to zero, creating a zero energy resonance, or zero energy ground state, depending, for example, on which a priori bound from Theorem 1.3 of Theorem 1.7 holds.

Specific examples of critical potentials in dimension one and two which are different from the zero potential seem to be rare. In the following we construct a family of potentials $V_{\alpha,d}$ in any dimension which are critical for $\alpha \geq 0$, having a zero energy resonance when $0 \leq \alpha \leq 1$ and a zero energy ground state when $\alpha > 1$, and which are not critical when $\alpha < 0$. To the best of our knowledge, our example is new.

Remark A.1. There are different definitions for a zero energy resonance available in the literature. One often calls ψ a zero energy resonance if it is a local positive eigenfunction of a Schrödinger operator H which is not square integrable on \mathbb{R}^d but its gradient $\nabla\psi$ is square integrable. We will follow this convention, except that we also allow that the L^2 -norm of $\nabla\psi$ is logarithmically divergent at infinity.

For $\alpha \in \mathbb{R}$ and $d \in \mathbb{N}$ define the potential $V_{\alpha,d}$ on \mathbb{R}^d by

$$V_{\alpha,d}(x) := \frac{4\alpha^2 - (d-2)^2}{4(1+|x|^2)} + \frac{1 - (\alpha + d/2)^2}{(1+|x|^2)^2} \quad (\text{A.1})$$

Clearly, $V_{\alpha,d}$ is bounded and goes to zero at infinity. Thus it is a Kato-class potential for all $d \geq 1$ and all $\alpha \geq 0$. In particular, $V_{\alpha,d}$ is both infinitesimally operator bounded, hence also infinitesimally form bounded, w.r.t. $-\Delta$. Therefore the Schrödinger operator $H_{\alpha,d} =$

$-\Delta + V_{\alpha,d}$ is a well defined self-adjoint operator on the domain $H^2(\mathbb{R}^d)$ with form domain $H^1(\mathbb{R}^d)$.

The key to understanding why the potentials $V_{\alpha,d}$ are critical for all $\alpha \geq 0$ and $d \geq 1$, not critical for $\alpha < 0$, and switch from having zero energy resonances to having zero energy ground states at $\alpha = 1$ is

Lemma A.2 (Ground state representation of $H_{\alpha,d}$). *Let $\alpha \in \mathbb{R}$, $d \geq 1$, and define*

$$\psi_{\alpha,d}(x) = (1 + |x|^2)^{(2-d)/4-\alpha/2} \quad (\text{A.2})$$

for $x \in \mathbb{R}^d$ and the measure

$$\mu_{\alpha,d}(B) = \int_B \psi_{\alpha,d}^2 dx \quad (\text{A.3})$$

on the Borel sets B in \mathbb{R}^d . Then the map $U_{\alpha,d} : L^2(\mathbb{R}^d, d\mu_{\alpha,d}) \rightarrow L^2(\mathbb{R}^d)$ given by

$$(U_{\alpha,d}\varphi) = \psi_{\alpha,d}\varphi \quad (\text{A.4})$$

is unitary with

$$U_{\alpha,d}^{-1}(H^1(\mathbb{R}^d)) = \{\varphi \in L^2(\mathbb{R}^d, d\mu_{\alpha,d}) : \nabla\varphi \in L^2(\mathbb{R}^d, d\mu_{\alpha,d})\}. \quad (\text{A.5})$$

Moreover, $U_{\alpha,d}H_{\alpha,d}U_{\alpha,d}^{-1} = -\Delta$ in the sense that for all $\psi \in H^1(\mathbb{R}^d)$, the form domain of $H_{\alpha,d}$,

$$\langle \psi, H_{\alpha,d}\psi \rangle = \langle \nabla\psi, \nabla\psi \rangle - \langle \psi, V_{\alpha,d}\psi \rangle = \int_{\mathbb{R}^d} |\nabla\varphi|^2 \psi_{\alpha,d}^2 dx \quad (\text{A.6})$$

where $\varphi = U_{\alpha,d}^{-1}\psi$.

Remark A.3. Lemma A.2 shows that the Schrödinger operator $H_{\alpha,d}$ is equivalent to the Dirichlet form $q(\varphi) = \langle \nabla\varphi, \nabla\varphi \rangle_{L^2(\mathbb{R}^d, d\mu_{\alpha,d})}$ on the weighted L^2 -space with measure $d\mu_{\alpha,d} = (1 + |x|^2)^{-(d-2)/2-\alpha} dx$. Note that this measure is finite if and only if $\alpha > 1$.

We give the proof of the lemma at the end of the appendix.

Theorem A.4. *Let $d \in \mathbb{N}$, $\alpha \in \mathbb{R}$, and $H_{\alpha,d} = -\Delta + V_{\alpha,d}$ be the self-adjoint Schrödinger operator with potential $V_{\alpha,d}$ given by (A.1). Then*

- a) $\sigma(H_{\alpha,d}) = \sigma_{\text{ess}}(H_{\alpha,d}) = [0, \infty)$.
- b) For all $\alpha \geq 0$ the potential $V_{\alpha,d}$ is critical, that is, the Schrödinger operator $H_{\alpha,d}$ has a virtual level.
- c) Zero is not an eigenvalue of $H_{\alpha,d}$ when $0 \leq \alpha \leq 1$. For $\alpha > 1$ zero is an eigenvalue. The zero energy resonance for $0 \leq \alpha \leq 1$, respectively ground state for $\alpha > 1$, is given by (A.2).
- d) For $\alpha < 0$, the potential $V_{\alpha,d}$ is subcritical, and zero is neither an eigenvalue nor a resonance.

Remark A.5. Using the early result of Kato, [28], see also [1, 49], the operator $H_{\alpha,d}$ has no strictly positive embedded eigenvalues. Since the potential $V_{\alpha,d}$ is short range, the spectrum of $H_{\alpha,d}$ is even purely absolutely continuous inside $(0, \infty)$, see [11, Theorem 5.10].

Proof. Using standard methods, [53], one sees that $\sigma_{ess}(H_{\alpha,d}) = \sigma_{ess}(-\Delta) = [0, \infty)$ since $V_{\alpha,d}$ is bounded and goes to zero at infinity. Moreover, the ground state representation (A.6) implies $\sigma(H_{\alpha,d}) \subset [0, \infty)$. Hence $\sigma(H_{\alpha,d}) = \sigma_{ess}(H_{\alpha,d}) = [0, \infty)$. This proves claim a).

Given $\varphi \in L^2(\mathbb{R}^d, d\mu_{\alpha,d})$ let $\psi := U_{\alpha,d}\varphi$. Taking $\varphi = 1$ gives $\psi = \psi_{\alpha,d} > 0$ which is in $H^1(\mathbb{R}^d)$ if and only if $\alpha > 1$. In this case (A.6) shows that $\psi_{\alpha,d}$ is the ground state of $H_{\alpha,d}$ corresponding to the eigenvalue zero. This proves the second claim in c). Lemma 2.6 also shows that the potential $V_{\alpha,d}$ is critical when $\alpha > 1$.

In addition, note that the right hand side of (A.6) is strictly positive unless φ is constant. Hence zero is not an eigenvalue of $H_{\alpha,d}$ when $0 \leq \alpha \leq 1$ because $\psi_{\alpha,d}$ is not square integrable in this case.

When $0 < \alpha \leq 1$ we take any $\varphi \in C_0^\infty(\mathbb{R})$ with $\varphi(t) = 1$ for $|t| \leq 1$, $\varphi(t) = 0$ for $|t| \geq 2$, and define

$$\varphi_R(x) = \varphi(|x|/R)$$

for $R > 0$. Then $|\nabla\varphi_R(x)| = R^{-1}|\varphi'(|x|/R)|$. Using $\psi_R = \psi_{\alpha,d}\varphi_R$ we get

$$\begin{aligned} \langle \psi_R, H_{\alpha,d}\psi_R \rangle &= R^{-2} \int |\varphi'(|x|/R)|^2 (1 + |x|^2)^{(2-d)/2-\alpha} dx \\ &\lesssim R^{-2} \int_R^{2R} (1 + r^2)^{(2-d)/2-\alpha} r^{d-1} dr \sim R^{-2} \int_R^{2R} (1 + r^2)^{-\alpha} r dr \lesssim R^{-2\alpha} \rightarrow 0 \end{aligned}$$

for $R \rightarrow \infty$ and $\alpha > 0$. Now let $W \geq 0$ have compact support, be infinitesimally form bounded w.r.t. $-\Delta$, and $W > 0$ on a set of positive Lebesgue measure. Since $\psi_R(x) \rightarrow (1 + |x|^2)^{(2-d)/4-\alpha/2}$ as $R \rightarrow \infty$ uniformly on compact sets we have

$$\lim_{R \rightarrow \infty} \langle \psi_R, (H_{\alpha,d} - \lambda W)\psi_R \rangle = -\lambda \int W(x)(1 + |x|^2)^{(2-d)/2-\alpha} dx < 0$$

for all $\lambda > 0$. Thus $\langle \psi_R, (H_{\alpha,d} - \lambda W)\psi_R \rangle < 0$ for all large enough $R > 0$. Since $\sigma_{ess}(H_{\alpha,d} - \lambda W) = [0, \infty)$, the Rayleigh Ritz principle shows that $H_{\alpha,d} - \lambda W$ has a negative eigenvalue for any $\lambda > 0$. Thus the potential $V_{\alpha,d}$ is critical. Clearly zero cannot be an eigenvalue nor a resonance, since then the potential $V_{\alpha,d}$ would have to be critical.

To see that $V_{0,d}$ is critical one needs to modify the ansatz function. Let $\delta > 0$ and set

$$\varphi_\delta(x) := \begin{cases} 1 & \text{if } |x| \leq 1 \\ (1 - \delta \ln |x|)_+ & \text{if } |x| > 1 \end{cases}$$

and $\psi_\delta = U_{\alpha,d}\varphi_\delta$. A straightforward calculation shows

$$\begin{aligned} \langle \psi_\delta, H_{0,d}\psi_\delta \rangle &= \delta^2 \int_{1 \leq |x| \leq e^{1/\delta}} (1 + |x|^2)^{(2-d)/2} |x|^{-2} dx \\ &\lesssim \delta^2 \int_1^{e^{1/\delta}} (1 + r^2)^{(2-d)/2} r^{d-3} dr \sim \delta^2 \int_1^{e^{1/\delta}} (1 + r^2)^{-1} r dr \\ &= \frac{\delta^2}{2} \ln(1 + e^{1/\delta}) \rightarrow 0 \quad \text{as } \delta \rightarrow 0. \end{aligned}$$

Thus $\lim_{\delta \rightarrow 0} \langle \psi_\delta, (H_{0,d} - \lambda W)\psi_\delta \rangle = -\lambda \int W(x)(1 + |x|^2)^{(2-d)/2} dx < 0$. As before this shows that $V_{0,d}$ is critical. Moreover, even though $\psi_{\alpha,d} \notin L^2(\mathbb{R}^d)$ when $0 \leq \alpha \leq 1$, its gradient $\nabla \psi_{\alpha,d}$ is in $L^2(\mathbb{R}^d)$ when $0 < \alpha \leq 1$ and the L^2 -norm of $\nabla \psi_{0,d}$ is only logarithmically divergent. Hence $\psi_{\alpha,d}$ is a zero energy resonance for $H_{\alpha,d}$ when $0 \leq \alpha \leq 1$.

Finally, we look at $V_{-\alpha,d}$ for $\alpha > 0$. A simple calculation shows

$$V_{-\alpha,d}(x) = V_{\alpha,d}(x) + 2\alpha d(1 + |x|^2)^{-2}.$$

Thus with $W(x) = (1 + |x|^2)^{-2} > 0$ and $\lambda = 2\alpha d > 0$ we have

$$\langle \psi, (H_{-\alpha,d} - \lambda W)\psi \rangle = \langle \psi, H_{\alpha,d}\psi \rangle \geq 0$$

for all $\psi \in H^1(\mathbb{R}^d)$, since $\sigma(H_{\alpha,d}) = [0, \infty)$ by part a). Hence $V_{-\alpha,d}$ is subcritical for $\alpha > 0$. ■

The family of potential $V_{\alpha,d}$ has several interesting properties summarized in

Lemma A.6 (Properties of $V_{\alpha,d}$). *Let $\alpha \in \mathbb{R}$ and $d \in \mathbb{N}$. Then*

- a) *In dimensions $d = 1, 2$ the potential $V_{\alpha,d}$ is non-trivial if $\alpha \neq |d-2|/2$ and in dimension $d \geq 3$ it is non-trivial if $\alpha \neq (2-d)/2$.*
- b) *In dimension $d = 1$ we have $V_{\alpha,1} > 0$ for $\alpha \leq -1/2$. If $-1/2 < \alpha < 1/2$ then $V_{\alpha,1} > 0$ near zero and it has a negative tail, i.e., $V_{\alpha,1}(x) < 0$ for large $|x|$. If $\alpha > 1/2$, then $V_{\alpha,1}$ is negative near zero and it has a positive tail.*
- c) *In dimension $d = 2$ we have $V_{\alpha,2} > 0$, i.e., $V_{\alpha,2}$ is purely repulsive for all $\alpha < 0$. For $\alpha > 0$ the potential $V_{\alpha,2}$ is negative near zero and has a positive tail.*
- d) *In dimension $d \geq 3$ we have $V_{\alpha,d} > 0$, i.e., the potential is repulsive, for $\alpha < (2-d)/2$. For $(2-d)/2 < \alpha \leq (d-2)/2$ we have $V_{\alpha,d} < 0$, i.e., the potential is attractive. If $\alpha > (d-2)/2$, then $V_{\alpha,d}$ is negative near zero and has a positive tail.*
- e) *For $d = 1$ the potential $V_{\alpha,1}$ is integrable and*

$$\int_{-\infty}^{\infty} V_{\alpha,1}(x) dx = \frac{\pi}{2}(\alpha - 1/2)^2 > 0 \quad \text{for } \alpha \neq 1/2.$$

- f) *For large enough R and all dimensions $d \geq 1$ the potentials $V_{\alpha,d}$ satisfy the bounds (1.6) for $0 \leq \alpha < 1$, respectively (1.8) for $\alpha = 1$, while they satisfy the complementary bound (1.7) for $0 < \epsilon < 4(\alpha^2 - 1)$ when $\alpha > 1$.*

Remark A.7. Claims b) and c) above are consistent with what is known about weakly coupled bound states in low dimensions. It is known that if $V \in L^1(\mathbb{R}^d)$ and $\int_{\mathbb{R}^d} V dx \leq 0$, where V is supposed to be non-trivial when $\int V dx = 0$, then the operator $-\Delta + \lambda V$ always has a negative bound state, no matter how small the coupling parameter $\lambda > 0$ is, when $d = 1, 2$. See, for example, [50] where this is proved under some additional assumptions, or [20] for the full result. In particular, this implies that critical potentials in one and two dimensions have to change sign and, if they are integrable, then $\int_{\mathbb{R}^d} V dx > 0$ unless V is trivial.

In addition, claim d) is consistent with our non-existence Theorem 1.3. Non-positive potentials cannot have a zero energy ground state in dimensions $d \leq 4$. They need to have a strong enough positive tail in order to be able to have zero energy bound states. Moreover, claim f) together with the fact that the potential $V_{\alpha,d}$ supports zero energy ground states if and only if $\alpha > 1$, see Theorem A.4, is consistent with our Theorems 1.3 and 1.7.

It is illuminating to plot $V_{\alpha,d}(x)$ for $|x| = r$ to explicitly see the behavior of $V_{\alpha,d}$ for various values of the parameters α and d .

Proof. The first claim a) is easy to check. To prove the rest, let $a_{\alpha,d} = \alpha^2 - (d-2)^2/4$ and $b_{\alpha,d} = 1 - (\alpha + d/2)^2$. Then

$$4V_{\alpha,d}(0) = 4(a_{\alpha,d} + b_{\alpha,d}) = -2d(d-2+2\alpha) > 0$$

if and only if $\alpha < (2-d)/2$. Moreover, unless $a_{\alpha,d} = 0$, the sign of $V_{\alpha,d}(x)$ for large $|x|$ is determined by the sign of $a_{\alpha,d}$. Since $a_{\alpha,d} > 0$ if and only if $|\alpha| > |d-2|/2$, it is straightforward to deduce the claims b), c), and d) from this.

Clearly, $V_{\alpha,1}$ is integrable. Using $\int_{-\infty}^{\infty} (1+x^2)^{-1} dx = \pi$ and $\int_{-\infty}^{\infty} (1+x^2)^{-2} dx = \pi/2$, claim e) follows from a simple calculation.

Since for large $|x|$ the second term in the definition of $V_{\alpha,d}$ is much smaller than the first, the last claim f) follows from a straightforward computation. \blacksquare

It remains to give the proof of the ground state representation.

Proof of Lemma A.2. Let $\gamma \in \mathbb{R}$ and set $\psi_\gamma(x) = (1+|x|^2)^{-\gamma/2}$ for $x \in \mathbb{R}^d$, which is a regularized version of $|x|^{-\gamma}$ used at the end of Section 2. When ψ and φ are related by

$$\psi = \psi_\gamma \varphi \tag{A.7}$$

then $\psi \in L^2(\mathbb{R}^d)$ is clearly equivalent to $\varphi \in L^2(\mathbb{R}^d, \psi_\gamma^2 dx)$ and the corresponding norms are the same. So the map $U_\gamma : L^2(\mathbb{R}^d, \psi_\gamma^2 dx) \rightarrow L^2(\mathbb{R}^d)$, $\varphi \mapsto \psi_\gamma \varphi$ preserves the corresponding norms. Its inverse is given by $U_\gamma^{-1}\psi = U_{-\gamma}\psi = \psi_\gamma^{-1}\psi$ and from this one easily checks that U_γ is a unitary map from the weighted space $L^2(\mathbb{R}^d, \psi_\gamma^2 dx)$ to $L^2(\mathbb{R}^d)$. This proves (A.4).

If $\psi \in L^2(\mathbb{R}^d)$ and $\varphi = U_\gamma^{-1}\psi \in L^2(\mathbb{R}^d, \psi_\gamma^2 dx)$, then we have, in the sense of distributions,

$$\nabla\psi = \varphi\nabla\psi_\gamma + \psi_\gamma\nabla\varphi = -\gamma\psi_\gamma(1+|x|^2)^{-1}x\varphi + \psi_\gamma\nabla\varphi = -\gamma(1+|x|^2)^{-1}x\psi + \psi_\gamma\nabla\varphi \tag{A.8}$$

since $\psi_\gamma \in C^\infty(\mathbb{R}^d)$. Clearly, $(1+|x|^2)^{-1}x$ is bounded on \mathbb{R}^d . Therefore, if $\varphi \in L^2(\mathbb{R}^d, \psi_\gamma^2 dx)$ and $\nabla\varphi \in L^2(\mathbb{R}^d, \psi_\gamma^2 dx)$ then (A.8) shows that $\nabla\psi \in L^2(\mathbb{R}^d)$. Hence, if φ and $\nabla\varphi$ are in $L^2(\mathbb{R}^d, \psi_\gamma^2 dx)$ then $\psi = U_\gamma\varphi$ is in $H^1(\mathbb{R}^d)$.

Conversely, if $\psi \in H^1(\mathbb{R}^d)$, then, as distributions, $\nabla\varphi = \gamma(1+|x|^2)^{\gamma/2-1}x\psi + (1+|x|^2)^{\gamma/2}\nabla\psi$, which shows that

$$\psi_\gamma\nabla\varphi = (1+|x|^2)^{-1}x\psi + \nabla\psi \in L^2(\mathbb{R}^d). \tag{A.9}$$

That is, if $\psi \in L^2(\mathbb{R}^d)$, then $\varphi = U_\gamma^{-1}\psi \in L^2(\mathbb{R}^d, \psi_\gamma^2 dx)$ and if, in addition, $\nabla\psi \in L^2(\mathbb{R}^d)$ then (A.9) shows that $\nabla\varphi \in L^2(\mathbb{R}^d, \psi_\gamma^2 dx)$. Altogether, this proves

$$U_\gamma^{-1}(H^1(\mathbb{R}^d)) = \{\varphi \in L^2(\mathbb{R}^d, \psi_\gamma^2 dx) : \nabla\varphi \in L^2(\mathbb{R}^d, \psi_\gamma^2 dx)\}$$

which is (A.5). Moreover, $C_0^\infty(\mathbb{R}^d)$ is dense in $H^1(\mathbb{R}^d)$ and since U_γ maps $C_0^\infty(\mathbb{R}^d)$ into itself it is also dense in $U_\gamma^{-1}(H^1(\mathbb{R}^d))$. So we only have to prove (A.6) for $\varphi \in C_0^\infty(\mathbb{R}^d)$.

Let $\gamma \in \mathbb{R}$ and $\psi = \psi_\gamma\varphi$ with $\varphi \in C_0^\infty(\mathbb{R}^d)$. Then, as already noticed before,

$$\nabla\psi(x) = -\gamma(1+|x|^2)^{-\gamma/2-1}x\varphi(x) + (1+|x|^2)^{-\gamma/2}\nabla\varphi(x),$$

hence

$$\begin{aligned} \langle \nabla \psi, \nabla \psi \rangle &= \langle \nabla \varphi, (1 + |x|^2)^{-\gamma} \nabla \varphi \rangle - 2\gamma \operatorname{Re}(\langle \nabla \varphi, (1 + |x|^2)^{-\gamma-1} x \varphi \rangle) \\ &\quad + \gamma^2 \langle \varphi, (1 + |x|^2)^{-\gamma-2} |x|^2 \varphi \rangle. \end{aligned} \quad (\text{A.10})$$

An integration by parts shows

$$\begin{aligned} \operatorname{Re}(\langle \nabla \varphi, (1 + |x|^2)^{-\gamma-1} x \varphi \rangle) &= -\operatorname{Re}(\langle \varphi, \nabla \cdot ((1 + |x|^2)^{-\gamma-1} x \varphi) \rangle) \\ &= 2(\gamma + 1) \langle \varphi, (1 + |x|^2)^{-\gamma-2} |x|^2 \varphi \rangle - d \langle \varphi, (1 + |x|^2)^{-\gamma-1} \varphi \rangle - \operatorname{Re}(\langle \varphi, (1 + |x|^2)^{-\gamma-1} x \nabla \varphi \rangle). \end{aligned}$$

Noticing that $\operatorname{Re}(\langle \varphi, (1 + |x|^2)^{-\gamma-1} x \nabla \varphi \rangle) = \operatorname{Re}(\langle \nabla \varphi, (1 + |x|^2)^{-\gamma-1} x \varphi \rangle)$ we get

$$2\gamma \operatorname{Re}(\langle \nabla \varphi, (1 + |x|^2)^{-\gamma-1} x \varphi \rangle) = 2\gamma(\gamma + 1) \langle \varphi, (1 + |x|^2)^{-\gamma-2} |x|^2 \varphi \rangle - d\gamma \langle \varphi, (1 + |x|^2)^{-\gamma-1} \varphi \rangle$$

and plugging this into (A.10) we arrive at

$$\begin{aligned} \langle \nabla \psi, \nabla \psi \rangle &= \langle \nabla \varphi, (1 + |x|^2)^{-\gamma} \nabla \varphi \rangle - 2\gamma(\gamma + 1) \langle \varphi, (1 + |x|^2)^{-\gamma-2} |x|^2 \varphi \rangle \\ &\quad + d\gamma \langle \varphi, (1 + |x|^2)^{-\gamma-1} \varphi \rangle + \gamma^2 \langle \varphi, (1 + |x|^2)^{-\gamma-2} |x|^2 \varphi \rangle \\ &= \langle \nabla \varphi, (1 + |x|^2)^{-\gamma} \nabla \varphi \rangle - \gamma(\gamma + 2 - d) \langle \varphi, (1 + |x|^2)^{-\gamma-1} \varphi \rangle \\ &\quad + \gamma(\gamma + 2) \langle \varphi, (1 + |x|^2)^{-\gamma-2} \varphi \rangle \\ &= \langle \nabla \varphi, (1 + |x|^2)^{-\gamma} \nabla \varphi \rangle - \gamma(\gamma + 2 - d) \langle \psi, (1 + |x|^2)^{-1} \psi \rangle \\ &\quad + \gamma(\gamma + 2) \langle \psi, (1 + |x|^2)^{-2} \psi \rangle. \end{aligned}$$

Choosing $\gamma = (d - 2)/2 + \alpha$ finishes the proof of Lemma A.2. ■

Remarks A.8. The proof of Lemma A.2 is clearly inspired by the proof of Hardy's inequality on $L^2(\mathbb{R}^d)$ for $d \geq 3$, where one considers $\psi(x) = |x|^{-\gamma/2} \varphi(x)$ for $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d \setminus \{0\})$ and optimizes in $\gamma > 0$. One needs to restrict to $\varphi \in \mathcal{C}_0^\infty(\mathbb{R}^d \setminus \{0\})$ due to the singularity of $|x|^{-\gamma/2}$ in zero. Since $\mathcal{C}_0^\infty(\mathbb{R}^d \setminus \{0\})$ is dense in $L^2(\mathbb{R}^d)$ only when $d \geq 3$, this leads to the well-known fact that Hardy's inequality only holds in dimensions $d \geq 3$.

Acknowledgements: Funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 258734477 – SFB 1173. This project also received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement MDFT No. 725528). Michal Jex also received financial support from the Ministry of Education, Youth and Sport of the Czech Republic under the Grant No. RVO 14000. Markus Lange was supported by NSERC of Canada and also acknowledges financial support from the European Research Council (ERC) under the European Union Horizon 2020 research and innovation programme (ERC StG MaMBoQ, grant agreement No. 802901). It is also a pleasure to thank the CIRM, Luminy, for the REB (research in residence) program, where part of this work was done.

REFERENCES

- [1] S. Agmon, *Lower bounds for solutions of Schrödinger equations*, J. Analyse Math. **23** (1970), 1–25. MR 276624 doi:10.1007/BF02795485
- [2] S. Agmon, *Bounds on exponential decay of eigenfunctions of Schrödinger operators*, Schrödinger Operators (Berlin, Heidelberg) (Sandro Graffi, ed.), Springer Berlin Heidelberg, 1985, pp. 1–38. MR 824986, doi:10.1007/BFb0080331

- [3] M. Aizenman and B. Simon, *Brownian motion and Harnack inequality for Schrödinger operators*, Comm. Pure Appl. Math. **35** (1982), no. 22, 209–273. MR 644024, doi:10.1002/cpa.3160350206.
- [4] S. Avramška-Lukarska, D. Hundertmark, and H. Kovařík, *Absence of positive eigenvalues of magnetic Schrödinger operators*, arXiv:2003.07294 (2020). <https://arxiv.org/abs/2003.07294>
- [5] S. Barth and A. Bitter, *On the virtual level of two-body interactions and applications to three-body systems in higher dimensions*, Journal of Mathematical Physics **60** (2019), no. 11, 113504. MR 4032166, doi:10.1063/1.5120366.
- [6] S. Barth and A. Bitter. *Decay rates of bound states at the spectral threshold of multi-particle Schrödinger operators*. Doc. Math. **25** (2020), 721–735. MR 4129671, doi:DOI:10.25537/dm.2020v25.721-735.
- [7] S. Barth, A. Bitter, and S. Vugalter, *Decay properties of zero-energy resonances of multi-particle Schrödinger operators and why the Efimov effect does not exist for systems of $n \geq 4$ particles*, Preprint arXiv:1910.04139 (2019). <https://arxiv.org/abs/1910.04139>.
- [8] S. Barth, A. Bitter, and S. Vugalter. *The absence of the Efimov effect in systems of one- and two-dimensional particles*. J. Math. Phys. 62(12):Paper No. 123502, 46, 2021. MR 4348075, doi:10.1063/5.0033524.
- [9] R. D. Benguria and C. Yarur, *Sharp condition on the decay of the potential for the absence of a zero-energy ground state of the Schrodinger equation*, J. of Phys. A: Mathematical and General **23** (1990), no. 9, 1513–1518. MR 1048781, <http://stacks.iop.org/0305-4470/23/1513>.
- [10] D. Bollé, F. Gesztesy, and W. Schweiger, *Scattering theory for long-range systems at threshold*, J. Math. Phys. **26** (1985), no. 7, 1661–1674. MR 793308, doi:10.1063/1.526963.
- [11] H. L. Cycon, R. G. Froese, W. Kirsch, and B. Simon, *Schrödinger operators with application to quantum mechanics and global geometry*, study ed., Texts and Monographs in Physics, Springer-Verlag, Berlin, 1987. MR 883643
- [12] P. Deift, W. Hunziker, B. Simon, and E. Vock, *Pointwise bounds on eigenfunctions and wave packets in N -body quantum systems. IV*, Comm. Math. Phys. **64** (1978/79), no. 1, 1–34. MR 516993 URL: <http://projecteuclid.org/euclid.cmp/1103904619>.
- [13] S. A. Denisov and A. Kiselev, *Spectral properties of Schrödinger operators with decaying potentials*, Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon’s 60th birthday, Proc. Sympos. Pure Math., vol. 76, Amer. Math. Soc., Providence, RI, 2007, pp. 565–589. MR 2307748 doi:10.1090/pspum/076.2/2307748.
- [14] J. Dereziński and E. Skibsted, *Quantum scattering at low energies*, J. Funct. Anal. **257** (2009), no. 6, 1828–1920. MR 2540993 doi:10.1016/j.jfa.2009.05.026.
- [15] W. G. Faris. Quadratic forms and essential self-adjointness. *Helv. Phys. Acta*, 45:1074–1088, 1972/73. MR 383964, <http://doi.org/10.5169/seals-114428>.
- [16] S. Fournais and E. Skibsted, *Zero energy asymptotics of the resolvent for a class of slowly decaying potentials*, Math. Z. **248** (2004), no. 3, 593–633. MR 2097376 doi:10.1007/s00209-004-0673-9.
- [17] R. L. Frank and B. Simon, *Eigenvalue bounds for Schrödinger operators with complex potentials. II*, J. Spectr. Theory **7** (2017), no. 3, 633–658. MR 3713021 doi:10.4171/JST/173.
- [18] H.-W. Goelden. *On non-degeneracy of the ground state of Schrödinger operators*. Math. Z. **155** (1977), no. 3, 239–247. doi:10.1007/BF02028443.
- [19] D. K. Gridnev and M. E. Garcia, *Rigorous conditions for the existence of bound states at the threshold in the two-particle case*, J. Phys. A **40** (2007), no. 30, 9003–9016. MR 2344533 MR 609535, doi:10.1088/1751-8113/40/30/022.
- [20] V. Hoang, D. Hundertmark, J. Richter, and S. Wugalter. *Quantitative bounds versus existence of weakly coupled bound states for Schrödinger type operators*. arXiv:arXiv:1610.09891, 40 pages, 2017. URL: <https://arxiv.org/abs/1610.09891>
- [21] T. Hoffmann-Ostenhof, M. Hoffmann-Ostenhof, and R. Ahlrichs, *“Schrödinger inequalities” and asymptotic behavior of many-electron densities*, Phys. Rev. A **18** (1978), 328–334. URL: <https://link.aps.org/doi/10.1103/PhysRevA.18.328>, doi:10.1103/PhysRevA.18.328.

- [22] D. Hundertmark, M. Jex, and M. Lange. *Quantum systems at The Brink: Helium-type systems*. *arXiv:1908.04883* (2021), 62 pages. URL: <https://arxiv.org/abs/1908.04883>
- [23] D. Hundertmark and Y.-R. Lee, *On non-local variational problems with lack of compactness related to non-linear optics*, *J. Nonlinear Sci.* **22** (2012), no. 1, 1–38. MR 2878650
- [24] A. D. Ionescu and D. Jerison, *On the absence of positive eigenvalues of Schrödinger operators with rough potentials*, *Geom. Funct. Anal.* **13** (2003), no. 5, 1029–1081. MR 2024415 doi:10.1007/s00039-003-0439-2.
- [25] A. Jensen and T. Kato, *Spectral properties of Schrödinger operators and time-decay of the wave functions*, *Duke Math. J.* **46** (1979), no. 3, 583–611. URL: <http://projecteuclid.org/euclid.dmj/1077313577>.
- [26] K. Jörgens and J. Weidmann, *Spectral properties of Hamiltonian operators*, *Lecture Notes in Mathematics*, Vol. 313, Springer-Verlag, Berlin-New York, 1973. MR 0492941
- [27] K. Kaleta and J. Lőrinczi, *Zero-energy bound state decay for non-local Schrödinger operators*, *Commun. Math. Phys.* **374** (2020), no. 3, 2151–2191. MR 4076095, doi:10.1007/s00220-019-03515-3.
- [28] T. Kato. *Growth properties of solutions of the reduced wave equation with a variable coefficient*. *Comm. Pure Appl. Math.* **12** (1959), 403–425. MR 108633, doi:10.1002/cpa.3160120302.
- [29] C. E. Kenig, *Restriction theorems, Carleman estimates, uniform Sobolev inequalities and unique continuation*, *Harmonic analysis and partial differential equations (El Escorial, 1987)*, *Lecture Notes in Math.*, vol. 1384, Springer, Berlin, 1989, pp. 69–90. MR 1013816 doi:10.1007/BFb0086794.
- [30] C. E. Kenig and N. Nadirashvili, *A counterexample in unique continuation*, *Math. Res. Lett.* **7** (2000), no. 5-6, 625–630. MR 1809288 doi:10.4310/MRL.2000.v7.n5.a8.
- [31] M. Klaus and Barry Simon, *Coupling constant thresholds in nonrelativistic quantum mechanics. I. Short-range two-body case*, *Ann. Physics* **130** (1980), no. 2, 251–281. MR 610664 doi:10.1016/0003-4916(80)90338-3.
- [32] I. Knowles, *On the location of eigenvalues of second-order linear differential operators*, *Proc. Roy. Soc. Edinburgh Sect. A* **80** (1978), no. 1-2, 15–22. MR 529565 doi:10.1017/S030821050001009X.
- [33] I. Knowles, *On the number of L^2 -solutions of second order linear differential equations*, *Proc. Roy. Soc. Edinburgh Sect. A* **80** (1978), no. 1-2, 1–13. MR 529564 doi:10.1017/S0308210500010088.
- [34] H. Koch and D. Tataru, *Sharp counterexamples in unique continuation for second order elliptic equations*, *J. Reine Angew. Math.* **542** (2002), 133–146. MR 1880829 doi:10.1515/crll.2002.003.
- [35] L. D. Landau and E. M. Lifshitz, *Quantum mechanics: non-relativistic theory. Course of Theoretical Physics, Vol. 3*. Addison-Wesley Series in Advanced Physics. Pergamon Press, Ltd., London-Paris; for U.S.A. and Canada: Addison-Wesley Publishing Company, Inc., Reading, Mass., 1958. Translated from the Russian by J. B. Sykes and J. S. Bell.
- [36] H. Leinfelder and C. G. Simader, *Schrödinger operators with singular magnetic vector potentials*, *Math. Z.* **176** (1981), no. 1, 1–19. MR 606167 doi:10.1007/BF01258900.
- [37] D. Lenz and P. Stollmann, *On the decomposition principle and a Persson type theorem for general regular Dirichlet forms*, *J. Spectr. Theory* **9** (2019), no. 3, 1089–1113. MR 4003551 doi:10.4171/JST/272.
- [38] E. H. Lieb, *Thomas-Fermi and related theories of atoms and molecules*, *Rev. Modern Phys.* **53** (1981), no. 4, 603–641. MR 629207 doi:10.1103/RevModPhys.53.603.
- [39] D. Lundholm. *Some spectral bounds for Schrödinger operators with Hardy-type potentials*. *arXiv:0911.3386*, (2009). URL: <https://arxiv.org/abs/0911.3386>.
- [40] D. Lundholm. *Zero-energy states in supersymmetric matrix models*. PhD dissertation, KTH (2010). URL: <http://urn.kb.se/resolve?urn=urn:nbn:se:kth:diva-12846>.
- [41] S. Nakamura, *Low energy asymptotics for Schrödinger operators with slowly decreasing potentials*, *Comm. Math. Phys.* **161** (1994), no. 1, 63–76. MR 1266070 URL: <http://projecteuclid.org/euclid.cmp/1104269792>.
- [42] R. G. Newton, *Nonlocal interactions; the generalized Levinson theorem and the structure of the spectrum*, *J. Math. Phys.* **18** (1977), no. 8, 1582–1588. MR 446169 doi:10.1063/1.523466.
- [43] J. F. Perez, F. A. B. Coutinho, and C. P. Malta. *Logarithmic corrections to the uncertainty principle and infinitude of the number of bound states of N -particle systems*. *J. Math. Phys.* **27** (1986), no. 6, 1537–1540. doi:10.1063/1.527115.

- [44] A. Persson. *Bounds for the discrete part of the spectrum of a semi-bounded Schrödinger operator*. Math. Scand. **8** (1960), 143–153. doi:10.7146/math.scand.a-10602.
- [45] A. G. Ramm, *Sufficient conditions for zero not to be an eigenvalue of the Schrödinger operator*, J. Math. Phys. **28** (1987), no. 6, 1341–1343. MR 890004 doi:10.1063/1.527817.
- [46] A. G. Ramm, *Conditions for zero not to be an eigenvalue of the Schrödinger operator: ii*, J. Math. Phys. **29** (1988), no. 6, 1431–1432. arXiv:https://doi.org/10.1063/1.527935, doi:10.1063/1.527935.
- [47] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV. Analysis of Operators*, Academic Press, New York-London, 1978. MR 0493421
- [48] C. G. Simader, *An elementary proof of Harnack’s inequality for Schrödinger operators and related topics*, Math. Z. **203** (1990), no. 1, 129–152. MR 1030712 doi:10.1007/BF02570727.
- [49] B. Simon, *On positive eigenvalues of one-body Schrödinger operators*, Comm. Pure Appl. Math. **22** (1969), 531–538. MR 247300 doi:10.1002/cpa.3160220405.
- [50] B. Simon. *The bound state of weakly coupled Schrödinger operators in one and two dimensions*. Ann. Physics **97** (1976), no. 2, 279–288. doi:10.1016/0003-4916(76)90038-5.
- [51] B. Simon, *Large time behavior of the L^p norm of Schrödinger semigroups*, J. Functional Analysis **40** (1981), no. 1, 66–83. MR 607592 doi:10.1016/0022-1236(81)90073-2.
- [52] B. Simon, *Schrödinger semigroups*, Bulletin (New Series) of the American Mathematical Society **7** (1982), no. 3, 447–526. https://doi.org/, doi:bams/1183549767.
- [53] G. Teschl, *Mathematical methods in quantum mechanics*, second ed., Graduate Studies in Mathematics, vol. 157, American Mathematical Society, Providence, RI, 2014, With applications to Schrödinger operators. MR 3243083 doi:10.1090/gsm/157.
- [54] J. von Neumann and E. P. Wigner, *Über merkwürdige diskrete Eigenwerte*, pp. 291–293, Springer Berlin Heidelberg, Berlin, Heidelberg, 1993. doi:10.1007/978-3-662-02781-3_19.
- [55] D. R. Yafaev, *The low energy scattering for slowly decreasing potentials*, Comm. Math. Phys. **85** (1982), no. 2, 177–196. MR 675998 http://projecteuclid.org/euclid.cmp/1103921410.

DEPARTMENT OF MATHEMATICS, INSTITUTE FOR ANALYSIS, KARLSRUHE INSTITUTE OF TECHNOLOGY, 76128 KARLSRUHE, GERMANY, AND DEPARTMENT OF MATHEMATICS, ALTGELD HALL, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, 1409 W. GREEN STREET, URBANA, IL 61801, USA

Email address: dirk.hundertmark@kit.edu

DEPARTMENT OF PHYSICS, FACULTY OF NUCLEAR SCIENCES AND PHYSICAL ENGINEERING, CZECH TECHNICAL UNIVERSITY IN PRAGUE, BŘEHOVÁ 7, 11519 PRAGUE, CZECH REPUBLIC, AND CEREMADE, DAUPHINE UNIVERSITY, PLACE DU MARÉCHAL DE LATTRE DE TASSIGNY, 75775 PARIS CEDEX 16, FRANCE

Email address: michal.jex@fjfi.cvut.cz

SISSA, MATHEMATICS AREA, VIA BONOMEA 265, 34136 TRIESTE, ITALY

Email address: mlang@sisssa.it