

VOLUME 29 NUMBER 1 January 2023

pISSN 2287-2728  
eISSN 2387-285X

# CLINICAL and MOLECULAR HEPATOLOGY

The forum for latest knowledge of hepatobiliary diseases



HCC incidence is decreasing in Korea  
but increasing in elderly

Early changes in biomarkers predict HBsAg response  
Baveno-VII predicts decompensation in cACLD

## Editorial

# New biomarkers of hepatitis B virus (HBV) infection: HBV RNA and HBV core-related antigen, new kids on the block?

Young-Suk Lim

Department of Gastroenterology, Liver Center, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

**Keywords:** Hepatitis B virus RNA; Hepatitis B virus core-related antigen; Hepatitis B; Biomarkers

See Article on Page 146

Through long-term treatment with nucleos(t)ide analogues (NUC), the prevention of disease progression to end-stage liver disease and reduction in the risk of hepatocellular carcinoma (HCC) is achievable in most patients with chronic hepatitis B (CHB). However, functional cure (hepatitis B surface antigen [HBsAg] seroclearance) is very rarely achievable even with long-term NUC treatment, and the optimal timing of treatment initiation and discontinuation remains debatable.

With recent advances in molecular analysis, several new biomarkers of the hepatitis B virus (HBV) have been identified, including quantification of HBsAg (qHBsAg), HBV RNA, and HBV core-related antigen (HBcrAg).<sup>1,2</sup> Integration of these biomarkers with the conventional ones, such as the hepatitis B e antigen (HBeAg) test and HBV DNA quantitation, may improve our understanding of the natural history of CHB and its response to antiviral therapy. All of qHBsAg, HBV RNA, and HBcrAg have been proposed as surrogate markers of HBV co-

valently closed circular DNA (cccDNA) activity. Studies have shown the potential utility of these novel biomarkers in a range of clinical settings, such as monitoring response to anti-viral therapy, predicting relapse after treatment cessation or estimating the risk of HCC.<sup>3,4</sup> However, HBsAg may be produced from both cccDNA and integrated viral genomes and its correlation with intrahepatic cccDNA is particularly weak in the HBeAg-negative patients. Therefore, attention has shifted to other serum biomarkers such as HBV RNA and core related antigen.

In this issue of Mak et al.,<sup>5</sup> a new role of HBV pre-genomic (pg) RNA and HBcrAg in predicting favourable HBsAg response (FHR; <100 IU/mL or HBsAg seroclearance) during median 17 years of NUC treatment. For HBeAg-positive patients, serum HBV pgRNA decline at week 4 was significantly greater for patients with FHR compared to non-FHR patients (5.49 vs. 4.32 log copies/mL, respectively). For HBeAg-negative patients, instead of increase in serum HBcrAg from baseline in non-FHR patients, FHR patients had median reduction in HBcrAg at week 4 (increment of 1.75 vs. reduction of 2.98 log U/mL). This may have a significance as a new study that

### Corresponding author : Young-Suk Lim

Department of Gastroenterology, Asan Medical Center, University of Ulsan College of Medicine, 88 Olympic-ro 43-gil, Songpa-gu, Seoul 05505, Korea  
Tel: +82-2-3010-5933, Fax: +82-2-485-5782, E-mail: limys@amc.seoul.kr  
<https://orcid.org/0000-0002-1544-577X>

**Editor:** Seung Up Kim, Yonsei University College of Medicine, Korea

**Received :** Nov. 20, 2022 / **Received :** Nov. 22, 2022 / **Accepted :** Nov. 23, 2022

showed the usefulness of HBV pgRNA and HBcrAg in predicting FHR during NUC treatment. However, its clinical implication is limited because this study cannot address whether early biomarker response can help to predict successful off-NA virological control. Nonetheless, the results of this study could be used as companion diagnostic tests in the clinical trials to develop new novel drugs to induce CHB functional cure.

It is noteworthy that the HBV RNA and HBcrAg were significantly lower in all time points among HBeAg-negative patients compared to HBeAg-positive patients, a finding that is consistent with previous reports. This is especially true for HBcrAg due to the poor detectability in HBeAg-negative patients because HBeAg is part of HBcrAg (which consists of HBcAg, HBeAg and p22cr). Therefore, the analyses were performed in HBeAg-positive and HBeAg-negative patients separately.

Overall, the use of novel biomarkers for HBV infection carries enormous potential, and it is possible that new biomarker-based models, used in combination with traditional ones, will become an integral part of our daily practice in near future as well as in the development of new drugs. The addition of HBV RNA and HBcrAg to our armamentarium as new biomarkers should be embraced and will be beneficial in our efforts to eliminate HBV. However, before using these biomarkers, the assays for these biomarkers should be standardized with their improvement in the detection sensitivity and

also availability.

### Conflicts of Interest

YS Lim is an advisory board member of Gilead Sciences and receives research funding from Gilead Sciences. No other disclosures are reported.

### REFERENCES

1. Wu JW, Kao JH, Tseng TC. Three heads are better than two: hepatitis B core-related antigen as a new predictor of hepatitis B virus-related hepatocellular carcinoma. *Clin Mol Hepatol* 2021;27:524-534.
2. Inoue T, Tanaka Y. Novel biomarkers for the management of chronic hepatitis B. *Clin Mol Hepatol* 2020;26:261-279.
3. Lok J, Dusheiko G, Carey I, Agarwal K. Review article: novel biomarkers in hepatitis B infection. *Aliment Pharmacol Ther* 2022;56:760-776.
4. Kramvis A, Chang KM, Dandri M, Farci P, Glebe D, Hu J, et al. A roadmap for serum biomarkers for hepatitis B virus: current status and future outlook. *Nat Rev Gastroenterol Hepatol* 2022;19:727-745.
5. Mak LY, Wong D, Kuchta A, Hilfiker M, Hamilton A, Chow N, et al. Hepatitis B virus pre-genomic RNA and hepatitis B core-related antigen reductions at week 4 predict favourable hepatitis B surface antigen response upon long-term nucleos(t)ide analogue in chronic hepatitis B. *Clin Mol Hepatol* 2023;29:146-162.

---

#### Abbreviations:

cccDNA, covalently closed circular DNA; CHB, chronic hepatitis B; FHR, favourable HBsAg response; HBcrAg, HBV core-related antigen; HBeAg, hepatitis B e antigen; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; NUC, nucleos(t)ide analogues; pg, pre-genomic; qHBsAg, quantification of HBsAg