

VOLUME 29 NUMBER 1 January 2023

pISSN 2287-2728  
eISSN 2387-285X

# CLINICAL and MOLECULAR HEPATOLOGY

The forum for latest knowledge of hepatobiliary diseases



HCC incidence is decreasing in Korea  
but increasing in elderly

Early changes in biomarkers predict HBsAg response  
Baveno-VII predicts decompensation in cACLD

## Review

# Wnt signaling in liver regeneration, disease, and cancer

Gengyi Zou<sup>1</sup> and Jae-Il Park<sup>1,2</sup>

<sup>1</sup>Department of Experimental Radiation Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX; <sup>2</sup>Genetics and Epigenetics Program, The University of Texas MD Anderson Cancer Center Graduate School of Biomedical Sciences, Houston, TX, USA

The liver exhibits the highest recovery rate from acute injuries. However, in chronic liver disease, the long-term loss of hepatocytes often leads to adverse consequences such as fibrosis, cirrhosis, and liver cancer. The Wnt signaling plays a pivotal role in both liver regeneration and tumorigenesis. Therefore, manipulating the Wnt signaling has become an attractive approach to treating liver disease, including cancer. Nonetheless, given the crucial roles of Wnt signaling in physiological processes, blocking Wnt signaling can also cause several adverse effects. Recent studies have identified cancer-specific regulators of Wnt signaling, which would overcome the limitation of Wnt signaling target approaches. In this review, we discussed the role of Wnt signaling in liver regeneration, precancerous lesion, and liver cancer. Furthermore, we summarized the basic and clinical approaches of Wnt signaling blockade and proposed the therapeutic prospects of cancer-specific Wnt signaling blockade for liver cancer treatment. (**Clin Mol Hepatol 2023;29:33-50**)

**Keywords:** WNT signaling pathway; Liver regeneration; Liver diseases; Hepatocellular carcinoma

## INTRODUCTION

Liver regeneration has been extensively studied.<sup>1-3</sup> *In vivo* studies have shown that partial hepatectomy or chemical injury activates extracellular and intracellular signaling pathways, leading to liver regeneration. Hepatocyte loss during chronic liver diseases triggers compensatory proliferation of the surviving hepatocytes.<sup>4-6</sup> Apart from liver regeneration in

physiological conditions, genotoxic risk factors might lead them to convert to neoplasia. Hepatitis virus, alcohol abuse, non-alcoholic fatty liver disease (NAFLD), and aflatoxin-B1 exposure are also the main etiological factors to induce the development of precancerous lesions in the liver. Liver cancer is one of the top 10 lethal cancers worldwide. Its estimated death rate in 2021 is 6% in males and 4% in females.<sup>7</sup> Liver cancer consists of hepatocellular carcinoma (HCC), cholan-

### Corresponding author : Gengyi Zou

Department of Experimental Radiation Oncology, The University of Texas MD Anderson Cancer Center, 6565 MD Anderson Blvd Unit 1054, Houston, TX 77030, USA

Tel: +1-713-792-3659, Fax: +1-713-794-5369, E-mail: gzou@mdanderson.org  
<https://orcid.org/0000-0002-5399-8674>

### Jae-Il Park

Department of Experimental Radiation Oncology, The University of Texas MD Anderson Cancer Center, 6565 MD Anderson Blvd. Unit 1052, Houston, TX 77030, USA

Tel: +1-713-792-3659, Fax: +1-713-794-5369, E-mail: jaeil@mdanderson.org  
<https://orcid.org/0000-0002-0737-2654>

giocarcinoma (CCA), hepatoblastoma (HB), and several other rare tumors (angiosarcoma, intraductal papillary neoplasm of the bile duct, and mucinous cystic neoplasm). HCC is the most common primary liver cancer frequently developed with chronic liver disease, such as cirrhosis caused by hepatitis virus infection.<sup>8</sup>

Among various signaling pathways associated with liver biology,<sup>9-12</sup> Wnt signaling is involved in all stages of liver disease progression, from liver injury to inflammation, fibrosis, cirrhosis, and tumorigenesis. Several Wnt ligands are secreted by various hepatic cells, including hepatocytes, stellate cells, Kupffer cells, biliary epithelial cells, and sinusoidal endothelial cells.<sup>13-16</sup> Based on the oncogenic roles of Wnt signaling in cancer, several components and regulators of Wnt signaling have been proposed as the druggable targets to improve the current therapeutic efficacy in the liver cancer treatment.<sup>17</sup>

Herein, we review the roles of Wnt signaling in liver regeneration and liver tumorigenesis and the therapeutic targets of Wnt signaling in liver cancer treatment.

## Wnt SIGNALING

Wnt signaling is evolutionarily conserved and orchestrates various cellular processes, including cell proliferation, differentiation, migration, polarity, stemness, and lineage plasticity.<sup>18,19</sup> Consequently, Wnt signaling plays a pivotal role in organogenesis, tissue homeostasis, tissue regeneration, and tumorigenesis.<sup>20-25</sup> The Wnt signaling is triggered by the binding of the Wnt ligands to the frizzled (FZD) receptors. The mammals have 19 Wnt ligands and 10 FZD receptors,<sup>26</sup> resulting in the complexity and specificity in Wnt signaling activation. Based on the involvement of  $\beta$ -catenin, a key component of Wnt signaling, Wnt signaling is generally classified


into canonical ( $\beta$ -catenin-mediated) and non-canonical ( $\beta$ -catenin-independent) Wnt signaling (Fig. 1). In the canonical Wnt/ $\beta$ -catenin pathway, the protein destruction complex (casein kinase 1 [CK1], glycogen synthase kinase 3 [GSK3], adenomatous polyposis coli [APC], and axis inhibition proteins [AXINs]) targets the  $\beta$ -catenin protein for degradation via CK1 and GSK3-mediated sequential phosphorylation at the N-terminus (Ser-45, Thr-41, Ser-37, and Ser-33) of  $\beta$ -catenin followed by  $\beta$ -TrCP, an E3 ligase, recruitment. Conversely, binding of the canonical Wnt ligands to the FZD receptors and LRP5/6 co-receptors activates dishevelled (DVL), which inhibits the protein destruction complex. As a result,  $\beta$ -catenin protein is stabilized and translocated into the nucleus to transactivate the canonical Wnt target genes by replacing the co-repressors associated with the T-cell factor/lymphoid enhancer-binding factor (TCF/LEF) with the co-activators. Non-canonical Wnt signaling pathways include the planar cell polarity pathway (involved in c-Jun N-terminal kinase [JNK] activation, small GTPase activation, and cytoskeletal rearrangement), and the Wnt/ $\text{Ca}^{2+}$  pathway (activating phospholipase C [PLC] and protein kinase C [PKC]).<sup>18,19</sup>

## Wnt SIGNALING IN LIVER REGENERATION

Upon partial hepatectomy or acute liver injury, the number of hepatocytes is drastically reduced. Various signaling pathways (epidermal growth factor [EGF], hepatocyte growth factor [HGF], Wnt/ $\beta$ -catenin, and Notch) stimulate the hepatocytes in the G0 phase to proliferate, compensating tissue loss and restoring the physiological functions of the liver.<sup>27-29</sup> During liver regeneration, endothelial cells under shear stress produce Wnts to activate Wnt/ $\beta$ -catenin signaling in hepatocytes. Additionally, the organ precisely senses the size of the

### Abbreviations:

AFB1, aflatoxin type B1; ALD, alcoholic liver disease; APC, adenomatous polyposis coli; AXIN, axis inhibition protein; CCA, cholangiocarcinoma;  $\text{CCl}_4$ , carbon tetrachloride; CEBPA, CCAAT enhancer-binding protein  $\alpha$ ; CK1, casein kinase 1; CREBBP/CBP, CREB binding protein; CV, central vein; DDC, 1,4-dihydro-2,4,6-trimethyl-pyridine-3,5-di carboxylate; DEN, diethylnitrosamine; DKK1, Dickkopf 1; DVL, dishevelled; E2F1, E2F transcription factor 1; EGF, epidermal growth factor; EMT, epithelial-mesenchymal transition; ERK, extracellular signal-regulated kinase; EtOH, ethanol; FZD, frizzled; GPC3, glypican-3; GSK3, glycogen synthase kinase 3; HB, hepatoblastoma; HBsAg, hepatitis B virus surface antigen; HBV, hepatitis B virus; HCC, hepatocellular carcinoma; HCV, hepatitis C virus; HGF, hepatocyte growth factor; Ig, immunoglobulin; IGF1, insulin-like growth factor 1; JNK, c-Jun N-terminal kinase; KO, knock-out; LGR5, leucine-containing repeat G-protein-coupled receptor 5; LOF, loss-of-function; LRP6, lipoprotein receptor-related protein 6; MAPK, mitogen-activated protein kinase; MCP, monocyte chemoattractant protein; MMP7, matrix metalloproteinase; mTOR, mammalian target of rapamycin; MVBs, multivesicular bodies; NAFLD, non-alcoholic fatty liver disease; NCOA2, nuclear receptor coactivator 2; NF- $\kappa$ B, nuclear factor kappa-light-chain-enhancer of activated B cells; peg-IFN, pegylated-Interferon- $\alpha$ 2a; PI3K, phosphoinositide 3-kinase; PKC, protein kinase C; PLC, phospholipase C; PORCN, porcupine; PPAR $\gamma$ , peroxisome proliferator-activated receptor  $\gamma$ ; RanBP3, Ran-binding protein 3; ROS, reactive oxygen species; SAL, salinomycin; SFRP1, secreted frizzled-related protein 1; sFZD7, secreted FZD7; SREBF, sterol regulatory element binding transcription factor; TCF/LEF, T-cell factor/lymphoid enhancer-binding factor; TMEM9, transmembrane protein 9; TNKS, tankyrase; VEGF, vascular endothelial growth factor; WFI1, Wnt inhibitory factor 1; WISP1, Wnt1-inducible-signaling pathway protein 1




**Figure 1.** Wnt signaling. Illustration of canonical and non-canonical Wnt signaling. The hallmark of the canonical Wnt/ $\beta$ -catenin pathway is the stabilization and nuclear translocation of  $\beta$ -catenin. In the absence of Wnt ligands, cytoplasmic  $\beta$ -catenin is degraded by the destruction complex (Axin, APC, GSK3 $\beta$ , and CK1 $\alpha$ ). Upon Wnt ligand binding to Frizzled receptors (FZDs) and LRP, the destruction complex is inhibited,  $\beta$ -catenin protein is stabilized in the cytosol and translocated into the nucleus. Nuclear  $\beta$ -catenin then recruits transcriptional coactivator CREBBP to transactivate target genes in conjunction with TCF/LEF transcription factors. Additionally, FZDs are ubiquitinated by ZNRF3 and RNF43 E3 ligases, which are inhibited by R-spondin binding to LGR5, increasing the cells' sensitivity to Wnt ligands. In Wnt/PCP signaling, Wnt ligands bind to FZDs or their co-receptors (ROR and RYK) to trigger a cascade reaction, involving the small GTPases RhoA and Ras-related C3 botulinum toxin substrate (Rac), then activating Rho-associated protein kinases (ROCKs) and JUN N-terminal kinases (JNK), respectively. These lead to cytoskeletal rearrangements and/or transcriptional responses such as ATF2. In Wnt/ $\text{Ca}^{2+}$  signaling, the activation of phospholipase C (PLC) triggers the release of  $\text{Ca}^{2+}$  from the endoplasmic reticulum (ER), which promotes the transcription of nuclear factor of activated T cells (NFAT) through several intermediate steps. Created with BioRender.com. LRP, lipoprotein receptor-related protein; LGR5, leucine-containing repeat G-protein-coupled receptor 5; RNF43, ring finger protein 43; ZNRF3, zinc and ring finger 3; GSK-3 $\beta$ , glycogen synthase kinase 3 $\beta$ ; CK1 $\alpha$ , casein kinase 1 $\alpha$ ; APC, adenomatous polyposis coli; CBP, CREB binding protein; TCF/LEF, T-cell factor/lymphoid enhancer-binding factor; ROR, receptor tyrosine kinase-like orphan receptor; RYK, receptor tyrosine kinase; ATF2, activating transcription factor 2; PIP2, phosphatidylinositol 4,5-bisphosphate; DAG, diacylglycerol; PKC, protein kinase C.

regenerating liver and adjusts its size to 100%.<sup>2</sup>

Several animal models (rat, mouse, and zebrafish) were utilized for liver regeneration study.<sup>30-33</sup> The partial hepatectomy is the classic strategy to create the murine liver regeneration model.<sup>30</sup> Carbon tetrachloride (CCl<sub>4</sub>) is a frequently used chemical to induce liver injury in rats and mice.<sup>34</sup> Meanwhile, several dietary-induced liver injury models are also commonly used.<sup>35</sup> Biliary injury and regeneration can be induced by the 1,4-dihydro-2,4,6-trimethyl-pyridine-3,5-dicarboxylate (DDC) diet.<sup>35</sup> Besides murine models, zebrafish emerged as a potent model for drug screening of liver regeneration.<sup>32,33</sup> Partial hepatectomy, drug-induced liver injury, and nitroreductase-mediated hepatocyte ablation were employed to estab-

lish the zebrafish liver injury model.<sup>32,33,36,37</sup>

Transient activation of the canonical Wnt signaling is indispensable for liver regeneration (Fig. 2).<sup>13,15,27</sup> In rat models, overexpressed Wnt1 and nuclear  $\beta$ -catenin are predominantly accumulated in remaining parenchymal cells after 70% partial hepatectomy. The level of  $\beta$ -catenin increased within 5 minutes after hepatectomy, accompanied by its nuclear translocation and subsequent target gene expression for hepatocyte proliferation.<sup>27</sup> Significantly, genetic ablation of  $\beta$ -catenin/Ctnnb1 impairs liver regeneration of mice from partial hepatectomy.<sup>38</sup> The liver-specific *Ctnnb1* knock-out (KO) delayed DNA synthesis and hepatocyte proliferation in mice after partial hepatectomy. Conversely, activation of



**Figure 2.** Wnt signaling in liver regeneration. In normal liver, most hepatocytes are polyploid with random chromosomal deletions. Upon liver injury, the increased narrow portal vein pressure stimulates the initiating signals for liver regeneration. The activation of Wnt signaling is crucial in liver regeneration. Moreover, in chronic liver injury, the ROS and lipid peroxide are the risk factors damaging the reproducing hepatocytes, leading to precancerous lesion development. Created with BioRender.com. ROS, reactive oxygen species.

Wnt/ $\beta$ -catenin signaling accelerates liver regeneration in the zebrafish model.<sup>39</sup> It was also shown that liver damage up-regulated leucine-containing repeat G-protein-coupled receptor 5 (LGR5) and AXIN2 in the hepatocytes.<sup>40</sup> LGR5 is a marker of actively dividing stem and progenitor cells in Wnt-driven self-renewing tissues.<sup>41</sup> LGR5 interacts with FZD and lipoprotein receptor-related protein 6 (LRP6) to enhance phosphorylation of LRP6, which in turn enhances the Wnt/ $\beta$ -catenin signaling.<sup>42</sup> While Lgr5 is not expressed in healthy adult livers, after liver damage, Lgr5+ cells appear near the bile ducts, consistent with strong activation of the Wnt signaling.<sup>41</sup> AXIN2 is another Wnt downstream target gene transactivated by  $\beta$ -catenin.<sup>43</sup> Like AXIN1, AXIN2 combined with other destruction complex components degrades  $\beta$ -catenin, serving as a negative feedback regulator of the Wnt signaling.<sup>44</sup>

Other than core components of Wnt signaling, additional regulators of Wnt signaling were implicated in liver regeneration. Recently, our group identified the transmembrane protein 9 (TMEM9) gene as an amplifier of Wnt/ $\beta$ -catenin signaling. TMEM9 is a type I transmembrane protein primarily localized in lysosomes and multivesicular bodies (MVBs). While the ablation of TMEM9 inhibits the activity of the Wnt/ $\beta$ -catenin signaling,  $\beta$ -catenin transactivates TMEM9, leading to hyperactivation of Wnt/ $\beta$ -catenin signaling.<sup>45</sup> Interestingly, TMEM9 is highly expressed in hepatocytes around the central vein (CV) of regenerating liver.<sup>46</sup> TMEM9 hyperactivates Wnt/ $\beta$ -catenin signaling to promote liver regeneration through

lysosomal degradation of APC protein.<sup>46</sup> *Tmem9* KO impairs CCl4-induced liver regeneration with downregulation of Wnt/ $\beta$ -catenin signaling.<sup>46</sup>


In addition to the role of Wnt/ $\beta$ -catenin signaling in regeneration, sustained activation of the Wnt signaling is associated with the progression of chronic liver diseases and liver tumorigenesis (Fig. 2). Additionally, reactive oxygen species (ROS) and lipid peroxide are the risk factors for the development of the precancerous lesion in the liver.<sup>47,48</sup> However, the crosstalk between Wnt signaling and ROS has not been fully revealed in the liver. It was reported that  $\beta$ -catenin can be further stabilized by ROS.<sup>49</sup> Meanwhile, lipid peroxidation products mainly generated by ROS activate the canonical Wnt pathway through oxidative stress.<sup>50</sup> Therefore, it is likely the potential crosstalk between Wnt signaling and ROS might contribute to liver cancer development.

Accumulating evidence suggests that many chronic liver diseases contribute to liver cancer development, described below.

## Wnt SIGNALING IN PRECANCEROUS LIVER LESION

### Hepatitis virus

Globally distributed hepatitis B virus (HBV) and hepatitis C virus (HCV) are the crucial triggers of HCC initiation. Both HBV



**Figure 3.** Wnt signaling in liver cancer. Dynamic activation of  $\beta$ -catenin and Wnt signaling-related gene mutations from risk factor exposure to final liver cancer. With the precancerous lesions induced by hepatitis virus, NAFLD, alcohol assumption, or aflatoxin-B1, genetic and epigenetic alteration (e.g., mutations in the *CTNNB1* or *AXIN1* genes) lead to the accumulation and nuclear translocation of  $\beta$ -catenin, resulting in initiating liver cancer development. Created with BioRender.com. NAFLD, non-alcoholic fatty liver disease.

and HCV can induce chronic infections and are essential pathogenic factors in cirrhosis and liver cancer (Fig. 3).<sup>51,52</sup> The epidemiological data show that more than 70% of patients with liver cancer have HBV infection, 10–20% have HCV infection, and a significant proportion of patients have both HBV and HCV infection.<sup>53–55</sup>

After infection, the DNA of HBV is integrated into the host genome, inducing genomic instability and transactivation of cancer-related genes, which culminates in the formation of early cancer cell clones. Mechanistically, HBV contributes to HCC development through direct and indirect means.<sup>56</sup> Direct mechanisms include virus mutations, HBV DNA integration, growth regulatory genes activation by HBV-encoded proteins.<sup>57</sup> Indirect mechanisms include the activation of cellular oncogenes associated with HBV DNA integration, genetic instability induced by viral integration or the regulatory protein HBx, and the development of liver disease mediated by immune enhancement due to viral proteins.<sup>58</sup>

Both hepatitis B virus surface antigen (HBsAg) and HBx modulate the expressions of genes involved in Wnt signaling activation. HBsAg activates the transcription factor LEF1 of the Wnt signaling.<sup>59</sup> The X protein encoded by the hepatitis B virus has a vital role in stimulating viral gene expression and replication, critical for maintaining chronic carrier status. HBx, a 17 kDa multifunctional protein, upregulates the expression of Wnt ligands (WNT1 and WNT3), the receptor (FZD2 and FZD7), a component of the destruction complex (GSK3 $\beta$ ), E-cadherin, and Wnt1-inducible-signaling pathway protein 1 (WISP1), a suppressor of Wnt antagonists (secreted frizzled-related protein 1 [SFRP1] and SFRP5). On the other hand, the

Wnt signaling key components ( $\beta$ -catenin and AXIN1) are highly mutated in HBV-associated HCC. Loss-of-function (LOF) mutations of the *AXIN1* are observed in HBV-HCC patients. In HBV and/or HCV-associated HCC patients, the most frequent mutation in the *CTNNB1* gene is enriched in the exon 3 encoding the N-terminal phosphorylation sites.<sup>60–62</sup> These aberrantly controlled genes in Wnt signaling subsequently promote and lead to the development of HCC.<sup>63–65</sup>

The oncogenic mechanism of HCV in liver cancer is mainly mediated by Wnt/ $\beta$ -catenin signaling hyperactivation via the core protein and two nonstructural proteins, NS3 and NS5A.<sup>66</sup> The core protein (HCV core antigen) is a significant component of HCV. It regulates hepatocyte transcription and promotes Wnt/ $\beta$ -catenin signaling by upregulating Wnt ligands (WNT1 and WNT3A), FZD receptors, and LRP5/6.<sup>67,68</sup> Additionally, at the early stage of HCV infection, the secreted Wnt antagonists, SFRP2 and Dickkopf 1 (DKK1), are downregulated by their promoter hypermethylation.<sup>69,70</sup> HCV core protein also promotes hypermethylation of the *CDH1* gene promoter region,<sup>71</sup> destabilizing the cadherin-catenin-actin complex for  $\beta$ -catenin release and activation.<sup>72</sup> NS5A stabilizes  $\beta$ -catenin via activating phosphoinositide 3-kinase (PI3K)/AKT, leading to GSK3 $\beta$  inactivation followed by inhibiting the protein-destruction complex-mediated  $\beta$ -catenin degradation for Wnt target gene activation. At the early stage of viral infection, HCV-activated Wnt/ $\beta$ -catenin signaling also promotes liver fibrosis by enhancing the activation and survival of hepatic stellate cells.<sup>17,73,74</sup>

## Alcohol abuse

Alcohol is a well-known risk factor for liver cancer. Alcoholic liver disease (ALD) is a chronic liver disease caused by long-term alcohol consumption (Fig. 3). ALD is characterized by the fatty liver at the beginning, then progressed to alcoholic hepatitis, liver fibrosis, and cirrhosis, which is pathologically associated with the precancerous lesions of HCC. *In vivo*, ethanol (EtOH) is metabolized into the reactive metabolite acetaldehyde, promoting liver tumorigenesis. Mice administered with the chemical carcinogen, diethylnitrosamine (DEN), for 7 weeks and the subsequent EtOH feeding for 16 weeks exhibited the increased total number of cancer foci and liver tumors.<sup>75</sup> Also, these tumors showed a 3- to 4-fold increase in the expression of proliferation markers and an increased expression of  $\beta$ -catenin, compared to non-tumor hepatocytes.<sup>75</sup> In a rat model of chronic liver disease, EtOH-treated liver was accompanied by the increased proliferation of hepatocytes, depletion of retinol and retinoic acid storage, augmented expression of phospho-GSK3 $\beta$  at the cell membrane, significant upregulation of soluble Wnt ligands (Wnt2 and Wnt7a), accumulation of nuclear  $\beta$ -catenin, and upregulation of  $\beta$ -catenin target genes (cyclin D1/CCND1, c-Myc/MYC, WISP1, and matrix metalloproteinase [MMP7]). These data suggest that long-term EtOH consumption activates the Wnt/ $\beta$ -catenin signaling and increases hepatocyte proliferation, promoting liver tumorigenesis.<sup>75</sup> Additionally, ROS accumulation, nuclear factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa$ B)-dependent vascular endothelial growth factor (VEGF) and monocyte chemoattractant protein (MCP)-1 upregulation, and activation of extracellular signal-regulated kinase (ERK)-mitogen-activated protein kinase (MAPK) signaling also contribute to EtOH-induced liver tumorigenesis.<sup>76-80</sup>

## NAFLD

The increasing prevalence of NAFLD was caused by an over-nourished lifestyle.<sup>81,82</sup> NAFLD is characterized by fat accumulation in the liver, evolving to end-stage liver diseases such as cirrhosis and HCC (Fig. 3).<sup>83</sup> The main risk factors of NAFLD include central obesity, overnutrition, insulin resistance, and metabolic syndrome.<sup>84</sup> In severe NAFLD, many tissue repair-related genes (TMEM204, FGFR2, matrix molecules, and matrix remodeling factors) were hypomethylated

at their promoters and overexpressed. Conversely, genes in specific metabolic pathways (lipid metabolism, cytochrome P450 family, multidrug resistance, and fatty acid anabolic pathways) were hypermethylated and silenced.<sup>85</sup> Hyperinsulinemia is one of the risk factors of NAFLD.<sup>86</sup> SOX17 plays a vital role in regulating insulin secretion. *Sox17* KO mice display high susceptibility to high-fat diet-induced hyperglycemia and diabetes.<sup>87</sup> SOX17 directly interacts with the TCF/LEF transcription factor to repress the transcription of Wnt signaling target genes. The methylation of the *SOX17* promoter is a frequent event in human cancers. Epigenetic silencing of *SOX17* contributes to the aberrant activation of Wnt/ $\beta$ -catenin signaling,<sup>88</sup> accelerating progression from NAFLD to HCC. Besides,  $\beta$ -catenin inhibits the expression of CCAAT enhancer-binding protein  $\alpha$  (CEBPA) and peroxisome proliferator-activated receptor  $\gamma$  (PPARG), which in turn inhibits the preadipocyte differentiation.<sup>89</sup> As the co-receptor of the Wnt/ $\beta$ -catenin signaling, LRP6 induces lipid accumulation in the liver via insulin-like growth factor 1 (IGF1)/AKT/mammalian target of rapamycin (mTOR)/sterol regulatory element binding transcription factor (SREBF) 1/2 signaling. Intriguingly, inhibiting the non-canonical Wnt signaling reduces lipid accumulation and inflammation.<sup>90</sup> Therefore, while reducing the effects of NAFLD risk factors, inhibition of the Wnt signaling is also essential for attenuating the development of NAFLD and preventing the initiation of HCC.

## Aflatoxin-B1 exposure

Among the aflatoxins, aflatoxin type B1 (AFB1) primarily targets the liver as a highly potent hepatotoxin and hepatocarcinogen (Fig. 3). AFB1 impairs DNA repair processes, resulting in severe DNA mutagenesis, and also inhibits DNA and RNA metabolism. This pathological event ultimately leads to excessive liver lipid accumulation, liver enlargement, bile duct epithelial hyperplasia, and liver cancer. The potency of aflatoxin to cause liver cancer is significantly enhanced in the presence of HBV infection. Under chronic HBV infection, cytochrome P450s could metabolize inactive AFB1 to mutagenic AFB1-8,9-epoxide. Also, the infection leads to hepatocyte necrosis and regeneration, producing oxygen and nitrogen reactive species and increasing the incidence of AFB1-induced mutagenesis.<sup>91</sup> Clinical studies have shown that *CTNNB1* mutations are present in approximately one-quarter of HCC in areas with low aflatoxin B1 exposure. Interestingly,

these *CTNNB1* mutations were similar to those previously reported in the human HCC.<sup>92</sup>

## Wnt SIGNALING IN LIVER CANCER

### HCC

HCC is a common and fatal malignancy worldwide.<sup>93</sup> Regardless of the risk factors mentioned above, aberrant hyperactivation of Wnt/ $\beta$ -catenin signaling is observed in 95% of HCCs.<sup>94</sup> The most common genetic mutations of the Wnt signaling in HCC are the gain-of-function mutations in the *CTNNB1* gene encoding  $\beta$ -catenin,<sup>61,95</sup> which is somewhat distinct from colorectal cancer where Wnt/ $\beta$ -catenin signaling hyperactivation is mainly driven by the *APC* gene inactivation.<sup>96</sup> Missense mutations of *CTNNB1* exon 3 were observed in 18.1% of HCC cases. Missense mutations at codons 32, 33, 38, or 45 of the *CTNNB1* gene lead to the unphosphorylation of the N-terminus of  $\beta$ -catenin for its stabilization, nuclear translocation, and target gene transactivation.<sup>60</sup> Secondly, the LOF mutations in the *AXIN1* gene were observed in 5–19% of HCC cases.<sup>97</sup> *CTNNB1* and *AXIN1* mutations occur in patients with advanced HCC (Fig. 3).<sup>98–100</sup> Importantly, hyperactivation of the Wnt signaling is considered a hallmark of advanced HCC.<sup>101</sup> It should also be noted that mutations in the *CTNNB1* and *AXIN1* genes lead to different HCC subtypes accompanied by distinct clinical and pathological features. *CTNNB1* mutations are associated with less aggressive HCC, including chromosomally stable and highly differentiated tumors,<sup>102</sup> with a better prognosis.<sup>95</sup> In contrast, *AXIN1* mutations occur more frequently in more aggressive HCC tumors characterized by hypodifferentiated tumor cells and chromatin instability.<sup>102</sup> Consistently, the HCC tumors with *CTNNB1* mutations or *AXIN1* mutations showed different target gene expression.<sup>61,95,103</sup>

### CCA

CCA is ranked as the second most common hepatobiliary cancer after HCC. CCA originates mainly from differentiated bile duct epithelial cells.<sup>104</sup> CCA is often diagnosed at an advanced stage with a poor prognosis. Current chemotherapy has not improved the survival rate of unresectable CCA patients. Clinical and preclinical studies have shown that activa-

tion of the Wnt/ $\beta$ -catenin signaling occurs throughout the initiation and progression of CCA. Wnt ligands (WNT2, WNT7b, and WNT10A) and TCF4 are upregulated in CCA, accompanied by nuclear translocation of  $\beta$ -catenin.<sup>105,106</sup> The progression of epithelial-mesenchymal transition (EMT) was observed in CCA, represented by the disrupted epithelial cell-cell junctions and mesenchymal characteristics.<sup>107–109</sup> Wnt/ $\beta$ -catenin signaling is one of the critical pathways promoting the EMT transition.<sup>110,111</sup> In CCA cells, suppression of Wnt/ $\beta$ -catenin signaling increased E-cadherin and downregulated vimentin,<sup>112,113</sup> suggesting that the Wnt/ $\beta$ -catenin signaling is associated with EMT during CCA tumorigenesis.  $\beta$ -catenin interacts with E-cadherin to form the cadherin-catenin-actin complex, maintaining epithelial cell adhesion, cytoskeleton, and integrity. During CCA development, the decreased E-cadherin releases  $\beta$ -catenin, resulting in  $\beta$ -catenin accumulation and nuclear translocation.<sup>111</sup> Then,  $\beta$ -catenin activates the transcription of *twist*, *snails*, and *ZEB1* to induce the EMT process in CCA cells.<sup>111</sup>

### HB

HB is a rare malignant tumor found in infants and children.<sup>114</sup> The preclinical and clinical studies showed the hyperactivation of Wnt/ $\beta$ -catenin signaling in HB. In HB cases,  $\beta$ -catenin was found to be increased in the cytoplasm and nucleus of the tumor cells.<sup>115,116</sup> While *CTNNB1* mutations are limited in the exon 3 in embryonal HB, the *CTNNB1* mutations in fetal HB encompass exon 3 and 4.<sup>117</sup> Meanwhile, missense, deletion, or insertion mutations in the *AXIN1* gene were detected in 8% of HB cases.<sup>118</sup>

## MANIPULATING Wnt SIGNALING

### Porcupine (PORCN)

PORCN is a membranous protein mainly localized in the endoplasmic reticulum. PORCN mediates the palmitoylation of Wnt ligands, an essential process for Wnt ligands secretion and ligand-frizzled receptor binding.<sup>119–121</sup> Genetic and pharmacological blockade of PORCN reduces palmitoylation and inhibits the secretion of Wnt ligands, suppressing Wnt signaling.<sup>122</sup> The clinical trials showed promising results of PORCN inhibitors in HCC treatment. ETC159, CGX1321, and RXC004



have entered phase I clinical trials, and IWP12 is still in the preclinical studies (Fig. 4).<sup>123</sup> In mouse models, *Porcn* KO induces embryonic lethality.<sup>124,125</sup> *Porcn* inhibition could cause adverse effects on bone homeostasis.<sup>126</sup>


## Wnt ligands

In physiological conditions, Wnt signaling is activated by binding of secreted Wnt ligands to LRP5/6 coreceptors and FZD receptors.<sup>127</sup> Thus, targeting Wnt ligands by chemicals or neutralizing antibodies efficiently inhibits Wnt signaling.

Based on the high expression of WNT1 in human HCC cell lines and tissues. Anti-WNT1 neutralizing antibody showed its growth inhibitory effect on HCC cell lines but not on normal hepatocytes, with reduced  $\beta$ -catenin's transcriptional activity (Fig. 4).<sup>128</sup>

## Wnt antagonists

SFRPs, WIFs, and DKKs are the secreted Wnt signaling antagonists.<sup>129,130</sup> SFRP-1 and Wnt inhibitory factor 1 (WIF1) inhibit Wnt signaling by directly binding to Wnt ligands.<sup>131</sup> The



**Figure 4.** Manipulating Wnt signaling. Illustration of components and processes of Wnt signal transduction as druggable targets for liver cancer treatment. See the text for detail. Created with BioRender.com. GPC3, glypican-3; LRP, lipoprotein receptor-related protein; FZD, frizzled; DKK1, Dickkopf 1; ROR, receptor tyrosine kinase-like orphan receptor; RYK, receptor tyrosine kinase; PORCN, porcupine; ER, endoplasmic reticulum; GSK-3 $\beta$ , glycogen synthase kinase 3 $\beta$ ; CK1 $\alpha$ , casein kinase 1 $\alpha$ ; APC, adenomatous polyposis coli; TCF, T-cell factor; CBP, CREB binding protein; LEF, lymphoid enhancer-binding factor; peg-IFN, pegylated-Interferon- $\alpha$ 2a; RanBP3, Ran-binding protein 3.

fusion proteins WIF1-Fc and SFRP1-Fc were constructed by adding the Fc fragment of human immunoglobulin (Ig) G1 to WIF1 and SFRP1, respectively (Fig. 4).<sup>132</sup> The fusion proteins exert potent anti-tumor activity by downregulating E2F transcription factor 1 (E2F1), cyclin D1, and c-Myc, increasing apoptosis of HCC cells and impairing tumor vascularization. DKK1 was initially considered a  $\beta$ -catenin-dependent tumor suppressor.<sup>130,133</sup> Several studies have shown that DKK1 promotes tumor cell proliferation, which may be due to DKK1-induced endocytosis of LRP and subsequent activation of the Wnt/PCP signaling pathway.<sup>134,135</sup> DKN-01 is a humanized monoclonal antibody targeting DKK1 in phase I/II clinical trial for HCC (Fig. 4). Phase I investigated the safety of DKN-01 as a single agent and in combination with sorafenib to treat HCC. Phase II explores the anti-tumor activity and safety of DKN-01 in patients with advanced HCC.

### FZD receptors

The FZD receptors are promising therapeutic targets for HCC. The anti-FZD antibody can effectively reduce the HCC tumor growth by blocking the activation of FZD receptors on the Wnt signaling.<sup>136</sup> FZD decoy receptor OMP-54F28 (ipafricapt) is a recombinant fusion protein that binds to a human IgG1 Fc fragment of FZD8,<sup>137,138</sup> which acts synergistically with chemotherapeutic agents (Fig. 4).<sup>139</sup> A phase Ib dose-escalation clinical trial evaluated the safety, tolerability, and pharmacokinetics of OMP-54F28 when combined with sorafenib. Secreted FZD7 (sFZD7) is the extracellular domain of FZD7, expressed and purified from *Escherichia coli*. sFZD7 binding to WNT3 decreased the transcriptional activity of  $\beta$ -catenin/TCF4 and inhibited the growth of HepG2, Hep40, and Huh7.<sup>140</sup> In combination with doxorubicin, sFZD7 inhibited the expression of c-Myc/MYC, Cyclin D1/CCND1, and Survivin/BIRC5, reduced the phosphorylation levels of AKT and ERK1/2, inhibited the growth of Huh7 xenograft tumors, and acted as a chemosensitizer.<sup>140</sup> OMP-54F28 is entering phase I clinical trials, while sFZD7 remains in preclinical studies (Fig. 4).

FZD antibody OMP-18R5 (vantictumab) is a monoclonal antibody directly binding to FZD receptors, which blocks the binding of Wnt ligands to FZD 1, 2, 5, 7, and 8,<sup>141</sup> which inhibits  $\beta$ -catenin-mediated transactivation (Fig. 4). In patient-derived xenograft models, OMP-18R5 combined with chemotherapeutic agents synergistically inhibited the development of several cancers.<sup>141,142</sup> However, like PORCN inhibitors, OMP-

18R5 has the same risk of impairing bone homeostasis.<sup>143</sup> In a dose-escalation clinical trial of OMP-18R5, one patient developed bone degeneration, controllable with zoledronic acid. The skeletal toxicity appeared to be manageable and reversible.<sup>144</sup>

### LRP co-receptors

Salinomycin (SAL), isolated from *Streptomyces albus*, is a monocarboxylic polyether ionophore antibiotic.<sup>145,146</sup> SAL blocks Wnt-induced LRP phosphorylation and leads to LRP protein degradation, destabilizing the Wnt/FZD/LRP complex and inhibiting the Wnt/ $\beta$ -catenin signaling (Fig. 4).<sup>147</sup> SAL effectively inhibits  $\beta$ -catenin expression in HepG2/C3a cell line.<sup>148</sup> SAL also inhibits the migration and invasiveness of liver cancer stem cells through the Wnt/ $\beta$ -catenin signaling suppression.<sup>149</sup>

### Tankyrase (TNKS)

TNKS mediates PARsylation and subsequent degradation of AXIN via the ubiquitin-proteasome pathway, which in turn disrupts the  $\beta$ -catenin destruction complex.<sup>150</sup> Subsequently, the released  $\beta$ -catenin enters the nucleus to transactivate Wnt target genes.<sup>151,152</sup> TNKS is overexpressed in many cancers, including HCC, gastric cancer, and colorectal cancer.<sup>153-155</sup> The TNKS inhibitors XAV939, WXL-8, and NVP-TNKS656, attenuated Wnt/ $\beta$ -catenin signaling and inhibited the growth of HCC cells (Fig. 4).<sup>155-157</sup> Moreover, TNKS inhibitors also suppressed HCC metastasis and invasion.<sup>157</sup> However, there are no relevant clinical trials for TNKS inhibitors in HCC.

### Nuclear export of $\beta$ -catenin

As shown in Table 1 and Supplementary Table 1, pegylated-Interferon- $\alpha$ 2a (peg-IFN), the first-line therapy for the HCV-infected,<sup>158</sup> attenuates the recurrence of HCC (Fig. 4).<sup>159</sup> Mechanistically, peg-IFN upregulates the expression of Ran-binding protein 3 (RanBP3),<sup>160</sup> which enhances the nuclear export of  $\beta$ -catenin.<sup>160</sup> Thus, it is likely that peg-IFN-induced  $\beta$ -catenin nuclear export is a mechanism delaying HCC and improving survival in HCV patients.

## β-catenin-mediated gene transactivation

The small molecule ICG-001 inhibits the interaction between β-catenin and CREB binding protein (CREBBP/CBP) for suppression of β-catenin-mediated gene transactivation (Fig. 4).<sup>161</sup> A phase Ib/IIa clinical trial of the ICG-001 derivative, PRI-724, targeting HCC has been terminated.<sup>162</sup> Similar to ICG-001, PMED-1 disrupts β-catenin-CREBBP interaction and suppresses β-catenin target gene activation.<sup>163</sup> PMED-1 inhibits

HCC cell proliferation but not normal human hepatocytes.<sup>163</sup>

PKF118-310, PKF115-584, and CGP049090 are small-molecule inhibitors targeting the β-catenin-TCF complex (Fig. 4).<sup>164</sup> These antagonists displayed the dose-dependent cytotoxicity in HepG2, Hep40, and Huh7 cell lines, with reduced cytotoxicity (10%) to normal hepatocytes. PKF118-310, PKF115-584, and CGP049090 downregulated β-catenin target genes (*MYC*, *CCND1*, and *Survivin/BIRC5*) and inhibited the growth of HepG2 xenografts.<sup>164,165</sup> Similar to the mechanism of PKF118-

**Table 1.** Targeting Wnt signaling in liver cancers

Agent	Target	Phase	Trial identifier	Type
DKN-01	DKK1	Phase I/II	NCT03645980	Protein
OMP-18R5	FZD1, 2, 5, 7, and 8	Phase I	NCT01345201	Protein
sFZD7	FZD7	Preclinical	NA	Protein
RHPDs	FZD7	Preclinical	NA	Protein
OMP-54F28	FZD8	Phase I	NCT02069145	Protein
Salinomycin	LRP5/6	Preclinical	NA	Natural compounds
CGX1321	PORCN	Phase I	NCT03507998	Small molecule inhibitors
IWP12	PORCN	Preclinical	NA	Small molecule inhibitors
ETC-159	PORCN	Phase I	NCT02521844	Small molecule inhibitors
RXC004	PORCN	Phase I	NCT03447470	Small molecule inhibitors
NVP-TNKS656	Tankyrase	Preclinical	NA	Small molecule inhibitors
XAV939/WXL-8	Tankyrase	Preclinical	NA	Small molecule inhibitors
CGP049090	TCF/β-catenin	Preclinical	NA	Natural compounds
PKF118-310	TCF/β-catenin	Preclinical	NA	Natural compounds
PKF115-584	TCF/β-catenin	Preclinical	NA	Natural compounds
FH535	TCF/β-catenin	Preclinical	NA	Small molecule inhibitors
Peg-IFN	TCF/β-catenin	Phase II	NCT00610389	Protein
WIF1-Fc and sFRP-Fc	Wnt ligands	Preclinical	NA	Protein
Anti-Wnt1	Wnt1	Preclinical	NA	Protein
CGK062	β-catenin phosphorylation	Preclinical	NA	Small molecule inhibitors
PMED-1	β-catenin/CBP	Preclinical	NA	Small molecule inhibitors
PRI-724	β-catenin/CBP	Phase I/II	NCT01302405	Small molecule inhibitors
Hydroxychloroquine	v-ATPase	Phase II	NCT03037437	Small molecule inhibitors
Chloroquine	v-ATPase	Preclinical	NA	Small molecule inhibitors
Bafilomycin	v-ATPase	Preclinical	NA	Small molecule inhibitors
Concanamycin	v-ATPase	Preclinical	NA	Small molecule inhibitors
CAR-GPC3 T cell	GPC3	Phase I	NCT02932956	Cells
Anti-GPC3 antibody	GPC3	Phase II	NCT01507168	Protein
CIK with anti-GPC3	GPC3	Phase II	NCT03146637	Cells

DKK1, Dickkopf 1; FZD, frizzed; NA, not available; PORCN, porcupine; TCF, T-cell factor; peg-IFN, pegylated-Interferon-α2a; CBP, CREB binding protein; v-ATPase, vacuolar-type ATPase.

310, PKF115-584, and CGP049090, FH535 inhibits  $\beta$ -catenin-mediated gene transactivation by interrupting the recruitment of nuclear receptor coactivator 2 (NCOA2)/GRIP1 to the  $\beta$ -catenin transcriptional complex.<sup>166</sup> It was shown that FH535 inhibits HCC cell proliferation by reducing cancer cell stemness.<sup>165</sup>

## $\beta$ -catenin phosphorylation

CGK062 promotes PKC $\alpha$ -mediated phosphorylation of  $\beta$ -catenin at Ser33/Ser37, which degrades  $\beta$ -catenin by the proteasome (Fig. 4).<sup>167</sup> Consistently, CGK062 inhibited the expression of  $\beta$ -catenin target genes (*CCND1*, *MYC*, and *AXIN2*) and suppressed the growth of Wnt/ $\beta$ -catenin-activated HCC cells.<sup>167</sup>

## Cancer-specific targeting of Wnt signaling

Given the pivotal role of Wnt signaling in the homeostasis and regeneration of multiple organs,<sup>168-170</sup> broad-spectrum Wnt signaling inhibitors cause detrimental effects on the normal cells and organs. Therefore, cancer-specific Wnt signaling regulators may be attractive for Wnt signaling blockade therapy. TMEM9, an amplifier of Wnt/ $\beta$ -catenin signaling, promotes lysosomal protein degradation via v-ATPase, resulting in APC downregulation.<sup>46</sup> TMEM9 is highly expressed in liver regeneration and HCC. Genetic ablation of *TMEM9* inhibits HCC tumorigenesis with downregulation of Wnt/ $\beta$ -catenin signaling.<sup>46</sup> Similarly, v-ATPase inhibitors, bafilomycin and concanamycin,<sup>171,172</sup> also inhibit Wnt/ $\beta$ -catenin signaling without toxicity to normal cells and animals (Fig. 4, Table 1).<sup>45,46</sup> Thus, molecular targeting of the TMEM9-v-ATPase axis can be used as cancer-specific Wnt/ $\beta$ -catenin blockade.

Glypican-3 (GPC3) is a proteoglycan binding to the FZD receptor and stimulates Wnt ligands-FZD interaction, resulting in the Wnt signaling activation (Fig. 4).<sup>173</sup> GPC-3 is specifically expressed in HCC but not in normal human liver tissue.<sup>174</sup> The ectopic expression of GPC3 promotes the proliferation of HCC cells.<sup>175</sup> HS20 (an anti-GPC3 monoclonal antibody) suppresses Wnt/ $\beta$ -catenin signaling via inhibiting the interaction of Wnt3a with the GPC3.<sup>176</sup> In xenograft mouse models, HS20 inhibited HCC progression without apparent concomitant toxicity.<sup>176</sup> To date, including CAR-GPC3 T cells or anti-GPC3 antibodies, 33 clinical trials related to GPC3 for HCC treatment were registered (<https://clinicaltrials.gov/>) (Table 1,

Supplementary Table 1).

## Concluding remarks

Wnt signaling activation plays a pivotal role in liver regeneration, metabolic zonation, liver diseases, and liver cancer. Aberrantly hyperactivated Wnt signaling promotes liver tumorigenesis and progression, often in conjunction with liver diseases. Although direct targeting of Wnt signaling sounds attractive as cancer therapy, given the crucial roles of Wnt signaling in tissue homeostasis and regeneration, severe adverse effects from Wnt blockade are inevitable. Nonetheless, an in-depth understanding of the biology of Wnt signaling in liver cancer and exploring cancer-specific Wnt signaling regulators are expected to identify molecular targets specific to liver cancer, which may overcome the current limitations of Wnt signaling inhibitors, and further improve therapeutic strategies of liver cancer treatment.

## Authors' contribution

G.Z. and J.I.P. wrote the manuscript.

## Acknowledgements

We apologize for not including all relevant studies in the field due to limited space. This work was supported by the Cancer Prevention and Research Institute of Texas (RP200315 to J.I.P.) and the National Cancer Institute (CA193297 and CA256207 to J.I.P.).

## Conflicts of Interest

The authors have no conflicts to disclose.

## SUPPLEMENTARY MATERIAL

Supplementary material is available at Clinical and Molecular Hepatology website (<http://www.e-cmh.org>).

## REFERENCES

1. Clemens MM, McGill MR, Apte U. Mechanisms and biomarkers of liver regeneration after drug-induced liver injury. *Adv Pharmacol* 2019;85:241-262.

2. Michalopoulos GK. Hepatostat: liver regeneration and normal liver tissue maintenance. *Hepatology* 2017;65:1384-1392.
3. Ko S, Russell JO, Molina LM, Monga SP. Liver progenitors and adult cell plasticity in hepatic injury and repair: knowns and unknowns. *Annu Rev Pathol* 2020;15:23-50.
4. Duncan AW, Taylor MH, Hickey RD, Hanlon Newell AE, Lenzi ML, Olson SB, et al. The ploidy conveyor of mature hepatocytes as a source of genetic variation. *Nature* 2010;467:707-710.
5. Anti M, Marra G, Rapaccini GL, Rumi C, Bussa S, Fadda G, et al. DNA ploidy pattern in human chronic liver diseases and hepatic nodular lesions. Flow cytometric analysis on echo-guided needle liver biopsy. *Cancer* 1994;73:281-288.
6. Duncan AW, Hanlon Newell AE, Smith L, Wilson EM, Olson SB, Thayer MJ, et al. Frequent aneuploidy among normal human hepatocytes. *Gastroenterology* 2012;142:25-28.
7. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. *CA Cancer J Clin* 2022;72:7-33.
8. Llovet JM, Zucman-Rossi J, Pikarsky E, Sangro B, Schwartz M, Sherman M, et al. Hepatocellular carcinoma. *Nat Rev Dis Primers* 2016;2:16018.
9. Tortelote GG, Reis RR, de Almeida Mendes F, Abreu JG. Complexity of the Wnt/ $\beta$ -catenin pathway: searching for an activation model. *Cell Signal* 2017;40:30-43.
10. Xia XH, Xiao CJ, Shan H. Facilitation of liver cancer SMCC7721 cell aging by siRNA 4 via inhibiting JAK2/STAT3 signal pathway. *Eur Rev Med Pharmacol Sci* 2017;21:1248-1253.
11. Chen J, Gingold JA, Su X. Immunomodulatory TGF- $\beta$  signaling in hepatocellular carcinoma. *Trends Mol Med* 2019;25:1010-1023.
12. Czauderna C, Castven D, Mahn FL, Marquardt JU. Context-dependent role of NF- $\kappa$ B signaling in primary liver cancer—from tumor development to therapeutic implications. *Cancers (Basel)* 2019;11:1053.
13. Russell JO, Monga SP. Wnt/ $\beta$ -catenin signaling in liver development, homeostasis, and pathobiology. *Annu Rev Pathol* 2018;13:351-378.
14. Cheng JH, She H, Han YP, Wang J, Xiong S, Asahina K, et al. Wnt antagonism inhibits hepatic stellate cell activation and liver fibrosis. *Am J Physiol Gastrointest Liver Physiol* 2008;294:G39-G49.
15. Zeng G, Awan F, Otruba W, Muller P, Apte U, Tan X, et al. Wnt'er in liver: expression of Wnt and frizzled genes in mouse. *Hepatology* 2007;45:195-204.
16. Ma R, Martínez-Ramírez AS, Borders TL, Gao F, Sosa-Pineda B. Metabolic and non-metabolic liver zonation is established non-synchronously and requires sinusoidal Wnts. *Elife* 2020;9:e46206.
17. Monga SP.  $\beta$ -catenin signaling and roles in liver homeostasis, injury, and tumorigenesis. *Gastroenterology* 2015;148:1294-1310.
18. Nusse R, Clevers H. Wnt/ $\beta$ -catenin signaling, disease, and emerging therapeutic modalities. *Cell* 2017;169:985-999.
19. Clevers H, Nusse R. Wnt/ $\beta$ -catenin signaling and disease. *Cell* 2012;149:1192-1205.
20. Scheibner K, Bakhti M, Bastidas-Ponce A, Lickert H. Wnt signaling: implications in endoderm development and pancreas organogenesis. *Curr Opin Cell Biol* 2019;61:48-55.
21. Steinhart Z, Angers S. Wnt signaling in development and tissue homeostasis. *Development* 2018;145:dev146589.
22. Hu S, Monga SP. Wnt/ $\beta$ -catenin signaling and liver regeneration: circuit, biology, and opportunities. *Gene Expr* 2021;20:189-199.
23. Zhan T, Rindtorff N, Boutros M. Wnt signaling in cancer. *Oncogene* 2017;36:1461-1473.
24. Raslan AA, Yoon JK. WNT signaling in lung repair and regeneration. *Mol Cells* 2020;43:774-783.
25. Girardi F, Le Grand F. Wnt signaling in skeletal muscle development and regeneration. *Prog Mol Biol Transl Sci* 2018;153:157-179.
26. Jackstadt R, Hodder MC, Sansom OJ. WNT and  $\beta$ -catenin in cancer: genes and therapy. *Annu Rev Cancer Biol* 2020;4:177-196.
27. Monga SP, Pediaditakis P, Mule K, Stolz DB, Michalopoulos GK. Changes in WNT/ $\beta$ -catenin pathway during regulated growth in rat liver regeneration. *Hepatology* 2001;33:1098-1109.
28. Stolz DB, Mars WM, Petersen BE, Kim TH, Michalopoulos GK. Growth factor signal transduction immediately after two-thirds partial hepatectomy in the rat. *Cancer Res* 1999;59:3954-3960.
29. Köhler C, Bell AW, Bowen WC, Monga SP, Fleig W, Michalopoulos GK. Expression of notch-1 and its ligand Jagged-1 in rat liver during liver regeneration. *Hepatology* 2004;39:1056-1065.
30. Higgins GM, Anderson RM. Experimental pathology of the liver. 1. Restoration of the liver of the white rat following partial surgical removal. *Arch Pathol* 1931;12:186-202.
31. Nikfarjam M, Malcontenti-Wilson C, Fanartzis M, Daruwalla J, Christophi C. A model of partial hepatectomy in mice. *J*

- Invest Surg 2004;17:291-294.
32. Sadler KC, Krahn KN, Gaur NA, Ukomadu C. Liver growth in the embryo and during liver regeneration in zebrafish requires the cell cycle regulator, *uhrf1*. *Proc Natl Acad Sci U S A* 2007;104:1570-1575.
  33. Curado S, Stainier DY, Anderson RM. Nitroreductase-mediated cell/tissue ablation in zebrafish: a spatially and temporally controlled ablation method with applications in developmental and regeneration studies. *Nat Protoc* 2008;3:948-954.
  34. Teschke R, Vierke W, Goldermann L. Carbon tetrachloride (CCl<sub>4</sub>) levels and serum activities of liver enzymes following acute CCl<sub>4</sub> intoxication. *Toxicol Lett* 1983;17:175-180.
  35. Preisegger KH, Factor VM, Fuchsbichler A, Stumptner C, Denk H, Thorgeirsson SS. Atypical ductular proliferation and its inhibition by transforming growth factor beta1 in the 3,5-diethoxycarbonyl-1,4-dihydrocollidine mouse model for chronic alcoholic liver disease. *Lab Invest* 1999;79:103-109.
  36. Oderberg IM, Goessling W. Partial hepatectomy in adult zebrafish. *J Vis Exp* 2021;(170):10.3791/62349.
  37. North TE, Babu IR, Vedder LM, Lord AM, Wishnok JS, Tannenbaum SR, et al. PGE<sub>2</sub>-regulated Wnt signaling and N-acetylcysteine are synergistically hepatoprotective in zebrafish acetaminophen injury. *Proc Natl Acad Sci U S A* 2010;107:17315-17320.
  38. Sekine S, Gutiérrez PJ, Lan BY, Feng S, Hebrok M. Liver-specific loss of beta-catenin results in delayed hepatocyte proliferation after partial hepatectomy. *Hepatology* 2007;45:361-368.
  39. Goessling W, North TE, Lord AM, Ceol C, Lee S, Weidinger G, et al. APC mutant zebrafish uncover a changing temporal requirement for wnt signaling in liver development. *Dev Biol* 2008;320:161-174.
  40. Sun T, Pikiolek M, Orsini V, Bergling S, Holwerda S, Morelli L, et al. AXIN2<sup>+</sup> pericentral hepatocytes have limited contributions to liver homeostasis and regeneration. *Cell Stem Cell* 2020;26:97-107.e6.
  41. Huch M, Dorrell C, Boj SF, van Es JH, Li VS, van de Wetering M, et al. In vitro expansion of single Lgr5<sup>+</sup> liver stem cells induced by Wnt-driven regeneration. *Nature* 2013;494:247-250.
  42. Carmon KS, Lin Q, Gong X, Thomas A, Liu Q. LGR5 interacts and cointernalizes with Wnt receptors to modulate Wnt/ $\beta$ -catenin signaling. *Mol Cell Biol* 2012;32:2054-2064.
  43. Jho EH, Zhang T, Domon C, Joo CK, Freund JN, Costantini F. Wnt/ $\beta$ -catenin/Tcf signaling induces the transcription of *Axin2*, a negative regulator of the signaling pathway. *Mol Cell Biol* 2002;22:1172-1183.
  44. Lustig B, Jerchow B, Sachs M, Weiler S, Pietsch T, Karsten U, et al. Negative feedback loop of Wnt signaling through upregulation of *conductin/axin2* in colorectal and liver tumors. *Mol Cell Biol* 2002;22:1184-1193.
  45. Jung YS, Jun S, Kim MJ, Lee SH, Suh HN, Lien EM, et al. TMEM9 promotes intestinal tumorigenesis through vacuolar-ATPase-activated Wnt/ $\beta$ -catenin signalling. *Nat Cell Biol* 2018;20:1421-1433.
  46. Jung YS, Stratton SA, Lee SH, Kim MJ, Jun S, Zhang J, et al. TMEM9-v-ATPase activates Wnt/ $\beta$ -catenin signaling via APC lysosomal degradation for liver regeneration and tumorigenesis. *Hepatology* 2021;73:776-794.
  47. Kudo Y, Sugimoto M, Arias E, Kasashima H, Cordes T, Linares JF, et al. PKC $\lambda$ /i loss induces autophagy, oxidative phosphorylation, and NRF2 to promote liver cancer progression. *Cancer Cell* 2020;38:247-262.e11.
  48. Wang Z, Li Z, Ye Y, Xie L, Li W. Oxidative stress and liver cancer: etiology and therapeutic targets. *Oxid Med Cell Longev* 2016;2016:7891574.
  49. Funato Y, Michiue T, Asashima M, Miki H. The thioredoxin-related redox-regulating protein nucleoredoxin inhibits Wnt-beta-catenin signalling through dishevelled. *Nat Cell Biol* 2006;8:501-508.
  50. Zhou T, Zhou KK, Lee K, Gao G, Lyons TJ, Kowluru R, et al. The role of lipid peroxidation products and oxidative stress in activation of the canonical wntless-type MMTV integration site (WNT) pathway in a rat model of diabetic retinopathy. *Diabetologia* 2011;54:459-468.
  51. Sun HC, Zhang W, Qin LX, Zhang BH, Ye QH, Wang L, et al. Positive serum hepatitis B e antigen is associated with higher risk of early recurrence and poorer survival in patients after curative resection of hepatitis B-related hepatocellular carcinoma. *J Hepatol* 2007;47:684-690.
  52. Rein DB, Wittenborn JS, Weinbaum CM, Sabin M, Smith BD, Lesesne SB. Forecasting the morbidity and mortality associated with prevalent cases of pre-cirrhotic chronic hepatitis C in the United States. *Dig Liver Dis* 2011;43:66-72.
  53. Yang JD, Kim WR, Coelho R, Mettler TA, Benson JT, Sanderson SO, et al. Cirrhosis is present in most patients with hepatitis B and hepatocellular carcinoma. *Clin Gastroenterol Hepatol* 2011;9:64-70.
  54. Roudot-Thoraval F. Epidemiology of hepatitis C virus infec-

- tion. *Clin Res Hepatol Gastroenterol* 2021;45:101596.
55. Mavilia MG, Wu GY. HBV-HCV coinfection: viral interactions, management, and viral reactivation. *J Clin Transl Hepatol* 2018;6:296-305.
  56. Levrero M, Zucman-Rossi J. Mechanisms of HBV-induced hepatocellular carcinoma. *J Hepatol* 2016;64(1 Suppl):S84-S101.
  57. Arbutnot P, Kew M. Hepatitis B virus and hepatocellular carcinoma. *Int J Exp Pathol* 2001;82:77-100.
  58. Neuveut C, Wei Y, Buendia MA. Mechanisms of HBV-related hepatocarcinogenesis. *J Hepatol* 2010;52:594-604.
  59. Tian X, Li J, Ma ZM, Zhao C, Wan DF, Wen YM. Role of hepatitis B surface antigen in the development of hepatocellular carcinoma: regulation of lymphoid enhancer-binding factor 1. *J Exp Clin Cancer Res* 2009;28:58.
  60. Javanmard D, Najafi M, Babaei MR, Karbalaie Niya MH, Esghaei M, Panahi M, et al. Investigation of CTNNB1 gene mutations and expression in hepatocellular carcinoma and cirrhosis in association with hepatitis B virus infection. *Infect Agent Cancer* 2020;15:37.
  61. Zucman-Rossi J, Benhamouche S, Godard C, Boyault S, Grimmer G, Balabaud C, et al. Differential effects of inactivated Axin1 and activated beta-catenin mutations in human hepatocellular carcinomas. *Oncogene* 2007;26:774-780.
  62. Galy O, Chemin I, Le Roux E, Villar S, Le Calvez-Kelm F, Lereau M, et al. Mutations in TP53 and CTNNB1 in relation to hepatitis B and C infections in hepatocellular carcinomas from Thailand. *Hepat Res Treat* 2011;2011:697162.
  63. Hsieh A, Kim HS, Lim SO, Yu DY, Jung G. Hepatitis B viral X protein interacts with tumor suppressor adenomatous polyposis coli to activate Wnt/ $\beta$ -catenin signaling. *Cancer Lett* 2011;300:162-172.
  64. Feitelson MA, Duan LX. Hepatitis B virus X antigen in the pathogenesis of chronic infections and the development of hepatocellular carcinoma. *Am J Pathol* 1997;150:1141-1157.
  65. Lau CC, Sun T, Ching AK, He M, Li JW, Wong AM, et al. Viral-human chimeric transcript predisposes risk to liver cancer development and progression. *Cancer Cell* 2014;25:335-349.
  66. Moradpour D, Penin F. Hepatitis C virus proteins: from structure to function. *Curr Top Microbiol Immunol* 2013;369:113-142.
  67. Fukutomi T, Zhou Y, Kawai S, Eguchi H, Wands JR, Li J. Hepatitis C virus core protein stimulates hepatocyte growth: correlation with upregulation of Wnt-1 expression. *Hepatology* 2005;41:1096-1105.
  68. Liu J, Ding X, Tang J, Cao Y, Hu P, Zhou F, et al. Enhancement of canonical Wnt/ $\beta$ -catenin signaling activity by HCV core protein promotes cell growth of hepatocellular carcinoma cells. *PLoS One* 2011;6:e27496.
  69. Umer M, Qureshi SA, Hashmi ZY, Raza A, Ahmad J, Rahman M, et al. Promoter hypermethylation of Wnt pathway inhibitors in hepatitis C virus - induced multistep hepatocarcinogenesis. *Virology* 2014;11:117.
  70. Quan H, Zhou F, Nie D, Chen Q, Cai X, Shan X, et al. Hepatitis C virus core protein epigenetically silences SFRP1 and enhances HCC aggressiveness by inducing epithelial-mesenchymal transition. *Oncogene* 2014;33:2826-2835.
  71. Ripoli M, Barbano R, Balsamo T, Piccoli C, Brunetti V, Coco M, et al. Hypermethylated levels of E-cadherin promoter in Huh-7 cells expressing the HCV core protein. *Virus Res* 2011;160:74-81.
  72. Jung YS, Wang W, Jun S, Zhang J, Srivastava M, Kim MJ, et al. Dereglulation of CRAD-controlled cytoskeleton initiates mucinous colorectal cancer via  $\beta$ -catenin. *Nat Cell Biol* 2018;20:1303-1314.
  73. Liu Y, El-Serag HB, Jiao L, Lee J, Moore D, Franco LM, et al. WNT signaling pathway gene polymorphisms and risk of hepatic fibrosis and inflammation in HCV-infected patients. *PLoS One* 2013;8:e84407.
  74. Takahara Y, Takahashi M, Zhang QW, Wagatsuma H, Mori M, Tamori A, et al. Serial changes in expression of functionally clustered genes in progression of liver fibrosis in hepatitis C patients. *World J Gastroenterol* 2008;14:2010-2022.
  75. Mercer KE, Hennings L, Ronis MJ. Alcohol consumption, Wnt/ $\beta$ -catenin signaling, and hepatocarcinogenesis. *Adv Exp Med Biol* 2015;815:185-195.
  76. Kim HS, Kim SJ, Bae J, Wang Y, Park SY, Min YS, et al. The p90rsk-mediated signaling of ethanol-induced cell proliferation in HepG2 cell line. *Korean J Physiol Pharmacol* 2016;20:595-603.
  77. Wang F, Yang JL, Yu KK, Xu M, Xu YZ, Chen L, et al. Activation of the NF- $\kappa$ B pathway as a mechanism of alcohol enhanced progression and metastasis of human hepatocellular carcinoma. *Mol Cancer* 2015;14:10.
  78. Yang CF, Zhong YJ, Ma Z, Li L, Shi L, Chen L, et al. NOX4/ROS mediate ethanol-induced apoptosis via MAPK signal pathway in L-02 cells. *Int J Mol Med* 2018;41:2306-2316.
  79. Zhang C, Ellis JL, Yin C. Inhibition of vascular endothelial growth factor signaling facilitates liver repair from acute ethanol-induced injury in zebrafish. *Dis Model Mech* 2016;9:1383-1396.

80. Mandrekar P, Ambade A, Lim A, Szabo G, Catalano D. An essential role for monocyte chemoattractant protein-1 in alcoholic liver injury: regulation of proinflammatory cytokines and hepatic steatosis in mice. *Hepatology* 2011;54:2185-2197.
81. Yuen MF, Tanaka Y, Fong DY, Fung J, Wong DK, Yuen JC, et al. Independent risk factors and predictive score for the development of hepatocellular carcinoma in chronic hepatitis B. *J Hepatol* 2009;50:80-88.
82. Wong VW, Chu WC, Wong GL, Chan RS, Chim AM, Ong A, et al. Prevalence of non-alcoholic fatty liver disease and advanced fibrosis in Hong Kong Chinese: a population study using proton-magnetic resonance spectroscopy and transient elastography. *Gut* 2012;61:409-415.
83. Bhalra N, Angulo P, van der Poorten D, Lee E, Hui JM, Saracco G, et al. The natural history of nonalcoholic fatty liver disease with advanced fibrosis or cirrhosis: an international collaborative study. *Hepatology* 2011;54:1208-1216.
84. Farrell GC, van Rooyen D, Gan L, Chitturi S. NASH is an inflammatory disorder: pathogenic, prognostic and therapeutic implications. *Gut Liver* 2012;6:149-171.
85. Murphy SK, Yang H, Moylan CA, Pang H, Dellinger A, Abdelmalek MF, et al. Relationship between methylome and transcriptome in patients with nonalcoholic fatty liver disease. *Gastroenterology* 2013;145:1076-1087.
86. Chettouh H, Lequoy M, Fartoux L, Vigouroux C, Desbois-Mouthon C. Hyperinsulinaemia and insulin signalling in the pathogenesis and the clinical course of hepatocellular carcinoma. *Liver Int* 2015;35:2203-2217.
87. Jonatan D, Spence JR, Method AM, Kofron M, Sinagoga K, Haataja L, et al. Sox17 regulates insulin secretion in the normal and pathologic mouse  $\beta$  cell. *PLoS One* 2014;9:e104675.
88. Jia Y, Yang Y, Liu S, Herman JG, Lu F, Guo M. SOX17 antagonizes WNT/ $\beta$ -catenin signaling pathway in hepatocellular carcinoma. *Epigenetics* 2010;5:743-749.
89. Ackers I, Malgor R. Interrelationship of canonical and non-canonical Wnt signalling pathways in chronic metabolic diseases. *Diab Vasc Dis Res* 2018;15:3-13.
90. Wang S, Song K, Srivastava R, Dong C, Go GW, Li N, et al. Nonalcoholic fatty liver disease induced by noncanonical Wnt and its rescue by Wnt3a. *FASEB J* 2015;29:3436-3445.
91. Kew MC. Synergistic interaction between aflatoxin B1 and hepatitis B virus in hepatocarcinogenesis. *Liver Int* 2003;23:405-409.
92. Devereux TR, Stern MC, Flake GP, Yu MC, Zhang ZQ, London SJ, et al. CTNNB1 mutations and beta-catenin protein accumulation in human hepatocellular carcinomas associated with high exposure to aflatoxin B1. *Mol Carcinog* 2001;31:68-73.
93. Yang JD, Hainaut P, Gores GJ, Amadou A, Plymoth A, Roberts LR. A global view of hepatocellular carcinoma: trends, risk, prevention and management. *Nat Rev Gastroenterol Hepatol* 2019;16:589-604.
94. Bengochea A, de Souza MM, Lefrançois L, Le Roux E, Galy O, Chemin I, et al. Common dysregulation of Wnt/frizzled receptor elements in human hepatocellular carcinoma. *Br J Cancer* 2008;99:143-150.
95. Boyault S, Rickman DS, de Reyniès A, Balabaud C, Rebouissou S, Jeannot E, et al. Transcriptome classification of HCC is related to gene alterations and to new therapeutic targets. *Hepatology* 2007;45:42-52.
96. Fodde R, Smits R, Clevers H. APC, signal transduction and genetic instability in colorectal cancer. *Nat Rev Cancer* 2001;1:55-67.
97. Khemlina G, Ikeda S, Kurzrock R. The biology of hepatocellular carcinoma: implications for genomic and immune therapies. *Mol Cancer* 2017;16:149.
98. Satoh S, Daigo Y, Furukawa Y, Kato T, Miwa N, Nishiwaki T, et al. AXIN1 mutations in hepatocellular carcinomas, and growth suppression in cancer cells by virus-mediated transfer of AXIN1. *Nat Genet* 2000;24:245-250.
99. Park JY, Park WS, Nam SW, Kim SY, Lee SH, Yoo NJ, et al. Mutations of beta-catenin and AXIN 1 genes are a late event in human hepatocellular carcinogenesis. *Liver Int* 2005;25:70-76.
100. Zhu M, Lu T, Jia Y, Luo X, Gopal P, Li L, et al. Somatic mutations increase hepatic clonal fitness and regeneration in chronic liver disease. *Cell* 2019;177:608-621.e12.
101. Marquardt JU, Seo D, Andersen JB, Gillen MC, Kim MS, Conner EA, et al. Sequential transcriptome analysis of human liver cancer indicates late stage acquisition of malignant traits. *J Hepatol* 2014;60:346-353.
102. Rebouissou S, Nault JC. Advances in molecular classification and precision oncology in hepatocellular carcinoma. *J Hepatol* 2020;72:215-229.
103. Abitbol S, Dahmani R, Coulouarn C, Ragazzon B, Mlecnik B, Senni N, et al. AXIN deficiency in human and mouse hepatocytes induces hepatocellular carcinoma in the absence of  $\beta$ -catenin activation. *J Hepatol* 2018;68:1203-1213.
104. Lazaridis KN, Gores GJ. Cholangiocarcinoma. In: Boyer TD, Manns MP, Sanyal AJ, eds. *Zakim and Boyer's Hepatology*.



- 6th ed. Philadelphia: Saunders, 2012:1032-1044.
105. Boulter L, Guest RV, Kendall TJ, Wilson DH, Wojtacha D, Robson AJ, et al. WNT signaling drives cholangiocarcinoma growth and can be pharmacologically inhibited. *J Clin Invest* 2015;125:1269-1285.
  106. Chen W, Liang J, Huang L, Cai J, Lei Y, Lai J, et al. Characterizing the activation of the Wnt signaling pathway in hilar cholangiocarcinoma using a tissue microarray approach. *Eur J Histochem* 2016;60:2536.
  107. Banales JM, Marin JJG, Lamarca A, Rodrigues PM, Khan SA, Roberts LR, et al. Cholangiocarcinoma 2020: the next horizon in mechanisms and management. *Nat Rev Gastroenterol Hepatol* 2020;17:557-588.
  108. Vaquero J, Guedj N, Clapéron A, Nguyen Ho-Boulidoires TH, Paradis V, Fouassier L. Epithelial-mesenchymal transition in cholangiocarcinoma: from clinical evidence to regulatory networks. *J Hepatol* 2017;66:424-441.
  109. Nitta T, Mitsuhashi T, Hatanaka Y, Miyamoto M, Oba K, Tsuchikawa T, et al. Prognostic significance of epithelial-mesenchymal transition-related markers in extrahepatic cholangiocarcinoma: comprehensive immunohistochemical study using a tissue microarray. *Br J Cancer* 2014;111:1363-1372.
  110. Chaw SY, Abdul Majeed A, Dalley AJ, Chan A, Stein S, Farah CS. Epithelial to mesenchymal transition (EMT) biomarkers--E-cadherin, beta-catenin, APC and vimentin--in oral squamous cell carcinogenesis and transformation. *Oral Oncol* 2012;48:997-1006.
  111. Valenta T, Hausmann G, Basler K. The many faces and functions of  $\beta$ -catenin. *EMBO J* 2012;31:2714-2736.
  112. Zhang GF, Qiu L, Yang SL, Wu JC, Liu TJ. Wnt/ $\beta$ -catenin signaling as an emerging potential key pharmacological target in cholangiocarcinoma. *Biosci Rep* 2020;40:BSR20193353.
  113. Xue W, Dong B, Zhao Y, Wang Y, Yang C, Xie Y, et al. Upregulation of TTYH3 promotes epithelial-to-mesenchymal transition through Wnt/ $\beta$ -catenin signaling and inhibits apoptosis in cholangiocarcinoma. *Cell Oncol (Dordr)* 2021;44:1351-1361.
  114. Carrillo-Reixach J, Torrens L, Simon-Coma M, Royo L, Domingo-Sàbat M, Abril-Fornaguera J, et al. Epigenetic footprint enables molecular risk stratification of hepatoblastoma with clinical implications. *J Hepatol* 2020;73:328-341.
  115. Gupta K, Rane S, Das A, Marwaha RK, Menon P, Rao KL. Relationship of  $\beta$ -catenin and postchemotherapy histopathologic changes with overall survival in patients with hepatoblastoma. *J Pediatr Hematol Oncol* 2012;34:e320-e328.
  116. Crippa S, Ancey PB, Vazquez J, Angelino P, Rougemont AL, Guettier C, et al. Mutant CTNNB1 and histological heterogeneity define metabolic subtypes of hepatoblastoma. *EMBO Mol Med* 2017;9:1589-1604.
  117. López-Terrada D, Gunaratne PH, Adesina AM, Pulliam J, Hoang DM, Nguyen Y, et al. Histologic subtypes of hepatoblastoma are characterized by differential canonical Wnt and notch pathway activation in DLK+ precursors. *Hum Pathol* 2009;40:783-794.
  118. Taniguchi K, Roberts LR, Aderca IN, Dong X, Qian C, Murphy LM, et al. Mutational spectrum of beta-catenin, AXIN1, and AXIN2 in hepatocellular carcinomas and hepatoblastomas. *Oncogene* 2002;21:4863-4871.
  119. Shah K, Panchal S, Patel B. Porcupine inhibitors: novel and emerging anti-cancer therapeutics targeting the Wnt signaling pathway. *Pharmacol Res* 2021;167:105532.
  120. Liu J, Pan S, Hsieh MH, Ng N, Sun F, Wang T, et al. Targeting Wnt-driven cancer through the inhibition of porcupine by LGK974. *Proc Natl Acad Sci U S A* 2013;110:20224-20229.
  121. Madan B, Ke Z, Harmston N, Ho SY, Frois AO, Alam J, et al. Wnt addiction of genetically defined cancers reversed by PORCN inhibition. *Oncogene* 2016;35:2197-2207.
  122. Proffitt KD, Madan B, Ke Z, Pendharkar V, Ding L, Lee MA, et al. Pharmacological inhibition of the Wnt acyltransferase PORCN prevents growth of WNT-driven mammary cancer. *Cancer Res* 2013;73:502-507.
  123. Wang W, Xu L, Liu P, Jairam K, Yin Y, Chen K, et al. Blocking Wnt secretion reduces growth of hepatocellular carcinoma cell lines mostly independent of  $\beta$ -catenin signaling. *Neoplasia* 2016;18:711-723.
  124. Barrott JJ, Cash GM, Smith AP, Barrow JR, Murtaugh LC. Deletion of mouse porcn blocks Wnt ligand secretion and reveals an ectodermal etiology of human focal dermal hypoplasia/Goltz syndrome. *Proc Natl Acad Sci U S A* 2011;108:12752-12757.
  125. Liu W, Shaver TM, Balasa A, Ljungberg MC, Wang X, Wen S, et al. Deletion of porcn in mice leads to multiple developmental defects and models human focal dermal hypoplasia (Goltz syndrome). *PLoS One* 2012;7:e32331.
  126. Funck-Brentano T, Nilsson KH, Brommage R, Henning P, Lerner UH, Koskela A, et al. Porcupine inhibitors impair trabecular and cortical bone mass and strength in mice. *J Endocrinol* 2018;238:13-23.
  127. He X, Semenov M, Tamai K, Zeng X. LDL receptor-related proteins 5 and 6 in Wnt/beta-catenin signaling: arrows point

- the way. *Development* 2004;131:1663-1677.
128. Wei W, Chua MS, Grepper S, So SK. Blockade of Wnt-1 signaling leads to anti-tumor effects in hepatocellular carcinoma cells. *Mol Cancer* 2009;8:76.
  129. Cruciat CM, Niehrs C. Secreted and transmembrane wnt inhibitors and activators. *Cold Spring Harb Perspect Biol* 2013;5:a015081.
  130. Niehrs C. Function and biological roles of the Dickkopf family of Wnt modulators. *Oncogene* 2006;25:7469-7481.
  131. Kawazoe M, Kaneko K, Nanki T. Glucocorticoid therapy suppresses Wnt signaling by reducing the ratio of serum Wnt3a to Wnt inhibitors, sFRP-1 and Wif-1. *Clin Rheumatol* 2021;40:2947-2954.
  132. Hu J, Dong A, Fernandez-Ruiz V, Shan J, Kawa M, Martínez-Ansó E, et al. Blockade of Wnt signaling inhibits angiogenesis and tumor growth in hepatocellular carcinoma. *Cancer Res* 2009;69:6951-6959.
  133. Glinka A, Wu W, Delius H, Monaghan AP, Blumenstock C, Niehrs C. Dickkopf-1 is a member of a new family of secreted proteins and functions in head induction. *Nature* 1998;391:357-362.
  134. Caneparo L, Huang YL, Staudt N, Tada M, Ahrendt R, Kazanskaya O, et al. Dickkopf-1 regulates gastrulation movements by coordinated modulation of Wnt/beta catenin and Wnt/PCP activities, through interaction with the Dally-like homolog Knypek. *Genes Dev* 2007;21:465-480.
  135. Cha SW, Tadjuidje E, Tao Q, Wylie C, Heasman J. Wnt5a and Wnt11 interact in a maternal Dkk1-regulated fashion to activate both canonical and non-canonical signaling in *Xenopus* axis formation. *Development* 2008;135:3719-3729.
  136. Nambotin SB, Lefrancois L, Sainsily X, Berthillon P, Kim M, Wands JR, et al. Pharmacological inhibition of frizzled-7 displays anti-tumor properties in hepatocellular carcinoma. *J Hepatol* 2011;54:288-299.
  137. Jimeno A, Gordon M, Chugh R, Messersmith W, Mendelson D, Dupont J, et al. A first-in-human phase I study of the anticancer stem cell agent ipafricept (OMP-54F28), a decoy receptor for wnt ligands, in patients with advanced solid tumors. *Clin Cancer Res* 2017;23:7490-7497.
  138. Moore KN, Gunderson CC, Sabbatini P, McMeekin DS, Mantia-Smaldone G, Burger RA, et al. A phase 1b dose escalation study of ipafricept (OMP54F28) in combination with paclitaxel and carboplatin in patients with recurrent platinum-sensitive ovarian cancer. *Gynecol Oncol* 2019;154:294-301.
  139. Hoey T. Development of FZD8-Fc (OMP-54F28), a Wnt signaling antagonist that inhibits tumor growth and reduces tumor initiating cell frequency. AACR Annual Meeting; 2013; Washington DC, USA.
  140. Wei W, Chua MS, Grepper S, So SK. Soluble frizzled-7 receptor inhibits Wnt signaling and sensitizes hepatocellular carcinoma cells towards doxorubicin. *Mol Cancer* 2011;10:16.
  141. Fischer MM, Cancilla B, Yeung VP, Cattaruzza F, Chartier C, Murriel CL, et al. WNT antagonists exhibit unique combinatorial antitumor activity with taxanes by potentiating mitotic cell death. *Sci Adv* 2017;3:e1700090.
  142. Gurney A, Axelrod F, Bond CJ, Cain J, Chartier C, Donigan L, et al. Wnt pathway inhibition via the targeting of frizzled receptors results in decreased growth and tumorigenicity of human tumors. *Proc Natl Acad Sci U S A* 2012;109:11717-11722.
  143. Davis SL, Cardin DB, Shahda S, Lenz HJ, Dotan E, O'Neil BH, et al. A phase 1b dose escalation study of Wnt pathway inhibitor vantictumab in combination with nab-paclitaxel and gemcitabine in patients with previously untreated metastatic pancreatic cancer. *Invest New Drugs* 2020;38:821-830.
  144. Smith DC, Rosen LS, Chugh R, Goldman J, Xu L, Kapoun AM, et al. First-in-human evaluation of the human monoclonal antibody vantictumab (OMP-18R5; antiFrizzled) targeting the WNT pathway in a phase I study for patients with advanced solid tumors. *J Clin Oncol* 2013;31(suppl):2540a.
  145. Gupta PB, Onder TT, Jiang G, Tao K, Kuperwasser C, Weinberg RA, et al. Identification of selective inhibitors of cancer stem cells by high-throughput screening. *Cell* 2009;138:645-659.
  146. Huczynski A. Salinomycin: a new cancer drug candidate. *Chem Biol Drug Des* 2012;79:235-238.
  147. Lu D, Choi MY, Yu J, Castro JE, Kipps TJ, Carson DA. Salinomycin inhibits Wnt signaling and selectively induces apoptosis in chronic lymphocytic leukemia cells. *Proc Natl Acad Sci U S A* 2011;108:13253-13257.
  148. Niwa AM, Semprebon SC, D'Epiro GFR, Marques LA, Zanetti TA, Mantovani MS. Salinomycin induces cell cycle arrest and apoptosis and modulates hepatic cytochrome P450 mRNA expression in HepG2/C3a cells. *Toxicol Mech Methods* 2022;32:341-351.
  149. Liu Q, Sun J, Luo Q, Ju Y, Song G. Salinomycin suppresses tumorigenicity of liver cancer stem cells and Wnt/beta-catenin signaling. *Curr Stem Cell Res Ther* 2021;16:630-637.
  150. Huang SM, Mishina YM, Liu S, Cheung A, Stegmeier F, Michaud GA, et al. Tankyrase inhibition stabilizes axin and antagonizes Wnt signalling. *Nature* 2009;461:614-620.
  151. Clevers H. Wnt/beta-catenin signaling in development and

- disease. *Cell* 2006;127:469-480.
152. Rubinfeld B, Albert I, Porfiri E, Fiol C, Munemitsu S, Polakis P. Binding of GSK3beta to the APC-beta-catenin complex and regulation of complex assembly. *Science* 1996;272:1023-1026.
  153. Ma Y, Zhang P, Zhang Q, Wang X, Miao Q, Lyu X, et al. Dihydroartemisinin suppresses proliferation, migration, the Wnt/ $\beta$ -catenin pathway and EMT via TNKS in gastric cancer. *Oncol Lett* 2021;22:688.
  154. Ma Z, Han C, Xia W, Wang S, Li X, Fang P, et al. circ5615 functions as a ceRNA to promote colorectal cancer progression by upregulating TNKS. *Cell Death Dis* 2020;11:356.
  155. Ma L, Wang X, Jia T, Wei W, Chua MS, So S. Tankyrase inhibitors attenuate WNT/ $\beta$ -catenin signaling and inhibit growth of hepatocellular carcinoma cells. *Oncotarget* 2015;6:25390-25401.
  156. Wu X, Luo F, Li J, Zhong X, Liu K. Tankyrase 1 inhibitor XAV939 increases chemosensitivity in colon cancer cell lines via inhibition of the Wnt signaling pathway. *Int J Oncol* 2016;48:1333-1340.
  157. Huang J, Qu Q, Guo Y, Xiang Y, Feng D. Tankyrases/ $\beta$ -catenin signaling pathway as an anti-proliferation and anti-metastatic target in hepatocarcinoma cell lines. *J Cancer* 2020;11:432-440.
  158. Ghany MG, Strader DB, Thomas DL, Seeff LB. Diagnosis, management, and treatment of hepatitis C: an update. *Hepatology* 2009;49:1335-1374.
  159. Lee D, Chung YH, Kim JA, Park WH, Jin YJ, Shim JH, et al. Safety and efficacy of adjuvant pegylated interferon therapy for metastatic tumor antigen 1-positive hepatocellular carcinoma. *Cancer* 2013;119:2239-2246.
  160. Thompson MD, Dar MJ, Monga SP. Pegylated interferon alpha targets Wnt signaling by inducing nuclear export of  $\beta$ -catenin. *J Hepatol* 2011;54:506-512.
  161. Deng J, Hua L, Han T, Tian M, Wang D, Tang H, et al. The CREB-binding protein inhibitor ICG-001: a promising therapeutic strategy in sporadic meningioma with NF2 mutations. *Neurooncol Adv* 2020;2:vdz055.
  162. Takahashi-Yanaga F, Kahn M. Targeting Wnt signaling: can we safely eradicate cancer stem cells? *Clin Cancer Res* 2010;16:3153-3162.
  163. Delgado ER, Yang J, So J, Leimgruber S, Kahn M, Ishitani T, et al. Identification and characterization of a novel small-molecule inhibitor of  $\beta$ -catenin signaling. *Am J Pathol* 2014;184:2111-2122.
  164. Wei W, Chua MS, Grepper S, So S. Small molecule antagonists of Tcf4/ $\beta$ -catenin complex inhibit the growth of HCC cells in vitro and in vivo. *Int J Cancer* 2010;126:2426-2436.
  165. Gedaly R, Galuppo R, Daily MF, Shah M, Maynard E, Chen C, et al. Targeting the Wnt/ $\beta$ -catenin signaling pathway in liver cancer stem cells and hepatocellular carcinoma cell lines with FH535. *PLoS One* 2014;9:e99272.
  166. Handeli S, Simon JA. A small-molecule inhibitor of Tcf/ $\beta$ -catenin signaling down-regulates PPARgamma and PPARdelta activities. *Mol Cancer Ther* 2008;7:521-529.
  167. Gwak J, Lee JH, Chung YH, Song GY, Oh S. Small molecule-based promotion of PKC $\alpha$ -mediated  $\beta$ -catenin degradation suppresses the proliferation of CRT-positive cancer cells. *PLoS One* 2012;7:e46697.
  168. Chen X, Yang J, Evans PM, Liu C. Wnt signaling: the good and the bad. *Acta Biochim Biophys Sin (Shanghai)* 2008;40:577-594.
  169. Chen D, Xie R, Shu B, Landay AL, Wei C, Reiser J, et al. Wnt signaling in bone, kidney, intestine, and adipose tissue and interorgan interaction in aging. *Ann N Y Acad Sci* 2019;1442:48-60.
  170. Behari J. The Wnt/ $\beta$ -catenin signaling pathway in liver biology and disease. *Expert Rev Gastroenterol Hepatol* 2010;4:745-756.
  171. Bowman EJ, Graham LA, Stevens TH, Bowman BJ. The bafilomycin/concanamycin binding site in subunit c of the v-ATPases from *Neurospora crassa* and *Saccharomyces cerevisiae*. *J Biol Chem* 2004;279:33131-33138.
  172. Huss M, Ingenhorst G, König S, Gassel M, Dröse S, Zeeck A, et al. Concanamycin A, the specific inhibitor of v-ATPases, binds to the v(o) subunit c. *J Biol Chem* 2002;277:40544-40548.
  173. Capurro M, Martin T, Shi W, Filmus J. Glypican-3 binds to frizzled and plays a direct role in the stimulation of canonical Wnt signaling. *J Cell Sci* 2014;127(Pt 7):1565-1575.
  174. Guo M, Zhang H, Zheng J, Liu Y. Glypican-3: a new target for diagnosis and treatment of hepatocellular carcinoma. *J Cancer* 2020;11:2008-2021.
  175. Capurro MI, Xiang YY, Lobe C, Filmus J. Glypican-3 promotes the growth of hepatocellular carcinoma by stimulating canonical Wnt signaling. *Cancer Res* 2005;65:6245-6254.
  176. Gao W, Kim H, Feng M, Phung Y, Xavier CP, Rubin JS, et al. Inactivation of Wnt signaling by a human antibody that recognizes the heparan sulfate chains of glypican-3 for liver cancer therapy. *Hepatology* 2014;60:576-587.