



Cardiovascular disease in asthma patients: from mechanisms to therapeutic implications

Authors: Mario Cazzola, Nicola A Hanania, Paola Rogliani, Maria Gabriella Matera

Article type: Review

Received: January 23, 2023

Accepted: February 2, 2023

Early publication date: February 5, 2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Cardiovascular disease in asthma patients: from mechanisms to therapeutic implications

Mario Cazzola¹, Nicola A Hanania², Paola Rogliani¹, Maria Gabriella Matera³

¹Chair of Respiratory Medicine, Department of Experimental Medicine, University of Rome Tor Vergata, Rome, Italy

²Section of Pulmonary and Critical Care Medicine, Baylor College of Medicine, Houston, TX, US

³Chair of Pharmacology, Department of Experimental Medicine, University of Campania Luigi Vanvitelli, Naples, Italy

Correspondence to:

Prof. Mario Cazzola, MD, PhD,

Chair of Respiratory Medicine, Department of Experimental Medicine, University of Rome Tor Vergata, Rome, Italy,

e-mail: mario.cazzola@uniroma2.it

ABSTRACT

Cardiovascular disease (CVD) is often associated with asthma, and asthma patients have an increased risk of CVD mortality. Our understanding of the bidirectional risk of CVD and asthma has been based on several observational studies. However, the specific pathogenetic mechanisms underlying the development of cardiovascular comorbidities in patients with asthma have not yet been fully determined. Such cardiovascular complications in patients with asthma have been attributed to airway and system inflammation present in both asthma and CVD. Indeed, there is evidence that mast cells, eosinophils, inflammatory cytokines, and immunoglobulin E increase in both lungs of patients with asthma, and in injured heart and vessels of CVD patients. These findings suggest that allergic asthma and CVD may share pathogenic pathways. Understanding these pathways is critical to the choice of pharmacological interventions. Currently, the most appropriate therapeutic approach lies in using the best available evidence to optimize the management of both asthma and CVD. Therapy should be optimized to take advantage of the favorable benefits that each medication may have on both organs while minimizing the likelihood of adverse effects on the lungs and heart. It is noteworthy that inhaled β_2 -agonists induce benefit in patients with acute decompensated heart failure. Furthermore, ICSs may reduce the risk of atherosclerosis. On the other side, asthma is

not an absolute contraindication to use cardio-selective β_1 -blockers, but these medications should be prescribed with caution especially if are necessary for acute cardiovascular events and alternative treatment options are unavailable. In addition, when aspirin intake causes the onset of hypersensitivity, P2Y₁₂ inhibitors (e.g., clopidogrel, prasugrel, and ticagrelor) are effective and safe treatment alternatives.

Key words: asthma, cardiovascular disease, comorbidity, mechanisms, therapeutic approaches

ASTHMA HETEROGENEITY

Asthma is a heterogeneous condition characterized by airway hyperresponsiveness, variable airflow obstruction that can become permanent over time, and chronic inflammation [1, 2]. However, not all of these characteristics are usually present, and one or two may prevail when large cohorts of asthma patients are analyzed [3]. Furthermore, asthma must at least be divided into two main endotypes according to their type 2 (T2) inflammation. It is defined as T2_{high}, characterized by upregulation of T2 immune pathways (ie, interleukin (IL)-4 and IL-13 gene sets) and eosinophilic airway inflammation, or T2_{low} (usually characterized by neutrophilic or paucigranulocytic airway inflammation) [4]. Patients with T2_{high} asthma are more allergic and have more eosinophils and airway hyperresponsiveness than patients with low expression [5]. T2_{low} asthma remains poorly understood. It encompasses extremely late-onset asthma, obesity-related asthma, smoking-related asthma, neutrophilic asthma, and asthmatic patients with mild inflammation [6].

Age at the onset of asthma has emerged as a crucial determinant in differentiating between endotypes [7]. Patients with early-onset (or childhood-onset) asthma are often atopic, with T2-predominant inflammation and a favorable prognosis. On the other hand, patients with adult-onset or late-onset asthma are frequently nonatopic, have a poor prognosis, and are more likely to have permanent airway obstruction. The absence of clinical allergy in late-onset asthma indicates that the T2 pathway is distinct from, and likely more complex than, the one associated with the early-onset allergic phenotype [8]. A T2 inflammatory process in the lung is achievable in the absence of mucosal-specific immunoglobulin (Ig)E and clinical allergic responses [8]. Some people have sputum neutrophilia mixed in with their eosinophilic process. This mixed inflammatory process suggests that other immune pathways interact with T2 immunity, including activation of IL-33 and IL-17 pathways [8].

Chest tightness, wheezing, and coughing can limit everyday activities, impair function, and disturb sleep [1, 2]. Asthma management aims not only to establish adequate control of

symptoms and relieve them when they occur but also to reduce the risk of asthma-related exacerbations and mortality, prolonged airflow limitation, and treatment-related adverse effects.

The basis of asthma management includes inhaled corticosteroids (ICS) and long-acting β_2 -agonists (LABAs) [1, 2]. However, long-acting muscarinic receptor antagonists (LAMAs), CysLT receptor antagonists (CysLTRAs), oral corticosteroid (OCS), and biological therapies such as omalizumab that targets IgE, preventing it from binding to receptors on basophils and mast cells, mepolizumab and reslizumab which are monoclonal antibodies (mAbs) that target interleukin-5 (IL-5), benralizumab which is an IL-5 receptor α (IL-5 R)-directed cytolytic mAb, or dupilumab which is an anti-IL-4 receptor α mAb that blocks both IL-4 and IL-13 signaling, are accessible as alternatives. In addition, growing evidence demonstrates the benefits of treating asthma with the triple regimen of ICS, LABA, and LAMA [9, 10].

Asthma guidelines and therapeutic strategies still provide treatment recommendations disregarding the complexity of asthma and advocate a progressive approach, increasing or de-escalating therapy by shifting to a higher or lower step, depending on the frequency and severity of the patient's symptoms [11].

However, it is critical to identify the particular component of asthma that drives symptoms in each patient prior to therapy selection [12]. Eosinophilic airway inflammation usually responds to standard treatment with corticosteroids and biologics, although biological therapies are suggested only in patients with severe asthma [1, 2]. $T2_{low}$ asthma is not responsive to treatment with ICSs, but instead, it is a predictor of response to antibiotics [6, 12]. Macrolides, including clarithromycin and azithromycin, have demonstrated their effectiveness [12]. No interventions have been evaluated for this specific inflammatory endotype, although preliminary evidence suggests therapies targeting the IL-6 pathway may be beneficial. Asthmatics with paucigranulocytic airway inflammation may not warrant anti-inflammatory treatment as symptoms in these patients may be driven solely by smooth muscle dysfunction (airway hyperresponsiveness) [6, 12]. Therefore, these patients may benefit from smooth muscle-directed therapies such as additional bronchodilators and LAMAs, mast-cell directed therapies, or in the most severe cases, bronchial thermoplasty [6, 12].

COMORBIDITIES AND ASTHMA

Adult-onset asthma may share metabolic and inflammatory components with other diseases such as obesity, metabolic syndrome, type 2 diabetes, cardiovascular disease (CVD), and psychiatric disorders [7]. Asthma, mainly in individuals with neutrophilic airway inflammation,

causes systemic inflammation, with an increase in circulating proinflammatory cytokines such as IL -6 and tumor necrosis factor- α (TNF- α), as well as high-sensitivity C-reactive protein (CRP) [13]. Systemic inflammation, a phenomenon common for many nonrespiratory disorders and asthma, constitutes a possible link between these disorders and at least certain endotypes of asthma [14].

Numerous comorbid illnesses, including gastroesophageal reflux, allergic rhinitis, obesity, depression, diabetes mellitus, and CVD, have been associated with asthma [15].

These comorbidities can affect the course of asthma and its severity as well as its treatment modality. Therefore, the identification of such comorbidities, a thorough evaluation of their effects, and appropriate treatment may reduce their impact must become a fundamental part of the approach to a patient with asthma [15].

Since asthma and CVD constitute major public health problems, there is genuine interest in determining the extent of any association between them and in understanding how one disease affects the other. Regrettably, epidemiological research on this relationship has produced contradictory evidence.

ARE ASTHMA AND CVD ASSOCIATED?

Some epidemiological studies found a strong link between asthma and CVD, particularly coronary artery disease (CAD), acute myocardial infarction (AMI), and arterial hypertension [16, 17]. Arterial hypertension shows the strongest association with asthma [18]. In a recent systematic review and meta-analysis that analyzed data from 5,493,776 subjects, the strength of association with asthma was weak for CAD, moderate for heart failure (HF), hypertension, cardiovascular comorbidities, and pulmonary hypertension, and strong for hypertensive cardiomyopathy [19]. In severe asthma, cardiovascular comorbidities, CAD, and HF were moderately associated, while hypertension was strongly associated. Furthermore, arterial hypertension has been associated with established measures of adult asthma severity, as evidenced by the use of more than 6 canisters of short-acting β -agonists (SABAs), history of emergency department visits or hospitalizations, and corticosteroid administration in a US case-control study [20].

The CVD risk has been reported to be 32% [21] to 42% [22] higher in asthmatics than in people without asthma. Analysis of 30 cohort studies comprising 4,157,823 participants showed that asthma and CVD had stronger links in women than in men (39% vs. 19%) and that patients with late-onset were at higher risk than patients with early-onset asthma (39% vs. 26%) [19]. Women have an increase in the prevalence rate of asthma, which is temporally correlated with

variations in estrogen levels [23]. However, overweight women (body mass index 25-30) with adult-onset asthma had a stronger association with CVD than those with adult-onset asthma who were not overweight [24]. Furthermore, asthma is associated with an increased risk of CVD among smokers [25] and among adults whose asthma is work-related [26].

However, several other investigations found only modest or no connections between asthma and CVD. According to the Atherosclerosis Risk in Communities (ARIC) study conducted in the United States, those who have ever had asthma have an adjusted relative risk of coronary heart disease of 0.87 (95% CI 0.66-1.14), while those who currently had asthma had an adjusted relative risk of 0.69 (95% CI 0.46-1.05) [27]. Although CVD and hypertension were more common in individuals with asthma than in the general population, an Italian population-based retrospective cross-sectional analysis revealed that there was only a weak correlation between asthma and these conditions [18]. A Polish epidemiological study revealed no increased prevalence of comorbidity between cardiovascular diseases and asthma, with only ischemic heart disease (IHD) showing a odds ratio (OR) > 1 [28]. Furthermore, asthma significantly reduced the risk of atherosclerosis (OR = 0.33) in the patients with asthma compared to controls. Asthma, particularly the uncontrolled eosinophilic phenotype, is a risk factor for CVD, not only in the classic form with coronary artery occlusion, but also in vasospastic episodes without occlusion [29].

Several population-based cohort studies have concluded that asthma is associated with an increased risk of CAD of up to about 30% [16, 30]. However, it has been reported that only those with asthma with an allergy have an elevated risk of CAD [31]. In any case, women with adult-onset asthma appear to have a higher risk of CAD [32].

In subjects with active asthma, defined as current users of asthma medications, the risk of acute myocardial infarction (AMI) is estimated to be 29% higher than in subjects without asthma, regardless of high-sensitivity (hs)CRP levels, smoking history, physical activity levels, or gender [33]. Inactive asthma, defined as the lack of asthma-related events within one year before AMI, does not increase the risk of AMI, although current asthma, characterized by the occurrence of any asthma-related episodes, including asthma symptoms (cough with wheezing, shortness of breath, and tightness in the chest), the use of asthma medications, unexpected office visits for asthma, emergency room visits for asthma, or hospitalization for asthma within a year before the index date of AMI, does [34]. Asthma exacerbations have been found to correlate with a significantly higher risk of AMI, particularly in the initial 1-week interval after acute exacerbation [35]. Data from the Korean Genome and Epidemiology Study prospective cohort

from 2004 to 2016 showed that asthma was mainly associated with IHD in the older age (≥ 53 years) group and the untreated asthma group [36].

Patients with asthma are at increased risk of all-cause and CVD mortality [23]. Chinese research using data from the National Health and Nutrition Examination Survey from 2001-2002 to 2013-2014 showed that people with current childhood-onset asthma had a slightly higher CVD death rate than those with current adulthood-onset asthma [37]. However, according to an Australian study, asthma in children had no effect on CVD mortality and CAD, HF, or major CVD events in adults [38]. Adult asthma is greatly influenced by factors intrinsic to the body, such as hormones and chest wall stiffness [30]. These inherent variables may increase the risk of CAD, but cigarette smoking also has an impact on adult asthma and is a risk factor for CAD [30]. In a sample of more than 400,000 Taiwanese adults, patients with active, but not inactive, asthma were more likely to die from CVD [39]. The risk of death from CAD was increased similarly. However, in a sensitivity analysis that excluded people with previous heart disease, only the relationship with CVD mortality persisted. Men had a higher correlation with active asthma than women regarding CVD mortality. Patients with asthma who had moderate to severe acute exacerbation during the previous year have higher mortality following CVD events [40].

PATHOGENETIC MECHANISMS LINKING ASTHMA TO CVD

Cardiovascular complications in patients with asthma can be attributed to the inflammation present in asthma and CVD. However, the specific pathogenetic mechanisms that can facilitate the development of cardiovascular comorbidities have not yet been fully determined.

Despite the limitations translating animal responses into humans, it should be noted that experimentally, acute and chronic allergic lung inflammation evoked in mice by sensitization and challenge with ovalbumin facilitated atheroma formation, regardless of whether lung inflammation occurred before, after, or concurrently with atherogenesis [41]. Pathological changes in the arteries have also been demonstrated in asthmatic patients [42]. Asthma is associated with pre-atherosclerotic vascular changes, such as increased arterial stiffness, and especially with a higher prevalence of overt atherosclerosis than in subjects without asthma. The severity of asthma is related to more pronounced pathological vascular alternations [42]. Furthermore, subjects with persistent asthma exhibited not only a more significant carotid plaque burden but also higher levels of serum inflammatory biomarkers (IL-6 and hsCRP) than those without asthma [43].

Chronic airway inflammation in asthma could contribute to systemic inflammation [44], suggesting that the vasculature is systemically exposed to inflammation generated in the lungs

[45] (figure 1). Female sex hormones have been shown to increase allergic lung inflammation in animal models [46]. This may suggest that systemic inflammatory responses to be triggered more strongly in women with asthma than in men, leading to a pro-inflammatory state that puts women with asthma at an increased risk of developing CVD [46].

Atherosclerosis has long been recognized as a chronic inflammatory disease characterized by inflammation and immunological dysfunction [47]. Several inflammatory cells, including macrophages, monocytes, lymphocytes, neutrophils, and mast cells, which are present in the walls of asthmatic bronchoalveolar and atherosclerotic vessel, play an important role in the pathogenesis of both diseases and perform comparable functions [48]. T1 cells appear to be the key driver driving atherosclerotic CVD [49]. Nevertheless, IgE concentrations are higher in coronary events [49]. However, it is unclear whether this is due to their involvement in the genesis of the coronary event or to the inflammatory response caused by the tissue damage that occurred during the event [50].

In any case, IgE-triggered mast cells and eosinophil leucocytes have been proposed as potential essential players [51]. Both types of cells, by producing chemokines and cytokines, particularly IL-4, IL-6, IL-9, IL-17A, and IL-33, but also interferon- γ and TNF- α [45], enhance vascular permeability and admission of low-density lipoprotein-cholesterol and inflammatory cells into the arterial wall. Furthermore, evidence shows that mast cells accumulate excessively in injured endothelium, which on the one hand stimulates the formation of foam cells and atherosclerotic plaques, and on the other hand, is an important factor in plaque degradation and the initiation of coagulation due to cytokines released from granules and proteolytic enzymes [52].

IL-6 is a crucial effector cytokine in the atherosclerotic T1 process, as it directly inhibits T-regulatory cells, allowing the proatherogenic T1 response to occur [53]. IL-6 also causes endothelial dysfunction, the first step in arterial damage and the production of atherosclerotic plaques, spreads inflammation, and predicts future atherosclerotic CVD events [54, 55].

The shared inflammatory mechanisms of asthma and CVD from the 5-lipoxygenase enzymatic pathway may contribute to long-term concomitant airway and atherosclerosis [56]. Cysteinyl-leukotrienes (CysLTs) operate as powerful smooth muscle constrictors in the airways, causing tissue edema and signaling eosinophil recruitment. Furthermore, they profoundly influence left ventricular function, heart rate, and coronary artery tone, causing myocardial depression when secreted by immunologically activated human cardiac mast cells. There is also evidence that CysLTs can increase the probability of thrombosis by activating of platelets and clotting factors with subsequent coronary thrombosis [57].

Human heart mast cells generate chymase and renin, which locally activate the angiotensin system and cause arteriolar vasoconstriction [58]. Furthermore, when immunologically activated, they also secrete histamine and platelet activating factor, with their well-known harmful effects on asthma and CVD [58].

Smooth muscle remodeling is another process that can be induced by inflammatory mediators. Hyperplasia and aberrant smooth muscle cell contraction leading to airway constriction in asthma are also recognized as characteristics of vascular remodeling and endothelial abnormalities seen in CVD [59].

Uncontrolled asthma, particularly in patients with a T2 inflammatory profile or significant eosinophilic activation, can promote a procoagulant state [16]. However, most patients with asthma do not experience thromboembolic events [16], although increasing evidence suggests that asthma is associated with greater risks of pulmonary embolism [60].

ASTHMA, CVD AND MUTUAL PHARMACOLOGICAL INTERFERENCES

There is a possibility that therapeutic approaches used to treat asthma can cause or worsen CVD [61, 62], but drugs used to treat CVD can also adversely affect asthma [62]. On the other hand, there is growing evidence that some asthma medications can improve CVD and that different CVD treatments can reduce the severity of asthma [62].

Inhaled β_2 -agonist treatment can increase the incidence of acute AMI, atrial fibrillation (AF), congestive HF, cardiac arrest, and sudden cardiac mortality in asthmatic patients [61]. This risk is higher in people with long QT syndrome [59]. The increased risk of AMI, which is not limited to β_2 -agonists but also involves ICSs, generally occurs primarily in the first 3 months, and then decreases [63]. Furthermore, long-term intensive use (more than 13 prescriptions in a year) of an ICS and a SABA has also been associated with an increased risk of AMI [62]. Some patients with asthma may have stronger sympathetic responses to β_2 -agonists contributing to their increased cardiovascular risk [64]. However, this possibility is negated by retrospectively collected data from the Taiwan National Health Insurance database [65]. The results showed that new bronchodilator users, but not previous users, had a reduced risk of coronary vasospastic angina, possibly because new users would be less susceptible to increased sympathetic activity and thus vasospasm. The ability of inhaled β_2 -agonists to induce benefit in patients with acute decompensated HF is important. In fact, they improve cardiovascular hemodynamics, increase cardiac output, reduce peripheral vascular resistance, and reduce pressure in the pulmonary circulation [61]. Furthermore, bronchoconstriction in patients with HF is frequently reversible with inhaled β_2 -agonists, and the resulting decrease in work of breathing might potentially

further reduce cardiac workload [61]. In any case, a recent nested case-control analysis conducted using the UK Clinical Practice Research Datalink of patients with asthma suggested that the use of SABA or ICS/LABA combination therapy was not associated with an increased risk of major adverse cardiovascular events (CVE) defined as the first occurrence of stroke, HF, AMI, arrhythmia, or cardiovascular death compared with ICS [66].

Similarly, other drugs used to treat asthma can adversely affect the heart. For example, ipratropium bromide, a short-acting muscarinic antagonist (SAMA), was associated with a 69% increase in the risk of arrhythmias in patients with asthma 12-24 years of age who used such high-dose SAMA (more than 114 mg) [67]. However, the risk was not statistically significant in low dose SAMA users (114 mg or less). Furthermore, it should be noted that LAMAs, such as tiotropium, glycopyrronium and umeclidinium, demonstrated safety and tolerability comparable to placebo when added to ICS therapy at different treatment steps in adults with symptomatic asthma [68].

Even at therapeutic blood concentrations, theophylline can cause tachycardia and serious arrhythmias, of which multifocal atrial tachycardia may herald sudden cardiac death [69].

Although not explicitly studied in patients with asthma, those receiving high-dose corticosteroid treatment (7.5 mg equivalent of prednisone per day) have an increased risk of developing CAD or HF [70]. Furthermore, the risk of AF occurrence with 7.5 mg equivalent of prednisone per day is six times higher than in patients who did not receive the drug [71]. Apparently, the risk of CVE when using oral doses of prednisolone >10 mg per day is higher during the first 30 days than during the longer period [72].

Interestingly, given higher levels of IgE in patients with CVD and their association with IHD [50], omalizumab has been reported to increase the risk of CVE, as well as arterial and venous thromboembolic events, in individuals with moderate to severe asthma [73]. However, the study that described these findings included individuals at a much higher CAD burden who received omalizumab. In fact, a subsequent pooled analysis of 25 omalizumab-related randomized controlled trials and 2 extension studies did not find evidence of any elevated cardiovascular risk [74]. Nonetheless, there are sporadic reports of cases in which omalizumab has been associated with arterial and venous thromboembolic events [75]. Furthermore, a French pharmacovigilance research demonstrated that omalizumab was associated with hypertension, ventricular arrhythmia, and venous thromboembolism to a significantly greater extent than mepolizumab, reslizumab and benralizumab [76].

Since inhaled bronchodilators must always be administered in combination with ICS in patients with asthma, ICS treatment was compared to ICS-LABA treatment using real-world data; no

significant differences were found between the two groups [77]. However, in women with asthma, the use of ICS is associated with significantly lower cardiovascular and all-cause mortality [78]. On the other hand, patients who had received much lower mean daily doses of ICS in the previous two years were reported to have greater atherosclerotic alterations [74]. ICSs may reduce the risk of atherosclerosis in two ways: directly through absorption across the lungs and exerting a direct impact on the artery wall, or indirectly by lowering the airway inflammatory response [79].

The documentation that CysLTs cause inflammatory changes and induction of immunothrombosis, and consequently, have also been implicated in CVD, is the reason why the use of a CysLT₁RA has been suggested to cause a reduction in AMI and the risk of atherosclerosis [57, 80]. Results of a recent observational retrospective 3-year study of 800 adult asthmatic patients suggest that montelukast, a potent and selective CysLT₁RA, may play a role in the prevention of incident myocardial ischemia events in the elderly asthmatic population [81]. However, although ICS + CysLT₁RA treatment in older adults is associated with a lower risk of CVEs than ICS + LABA treatment, its risk of asthma exacerbations is higher [82].

β -Blockers are frequently used in the treatment of CVD, but they can deleteriously impact asthma patients. Over time, evidence has accumulated on their ability to cause bronchoconstriction, neutralize the effectiveness of β_2 -agonists, and induce moderate and severe asthma exacerbations [83, 84]. However, this unfavorable impact on the heart is mainly observed with non-selective β -blockers while cardio-selective β -blockers, which are 13 to 19 times more potent in blocking β_1 -adrenoceptors (ARs) than β_2 -ARs [85], are better tolerated than non-selective β -blockers, although not completely risk-free mainly in patients with unstable asthma or those with severe airway obstruction [86]. Therefore, asthma is not an absolute contraindication, but these medications should still be prescribed with caution especially if cardio-selective β_1 -blockers are necessary for acute CVEs and alternative treatment options are unavailable [86]. The choice could possibly fall on drugs more suitable for concomitant CVD and asthma, such as nebivolol, a selective for β_1 -AR with nitric oxide-mediated vasodilatory effects and is metabolically neutral, or celiprolol, a β -blocker possessing strong β_1 -AR antagonist and mild β_2 -AR agonist properties, but even in this case the risk of bronchoconstriction, even if of modest magnitude, cannot be ruled out [87]. Chronic use of a nonselective β -blocker with β_2 -AR inverse agonist characteristics, such as nadolol, which exert their beneficial effects on airway epithelial cells and immune cells by inhibiting constitutive pro-inflammatory signaling through non-canonical β -arrestin-2-mediated signaling may be

another choice [88]. In fact, it is well tolerated, may have favorable effects on airway hyperresponsiveness, and does not neutralize the effectiveness of salbutamol, according to two small clinical studies that were not supported by larger research [89, 90].

Statin use is the cornerstone of CVD prevention [91]. Furthermore, even in smokers with asthma, they improve the anti-inflammatory effects of ICSs [92]. However, a Cochrane review published in 2020 concluded that statin therapy was unsuccessful for asthma [93]. Nevertheless, a recent meta-analysis of 8 observational studies and 11 RCTs on asthma reveals that statins can reduce exacerbations and improve asthma management [94]. Furthermore, statins have been shown to improve asthma symptoms and lower levels of hsCRP, sputum eosinophil ratio, and IL-6 without affecting lung function in people with asthma [95].

Due to its role in the development of pro-inflammatory mediators in the lungs, the renin-angiotensin system may have a role in the pathophysiology of asthma. There is substantial meta-analytical evidence linking high circulating ACE concentrations with the onset of asthma [96]. The high levels of angiotensin II and renin found in patients with severe asthma during exacerbations compared to those without exacerbation cause bronchoconstriction, smooth muscle cell hyperplasia, and lung inflammation [97]. As a result, angiotensin converting enzyme (ACE) inhibitors and angiotensin II type 1 (AT1) receptor blockers may be beneficial for people with asthma. However, contrary to expectations, ACE inhibitors were linked to higher asthma morbidity in patients with hypertension and asthma, including increased use of SABAs, systemic corticosteroids, and emergency department visits or hospitalizations [20]. These events are quite rare, although potentially serious, and generally occur in the first few weeks of treatment [98]. On the contrary, a small reduction in airway hyperresponsiveness was observed with AT1 receptor blockers [99].

There is no evidence that calcium channel blockers worsen asthma when administered orally [62]. On the contrary, they may be useful for improving lung function, particularly in exercise-induced asthma [100]. Additionally, calcium channel blockers reduce the annual decline in lung function in asthmatic patients. This effect is thought to be caused by their inhibitory action on inflammation and airway remodeling [101].

Aspirin is the cornerstone for treating ischemic heart disease and secondary prevention of ischemic and valvular heart disease. Aspirin intake can cause the onset of hypersensitivity, known as aspirin-exacerbated respiratory disease (AERD), characterized by symptoms such as rhinorrhea, sneezing, nasal congestion, ocular tearing, bronchospasm, skin flushing, hives, hypotension [102]. After aspirin administration, changes in arachidonic acid metabolism may cause an imbalance between pro-inflammatory and anti-inflammatory substances, resulting in

an overproduction of CysLTs, IL-33/Thymic stromal lymphopoietin, and prostaglandin (PG)D₂ and diminished production (or increased catabolism) of PGE₂ in the respiratory tract [103]. AERD affects a considerable number of patients with asthma, with a prevalence of about 7% [104], and becomes higher in those who also have nasal polyposis, with estimates ranging from 30 to 40% [105].

Antithrombotic agents that do not include aspirin, such as P2Y₁₂ inhibitors (e.g., clopidogrel, prasugrel, and ticagrelor) are treatment alternatives for people who are allergic to aspirin [106]. Clopidogrel was recommended in 2014 by the American College of Cardiology Foundation/American Heart Association guidelines for the management of patients with non-ST-elevation acute coronary syndromes in those with aspirin hypersensitivity, with a loading dosage followed by daily maintenance [107]. Compared to aspirin alone, P2Y₁₂ inhibitor monotherapy for secondary prevention is linked with a considerable reduction in atherothrombotic events without an increased risk of serious bleeding [108]. However, dual antiplatelet medication is suggested during acute coronary syndromes, particularly after percutaneous coronary intervention [109]. Aspirin desensitization is possible for these individuals; however, many “standard” desensitization methods take several days, rendering them problematic in the emergency context [110].

Nevertheless, patients with AERD who are on a stable asthma regimen and have a baseline FEV₁ of at least 70% predicted can be safely desensitized to aspirin using a 90-minute dose escalation protocol, beginning at 40.5 mg, and identifying desensitization as tolerance of the repeated provocation dose and at least one subsequent aspirin dose, bringing total cumulative daily dose to ≥ 325 mg [111]. This procedure is often performed in a single day. Furthermore, there is some evidence that treatment with omalizumab can help reduce respiratory symptoms during aspirin desensitization and potentially restore aspirin tolerance without the need for aspirin desensitization [112].

CONCLUSION

In summary, accumulating evidence has emerged to suggest associations between asthma and CVD. As a result, our understanding of the bidirectional risk of CVD and asthma has increased, although our knowledge about this association and its mechanisms remain incomplete. Indeed, the clinical evidence supporting the hypothesis that asthma confers an elevated CVD risk in patients is not univocal, and the specific biological mechanisms that may facilitate the development of cardiovascular comorbidities are unclear because they often overlap and sometimes conflict with each other. Understanding these biological mechanisms is critical for

choosing pharmacological interventions. Currently, the most appropriate therapeutic approach lies in using the best available evidence to optimize the management of both asthma and CVD. In any case, it will be necessary to conduct specific clinical trials studying the effect on clinical outcomes of drugs known to be helpful for both CVD and asthma.

Article information

Conflict of interest: None declared.

Funding: None.

Open access: This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at kardiologiapolska@ptkardio.pl.

REFERENCES

1. Cloutier MM, Baptist AP, Blake KV, et al. 2020 Focused Updates to the Asthma Management Guidelines: A Report from the National Asthma Education and Prevention Program Coordinating Committee Expert Panel Working Group. *J Allergy Clin Immunol.* 2020; 146(6): 1217–1270, doi: [10.1016/j.jaci.2020.10.003](https://doi.org/10.1016/j.jaci.2020.10.003), indexed in Pubmed: [33280709](https://pubmed.ncbi.nlm.nih.gov/33280709/).
2. Reddel HK, Bacharier LB, Bateman ED, et al. Global Initiative for Asthma Strategy 2021: Executive Summary and Rationale for Key Changes. *Am J Respir Crit Care Med.* 2022; 205(1): 17–35, doi: [10.1164/rccm.202109-2205PP](https://doi.org/10.1164/rccm.202109-2205PP), indexed in Pubmed: [34658302](https://pubmed.ncbi.nlm.nih.gov/34658302/).
3. Lötvall J, Akdis CA, Bacharier LB, et al. Asthma endotypes: a new approach to classification of disease entities within the asthma syndrome. *J Allergy Clin Immunol.* 2011; 127(2): 355–360, doi: [10.1016/j.jaci.2010.11.037](https://doi.org/10.1016/j.jaci.2010.11.037), indexed in Pubmed: [21281866](https://pubmed.ncbi.nlm.nih.gov/21281866/).
4. Fahy JV. Type 2 inflammation in asthma--present in most, absent in many. *Nat Rev Immunol.* 2015; 15(1): 57–65, doi: [10.1038/nri3786](https://doi.org/10.1038/nri3786), indexed in Pubmed: [25534623](https://pubmed.ncbi.nlm.nih.gov/25534623/).
5. Woodruff PG, Modrek B, Choy DF, et al. T-helper type 2-driven inflammation defines major subphenotypes of asthma. *Am J Respir Crit Care Med.* 2009; 180(5): 388–395, doi: [10.1164/rccm.200903-0392OC](https://doi.org/10.1164/rccm.200903-0392OC), indexed in Pubmed: [19483109](https://pubmed.ncbi.nlm.nih.gov/19483109/).

6. Fitzpatrick AM, Chipps BE, Holguin F, et al. T2-"Low" Asthma: Overview and Management Strategies. *J Allergy Clin Immunol Pract.* 2020; 8(2): 452–463, doi: [10.1016/j.jaip.2019.11.006](https://doi.org/10.1016/j.jaip.2019.11.006), indexed in Pubmed: [32037109](https://pubmed.ncbi.nlm.nih.gov/32037109/).
7. Ilmarinen P, Tuomisto LE, Kankaanranta H. Phenotypes, Risk Factors, and Mechanisms of Adult-Onset Asthma. *Mediators Inflamm.* 2015; 2015: 514868, doi: [10.1155/2015/514868](https://doi.org/10.1155/2015/514868), indexed in Pubmed: [26538828](https://pubmed.ncbi.nlm.nih.gov/26538828/).
8. Wenzel SE. Asthma phenotypes: the evolution from clinical to molecular approaches. *Nat Med.* 2012; 18(5): 716–725, doi: [10.1038/nm.2678](https://doi.org/10.1038/nm.2678), indexed in Pubmed: [22561835](https://pubmed.ncbi.nlm.nih.gov/22561835/).
9. Cazzola M, Calzetta L, Rinaldi B, et al. Clinical characteristics, treatment patterns and adherence in patients with asthma on multiple inhaler triple therapy: a review of findings. *Expert Rev Respir Med.* 2022; 16(11-12): 1205–1212, doi: [10.1080/17476348.2023.2167715](https://doi.org/10.1080/17476348.2023.2167715), indexed in Pubmed: [36629483](https://pubmed.ncbi.nlm.nih.gov/36629483/).
10. Cazzola M, Braido F, Calzetta L, et al. The 5T approach in asthma: Triple Therapy Targeting Treatable Traits. *Respir Med.* 2022; 200: 106915, doi: [10.1016/j.rmed.2022.106915](https://doi.org/10.1016/j.rmed.2022.106915), indexed in Pubmed: [35753188](https://pubmed.ncbi.nlm.nih.gov/35753188/).
11. Cazzola M, Matera MG, Rogliani P, et al. Pharmacodynamic and pharmacokinetic assessment of fluticasone furoate + vilanterol for the treatment of asthma. *Expert Opin Drug Metab Toxicol.* 2016; 12(7): 813–822, doi: [10.1080/17425255.2016.1192125](https://doi.org/10.1080/17425255.2016.1192125), indexed in Pubmed: [27253498](https://pubmed.ncbi.nlm.nih.gov/27253498/).
12. Svenningsen S, Nair P. Asthma Endotypes and an Overview of Targeted Therapy for Asthma. *Front Med (Lausanne).* 2017; 4: 158, doi: [10.3389/fmed.2017.00158](https://doi.org/10.3389/fmed.2017.00158), indexed in Pubmed: [29018800](https://pubmed.ncbi.nlm.nih.gov/29018800/).
13. Wood LG, Baines KJ, Fu J, et al. The neutrophilic inflammatory phenotype is associated with systemic inflammation in asthma. *Chest.* 2012; 142(1): 86–93, doi: [10.1378/chest.11-1838](https://doi.org/10.1378/chest.11-1838), indexed in Pubmed: [22345378](https://pubmed.ncbi.nlm.nih.gov/22345378/).
14. Kankaanranta H, Kauppi P, Tuomisto LE, et al. Emerging Comorbidities in Adult Asthma: Risks, Clinical Associations, and Mechanisms. *Mediators Inflamm.* 2016; 2016: 3690628, doi: [10.1155/2016/3690628](https://doi.org/10.1155/2016/3690628), indexed in Pubmed: [27212806](https://pubmed.ncbi.nlm.nih.gov/27212806/).
15. Cazzola M, Rogliani P, Ora J, et al. Asthma and comorbidities: recent advances. *Pol Arch Intern Med.* 2022; 132(4), doi: [10.20452/pamw.16250](https://doi.org/10.20452/pamw.16250), indexed in Pubmed: [35485651](https://pubmed.ncbi.nlm.nih.gov/35485651/).
16. Argueta F, Alviar C, Peters J, et al. Chronic asthma and the risk of cardiovascular disease. *Cardiovascular Complications of Respiratory Disorders.* Sheffield, European Respiratory Society. 2020: 82–95, doi: [10.1183/2312508x.10027519](https://doi.org/10.1183/2312508x.10027519).

17. Guo J, Zhang Y, Liu T, et al. Allergic asthma is a risk factor for human cardiovascular diseases. *Nat Cardiovas Res.* 2022; 1(5): 417–430, doi: [10.1038/s44161-022-00067-z](https://doi.org/10.1038/s44161-022-00067-z).
18. Cazzola M, Calzetta L, Bettoncelli G, et al. Cardiovascular disease in asthma and COPD: a population-based retrospective cross-sectional study. *Respir Med.* 2012; 106(2): 249–256, doi: [10.1016/j.rmed.2011.07.021](https://doi.org/10.1016/j.rmed.2011.07.021), indexed in Pubmed: [21856140](https://pubmed.ncbi.nlm.nih.gov/21856140/).
19. Rogliani P, Laitano R, Ora J, et al. Strength of association between comorbidities and asthma: a meta-analysis. *Eur Respir Rev.* ; 2023(in press).
20. Christiansen SC, Schatz M, Yang SJ, et al. Hypertension and Asthma: A Comorbid Relationship. *J Allergy Clin Immunol Pract.* 2016; 4(1): 76–81, doi: [10.1016/j.jaip.2015.07.009](https://doi.org/10.1016/j.jaip.2015.07.009), indexed in Pubmed: [26342745](https://pubmed.ncbi.nlm.nih.gov/26342745/).
21. Zhang B, Li ZF, An ZY, et al. Association Between Asthma and All-Cause Mortality and Cardiovascular Disease Morbidity and Mortality: A Meta-Analysis of Cohort Studies. *Front Cardiovasc Med.* 2022; 9: 861798, doi: [10.3389/fcvm.2022.861798](https://doi.org/10.3389/fcvm.2022.861798), indexed in Pubmed: [35369308](https://pubmed.ncbi.nlm.nih.gov/35369308/).
22. Hekking PP, Amelink M, Wener RR, et al. Comorbidities in Difficult-to-Control Asthma. *J Allergy Clin Immunol Pract.* 2018; 6(1): 108–113, doi: [10.1016/j.jaip.2017.06.008](https://doi.org/10.1016/j.jaip.2017.06.008), indexed in Pubmed: [28734857](https://pubmed.ncbi.nlm.nih.gov/28734857/).
23. Onufrak SJ, Abramson JL, Austin HD, et al. Relation of adult-onset asthma to coronary heart disease and stroke. *Am J Cardiol.* 2008; 101(9): 1247–1252, doi: [10.1016/j.amjcard.2007.12.024](https://doi.org/10.1016/j.amjcard.2007.12.024), indexed in Pubmed: [18435952](https://pubmed.ncbi.nlm.nih.gov/18435952/).
24. Lee HMu, Truong ST, Wong ND. Association of adult-onset asthma with specific cardiovascular conditions. *Respir Med.* 2012; 106(7): 948–953, doi: [10.1016/j.rmed.2012.02.017](https://doi.org/10.1016/j.rmed.2012.02.017), indexed in Pubmed: [22445771](https://pubmed.ncbi.nlm.nih.gov/22445771/).
25. Pollevick ME, Xu KY, Mhango G, et al. The Relationship Between Asthma and Cardiovascular Disease: An Examination of the Framingham Offspring Study. *Chest.* 2021; 159(4): 1338–1345, doi: [10.1016/j.chest.2020.11.053](https://doi.org/10.1016/j.chest.2020.11.053), indexed in Pubmed: [33316236](https://pubmed.ncbi.nlm.nih.gov/33316236/).
26. Dodd KE, Blackley DJ, Mazurek JM. Cardiovascular Disease Among Adults With Work-Related Asthma, 2012-2017. *Am J Prev Med.* 2023; 64(2): 194–203, doi: [10.1016/j.amepre.2022.09.010](https://doi.org/10.1016/j.amepre.2022.09.010), indexed in Pubmed: [36371324](https://pubmed.ncbi.nlm.nih.gov/36371324/).
27. Schanen JG, Iribarren C, Shahar E, et al. Asthma and incident cardiovascular disease: the Atherosclerosis Risk in Communities Study. *Thorax.* 2005; 60(8): 633–638, doi: [10.1136/thx.2004.026484](https://doi.org/10.1136/thx.2004.026484), indexed in Pubmed: [16061703](https://pubmed.ncbi.nlm.nih.gov/16061703/).

28. Panek M, Mokros Ł, Pietras T, et al. The epidemiology of asthma and its comorbidities in Poland--Health problems of patients with severe asthma as evidenced in the Province of Lodz. *Respir Med.* 2016; 112: 31–38, doi: [10.1016/j.rmed.2016.01.009](https://doi.org/10.1016/j.rmed.2016.01.009), indexed in Pubmed: [26852088](https://pubmed.ncbi.nlm.nih.gov/26852088/).
29. Picado C, Pernigotti A, Arismendi E. Asthma: A New Cardiovascular Risk Factor? *Arch Bronconeumol (Engl Ed).* 2019; 55(7): 353–354, doi: [10.1016/j.arbres.2018.10.016](https://doi.org/10.1016/j.arbres.2018.10.016), indexed in Pubmed: [30559032](https://pubmed.ncbi.nlm.nih.gov/30559032/).
30. Wang L, Gao S, Yu M, et al. Association of asthma with coronary heart disease: A meta analysis of 11 trials. *PLoS One.* 2017; 12(6): e0179335, doi: [10.1371/journal.pone.0179335](https://doi.org/10.1371/journal.pone.0179335), indexed in Pubmed: [28609456](https://pubmed.ncbi.nlm.nih.gov/28609456/).
31. Ingebrigtsen TS, Marott JL, Vestbo J, et al. Coronary heart disease and heart failure in asthma, COPD and asthma-COPD overlap. *BMJ Open Respir Res.* 2020; 7(1), doi: [10.1136/bmjresp-2019-000470](https://doi.org/10.1136/bmjresp-2019-000470), indexed in Pubmed: [33371008](https://pubmed.ncbi.nlm.nih.gov/33371008/).
32. Liu H, Fu Y, Wang K. Asthma and risk of coronary heart disease: A meta-analysis of cohort studies. *Ann Allergy Asthma Immunol.* 2017; 118(6): 689–695, doi: [10.1016/j.anai.2017.03.012](https://doi.org/10.1016/j.anai.2017.03.012), indexed in Pubmed: [28433577](https://pubmed.ncbi.nlm.nih.gov/28433577/).
33. Cepelis A, Brumpton BM, Laugsand LE, et al. Asthma, asthma control and risk of acute myocardial infarction: HUNT study. *Eur J Epidemiol.* 2019; 34(10): 967–977, doi: [10.1007/s10654-019-00562-x](https://doi.org/10.1007/s10654-019-00562-x), indexed in Pubmed: [31512117](https://pubmed.ncbi.nlm.nih.gov/31512117/).
34. Bang DW, Wi CI, Kim ENa, et al. Asthma Status and Risk of Incident Myocardial Infarction: A Population-Based Case-Control Study. *J Allergy Clin Immunol Pract.* 2016; 4(5): 917–923, doi: [10.1016/j.jaip.2016.02.018](https://doi.org/10.1016/j.jaip.2016.02.018), indexed in Pubmed: [27157653](https://pubmed.ncbi.nlm.nih.gov/27157653/).
35. Raita Y, Camargo CA, Faridi MK, et al. Risk of Acute Myocardial Infarction and Ischemic Stroke in Patients with Asthma Exacerbation: A Population-Based, Self-Controlled Case Series Study. *J Allergy Clin Immunol Pract.* 2020; 8(1): 188–194.e8, doi: [10.1016/j.jaip.2019.06.043](https://doi.org/10.1016/j.jaip.2019.06.043), indexed in Pubmed: [31323338](https://pubmed.ncbi.nlm.nih.gov/31323338/).
36. Wee JH, Park MW, Min C, et al. Association between asthma and cardiovascular disease. *Eur J Clin Invest.* 2021; 51(3): e13396, doi: [10.1111/eci.13396](https://doi.org/10.1111/eci.13396), indexed in Pubmed: [32888313](https://pubmed.ncbi.nlm.nih.gov/32888313/).
37. He X, Cheng G, He Lu, et al. Adults with current asthma but not former asthma have higher all-cause and cardiovascular mortality: a population-based prospective cohort study. *Sci Rep.* 2021; 11(1): 1329, doi: [10.1038/s41598-020-79264-4](https://doi.org/10.1038/s41598-020-79264-4), indexed in Pubmed: [33446724](https://pubmed.ncbi.nlm.nih.gov/33446724/).

38. Ng CA, Knuiman MW, Murray K, et al. Childhood asthma and cardiovascular morbidity and mortality in adulthood: The Busselton Health Study. *Pediatr Pulmonol.* 2021; 56(7): 1915–1923, doi: [10.1002/ppul.25386](https://doi.org/10.1002/ppul.25386), indexed in Pubmed: [33819390](https://pubmed.ncbi.nlm.nih.gov/33819390/).
39. Strand LB, Tsai MK, Wen CP, et al. Is having asthma associated with an increased risk of dying from cardiovascular disease? A prospective cohort study of 446 346 Taiwanese adults. *BMJ Open.* 2018; 8(5): e019992, doi: [10.1136/bmjopen-2017-019992](https://doi.org/10.1136/bmjopen-2017-019992), indexed in Pubmed: [29858410](https://pubmed.ncbi.nlm.nih.gov/29858410/).
40. Yu S, Lee SC, Hong JH, et al. Cardiovascular and cerebrovascular mortality in patients with preceding asthma exacerbation. *Respirology.* 2023 [Epub ahead of print], doi: [10.1111/resp.14444](https://doi.org/10.1111/resp.14444), indexed in Pubmed: [36596607](https://pubmed.ncbi.nlm.nih.gov/36596607/).
41. Liu CL, Wang Yi, Liao M, et al. Allergic lung inflammation promotes atherosclerosis in apolipoprotein E-deficient mice. *Transl Res.* 2016; 171: 1–16, doi: [10.1016/j.trsl.2016.01.008](https://doi.org/10.1016/j.trsl.2016.01.008), indexed in Pubmed: [26898714](https://pubmed.ncbi.nlm.nih.gov/26898714/).
42. Tuleta I, Skowasch D, Aurich F, et al. Asthma is associated with atherosclerotic artery changes. *PLoS One.* 2017; 12(10): e0186820, doi: [10.1371/journal.pone.0186820](https://doi.org/10.1371/journal.pone.0186820), indexed in Pubmed: [29073174](https://pubmed.ncbi.nlm.nih.gov/29073174/).
43. Tattersall MC, Dasiewicz AS, McClelland RL, et al. Persistent Asthma Is Associated With Carotid Plaque in MESA. *J Am Heart Assoc.* 2022; 11(23): e026644, doi: [10.1161/JAHA.122.026644](https://doi.org/10.1161/JAHA.122.026644), indexed in Pubmed: [36416156](https://pubmed.ncbi.nlm.nih.gov/36416156/).
44. Jousilahti P, Salomaa V, Hakala K, et al. The association of sensitive systemic inflammation markers with bronchial asthma. *Ann Allergy Asthma Immunol.* 2002; 89(4): 381–385, doi: [10.1016/S1081-1206\(10\)62039-X](https://doi.org/10.1016/S1081-1206(10)62039-X), indexed in Pubmed: [12392382](https://pubmed.ncbi.nlm.nih.gov/12392382/).
45. Gurgone D, McShane L, McSharry C, et al. Cytokines at the Interplay Between Asthma and Atherosclerosis? *Front Pharmacol.* 2020; 11: 166, doi: [10.3389/fphar.2020.00166](https://doi.org/10.3389/fphar.2020.00166), indexed in Pubmed: [32194407](https://pubmed.ncbi.nlm.nih.gov/32194407/).
46. Card JW, Zeldin DC. Hormonal influences on lung function and response to environmental agents: lessons from animal models of respiratory disease. *Proc Am Thorac Soc.* 2009; 6(7): 588–595, doi: [10.1513/pats.200904-020RM](https://doi.org/10.1513/pats.200904-020RM), indexed in Pubmed: [19934354](https://pubmed.ncbi.nlm.nih.gov/19934354/).
47. Gisterå A, Hansson GK. The immunology of atherosclerosis. *Nat Rev Nephrol.* 2017; 13(6): 368–380, doi: [10.1038/nrneph.2017.51](https://doi.org/10.1038/nrneph.2017.51), indexed in Pubmed: [28392564](https://pubmed.ncbi.nlm.nih.gov/28392564/).

48. Liu CL, Zhang JY, Shi GP. Interaction between allergic asthma and atherosclerosis. *Transl Res.* 2016; 174: 5–22, doi: [10.1016/j.trsl.2015.09.009](https://doi.org/10.1016/j.trsl.2015.09.009), indexed in Pubmed: [26608212](https://pubmed.ncbi.nlm.nih.gov/26608212/).
49. Libby P, Lichtman AH, Hansson GK. Immune effector mechanisms implicated in atherosclerosis: from mice to humans. *Immunity.* 2013; 38(6): 1092–1104, doi: [10.1016/j.immuni.2013.06.009](https://doi.org/10.1016/j.immuni.2013.06.009), indexed in Pubmed: [23809160](https://pubmed.ncbi.nlm.nih.gov/23809160/).
50. Kounis NG, Hahalis G. Serum IgE levels in coronary artery disease. *Atherosclerosis.* 2016; 251: 498–500, doi: [10.1016/j.atherosclerosis.2016.05.045](https://doi.org/10.1016/j.atherosclerosis.2016.05.045), indexed in Pubmed: [27288268](https://pubmed.ncbi.nlm.nih.gov/27288268/).
51. Winder B, Kiechl SJ, Gruber NM, et al. The association of allergic asthma and carotid intima-media thickness in adolescence: data of the prospective early vascular ageing (EVA)-Tyrol cohort study. *BMC Cardiovasc Disord.* 2022; 22(1): 11, doi: [10.1186/s12872-021-02452-1](https://doi.org/10.1186/s12872-021-02452-1), indexed in Pubmed: [35042472](https://pubmed.ncbi.nlm.nih.gov/35042472/).
52. Bergmann K, Sypniewska G. Is there an association of allergy and cardiovascular disease? *Biochem Med (Zagreb).* 2011; 21(3): 210–218, doi: [10.11613/bm.2011.030](https://doi.org/10.11613/bm.2011.030), indexed in Pubmed: [22420234](https://pubmed.ncbi.nlm.nih.gov/22420234/).
53. Kimura A, Kishimoto T. IL-6: regulator of Treg/Th17 balance. *Eur J Immunol.* 2010; 40(7): 1830–1835, doi: [10.1002/eji.201040391](https://doi.org/10.1002/eji.201040391), indexed in Pubmed: [20583029](https://pubmed.ncbi.nlm.nih.gov/20583029/).
54. Ridker PM, Rifai N, Stampfer MJ, et al. Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men. *Circulation.* 2000; 101(15): 1767–1772, doi: [10.1161/01.cir.101.15.1767](https://doi.org/10.1161/01.cir.101.15.1767), indexed in Pubmed: [10769275](https://pubmed.ncbi.nlm.nih.gov/10769275/).
55. Spagnoli LG, Bonanno E, Sangiorgi G, et al. Role of inflammation in atherosclerosis. *J Nucl Med.* 2007; 48(11): 1800–1815, doi: [10.2967/jnumed.107.038661](https://doi.org/10.2967/jnumed.107.038661), indexed in Pubmed: [17942804](https://pubmed.ncbi.nlm.nih.gov/17942804/).
56. Tattersall MC, Guo M, Korcarz CE, et al. Asthma predicts cardiovascular disease events: the multi-ethnic study of atherosclerosis. *Arterioscler Thromb Vasc Biol.* 2015; 35(6): 1520–1525, doi: [10.1161/ATVBAHA.115.305452](https://doi.org/10.1161/ATVBAHA.115.305452), indexed in Pubmed: [25908767](https://pubmed.ncbi.nlm.nih.gov/25908767/).
57. Sutton SS, Magagnoli J, Cummings TH, et al. Leukotriene receptor antagonism with montelukast as a possible therapeutic for venous thromboembolism prophylaxis: An observational study. *Prostaglandins Other Lipid Mediat.* 2022; 161: 106649, doi: [10.1016/j.prostaglandins.2022.106649](https://doi.org/10.1016/j.prostaglandins.2022.106649), indexed in Pubmed: [35595009](https://pubmed.ncbi.nlm.nih.gov/35595009/).

58. Triggiani M, Patella V, Staiano RI, et al. Allergy and the cardiovascular system. *Clin Exp Immunol.* 2008; 153 Suppl 1(Suppl 1): 7–11, doi: [10.1111/j.1365-2249.2008.03714.x](https://doi.org/10.1111/j.1365-2249.2008.03714.x), indexed in Pubmed: [18721322](https://pubmed.ncbi.nlm.nih.gov/18721322/).
59. Christiansen SC, Zuraw BL, Christiansen SC, et al. Treatment of Hypertension in Patients with Asthma. *N Engl J Med.* 2019; 381(11): 1046–1057, doi: [10.1056/NEJMra1800345](https://doi.org/10.1056/NEJMra1800345), indexed in Pubmed: [31509675](https://pubmed.ncbi.nlm.nih.gov/31509675/).
60. Lee PH, Fu PK. Pulmonary Embolism and Severe Asthma: Case Report and Literature Review. *Medicina (Kaunas).* 2019; 55(10), doi: [10.3390/medicina55100647](https://doi.org/10.3390/medicina55100647), indexed in Pubmed: [31561622](https://pubmed.ncbi.nlm.nih.gov/31561622/).
61. Cazzola M, Rogliani P, Calzetta L, et al. Bronchodilators in subjects with asthma-related comorbidities. *Respir Med.* 2019; 151: 43–48, doi: [10.1016/j.rmed.2019.04.001](https://doi.org/10.1016/j.rmed.2019.04.001), indexed in Pubmed: [31047116](https://pubmed.ncbi.nlm.nih.gov/31047116/).
62. Ora J, Cavalli F, Cazzola M. Management of patients with asthma or COPD and cardiovascular disease: risks *versus* benefits. *Cardiovascular Complications of Respiratory Disorders (ERS Monograph)*. Sheffield, European Respiratory Society, 2020: 66–81, doi: [10.1183/2312508x.10027419](https://doi.org/10.1183/2312508x.10027419).
63. Zhang B, de Vries F, Setakis E, et al. The pattern of risk of myocardial infarction in patients taking asthma medication: a study with the General Practice Research Database. *J Hypertens.* 2009; 27(7): 1485–1492, doi: [10.1097/HJH.0b013e32832af68d](https://doi.org/10.1097/HJH.0b013e32832af68d), indexed in Pubmed: [19491706](https://pubmed.ncbi.nlm.nih.gov/19491706/).
64. Moore LE, Kapoor K, Byers BW, et al. Acute effects of salbutamol on systemic vascular function in people with asthma. *Respir Med.* 2019; 155: 133–140, doi: [10.1016/j.rmed.2019.07.018](https://doi.org/10.1016/j.rmed.2019.07.018), indexed in Pubmed: [31349187](https://pubmed.ncbi.nlm.nih.gov/31349187/).
65. Hung MJ, Mao CT, Hung MY, et al. Impact of Asthma on the Development of Coronary Vasospastic Angina: A Population-Based Cohort Study. *Medicine (Baltimore).* 2015; 94(42): e1880, doi: [10.1097/MD.0000000000001880](https://doi.org/10.1097/MD.0000000000001880), indexed in Pubmed: [26496346](https://pubmed.ncbi.nlm.nih.gov/26496346/).
66. Amegadzie JE, Gamble JM, Farrell J, et al. Association between Inhaled β -agonists Initiation and Risk of Major Adverse Cardiovascular Events: A Population-based Nested Case-Control Study. *Int J Chron Obstruct Pulmon Dis.* 2022; 17: 1205–1217, doi: [10.2147/COPD.S358927](https://doi.org/10.2147/COPD.S358927), indexed in Pubmed: [35645559](https://pubmed.ncbi.nlm.nih.gov/35645559/).
67. Adimadhyam S, Schumock GT, Walton S, et al. Risk of arrhythmias associated with ipratropium bromide in children, adolescents, and young adults with asthma: a nested case-control study. *Pharmacotherapy.* 2014; 34(4): 315–323, doi: [10.1002/phar.1336](https://doi.org/10.1002/phar.1336), indexed in Pubmed: [23918239](https://pubmed.ncbi.nlm.nih.gov/23918239/).

68. Cazzola M, Rogliani P, Matera MG. The latest on the role of LAMAs in asthma. *J Allergy Clin Immunol.* 2020; 146(6): 1288–1291, doi: [10.1016/j.jaci.2020.06.014](https://doi.org/10.1016/j.jaci.2020.06.014), indexed in Pubmed: [32603663](https://pubmed.ncbi.nlm.nih.gov/32603663/).
69. Bittar G, Friedman HS. The arrhythmogenicity of theophylline. A multivariate analysis of clinical determinants. *Chest.* 1991; 99(6): 1415–1420, doi: [10.1378/chest.99.6.1415](https://doi.org/10.1378/chest.99.6.1415), indexed in Pubmed: [2036824](https://pubmed.ncbi.nlm.nih.gov/2036824/).
70. Wei Li, MacDonald TM, Walker BR. Taking glucocorticoids by prescription is associated with subsequent cardiovascular disease. *Ann Intern Med.* 2004; 141(10): 764–770, doi: [10.7326/0003-4819-141-10-200411160-00007](https://doi.org/10.7326/0003-4819-141-10-200411160-00007), indexed in Pubmed: [15545676](https://pubmed.ncbi.nlm.nih.gov/15545676/).
71. van der Hoof CS, Heeringa J, Brusselle GG, et al. Corticosteroids and the risk of atrial fibrillation. *Arch Intern Med.* 2006; 166(9): 1016–1020, doi: [10.1001/archinte.166.9.1016](https://doi.org/10.1001/archinte.166.9.1016), indexed in Pubmed: [16682576](https://pubmed.ncbi.nlm.nih.gov/16682576/).
72. Varas-Lorenzo C, Rodriguez LA, Maguire A, et al. Use of oral corticosteroids and the risk of acute myocardial infarction. *Atherosclerosis.* 2007; 192(2): 376–383, doi: [10.1016/j.atherosclerosis.2006.05.019](https://doi.org/10.1016/j.atherosclerosis.2006.05.019), indexed in Pubmed: [16787647](https://pubmed.ncbi.nlm.nih.gov/16787647/).
73. Iribarren C, Rahmaoui A, Long AA, et al. Cardiovascular and cerebrovascular events among patients receiving omalizumab: Results from EXCELS, a prospective cohort study in moderate to severe asthma. *J Allergy Clin Immunol.* 2017; 139(5): 1489–1495.e5, doi: [10.1016/j.jaci.2016.07.038](https://doi.org/10.1016/j.jaci.2016.07.038), indexed in Pubmed: [27639934](https://pubmed.ncbi.nlm.nih.gov/27639934/).
74. Iribarren C, Rothman KJ, Bradley MS, et al. Cardiovascular and cerebrovascular events among patients receiving omalizumab: Pooled analysis of patient-level data from 25 randomized, double-blind, placebo-controlled clinical trials. *J Allergy Clin Immunol.* 2017; 139(5): 1678–1680, doi: [10.1016/j.jaci.2016.12.953](https://doi.org/10.1016/j.jaci.2016.12.953), indexed in Pubmed: [28108337](https://pubmed.ncbi.nlm.nih.gov/28108337/).
75. Oblitas CM, Galeano-Valle F, Vela-De La Cruz L, et al. Omalizumab as a Provoking Factor for Venous Thromboembolism. *Drug Target Insights.* 2019; 13: 1177392819861987, doi: [10.1177/1177392819861987](https://doi.org/10.1177/1177392819861987), indexed in Pubmed: [31320796](https://pubmed.ncbi.nlm.nih.gov/31320796/).
76. Quinta JB, Montastruc F, Sommet A, et al. Cardiovascular adverse effects of anti-IL-5/IL-5R α therapies: A real-world study. *J Allergy Clin Immunol Pract.* 2021; 9(3): 1411–1413, doi: [10.1016/j.jaip.2020.12.031](https://doi.org/10.1016/j.jaip.2020.12.031), indexed in Pubmed: [33412311](https://pubmed.ncbi.nlm.nih.gov/33412311/).
77. Park J, Kang H, Hong S, et al. Comparing Inhaled Corticosteroids and Long-Acting Beta Agonists Combination To Inhaled Corticosteroids on Risk of Ischemic

- Cardiovascular Diseases in Patients with Asthma. *Value in Health*. 2017; 20(9): A640, doi: [10.1016/j.jval.2017.08.1462](https://doi.org/10.1016/j.jval.2017.08.1462).
78. Camargo CA, Barr RG, Chen R, et al. Prospective study of inhaled corticosteroid use, cardiovascular mortality, and all-cause mortality in asthmatic women. *Chest*. 2008; 134(3): 546–551, doi: [10.1378/chest.07-3126](https://doi.org/10.1378/chest.07-3126), indexed in Pubmed: [18641096](https://pubmed.ncbi.nlm.nih.gov/18641096/).
 79. Podgórski M, Kupczyk M, Grzelak P, et al. Inhaled Corticosteroids in Asthma: Promoting or Protecting Against Atherosclerosis? *Med Sci Monit*. 2017; 23: 5337–5344, doi: [10.12659/msm.904469](https://doi.org/10.12659/msm.904469), indexed in Pubmed: [29120994](https://pubmed.ncbi.nlm.nih.gov/29120994/).
 80. Hoxha M, Lewis-Mikhael AM, Bueno-Cavanillas A. Potential role of leukotriene receptor antagonists in reducing cardiovascular and cerebrovascular risk: A systematic review of human clinical trials and in vivo animal studies. *Biomed Pharmacother*. 2018; 106: 956–965, doi: [10.1016/j.biopha.2018.07.033](https://doi.org/10.1016/j.biopha.2018.07.033), indexed in Pubmed: [30119268](https://pubmed.ncbi.nlm.nih.gov/30119268/).
 81. Hoxha M, Tedesco CC, Quaglin S, et al. Montelukast Use Decreases Cardiovascular Events in Asthmatics. *Front Pharmacol*. 2020; 11: 611561, doi: [10.3389/fphar.2020.611561](https://doi.org/10.3389/fphar.2020.611561), indexed in Pubmed: [33519477](https://pubmed.ncbi.nlm.nih.gov/33519477/).
 82. Altawalbeh SM, Thorpe CT, Zgibor JC, et al. Antileukotriene Agents Versus Long-Acting Beta-Agonists in Older Adults with Persistent Asthma: A Comparison of Add-On Therapies. *J Am Geriatr Soc*. 2016; 64(8): 1592–1600, doi: [10.1111/jgs.14235](https://doi.org/10.1111/jgs.14235), indexed in Pubmed: [27351988](https://pubmed.ncbi.nlm.nih.gov/27351988/).
 83. Pité H, da Cruz MB, Morais-Almeida M. Obstructive lung diseases and beta-blockers: Where do we stand? *Eur J Intern Med*. 2016; 34: e32–e33, doi: [10.1016/j.ejim.2016.04.024](https://doi.org/10.1016/j.ejim.2016.04.024), indexed in Pubmed: [27179413](https://pubmed.ncbi.nlm.nih.gov/27179413/).
 84. Morales DR, Jackson C, Lipworth BJ, et al. Adverse respiratory effect of acute β -blocker exposure in asthma: a systematic review and meta-analysis of randomized controlled trials. *Chest*. 2014; 145(4): 779–786, doi: [10.1378/chest.13-1235](https://doi.org/10.1378/chest.13-1235), indexed in Pubmed: [24202435](https://pubmed.ncbi.nlm.nih.gov/24202435/).
 85. Baker JG, Wilcox RG. β -Blockers, heart disease and COPD: current controversies and uncertainties. *Thorax*. 2017; 72(3): 271–276, doi: [10.1136/thoraxjnl-2016-208412](https://doi.org/10.1136/thoraxjnl-2016-208412), indexed in Pubmed: [27927840](https://pubmed.ncbi.nlm.nih.gov/27927840/).
 86. Tiotiu A, Novakova P, Kowal K, et al. Beta-blockers in asthma: myth and reality. *Expert Rev Respir Med*. 2019; 13(9): 815–822, doi: [10.1080/17476348.2019.1649147](https://doi.org/10.1080/17476348.2019.1649147), indexed in Pubmed: [31352857](https://pubmed.ncbi.nlm.nih.gov/31352857/).

87. Cazzola M, Noschese P, D'Amato M, et al. Comparison of the effects of single oral doses of nebivolol and celiprolol on airways in patients with mild asthma. *Chest*. 2000; 118(5): 1322–1326, doi: [10.1378/chest.118.5.1322](https://doi.org/10.1378/chest.118.5.1322), indexed in Pubmed: [11083681](https://pubmed.ncbi.nlm.nih.gov/11083681/).
88. Dickey BF, Walker JKL, Hanania NA, et al. beta-Adrenoceptor inverse agonists in asthma. *Curr Opin Pharmacol*. 2010; 10(3): 254–259, doi: [10.1016/j.coph.2010.03.002](https://doi.org/10.1016/j.coph.2010.03.002), indexed in Pubmed: [20399707](https://pubmed.ncbi.nlm.nih.gov/20399707/).
89. Hanania NA, Singh S, El-Wali R, et al. The safety and effects of the beta-blocker, nadolol, in mild asthma: an open-label pilot study. *Pulm Pharmacol Ther*. 2008; 21(1): 134–141, doi: [10.1016/j.pupt.2007.07.002](https://doi.org/10.1016/j.pupt.2007.07.002), indexed in Pubmed: [17703976](https://pubmed.ncbi.nlm.nih.gov/17703976/).
90. Hanania NA, Mannava B, Franklin AE, et al. Response to salbutamol in patients with mild asthma treated with nadolol. *Eur Respir J*. 2010; 36(4): 963–965, doi: [10.1183/09031936.00003210](https://doi.org/10.1183/09031936.00003210), indexed in Pubmed: [20889466](https://pubmed.ncbi.nlm.nih.gov/20889466/).
91. Cardet JC, Bulkhi AA, Lockey RF. Nonrespiratory Comorbidities in Asthma. *J Allergy Clin Immunol Pract*. 2021; 9(11): 3887–3897, doi: [10.1016/j.jaip.2021.08.027](https://doi.org/10.1016/j.jaip.2021.08.027), indexed in Pubmed: [34492402](https://pubmed.ncbi.nlm.nih.gov/34492402/).
92. Thomson NC, Charron CE, Chaudhuri R, et al. Atorvastatin in combination with inhaled beclometasone modulates inflammatory sputum mediators in smokers with asthma. *Pulm Pharmacol Ther*. 2015; 31: 1–8, doi: [10.1016/j.pupt.2015.01.001](https://doi.org/10.1016/j.pupt.2015.01.001), indexed in Pubmed: [25595138](https://pubmed.ncbi.nlm.nih.gov/25595138/).
93. Naing C, Ni H. Statins for asthma. *Cochrane Database Syst Rev*. 2020; 7(7): CD013268, doi: [10.1002/14651858.CD013268.pub2](https://doi.org/10.1002/14651858.CD013268.pub2), indexed in Pubmed: [32668027](https://pubmed.ncbi.nlm.nih.gov/32668027/).
94. Sunata K, Kabata H, Kuno T, et al. The effect of statins for asthma. A systematic review and meta-analysis. *J Asthma*. 2022; 59(4): 801–810, doi: [10.1080/02770903.2021.1879850](https://doi.org/10.1080/02770903.2021.1879850), indexed in Pubmed: [33504228](https://pubmed.ncbi.nlm.nih.gov/33504228/).
95. Zhang QX, Zhang HF, Lu XT, et al. Statins improve asthma symptoms by suppressing inflammation: a meta-analysis based on RCTs. *Eur Rev Med Pharmacol Sci*. 2022; 26(22): 8401–8410, doi: [10.26355/eurev_202211_30376](https://doi.org/10.26355/eurev_202211_30376), indexed in Pubmed: [36459023](https://pubmed.ncbi.nlm.nih.gov/36459023/).
96. Hui Q, Hao Y, Ye F, et al. Genetically high angiotensin-converting enzyme concentrations causally increase asthma risk: A meta-analysis using Mendelian randomization. *Front Med (Lausanne)*. 2022; 9: 941944, doi: [10.3389/fmed.2022.941944](https://doi.org/10.3389/fmed.2022.941944), indexed in Pubmed: [36419791](https://pubmed.ncbi.nlm.nih.gov/36419791/).

97. Gregório JF, Rodrigues-Machado Md, Santos RA, et al. Asthma: role of the angiotensin-(1-7)/Mas (MAS1) pathway in pathophysiology and therapy. *Br J Pharmacol*. 2021; 178(22): 4428–4439, doi: [10.1111/bph.15619](https://doi.org/10.1111/bph.15619), indexed in Pubmed: [34235725](https://pubmed.ncbi.nlm.nih.gov/34235725/).
98. Lunde H, Hedner T, Samuelsson O, et al. Dyspnoea, asthma, and bronchospasm in relation to treatment with angiotensin converting enzyme inhibitors. *BMJ*. 1994; 308(6920): 18–21, doi: [10.1136/bmj.308.6920.18](https://doi.org/10.1136/bmj.308.6920.18), indexed in Pubmed: [8298346](https://pubmed.ncbi.nlm.nih.gov/8298346/).
99. Myou S, Fujimura M, Kamio Y, et al. Effect of losartan, a type 1 angiotensin II receptor antagonist, on bronchial hyperresponsiveness to methacholine in patients with bronchial asthma. *Am J Respir Crit Care Med*. 2000; 162(1): 40–44, doi: [10.1164/ajrccm.162.1.9907127](https://doi.org/10.1164/ajrccm.162.1.9907127), indexed in Pubmed: [10903217](https://pubmed.ncbi.nlm.nih.gov/10903217/).
100. Chiu KYu, Li JG, Lin Y. Calcium channel blockers for lung function improvement in asthma: A systematic review and meta-analysis. *Ann Allergy Asthma Immunol*. 2017; 119(6): 518–523.e3, doi: [10.1016/j.anai.2017.08.013](https://doi.org/10.1016/j.anai.2017.08.013), indexed in Pubmed: [29032888](https://pubmed.ncbi.nlm.nih.gov/29032888/).
101. Tsuzuki R, To M, Yamawaki S, et al. Inhibitory effect of calcium channel blockers on the deterioration of lung function in adult-onset asthma. *Ann Allergy Asthma Immunol*. 2021; 126(6): 731–733, doi: [10.1016/j.anai.2021.02.029](https://doi.org/10.1016/j.anai.2021.02.029), indexed in Pubmed: [33705916](https://pubmed.ncbi.nlm.nih.gov/33705916/).
102. Li KL, Lee AY, Abuzeid WM. Aspirin Exacerbated Respiratory Disease: Epidemiology, Pathophysiology, and Management. *Med Sci (Basel)*. 2019; 7(3), doi: [10.3390/medsci7030045](https://doi.org/10.3390/medsci7030045), indexed in Pubmed: [30884882](https://pubmed.ncbi.nlm.nih.gov/30884882/).
103. Laidlaw TM, Boyce JA. Updates on immune mechanisms in aspirin-exacerbated respiratory disease. *J Allergy Clin Immunol*. 2022 [Epub ahead of print], doi: [10.1016/j.jaci.2022.08.021](https://doi.org/10.1016/j.jaci.2022.08.021), indexed in Pubmed: [36184313](https://pubmed.ncbi.nlm.nih.gov/36184313/).
104. Rajan JP, Wineinger NE, Stevenson DD, et al. Prevalence of aspirin-exacerbated respiratory disease among asthmatic patients: A meta-analysis of the literature. *J Allergy Clin Immunol*. 2015; 135(3): 676–81.e1, doi: [10.1016/j.jaci.2014.08.020](https://doi.org/10.1016/j.jaci.2014.08.020), indexed in Pubmed: [25282015](https://pubmed.ncbi.nlm.nih.gov/25282015/).
105. Jenkins C, Costello J, Hodge L. Systematic review of prevalence of aspirin induced asthma and its implications for clinical practice. *BMJ*. 2004; 328(7437): 434, doi: [10.1136/bmj.328.7437.434](https://doi.org/10.1136/bmj.328.7437.434), indexed in Pubmed: [14976098](https://pubmed.ncbi.nlm.nih.gov/14976098/).
106. Thakker RA, Salazar L, Jazar DA, et al. Coronary Artery Disease and Aspirin Intolerance: Background and Insights on Current Management. *Cardiol Ther*. 2022; 11(2): 175–183, doi: [10.1007/s40119-022-00255-9](https://doi.org/10.1007/s40119-022-00255-9), indexed in Pubmed: [35344187](https://pubmed.ncbi.nlm.nih.gov/35344187/).

107. Amsterdam EA, Wenger NK, Brindis RG, et al. 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014; 130(25): e344–e426, doi: [10.1161/CIR.000000000000134](https://doi.org/10.1161/CIR.000000000000134), indexed in Pubmed: [25249585](https://pubmed.ncbi.nlm.nih.gov/25249585/).
108. Aggarwal D, Bhatia K, Chunawala ZS, et al. P2Y inhibitor versus aspirin monotherapy for secondary prevention of cardiovascular events: meta-analysis of randomized trials. *Eur Heart J Open*. 2022; 2(2): oeac019, doi: [10.1093/ehjopen/oeac019](https://doi.org/10.1093/ehjopen/oeac019), indexed in Pubmed: [35919116](https://pubmed.ncbi.nlm.nih.gov/35919116/).
109. Correction to: 2016 ACC/AHA Guideline Focused Update on Duration of Dual Antiplatelet Therapy in Patients With Coronary Artery Disease: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines: An Update of the 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention, 2011 ACCF/AHA Guideline for Coronary Artery Bypass Graft Surgery, 2012 ACC/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the Diagnosis and Management of Patients With Stable Ischemic Heart Disease, 2013 ACCF/AHA Guideline for the Management of ST-Elevation Myocardial Infarction, 2014 AHA/ACC Guideline for the Management of Patients With Non-ST-Elevation Acute Coronary Syndromes, and 2014 ACC/AHA Guideline on Perioperative Cardiovascular Evaluation and Management of Patients Undergoing Noncardiac Surgery. *Circulation*. 2016; 134(10): e192–e194, doi: [10.1161/CIR.0000000000000452](https://doi.org/10.1161/CIR.0000000000000452), indexed in Pubmed: [27601566](https://pubmed.ncbi.nlm.nih.gov/27601566/).
110. Pattanaik D, Lieberman P, Das P. Aspirin sensitivity and coronary artery disease: implications for the practicing cardiologist. *Future Cardiol*. 2012; 8(4): 555–562, doi: [10.2217/fca.12.38](https://doi.org/10.2217/fca.12.38), indexed in Pubmed: [22871195](https://pubmed.ncbi.nlm.nih.gov/22871195/).
111. DeGregorio GA, Singer J, Cahill KN, et al. A 1-Day, 90-Minute Aspirin Challenge and Desensitization Protocol in Aspirin-Exacerbated Respiratory Disease. *J Allergy Clin Immunol Pract*. 2019; 7(4): 1174–1180, doi: [10.1016/j.jaip.2018.10.032](https://doi.org/10.1016/j.jaip.2018.10.032), indexed in Pubmed: [30391549](https://pubmed.ncbi.nlm.nih.gov/30391549/).
112. Woo SD, Luu QQ, Park HS. NSAID-Exacerbated Respiratory Disease (NERD): From Pathogenesis to Improved Care. *Front Pharmacol*. 2020; 11: 1147, doi: [10.3389/fphar.2020.01147](https://doi.org/10.3389/fphar.2020.01147), indexed in Pubmed: [32848759](https://pubmed.ncbi.nlm.nih.gov/32848759/).

Figure 1. Some pathogenetic mechanisms linking asthma to cardiovascular disease

