



KARDIOLOGIA POLSKA

Polish Heart Journal
The Official Peer-reviewed Journal
of the Polish Cardiac Society
since 1957

Online first

This is a provisional PDF only. Copyedited and fully
formatted version will be made available soon

ISSN 0022-9032

e-ISSN 1897-4279

Transcatheter closure of giant left atrial appendage aneurysm

Authors: Witold Streb, Stanisław Morawski, Tomasz Podolecki, Zbigniew Kalarus

Article type: Clinical vignette

Received: December 7, 2022

Accepted: January 22, 2023

Early publication date: January 27, 2023

This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Transcatheter closure of giant left atrial appendage aneurysm

Short title: Closure of left atrial appendage aneurysm

Witold Streb^{1,2}, Stanisław Morawski², Tomasz Podolecki^{1,2}, Zbigniew Kalarus^{1,2}

¹Department of Cardiology, Congenital Heart Diseases and Electrotherapy, Faculty of Medical Sciences in Zabrze, Medical University of Silesia, Katowice, Poland

²Silesian Center for Heart Diseases in Zabrze, Zabrze, Poland

Correspondence to:

Witold Streb, MD, PhD,

Silesian Center for Heart Diseases in Zabrze,

Curie-Skłodowskiej 9, 41–800 Zabrze, Poland,

phone: +48 373 36 48,

e-mail: w.streb@sccs.pl

A 67-year-old female patient diagnosed with left atrial appendage (LAA) aneurysm and persistent atrial fibrillation was referred for expanded diagnosis and treatment. The patient had a several-year paroxysmal atrial fibrillation history associated with troublesome symptoms (European Heart Rhythm Association [EHRA] class 2b). For this reason, she underwent successful electrical cardioversions in May 2021 and March 2022. One day after the second cardioversion, an ischemic stroke occurred despite using non-vitamin K antagonist oral anticoagulants (NOAC). Treatment at that time included mechanical thrombectomy, which prevented permanent neurological deficits. As part of further stroke prevention, given the ineffectiveness of NOAC, it was decided to include Warfarin. During further follow-up, the patient developed a recurrence of arrhythmia. For this reason, the patient was qualified for cryoablation of the pulmonary veins.

Transesophageal echocardiography (TEE) before cryoablation showed an LAA aneurysm with a longitudinal dimension of 63 × 40 × 39 mm with a distal thrombus and sludge, with a preserved LAA neck structure (width 23 × 31 mm, length 13 mm). The dimensions of the LAA

aneurysm assessed by a computed tomography (CT) scan were $4.59 \times 4.66 \times 4.9$ cm. Contrast defects indicating the presence of a thrombus in the lumen of the aneurysm were seen.

The heart team qualified the patient for endovascular elimination of the LAA aneurysm, with cryoablation performed in the second stage. The LAA closure procedure was performed using a 34 mm Amplatzer Amulet occlude and Sentinel Cerebral Protection System. Despite the use of the neuroprotection system, the implantation technique was modified by forgoing the LAA angiography and relying on the so-called touchless technique. It involves avoiding the insertion of catheters deep into the lumen of the LAA lobe, which in this case was facilitated by the size of the aneurysm and the presence of a thrombus in the top of the LAA lobe. These modifications to the implantation technique reduce the risk that the thrombus will migrate beyond the LAA. After the occluder's expansion in the LAA's neck, no flow was recorded in LAA. The device's disk completely covered the entrance to LAA, protecting not only the flow of blood into the LAA but also guarding the occluder against possible displacement into the lumen of the aneurysm after its release. After implantation of the occluder, the neuroprotection system was withdrawn, and its filters were dissected and flushed with saline. In the washings, only two approximately 2-millimetre structures were found, which may correspond to small fragments of thrombus. The procedure itself and the post-operative period went without complications. The patient was discharged the next day with a recommendation to take acetylsalicylic acid and clopidogrel until a follow-up visit was scheduled 6 weeks after the procedure.

The TEE examination performed at the follow-up visit showed no dislocation of the occluder or the presence of a thrombus or leakage on its surface. CT scans also verified the results of the TEE study. According to a previous qualification, successful cryoablation of all pulmonary veins was performed.

LAA aneurysm is one of the infrequent anomalies. Only single cases have been described in the literature, based on which both congenital and acquired etiology of this condition can be inferred. LAA aneurysms are related to factors such as trauma to the LAA wall, an inflammatory process in the left atrial wall, abnormal vascularization, tuberculosis, or valvular defects [1].

Most cases, the clinical symptoms associated with LAA aneurysms appear after age 30 as the lumen enlarges. In 34% of the cases described, the diagnosis occurred incidentally in asymptomatic individuals [2]. The most common symptoms associated with LAA aneurysms are supraventricular arrhythmias, strokes, shortness of breath, chest pain and cough [3]. Cases of cardiac tamponade have also been described [4]. Because of the complications associated with LAA aneurysms, most notably the risk of the central nervous system and peripheral

embolization, LAA aneurysms are often removed by cardiac surgery [5]. To the authors' knowledge, the case report presented here is the first case of transcatheter closure of an LAA aneurysm.

Article information

Conflict of interest: ZK and WS are proctors of Abbott company.

Funding: None.

Open access: This article is available in open access under Creative Common Attribution-Non-Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) license, allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially. For commercial use, please contact the journal office at kardiologiapolska@ptkardio.pl.

REFERENCES

1. Fakhri G, Obeid M, El Rassi I, et al. Large congenital left atrial wall aneurysm: An updated and comprehensive review of the literature. *Echocardiography*. 2020; 37(6): 965–970, doi: [10.1111/echo.14687](https://doi.org/10.1111/echo.14687), indexed in Pubmed: [32428351](https://pubmed.ncbi.nlm.nih.gov/32428351/).
2. Hoffmann U, Hamed N, Herold C, et al. Radiological signs of a left atrial aneurysm. *Eur Radiol*. 2000; 10(8): 1332–1334, doi: [10.1007/s003309900307](https://doi.org/10.1007/s003309900307), indexed in Pubmed: [10939502](https://pubmed.ncbi.nlm.nih.gov/10939502/).
3. Aryal MR, Hakim FA, Ghimire S, et al. Left atrial appendage aneurysm: a systematic review of 82 cases. *Echocardiography*. 2014; 31(10): 1312–1318, doi: [10.1111/echo.12667](https://doi.org/10.1111/echo.12667), indexed in Pubmed: [24976376](https://pubmed.ncbi.nlm.nih.gov/24976376/).
4. Kaliciński ZM, Orłowska H, Werner B. Images in congenital heart disease. Symptomatic left atrial aneurysm in a neonate. *Cardiol Young*. 2001; 11(6): 654–655, doi: [10.1017/s1047951101001032](https://doi.org/10.1017/s1047951101001032), indexed in Pubmed: [11813919](https://pubmed.ncbi.nlm.nih.gov/11813919/).
5. Zhao J, Ge Y, Yan H, et al. Treatment of congenital aneurysms of the left atrium and left atrial appendage. *Tex Heart Inst J*. 1999; 26(2): 136–139, indexed in Pubmed: [10397438](https://pubmed.ncbi.nlm.nih.gov/10397438/).




Figure 1. **A.** Visualization of LAA aneurysm on CT scan. **B.** Visible thrombus in LAA aneurysm on CT with contrast (asterisk). **C.** Four-chamber view showing LAA aneurysm on TEE. **D.** Fluoroscopy image showing the occluder before release. **E.** TEE view of LAA aneurysm after closure with Amplatzer Amulet occlude. **F.** Image of LAA aneurysm closed with an occluder on CT scan after 6 weeks (study without contrast)

Abbreviations: LA, left atrium; LV, left ventricle; RV, right ventricle; arrows indicate LAA aneurysm; TEE, transesophageal echocardiography