



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

C. R. Acad. Sci. Paris, Ser. I 336 (2003) 785–790



Mathematical Problems in Mechanics/Mathematical Physics

Binary quantum collision operators conserving mass momentum and energy

Pierre Degond^a, Christian Ringhofer^b

^a MIP (UMR CNRS 5640), Université Paul Sabatier, 118, route de Narbonne, 31062 Toulouse cedex 4, France

^b Department of Mathematics, Arizona State University, Tempe, AZ 85287-1804, USA

Received and accepted 12 March 2003

Presented by Philippe G. Ciarlet

Abstract

In this Note, we generalize the Boltzmann collision operator modeling binary particle–particle collisions to a quantum framework using nonlocal quantum entropy principles. *To cite this article: P. Degond, C. Ringhofer, C. R. Acad. Sci. Paris, Ser. I 336 (2003).*

© 2003 Académie des sciences. Published by Éditions scientifiques et médicales Elsevier SAS. All rights reserved.

Résumé

Opérateurs de collisions quantiques conservant la masse, l'impulsion et l'énergie. Dans cette Note, nous généralisons l'opérateur de collision de Boltzmann modélisant les collisions binaires particule–particule au cadre quantique, en utilisant un principe non-local de minimisation d'entropie quantique. *Pour citer cet article: P. Degond, C. Ringhofer, C. R. Acad. Sci. Paris, Ser. I 336 (2003).*

© 2003 Académie des sciences. Published by Éditions scientifiques et médicales Elsevier SAS. All rights reserved.

Version française abrégée

L'objet de cet article est la dérivation d'opérateurs de collisions pour l'équation de Boltzmann quantique (1) où $w(r, p, t)$ est la distribution de Wigner, $r \in \mathbb{R}^3$ dénotant la position, $p \in \mathbb{R}^3$, l'impulsion et t , le temps, $V(r, t)$ est le potentiel et l'opérateur $V(r + \frac{\sigma\hbar}{2i}\nabla_p, t)$ dans (1) doit être compris dans le sens des opérateurs pseudo-différentiels, et $q(w, w)$ est un opérateur nonlinéaire modélisant les interactions binaires entre les particules. Le but de cette Note est de proposer une expression adéquate de $q(w, w)$. Différentes expressions de q sont disponibles dans la littérature pour modéliser des collisions avec l'environnement [1–3,7]. Ici, nous nous intéressons aux interactions binaires. A notre connaissance, l'expression proposée est la première permettant de satisfaire les deux principes énoncés ci-dessous.

Par analogie avec le cas classique, nous dérivons une forme générale de l'opérateur en se fondant sur deux principes : (i) q conserve localement un ensemble de grandeurs macroscopiques (par ex. la masse, l'impulsion et

E-mail addresses: degond@mip.ups-tlse.fr (P. Degond), ringhofer@asu.edu (C. Ringhofer).

l'énergie); (ii) q dissipe l'entropie. La différence avec le cas classique est que l'entropie est définie globalement comme étant la trace d'un opérateur, et non localement.

La contrainte (i) combinée à des considérations de symétrie et de microréversibilité conduit à une expression de l'opérateur de la forme (4) dans laquelle l'opérateur de taux de transition K satisfait (5). La contrainte (ii) est satisfaite dès que le taux de transition K s'écrit sous la forme (8) où S est la section efficace de collision et $A[w]$ est un opérateur défini par (12). Cet opérateur est non-local à la fois en impulsion et en position en raison de la non localité de l'opérateur W défini par (3) et de son inverse W^{-1} . On en déduit donc une expression de q de la forme (13).

Dans le cas où les quantités conservées se réduisent aux quantités hydrodynamiques habituelles (masse, impulsion et énergie), l'opérateur q ainsi défini admet pour états d'équilibres (éléments du noyau de q) les maxwelliennes quantiques, qui sont de la forme (15). On notera que là encore, du fait de la non-localité de l'opérateur W , les relations entre les paramètres de la Maxwellienne quantique (les fonctions $a_m(r, t)$), et ses moments (la densité, impulsion et énergie) est non-locale en espace. Cette caractéristique a déjà été notée et exploitée dans [5,6].

1. Introduction

This paper is concerned with the derivation of collision operators for the quantum Boltzmann equation

$$\partial_t w + \frac{1}{m_*} \nabla_r \cdot (w p) - \frac{i}{\hbar} \sum_{\sigma=\pm 1} \sigma V \left(r + \frac{\sigma \hbar}{2i} \nabla_p, t \right) w = q(w, w), \quad (1)$$

where $w(r, p, t)$ denotes the Wigner function of a mixed quantum state, $r \in \mathbb{R}^3$ denotes position and $p \in \mathbb{R}^3$ denotes momentum. The function $V(r, t)$ denotes the potential and the operator $V(r + \frac{\sigma \hbar}{2i} \nabla_p, t)$ in (1) is understood in the sense of pseudo differential operators [8]. $q(w, w)$ is a bilinear operator modeling binary collisions between particles. The Wigner equation (1) is obtained from the Heisenberg equation for the density matrix $\rho(x, y, t)$

$$i\hbar \partial_t \rho = [H, \rho] + i\hbar Q(\rho, \rho), \quad (2)$$

where $[H, \rho]$ denotes the usual commutator of ρ with the Hamiltonian $H = -\frac{\hbar^2}{2m_*} \Delta + V$, through the Wigner–Weyl transform

$$w = W[\rho](r, p, t) = (2\pi)^{-3} \int \rho \left(r - \frac{\hbar}{2} \eta, r + \frac{\hbar}{2} \eta, t \right) \exp(i\eta \cdot p) d\eta. \quad (3)$$

The form of the collision operator Q , which is related to q in the Wigner picture (1) via $q(W[\rho], W[\rho]) = W[Q(\rho, \rho)]$, is the subject of this paper. Various approaches exist to modeling collisions of particles with a background in a quantum mechanical framework [1–3,7]. Since they model collisions with a background they result in linear collision operators. We, on the other hand, are concerned with particle–particle collisions, and therefore the collision operators in (1) and (2) have to be nonlinear. In analogy to the classical case, we will derive the general form of the collision operator q from two requirements, namely

- $q(w, w)$ locally conserves a given set of quantities $\kappa_m(p)$, $m = 0, \dots, M$, where we always assume $\kappa_0(p) = 1$, i.e., we assume at least mass conservation.
- $q(w, w)$, or in the density matrix picture (2) $Q(\rho, \rho)$, dissipates the logarithmic quantum entropy.

Based on these two assumptions we can derive the form of $Q(\rho, \rho)$. The derivation is analogous to the one leading to the classical Boltzmann operator [4], except that the quantum entropy is defined as a global quantity, namely the trace of an operator, instead of the simple logarithm of the density function.

2. The quantum collision operator

In this section we derive the form of the collision operator from the two requirements above. The way the local conservation properties are used in Section 2.1 is precisely the same as for the classical Boltzmann collision operator and can be found cf. in [4]. The quantum nature of the collision operator enters through the use of the quantum entropy in Section 2.2.

2.1. Conserved quantities

We start by using the assumption that $q(w, w)$ locally conserves mass, i.e., $\kappa_0 = 1$ holds, or

$$\int q(w, w)(r, p, t) dp = 0, \quad \forall w, r, t,$$

holds. This implies that we can write the collision operator q in weak form as

$$\begin{aligned} & \int \phi(p)q(w, w)(r, p, t) dp \\ &= \frac{1}{2} \int K(w, w, t, r, p, p_1, p', p'_1)[\phi(p') + \phi(p'_1) - \phi(p) - \phi(p_1)] dp dp_1 dp' dp'_1, \end{aligned} \tag{4}$$

for any test function $\phi(p)$, where K is an operator acting on the Wigner function w satisfying the symmetry relations

$$K(w, w, t, r, p, p_1, p', p'_1) = K(\dots, p_1, p, p', p'_1) = K(\dots, p, p_1, p'_1, p') = K(\dots, p', p'_1, p, p_1). \tag{5}$$

The first two equalities in (5) express particle indiscernability. The last one is the so-called microreversibility property, which states that the direct and reverse collision processes are equally probable. Clearly, the form (4) implies local mass conservation since the integral vanishes for constant test functions ϕ . On the other hand, any mass conserving operator q can be straightforwardly written in the form (4). Next we use the additional conservation properties, i.e., the integral in (4) has to vanish for all Wigner functions w when choosing $\kappa_m, m = 1, \dots, M$, as test functions. So

$$\begin{aligned} & \int K(w, w, t, r, p, p_1, p', p'_1)[\kappa_m(p') + \kappa_m(p'_1) - \kappa_m(p) - \kappa_m(p_1)] dp dp_1 dp' dp'_1 = 0, \\ & m = 1, \dots, M, \quad \forall w, r, t, \end{aligned}$$

has to hold. Note, that the index m only ranges from $m = 1, \dots, M$ since $\kappa_0 = 1$ is automatically conserved by writing q in the form (4). This implies that K is supported only on the set of p, p_1, p', p'_1 for which $\kappa_m(p') + \kappa_m(p'_1) - \kappa_m(p) - \kappa_m(p_1) = 0$ holds for $m = 1, \dots, M$. Therefore we can write

$$K(w, w, t, r, p, p_1, p', p'_1) = K_1(w, w, t, r, p, p_1, p', p'_1) \prod_{m=1}^M \delta(\kappa_m(p') + \kappa_m(p'_1) - \kappa_m(p) - \kappa_m(p_1)), \tag{6}$$

where the operator K_1 satisfies the same symmetry relations (5) as K . Usually, momentum and energy will be conserved in addition to mass. So $M = 4, \kappa_i(p) = p_i, \kappa_4(p) = \frac{|p|^2}{2m^*}$ will hold.

2.2. Dissipation of the logarithmic quantum entropy

As an additional requirement we impose that the operator Q in the Heisenberg equation (2) dissipates the quantum entropy [9], i.e., that

$$\text{Tr}[\ln_M(\rho)Q(\rho, \rho)] \leq 0, \quad \forall \rho,$$

holds for all density matrices ρ . Here, we denote by $\ln_M(\rho)$ the logarithm of a selfadjoint operator (or density matrix), defined in the usual way via its spectral decomposition. The symbol Tr denotes the trace of the operator, so $\text{Tr}[\rho_1\rho_2] = \int \rho_1(x, y)\rho_2(y, x) dx dy$ holds. A direct calculation, using the definition (3) of the Wigner transform, yields that the trace of the product of two density matrices translates in the Wigner picture to

$$\text{Tr}[\rho_1\rho_2] = \int W[\rho_1](r, p, t)W[\rho_2](r, p, t) dr dp.$$

Thus, we require that

$$\int W[\ln_M(\rho)]q(w, w) dr dp \leq 0$$

holds for all density matrices ρ with $w = W[\rho]$. Note, that, other than in the classical case, the entropy is a global quantity. Using the weak formulation (4) of the collision operator q this gives

$$\int K(w, w, t, r, p, p_1, p', p'_1)(f' + f'_1 - f - f_1) dr dp dp_1 dp'_1 \leq 0, \quad f = W[\ln_M(\rho)], \quad w = W(\rho), \quad (7)$$

where we write, for short $f' = f(r, p', t)$, $f_1 = f(r, p_1, t)$ and so on. In analogy to the classical case, we write the operator K in (4) as

$$K(w, w, t, r, p, p_1, p', p'_1) = A[w](r, p, t)A[w](r, p_1, t)S(r, p, p_1, p', p'_1) \quad (8)$$

with A some operator acting on the Wigner function w and the scattering cross section S satisfying the symmetry conditions

$$S(r, p, p_1, p', p'_1) = S(r, p_1, p, p', p'_1) = S(r, p, p_1, p'_1, p') = S(r, p', p'_1, p, p_1). \quad (9)$$

In addition, because of the conservation properties discussed in Section 2.1, the scattering cross section S has to be of the form

$$S(r, p, p_1, p', p'_1) = S_1(r, p, p_1, p', p'_1) \prod_{m=1}^M \delta(\kappa_m(p') + \kappa_m(p'_1) - \kappa_m(p) - \kappa_m(p_1)), \quad (10)$$

in order to satisfy (6). The additional symmetry of S in (9) allows us to write (7) as

$$\frac{1}{2} \int (gg_1 - g'g'_1)S(r, p, p_1, p', p'_1)(f' + f'_1 - f - f_1) dr dp dp_1 dp'_1 \leq 0, \\ g = A[w], \quad f = W[\ln_M(\rho)], \quad w = W(\rho). \quad (11)$$

Assuming that the scattering cross section S_1 is strictly positive, the inequality (11) can now be achieved by setting $g = e^f$ or

$$A[w] = A[W[\rho]] = e^{W[\ln_M(\rho)]} = e^{W[\ln_M(W^{-1}(w))]} \quad (12)$$

This gives for the collision operator $q(w, w)$ in its strong formulation

$$q(w, w)(r, p, t) = \int S(r, p, p_1, p', p'_1)(g'g'_1 - gg_1) dp'_1 dp_1, \quad g(r, p, t) = e^{W[\ln_M(W^{-1}[w])]}, \quad (13)$$

where the inverse Wigner transform W^{-1} is given by

$$W^{-1}[w](x, y, t) = \int w\left(\frac{x+y}{2}, p, t\right) \exp\left[\frac{i}{\hbar} p \cdot (x-y)\right] dp, \quad (14)$$

and the kernel S is given by (10) with some scattering cross section S_1 . Note, that the exponential function in (13) is the usual exponential function while the logarithm is the logarithm of an operator.

3. Equilibria and local Maxwellians

We now investigate the kernel of the collision operator $q(w, w)$, given in (13). If the density matrix ρ_0 is such that $q(w_0, w_0) = 0$ holds for $w_0 = W[\rho_0]$, then obviously

$$\int W[\ln_M(\rho_0)]q(w_0, w_0) \, dr \, dp = 0$$

holds as well. In this case, using (11), we obtain

$$\frac{1}{2} \int (gg_1 - g'g'_1)S(r, p, p_1, p', p'_1)(f' + f'_1 - f - f_1) \, dp \, dp_1 \, dp' \, dp'_1 = 0$$

with $g = e^{W[\ln_M(\rho_0)]}$, $f = W[\ln_M(\rho_0)] = \ln(g)$.

Using the form (10) of the scattering cross section S , and the fact that $S_1 > 0$, holds we obtain

$$(e^{f'+f'_1} - e^{f+f_1})(f' + f'_1 - f - f_1) \prod_{m=1}^M \delta(\kappa_m(p') + \kappa_m(p'_1) - \kappa_m(p) - \kappa_m(p_1)) = 0 \quad \forall p, p_1, p', p'_1.$$

According to a celebrated result of Boltzmann, in the case where the conserved quantities are the hydrodynamic ones, i.e., $M = 4$, $\kappa_i(p) = p_i$, $\kappa_4(p) = |p|^2/(2m_*)$, this implies that f is a linear combination of the quantities $\kappa_m(p)$, $m = 0, \dots, 4$, with coefficients which can depend on the position r and the time t . Note that such a result is not known if more than just the hydrodynamic quantities are conserved. Thus, in this case, we obtain

$$f(r, p, t) = \sum_{m=0}^M a_m(r, t)\kappa_m(p), \quad \text{and} \quad \rho_0 = \exp_M(W^{-1}[f]), \quad w_0 = W[\exp_M(W^{-1}[f])], \tag{15}$$

for the equilibrium density matrix and Wigner function ρ_0 and w_0 . On the other hand, a Wigner function of the form (15) clearly is in the kernel of the operator q defined by (13), and therefore the kernel of q consists precisely of Wigner functions of the form (15). Here \exp_M denotes the matrix exponential defined via the spectral decomposition of ρ and W and W^{-1} are defined by (3) and (14) respectively. (15) is the quantum equivalent of the classical local Maxwellian and has been used in [5,6] for the derivation of hydrodynamic closures of the quantum Boltzmann equation.

4. Conclusions

Based on the assumptions that binary collisions locally conserve a given set of quantities and dissipate the global logarithmic quantum entropy, we have derived a quantum version of the Boltzmann collision operator. The nonlocal nature of quantum collisions is reflected by the fact that, although its conservation properties are local, the operator itself is spatially nonlocal because of the operator logarithm in (13). The quantum collision operator will reduce to the usual Boltzmann operator in the classical limit, since the operator $A[w] = e^{W[\ln_M(W^{-1}[w])]}$ can be expected to reduce to the identity in this limit.

Acknowledgements

Work supported by NSF award DECS-0218008. Support by the European network HYKE, funded by the EC as contract HPRN-CT-2002-00282, is also acknowledged.

References

- [1] P. Argyres, Quantum kinetic equations for electrons in high electric and phonon fields, *Phys. Lett. A* 171 (1992).
- [2] A. Arnold, J. Lopez, P. Markowich, J. Soler, An analysis of quantum Fokker–Planck models: A Wigner function approach, Preprint, 2002.
- [3] J. Barker, D. Ferry, Self-scattering path-variable formulation of high-field, time-dependent, quantum kinetic equations for semiconductor transport in the finite collision–duration regime, *Phys. Rev. Lett.* 42 (1997).
- [4] C. Cercignani, The Boltzmann Equation and Its Applications, in: *Appl. Math. Sci.*, Vol. 67, Springer-Verlag, 1988.
- [5] P. Degond, C. Ringhofer, Quantum moment hydrodynamics and the entropy principle, *J. Stat. Phys.* (2002), submitted. Preprint available at URL: <http://math.la.asu.edu/~chris>.
- [6] P. Degond, C. Ringhofer, A note on quantum moment hydrodynamics and the entropy principle, *C. R. Acad. Sci. Paris, Ser. I* 335 (2002) 967–972.
- [7] F. Fromlet, P. Markowich, C. Ringhofer, A Wignerfunction approach to phonon scattering, *VLSI Design* 9 (1999) 339–350.
- [8] M. Shubin, *Pseudodifferential Operators and Spectral Theory*, Springer, 1980.
- [9] D. Zubarev, V. Morozov, G. Röpke, *Statistical Mechanics of Nonequilibrium Processes*, Vol. 1, Basic Concepts, Kinetic Theory, Akademie-Verlag, Berlin, 1996.