



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Group theory/Algebraic geometry

# The Lie algebra of type $G_2$ is rational over its quotient by the adjoint action $\star$



## Rationalité de l'algèbre de Lie de type $G_2$ sur son quotient par l'action adjointe

Dave Anderson<sup>a</sup>, Mathieu Florence<sup>b</sup>, Zinovy Reichstein<sup>c</sup><sup>a</sup> Instituto Nacional de Matemática Pura e Aplicada, Rio de Janeiro, RJ 22460-320, Brazil<sup>b</sup> Institut de mathématiques de Jussieu, Université Paris-6, 4, place Jussieu, 75005 Paris, France<sup>c</sup> Department of Mathematics, University of British Columbia, BC V6T 1Z2, Canada

## ARTICLE INFO

## Article history:

Received 27 August 2013

Accepted 24 October 2013

Presented by Jean-Pierre Serre

## ABSTRACT

Let  $G$  be a split simple group of type  $G_2$  over a field  $k$ , and let  $\mathfrak{g}$  be its Lie algebra. Answering a question of J.-L. Colliot-Thélène, B. Konyavskiĭ, V.L. Popov, and Z. Reichstein, we show that the function field  $k(\mathfrak{g})$  is generated by algebraically independent elements over the field of adjoint invariants  $k(\mathfrak{g})^G$ .

© 2013 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

## R É S U M É

Soit  $G$  un groupe algébrique simple et déployé de type  $G_2$  sur un corps  $k$ . Soit  $\mathfrak{g}$  son algèbre de Lie. On démontre que le corps des fonctions  $k(\mathfrak{g})$  est transcendant pur sur le corps  $k(\mathfrak{g})^G$  des invariants adjoints. Ceci répond par l'affirmative à une question posée par J.-L. Colliot-Thélène, B. Konyavskiĭ, V.L. Popov et Z. Reichstein.

© 2013 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

## 1. Introduction

Let  $G$  be a split connected reductive group over a field  $k$  and let  $\mathfrak{g}$  be the Lie algebra of  $G$ . We will be interested in the following natural question:

**Question 1.** Is the function field  $k(\mathfrak{g})$  *purely transcendental* over the field of invariants  $k(\mathfrak{g})^G$  for the adjoint action of  $G$  on  $\mathfrak{g}$ ? That is, can  $k(\mathfrak{g})$  be generated over  $k(\mathfrak{g})^G$  by algebraically independent elements?

In [5], the authors reduce this question to the case where  $G$  is simple, and show that in the case of simple groups, the answer is affirmative for split groups of types  $A_n$  and  $C_n$ , and negative for all other types except possibly for  $G_2$ . The standing assumption in [5] is that  $\text{char}(k) = 0$ , but here we work in arbitrary characteristic.

The purpose of this note is to settle **Question 1** for the remaining case  $G = G_2$ .

$\star$  D.A. was partially supported by NSF Grant DMS-0902967. Z.R. was partially supported by National Sciences and Engineering Research Council of Canada Grant No. 250217-2012.

E-mail addresses: [dave@impa.br](mailto:dave@impa.br) (D. Anderson), [florence@math.jussieu.fr](mailto:florence@math.jussieu.fr) (M. Florence), [reichst@math.ubc.ca](mailto:reichst@math.ubc.ca) (Z. Reichstein).

**Theorem 2.** *Let  $k$  be an arbitrary field and  $G$  be the simple split  $k$ -group of type  $G_2$ . Then  $k(\mathfrak{g})$  is purely transcendental over  $k(\mathfrak{g})^G$ .*

Under the same hypothesis, and also assuming  $\text{char}(k) = 0$ , it follows from [Theorem 2](#) and [\[5, Theorem 4.10\]](#) that the field extension  $k(G)/k(G)^G$  is also purely transcendental, where  $G$  acts on itself by conjugation.

Apart from settling the last case left open in [\[5\]](#), we were motivated by the (still mysterious) connection between [Question 1](#) and the Gelfand–Kirillov (GK) conjecture [\[9\]](#). In this context,  $\text{char}(k) = 0$ . A. Premet [\[11\]](#) recently showed that the GK conjecture fails for simple Lie algebras of any type other than  $A_n$ ,  $C_n$  and  $G_2$ . His paper relies on the negative results of [\[5\]](#) and their characteristic  $p$  analogues ([\[11\]](#), see also [\[5, Theorem 6.3\]](#)). It is not known whether a positive answer to [Question 1](#) for  $\mathfrak{g}$  implies the GK conjecture for  $\mathfrak{g}$ . The GK conjecture has been proved for algebras of type  $A_n$  (see [\[9\]](#)), but remains open for types  $C_n$  and  $G_2$ . While [Theorem 2](#) does not settle the GK conjecture for type  $G_2$ , it puts the remaining two open cases—for algebras of type  $C_n$  and  $G_2$ —on equal footing vis-à-vis [Question 1](#).

## 2. Twisting

Temporarily, let  $W$  be a linear algebraic group over a field  $k$ . (In the sequel,  $W$  will be the Weyl group of  $G$ ; in particular, it will be finite and smooth.) We refer to [\[7, Section 3\]](#), [\[8, Section 2\]](#), or [\[5, Section 2\]](#) for details about the following facts.

Let  $X$  be a quasi-projective variety with a (right)  $W$ -action defined over  $k$ , and let  $\zeta$  be a (left)  $W$ -torsor over  $k$ . The diagonal left action of  $W$  on  $X \times_{\text{Spec}(k)} \zeta$  (by  $g \cdot (x, z) = (xg^{-1}, gz)$ ) makes  $X \times_{\text{Spec}(k)} \zeta$  into the total space of a  $W$ -torsor  $X \times_{\text{Spec}(k)} \zeta \rightarrow B$ . The base space  $B$  of this torsor is usually called the *twist* of  $X$  by  $\zeta$ . We denote it by  ${}^\zeta X$ .

It is easy to see that if  $\zeta$  is trivial then  ${}^\zeta X$  is  $k$ -isomorphic to  $X$ . Hence,  ${}^\zeta X$  is a  $k$ -form of  $X$ , i.e.,  $X$  and  ${}^\zeta X$  become isomorphic over an algebraic closure of  $k$ .

The twisting construction is functorial in  $X$ : a  $W$ -equivariant morphism  $X \rightarrow Y$  (or rational map  $X \dashrightarrow Y$ ) induces a  $k$ -morphism  ${}^\zeta X \rightarrow {}^\zeta Y$  (resp., rational map  ${}^\zeta X \dashrightarrow {}^\zeta Y$ ).

## 3. The split group of type $G_2$

We fix notation and briefly review the basic facts, referring to [\[13\]](#), [\[1\]](#), or [\[2\]](#) for more details. Over any field  $k$ , a simple split group  $G$  of type  $G_2$  has a faithful seven-dimensional representation  $V$ . Following [\[2, \(3.11\)\]](#), one can fix a basis  $f_1, \dots, f_7$ , with dual basis  $X_1, \dots, X_7$ , so that  $G$  preserves the nonsingular quadratic norm  $N = X_1X_7 + X_2X_6 + X_3X_5 + X_4^2$ . (See [\[1, §6.1\]](#) for the case  $\text{char}(k) = 2$ . In this case  $V$  is not irreducible, since the subspace spanned by  $f_4$  is invariant; the quotient  $V/(k \cdot f_4)$  is the minimal irreducible representation. However, irreducibility will not be necessary in our context.) The corresponding embedding  $G \hookrightarrow \text{GL}_7$  yields a split maximal torus and Borel subgroup  $T \subset B \subset G$ , by intersecting with diagonal and upper-triangular matrices. Explicitly, the maximal torus is:

$$T = \text{diag}(t_1, t_2, t_1t_2^{-1}, 1, t_1^{-1}t_2, t_2^{-1}, t_1^{-1}); \quad (1)$$

cf. [\[2, Lemma 3.13\]](#).

The Weyl group  $W = N(T)/T$  is isomorphic to the dihedral group with 12 elements, and the surjection  $N(T) \rightarrow W$  splits. The inclusion  $G \hookrightarrow \text{GL}_7$  thus gives rise to an inclusion  $N(T) = T \rtimes W \hookrightarrow D \rtimes S_7$ , where  $D \subset \text{GL}_7$  is the subgroup of diagonal matrices. On the level of the dual basis  $X_1, \dots, X_7$ , we obtain an isomorphism  $W \cong S_3 \times S_2$  realized as follows:  $S_3$  permutes the three ordered pairs  $(X_1, X_7)$ ,  $(X_6, X_2)$ , and  $(X_5, X_3)$ , and  $S_2$  exchanges the two ordered triples  $(X_1, X_5, X_6)$  and  $(X_7, X_3, X_2)$ . The variable  $X_4$  is fixed by  $W$ . For details, see [\[2, §A.3\]](#).

The subgroup  $P \subset G$  stabilizing the isotropic line spanned by  $f_1$  is a maximal standard parabolic, and the corresponding homogeneous space  $P \backslash G$  is isomorphic to the five-dimensional quadric  $\mathcal{Q} \subset \mathbb{P}(V)$  defined by the vanishing of the norm, i.e., by the equation:

$$X_1X_7 + X_2X_6 + X_3X_5 + X_4^2 = 0. \quad (2)$$

Note that the quadric  $\mathcal{Q}$  is endowed with an action of  $T$ . An easy tangent space computation shows that  $P$  is smooth regardless of the characteristic of  $k$ .

**Lemma 3.** *The group  $P$  is special, i.e.,  $H^1(l, P) = \{1\}$  for every field extension  $l/k$ . Moreover,  $P$  is rational, as a variety over  $k$ .*

**Proof.** Since the split group of type  $G_2$  is defined over the prime field, we may replace  $k$  by the prime field for the purpose of proving this lemma, and in particular, we may assume  $k$  is perfect. We begin by briefly recalling a construction of Chevalley [\[4\]](#). The isotropic line  $E_1 \subset V$  stabilized by  $P$  is spanned by  $f_1$ , and  $P$  also preserves an isotropic 3-space  $E_3$  spanned by  $f_1, f_2, f_3$ ; see, e.g., [\[2, §2.2\]](#). There is a corresponding map  $P \rightarrow \text{GL}(E_3/E_1) \cong \text{GL}_2$ , which is a split surjection thanks to the block matrix described in [\[10, p. 13\]](#) as the image of “ $B$ ” in  $\text{GL}_7$ . The kernel is unipotent, and we have a split exact sequence corresponding to the Levi decomposition:

$$1 \rightarrow R_u(P) \rightarrow P \rightarrow \text{GL}_2 \rightarrow 1. \quad (3)$$

Combining the exact sequence in cohomology induced by (3) with the fact that both  $R_u(P)$  and  $GL_2$  are special (see [12, pp. 122 and 128]), shows that  $P$  is special.

Since  $P$  is isomorphic to  $R_u(P) \times GL_2$  as a variety over  $k$ , and  $P$  is smooth, so is  $R_u(P)$ . A smooth connected unipotent group over a perfect field is rational [6, IV, §2(3.10)]; therefore  $R_u(P)$  is  $k$ -rational, and so is  $P$ .  $\square$

**4. Proof of Theorem 2**

Keep the notation of the previous section. By a  $W$ -model (of  $k(\mathcal{Q})^T$ ), we mean a quasi-projective  $k$ -variety  $Y$ , endowed with a right action of  $W$ , together with a dominant  $W$ -equivariant  $k$ -rational map  $\mathcal{Q} \dashrightarrow Y$  which, on the level of function fields, identifies  $k(Y)$  with  $k(\mathcal{Q})^T$ . Such a map  $\mathcal{Q} \dashrightarrow Y$  is called a ( $W$ -equivariant) rational quotient map. A  $W$ -model is unique up to a  $W$ -equivariant birational isomorphism; we will construct an explicit one below.

We reduce Theorem 2 to a statement about rationality of a twisted  $W$ -model, in two steps. The first holds for general split connected semisimple groups  $G$ .

**Proposition 4.** *Let  $G$  be a split connected semisimple group over  $k$ , with split maximal  $k$ -torus  $T$ . Let  $K = k(\mathfrak{t})^W$ ,  $L = k(\mathfrak{t})$ , and let  $\zeta$  be the  $W$ -torsor corresponding to the field extension  $L/K$ . If the twisted variety  ${}^\zeta(G_K/T_K)$  is rational over  $K$ , then  $k(\mathfrak{g})$  is purely transcendental over  $k(\mathfrak{g})^G$ .*

**Proof.** Consider the  $(G \times W)$ -equivariant morphism:

$$f : G/T \times_{\text{Spec}(k)} \mathfrak{t} \rightarrow \mathfrak{g}$$

given by  $(\bar{a}, t) \mapsto \text{Ad}(at)$ , where  $\mathfrak{t}$  is the Lie algebra of  $T$ ,  $\bar{a} \in G/T$  is the class of  $a \in G$ , modulo  $T$ . Here  $G$  acts on  $G/T \times \mathfrak{t}$  by translations on the first factor (and trivially on  $\mathfrak{t}$ ), and via the adjoint representation on  $\mathfrak{g}$ . The Weyl group  $W$  naturally acts on  $\mathfrak{t}$  and  $G/T$  (on the right), diagonally on  $G/T \times \mathfrak{t}$ , and trivially on  $\mathfrak{g}$ .

The image of  $f$  contains the semisimple locus in  $\mathfrak{g}$ , so  $f$  is dominant and induces an inclusion  $f^*: k(\mathfrak{g}) \hookrightarrow k(G/T \times \mathfrak{t})$ . Clearly  $f^*k(\mathfrak{g}) \subset k(G/T \times \mathfrak{t})^W$ . We will show that in fact:

$$f^*k(\mathfrak{g}) = k(G/T \times \mathfrak{t})^W. \tag{4}$$

Write  $\bar{k}$  for an algebraic closure of  $k$ , and note that the preimage of a  $\bar{k}$ -point of  $\mathfrak{g}$  in general position is a single  $W$ -orbit in  $(G/T \times \mathfrak{t})_{\bar{k}}$ . To establish (4), it remains to check that  $f$  is smooth at a general point  $(g, x)$  of  $G/T \times \mathfrak{t}$ . (In particular, when  $\text{char}(k) = 0$  nothing more is needed.) To carry out this calculation, we may assume without loss of generality that  $k$  is algebraically closed and (since  $f$  is  $G$ -equivariant)  $g = 1$ . Since  $\dim(G/T \times \mathfrak{t}) = \dim(\mathfrak{g})$ , it suffices to show that the differential:

$$df : T_{(1,x)}(G/T \times \mathfrak{t}) \rightarrow T_x(\mathfrak{g})$$

is surjective, for any regular semisimple element  $x \in \mathfrak{t}$ . Equivalently, we want to show that  $[x, \mathfrak{g}] + \mathfrak{t} = \mathfrak{g}$ . Since  $x$  is regular, we have  $\dim([x, \mathfrak{g}]) + \dim \mathfrak{t} = \dim \mathfrak{g}$ . Thus it remains to show that  $[x, \mathfrak{g}] \cap \mathfrak{t} = 0$ . To see this, suppose  $[x, y] \in \mathfrak{t}$  for some  $y \in \mathfrak{g}$ . Since  $x$  is semisimple, we can write  $y = \sum_{i=1}^r y_{\lambda_i}$ , where  $y_{\lambda}$  is an eigenvector for  $\text{ad}(x)$  with eigenvalue  $\lambda$ , and  $\lambda_1, \dots, \lambda_r$  are distinct. Then  $[x, y] = \sum_{i=1}^r \lambda_i y_{\lambda_i} \in \mathfrak{t}$  is an eigenvector for  $\text{ad}(x)$  with eigenvalue 0. Remembering that eigenvectors of  $\text{ad}(x)$  with distinct eigenvalues are linearly independent, we conclude that  $[x, y] = 0$ . This completes the proof of (4).

It is easy to see  $k(G/T \times \mathfrak{t})^{G \times W} = k(\mathfrak{t})^W$ . Summarizing,  $f^*$  induces a diagram:

$$\begin{array}{ccc} k(G/T \times_{\text{Spec}(k)} \mathfrak{t})^W & \xrightarrow{\sim} & k(\mathfrak{g}) \\ \downarrow & & \downarrow \\ k(\mathfrak{t})^W & \xrightarrow{\sim} & k(\mathfrak{g})^G, \end{array}$$

where the top row is the  $G$ -equivariant isomorphism (4), and the bottom row is obtained from the top by taking  $G$ -invariants. Note that:

$$k(G/T \times_{\text{Spec}(k)} \mathfrak{t}) \simeq K((G/T)_K \times_{\text{Spec}(K)} \text{Spec} L),$$

where  $\simeq$  denotes a  $G$ -equivariant isomorphism of fields. (Recall that  $G$  acts trivially on  $\mathfrak{t}$  and hence also on  $L$  and  $K$ .) Thus the field extension on the left side of our diagram can be rewritten as  $K({}^\zeta(G_K/T_K))/K$ , where  $\zeta$  is the  $W$ -torsor  $\text{Spec}(L) \rightarrow \text{Spec}(K)$ . By assumption, this field extension is purely transcendental; the diagram shows it is isomorphic to  $k(\mathfrak{g})/k(\mathfrak{g})^G$ .  $\square$

For the second reduction, we return to the assumptions of Section 3.

**Proposition 5.** Let  $G$  be a split simple group of type  $G_2$ , with maximal torus  $T$  and Weyl group  $W$ , and let  $\mathcal{Q}$  be the quadric defined in Section 3. Suppose that for a given  $W$ -model  $Y$  of  $k(\mathcal{Q})^T$ , and for some  $W$ -torsor  $\zeta$  over some field  $K/k$ , the twisted variety  ${}^\zeta(Y_K)$  is rational over  $K$ . Then the twisted variety  ${}^\zeta(G_K/T_K)$  is rational over  $K$ .

**Proof.** For the purpose of this proof, we may view  $K$  as a new base field and replace it with  $k$ .

We claim that the left action of  $P$  on  $G/T$  is generically free. Since  $G$  has trivial center, the (characteristic-free) argument at the beginning of the proof of [5, Lemma 9.1] shows that in order to establish this claim it suffices to show that the right  $T$ -action on  $\mathcal{Q} = P \backslash G$  is generically free. The latter action, given by restricting the linear action (1) of  $T$  on  $\mathbb{P}^6$  to the quadric  $\mathcal{Q}$  given by (2), is clearly generically free.

Let  $Y$  be a  $W$ -model. The  $W$ -equivariant rational map  $G/T \dashrightarrow Y$  induced by the projection  $G \rightarrow P \backslash G = \mathcal{Q}$  is a rational quotient map for the left  $P$ -action on  $G/T$ ; cf. [5, p. 458]. Since the  $P$ -action is generically free, this map is a  $P$ -torsor over the generic point of  $Y$ ; see [3, Theorem 4.7]. By the functoriality of the twisting operation, after twisting by a  $W$ -torsor  $\zeta$ , we obtain a rational map  ${}^\zeta(G/T) \dashrightarrow {}^\zeta Y$ , which is a  $P$ -torsor over the generic point of  ${}^\zeta Y$ . This torsor has a rational section, since  $P$  is special; see Lemma 3. In particular,  ${}^\zeta(G/T)$  is  $k$ -birationally isomorphic to  $P \times {}^\zeta Y$ . Since  $P$  is  $k$ -rational (once again, by Lemma 3),  ${}^\zeta(G/T)$  is rational over  ${}^\zeta Y$ . Since  ${}^\zeta Y$  is rational over  $k$ , we conclude that so is  ${}^\zeta(G/T)$ , as desired.  $\square$

It remains to show that the hypothesis of Proposition 5 holds. As before, we may replace the field  $K$  with  $k$ . The following lemma completes the proof of Theorem 2.

**Lemma 6.** Let  $Y$  be a  $W$ -model for  $k(\mathcal{Q})^T$ . The twisted variety  ${}^\zeta Y$  is rational over  $k$ , for every  $W$ -torsor  $\zeta$  over  $k$ .

**Proof.** We begin by constructing an explicit  $W$ -model. The affine open subset  $\mathcal{Q}^{\text{aff}} = \{x_1x_7 + x_2x_6 + x_3x_5 + 1 = 0\} \subset \mathbb{A}^6$  (where  $x_4 \neq 0$ ) is  $N(T)$ -invariant. Here the affine coordinates on  $\mathbb{A}^6$  are  $x_i := X_i/X_4$ , for  $i \neq 4$ . The field of rational functions invariant for the  $T$ -action on  $\mathcal{Q}^{\text{aff}}$  is  $k(y_1, y_2, y_3, z_1, z_2)$ , where the variables

$$y_1 = x_1x_7, \quad y_2 = x_2x_6, \quad y_3 = x_3x_5, \quad z_1 = x_1x_5x_6, \quad \text{and} \quad z_2 = x_2x_3x_7$$

are subject to the relations  $y_1 + y_2 + y_3 + 1 = 0$  and  $y_1y_2y_3 = z_1z_2$ . Thus we may choose as a  $W$ -model the affine subvariety  $\Lambda_1$  of  $\mathbb{A}^5$  given by these two equations, where  $W = S_2 \times S_3$  acts on the coordinates as follows:  $S_2$  permutes  $z_1, z_2$ , and  $S_3$  permutes  $y_1, y_2, y_3$ . (Recall the  $W$ -action defined in Section 3, and note that the field  $k(\mathcal{Q})$  is recovered by adjoining the classes of variables  $x_1$  and  $x_2$ .) We claim that  $\Lambda_1$  is  $W$ -equivariantly birationally isomorphic to

$$\begin{aligned} \Lambda_2 &= \{(Y_1 : Y_2 : Y_3 : Z_0 : Z_1 : Z_2) : Y_1 + Y_2 + Y_3 + Z_0 = 0 \text{ and } Y_1Y_2Y_3 = Z_1Z_2Z_0\} \subset \mathbb{P}^5, \\ \Lambda_3 &= \{(Y_1 : Y_2 : Y_3 : Z_1 : Z_2) : Y_1Y_2Y_3 + (Y_1 + Y_2 + Y_3)Z_1Z_2 = 0\} \subset \mathbb{P}^4, \quad \text{and} \\ \Lambda_4 &= \{(Y_1 : Y_2 : Y_3 : Z_1 : Z_2) : Z_1Z_2 + Y_2Y_3 + Y_1Y_3 + Y_1Y_2 = 0\} \subset \mathbb{P}^4, \end{aligned}$$

where  $W$  acts on the projective coordinates  $Y_1, Y_2, Y_3, Z_1, Z_2, Z_0$  as follows:  $S_2$  permutes  $Z_1, Z_2$ ,  $S_3$  permutes  $Y_1, Y_2, Y_3$ , and every element of  $W$  fixes  $Z_0$ . Note that  $\Lambda_2 \subset \mathbb{P}^5$  is the projective closure of  $\Lambda_1 \subset \mathbb{A}^5$ ; hence, using  $\simeq$  to denote  $W$ -equivariant birational equivalence, we have  $\Lambda_1 \simeq \Lambda_2$ . The isomorphism  $\Lambda_2 \simeq \Lambda_3$  is obtained by eliminating  $Z_0$  from the system of equations defining  $\Lambda_2$ . Finally, the isomorphism  $\Lambda_3 \simeq \Lambda_4$  comes from the Cremona transformation  $\mathbb{P}^4 \dashrightarrow \mathbb{P}^4$  given by  $Y_i \rightarrow 1/Y_i$  and  $Z_j \rightarrow 1/Z_j$  for  $i = 1, 2, 3$  and  $j = 1, 2$ .

Let  $\zeta$  be a  $W$ -torsor over  $k$ . It remains to be shown that  ${}^\zeta \Lambda_4$  is  $k$ -rational. Since  $\Lambda_4$  is a  $W$ -equivariant quadric hypersurface in  $\mathbb{P}^4$ , and the  $W$ -action on  $\mathbb{P}^4$  is induced by a linear representation  $W \rightarrow \text{GL}_5$ , Hilbert's Theorem 90 tells us that  ${}^\zeta \mathbb{P}^4$  is  $k$ -isomorphic to  $\mathbb{P}^4$ , and  ${}^\zeta \Lambda_4$  is isomorphic to a quadric hypersurface in  $\mathbb{P}^4$  defined over  $k$ ; see [7, Lemma 10.1]. It is easily checked that  $\Lambda_4$  is smooth over  $k$ , and therefore so is  ${}^\zeta \Lambda_4$ . The zero-cycle of degree 3 given by  $(1 : 0 : 0 : 0 : 0) + (0 : 1 : 0 : 0 : 0) + (0 : 0 : 1 : 0 : 0)$  in  $\Lambda_4$  is  $W$ -invariant, so it defines a zero-cycle of degree 3 in  ${}^\zeta \Lambda_4$ . By Springer's theorem, the smooth quadric  ${}^\zeta \Lambda_4$  has a  $k$ -rational point, hence is  $k$ -rational.  $\square$

## Acknowledgement

We are grateful to J.-L. Colliot-Thélène for stimulating conversations.

## References

- [1] D. Anderson, Degeneracy loci and  $G_2$  flags, PhD thesis, University of Michigan, 2009, <http://deepblue.lib.umich.edu/handle/2027.42/62415>.
- [2] D. Anderson, Chern class formulas for  $G_2$  Schubert loci, *Trans. Amer. Math. Soc.* 363 (12) (2011) 6615–6646.
- [3] G. Berhuy, G. Favi, Essential dimension: A functorial point of view (after A. Merkurjev), *Doc. Math.* 8 (2003) 279–330.
- [4] C. Chevalley, Les groupes de type  $G_2$ , in: Séminaire Claude Chevalley, tome 2 (1956–1958), reprinted in: *Classification des groupes algébriques semi-simples*, Springer, 2005.
- [5] J.-L. Colliot-Thélène, B. Kunyavskii, V.L. Popov, Z. Reichstein, Is the function field of a reductive Lie algebra purely transcendental over the field of invariants for the adjoint action? *Compos. Math.* 147 (2011) 428–466.
- [6] M. Demazure, P. Gabriel, *Groupes algébriques*, tome I, Masson & Cie, Paris, 1970.

- [7] A. Duncan, Z. Reichstein, Versality of algebraic group actions and rational points on twisted varieties, *J. Alg. Geom.* (2013), in press, arXiv:1109.6093.
- [8] M. Florence, On the essential dimension of cyclic  $p$ -groups, *Invent. Math.* 171 (1) (2008) 175–189.
- [9] I.M. Gelfand, A.A. Kirillov, Sur les corps liés aux algèbres enveloppantes des algèbres de Lie, *IHES Publ. Math.* 31 (1966) 5–19.
- [10] J. Heinloth, Bounds for Behrend's conjecture on the canonical reduction, *Int. Math. Res. Not. IMRN* 14 (2008) rnn045, 17 pp.
- [11] A. Premet, Modular Lie algebras and the Gelfand–Kirillov conjecture, *Invent. Math.* 181 (2) (2010) 395–420.
- [12] J.-P. Serre, *Galois Cohomology*, Springer-Verlag, Berlin, 1997.
- [13] T.A. Springer, F.D. Veldkamp, *Octonions, Jordan Algebras, and Exceptional Groups*, Springer, 2000.