



Number theory

Plus and minus logarithms and Amice transform

*Les logarithmes plus et moins et la transformation d'Amice*

Cédric Dion, Antonio Lei

Département de mathématiques et de statistique, Université Laval, pavillon Alexandre-Vachon, 1045, avenue de la Médecine, Québec, QC, G1V 0A6, Canada

ARTICLE INFO

Article history:

Received 18 August 2017

Accepted 20 September 2017

Available online 25 September 2017

Presented by Jean-Pierre Serre

ABSTRACT

We give a new description of Pollack's plus and minus p -adic logarithms \log_p^\pm in terms of distributions. In particular, if μ_\pm denote the pre-images of \log_p^\pm under the Amice transform, we give explicit formulae for the values $\mu_\pm(a + p^n\mathbb{Z}_p)$ for all $a \in \mathbb{Z}_p$ and all integers $n \geq 1$. Our formulae imply that the distribution μ_- agrees with a distribution studied by Koblitz in 1977. Furthermore, we show that a similar description exists for Loeffler's two-variable analogues of these plus and minus logarithms.

© 2017 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

Nous donnons une nouvelle description des logarithmes p -adiques plus et moins définis par Pollack en termes de distributions. En particulier, si μ_\pm dénote la pré-image de \log_p^\pm sous la transformation d'Amice, nous donnons des formules explicites pour les valeurs $\mu_\pm(a + p^n\mathbb{Z}_p)$ pour tout $a \in \mathbb{Z}_p$ et tout entier $n \geq 1$. Nos formules impliquent que la distribution μ_- correspond à une distribution étudiée par Koblitz en 1977. Par ailleurs, nous montrons qu'il existe une description similaire, due à Loeffler, pour des analogues à deux variables de ces logarithmes plus et moins.

© 2017 Académie des sciences. Published by Elsevier Masson SAS. All rights reserved.

1. Introduction

Let p a fixed prime and E an elliptic curve with good supersingular reduction at p . Amice-Vélu [1] and Visik [12] constructed two p -adic L -functions, L_α and L_β , corresponding to the two roots (α and β) of the Hecke polynomial $X^2 - a_p X + p$. These p -adic L -functions can be regarded as distributions on the Galois group \mathbb{Z}_p^\times . When evaluated at Dirichlet characters of p -power conductor, these distributions interpolate the complex L -values of E twisted by these characters.

Unlike the good ordinary case, where the p -adic L -function can be used to formulate the Iwasawa main conjecture of E at p , the p -adic L -functions in the supersingular case are less suited for the study of Iwasawa theory. The main reason

E-mail addresses: cedric.dion.1@ulaval.ca (C. Dion), antonio.lei@mat.ulaval.ca (A. Lei).

is that they turn out to be unbounded distributions, rather than bounded measures on \mathbb{Z}_p^\times . In the case where $a_p = 0$, Pollack [11] resolved this issue by defining the plus/minus logarithms, which we denote by \log_p^\pm , and proved that there exist two bounded measures, L_+ and L_- , such that

$$L_\lambda = \log_p^+ L_+ + \lambda \log_p^- L_-, \tag{1}$$

for $\lambda \in \{\alpha, \beta\}$.

Kobayashi [7] formulated an Iwasawa main conjecture using the p -adic L -functions constructed by Pollack and showed that one inclusion of this conjecture holds true. More recently, Wan [13] has announced a proof for the other inclusion of the conjecture.

In [11], \log_p^\pm are defined to be power series given by infinite products of cyclotomic polynomials. In this article, we show that these power series may be described in terms of distributions. We define the following subsets of \mathbb{Z}_p :

$$S_n^+ := \{a \in \mathbb{Z}_p : \text{all even powers of } p \text{ vanish in the } p\text{-adic expansion of } a \text{ modulo } p^n\};$$

$$S_n^- := \{a \in \mathbb{Z}_p : \text{all odd powers of } p \text{ vanish in the } p\text{-adic expansion of } a \text{ modulo } p^n\}.$$

Our main result describes the distributions attached to \log_p^\pm in terms of these S_n^\pm .

Theorem 1.1 (Corollary 3.7). *Let μ_\pm be the distributions whose Amice transforms equal \log_p^\pm . Then, μ_+ is characterized by*

$$\mu_+(a + p^n \mathbb{Z}_p) = \begin{cases} p^{-\lfloor (n+2)/2 \rfloor} & \text{if } a \in S_n^+, \\ 0 & \text{otherwise.} \end{cases}$$

The distribution μ_- is characterized by

$$\mu_-(a + p^n \mathbb{Z}_p) = \begin{cases} p^{-\lfloor (n+3)/2 \rfloor} & \text{if } a \in S_n^-, \\ 0 & \text{otherwise.} \end{cases}$$

It is a straightforward exercise to verify that the values given in Theorem 1.1 are additive. In fact, the values of $\mu_-(a + p^n \mathbb{Z}_p)$ are precisely those given in [8, Chapter II, §3, Exercise 7] (after multiplying by p).

Since the works of Kobayashi [7] and Pollack [11], many attempts to generalize the plus/minus Iwasawa theory have been made. For example, Darmon–Iovita [3], Longo–Vigni [10], Castella–Wan [2] have studied the behaviour of an elliptic curve with supersingular reduction at p over the anticyclotomic \mathbb{Z}_p -extension of an imaginary quadratic field. Iovita–Pollack [4] have done something similar for more general \mathbb{Z}_p -extensions of a number field. In all these works, the plus and minus logarithms play an important role.

Another direction that has been taken was the study of two-variable plus/minus Iwasawa theory as introduced by Loeffler [9] and Kim [6,5]. More specifically, they studied the p -adic L -functions and Selmer groups of E over the \mathbb{Z}_p^2 -extension of an imaginary quadratic field K where p splits. This leads to the construction of four 2-variable p -adic L -functions, say $L_{\lambda, \nu}$, where $\lambda, \nu \in \{\alpha, \beta\}$. The two variables correspond to the two primes of K lying above p . A similar decomposition to (1) exists, namely,

$$L_{\lambda, \nu} = \log_p^{++} L_{++} + \lambda \log_p^{-+} L_{-+} + \nu \log_p^{+-} L_{+-} + \log_p^{--} L_{--},$$

where $\log_p^{\pm\pm}$ are some two-variable plus/minus logarithms, defined by the product of two one-variable plus/minus logarithms, and $L_{\pm\pm}$ are some bounded two-variable p -adic L -functions. We shall show that we may deduce from Theorem 1.1 a similar description of the two-variable plus/minus logarithms in terms of some explicit distributions on \mathbb{Z}_p^2 (see Corollary 4.6 below).

2. Definition and properties of Pollack’s plus/minus logarithms

In this section, we recall the definition and some basic properties of Pollack’s plus/minus logarithms from [11]. We begin by introducing some notation.

Let $\Phi_n(T) = \sum_{t=0}^{p-1} T^{p^{n-1}t}$ be the p^n -th cyclotomic polynomial. We shall also write ζ_k for a primitive p^k -th root of unity.

$$\log_p^+(T) := \frac{1}{p} \prod_{n=1}^{\infty} \frac{\Phi_{2n}(1+T)}{p},$$

$$\log_p^-(T) := \frac{1}{p} \prod_{n=1}^{\infty} \frac{\Phi_{2n-1}(1+T)}{p}.$$

Pollack showed that the products above converge and give two power series in $\mathbb{Q}_p[[T]]$. Furthermore, both \log_p^\pm converge on the open unit disc $\{x \in \mathbb{C}_p : |x|_p < 1\}$. In fact, they are related to the usual p -adic logarithm via the formula

$$\log_p^+(T) \log_p^-(T) = \frac{\log_p(1 + T)}{p^2 T},$$

hence their names. These logarithms satisfy the following interpolation formulae.

Lemma 2.1. *Let $n \geq 1$ be an integer. Then,*

$$\log_p^+(\zeta_n - 1) = \begin{cases} 0 & 2|n, \\ p^{-(n+1)/2} \prod_{j=1}^{(n-1)/2} \Phi_{2j}(\zeta_n) & 2 \nmid n, \end{cases}$$

$$\log_p^-(\zeta_n - 1) = \begin{cases} p^{-n/2-1} \prod_{j=1}^{n/2} \Phi_{2j-1}(\zeta_n) & 2|n, \\ 0 & 2 \nmid n. \end{cases}$$

Proof. This is [11, Lemma 4.7]. \square

Remark 2.2. Let $k < n$ be two odd positive integers, then

$$p^{-(k+1)/2} \prod_{j=1}^{(k-1)/2} \Phi_{2j}(\zeta_k) = p^{-(n+1)/2} \prod_{j=1}^{(n-1)/2} \Phi_{2j}(\zeta_k).$$

This is because every term with index greater than $(k - 1)/2$ on the right-hand side equals p . So, it does not change the value of the product as it gets cancelled by the extra factors of p in $p^{-(n+1)/2}$. In other words, we may write

$$\log_p^+(\zeta_k - 1) = p^{-(n+1)/2} \prod_{j=1}^{(n-1)/2} \Phi_{2j}(\zeta_k).$$

Similarly, we have

$$\log_p^-(\zeta_k - 1) = p^{-n/2-1} \prod_{j=1}^{n/2} \Phi_{2j-1}(\zeta_k)$$

if $k < n$ are two even positive integers.

Remark 2.3. It is proved in [11, Lemma 4.5] that both \log_p^\pm are $o(\log_p)$. Hence, they are uniquely determined by the interpolation formulae given in Lemma 2.1.

3. Proof of Theorem 1.1

In this section, we give a detailed proof of the main result of this article (Theorem 1.1). We begin by some preliminary lemmas.

The original definition of the plus and minus logarithms are given by infinite products of even and odd p -power cyclotomic polynomials, respectively. We shall express the truncated product of these polynomials in the form $\sum a_i x^i$.

For an integer $n \geq 1$, we define

$$R_n^+ = \left\{ \sum_{l=0}^{n-1} a_l p^{2l+1} : a_l \in \{0, 1, \dots, p-1\} \right\},$$

$$R_n^- = \left\{ \sum_{l=0}^{n-1} a_l p^{2l} : a_l \in \{0, 1, \dots, p-1\} \right\}.$$

These sets, which are related to the sets S_n^\pm given in the introduction, will play an important role in our proof of Theorem 1.1.

Lemma 3.1. *Let $n \geq 1$. We have the formula*

$$\prod_{j=1}^n \Phi_{2j}(x) = \sum_{i \in R_n^+} x^i.$$

Proof. We prove this formula by induction on n . For the base case $n = 1$, the sum on the right-hand side of the formula is simply the definition of the cyclotomic polynomial $\Phi_2(x)$.

We now suppose that the lemma holds for n and we shall show that it also holds for $n + 1$. Our inductive hypothesis implies that

$$\begin{aligned} \prod_{j=1}^{n+1} \Phi_{2j}(x) &= \Phi_{2(n+1)}(x) \prod_{j=1}^n \Phi_{2j}(x) \\ &= \left(x^{p^{2n+1}(p-1)} + \dots + x^{p^{2n+1}} + 1 \right) \sum_{i \in R_n^+} x^i \\ &= \sum_{j=0}^{p-1} \sum_{i \in R_n^+} x^{i+jp^{2n+1}}. \end{aligned}$$

But it is clear from the definition that

$$R_{n+1}^+ = \left\{ i + jp^{2n+1} : i \in R_n^+, 0 \leq j \leq p - 1 \right\}.$$

This completes the induction. \square

We have the following analogous result for the product of odd p -power cyclotomic polynomials.

Lemma 3.2. *Let $n \geq 1$ be an integer. Then,*

$$\prod_{j=1}^n \Phi_{2j-1}(x) = \sum_{i \in R_n^-} x^i.$$

Proof. The proof is analogous to that of Lemma 3.1. \square

We now recall the definition of the Amice transform.

Definition 3.3. The Amice transform of a distribution μ on \mathbb{Z}_p is defined to be

$$A_\mu(T) = \int_{\mathbb{Z}_p} (1 + T)^x \mu(x).$$

Definition 3.4. Let μ_+ be the distribution associated with \log_p^+ and let μ_- be that associated with \log_p^- .

If $z \in \mathbb{C}_p$ satisfies $|z|_p < 1$, it is immediate from the definition of μ_\pm that $\int_{\mathbb{Z}_p} z^x \mu_\pm(x) = \log_p^\pm(z - 1)$.

Proposition 3.5. *Let $n \geq 1$ be an integer and fix $a \in \mathbb{Z}_p$. The distribution μ_+ satisfies*

$$\mu_+(a + p^n \mathbb{Z}_p) = \frac{1}{p^{\lfloor (3n+2)/2 \rfloor}} \sum_{\zeta \in \mu_{p^n}} \zeta^{-a} \prod_{j=1}^{\lfloor n/2 \rfloor} \Phi_{2j}(\zeta)$$

and the distribution μ_- satisfies

$$\mu_-(a + p^n \mathbb{Z}_p) = \frac{1}{p^{\lfloor (3n+1)/2 \rfloor + 1}} \sum_{\zeta \in \mu_{p^n}} \zeta^{-a} \prod_{j=1}^{\lfloor (n+1)/2 \rfloor} \Phi_{2j-1}(\zeta).$$

Proof. We shall only give the proof for μ_+ since that for μ_- is similar. Let us write $\chi_{a+p^n \mathbb{Z}_p}$ for the characteristic function of $a + p^n \mathbb{Z}_p$. It can be decomposed as

$$\chi_{a+p^n \mathbb{Z}_p}(x) = \frac{1}{p^n} \sum_{\zeta \in \mu_{p^n}} \zeta^{x-a}.$$

By linearity, we have

$$\begin{aligned} \mu_+(a + p^n \mathbb{Z}_p) &= \int_{\mathbb{Z}_p} \chi_{a+p^n \mathbb{Z}_p}(x) \mu_+(x) \\ &= \sum_{\zeta \in \mu_{p^n}} \frac{\int_{\mathbb{Z}_p} \zeta^{x-a} \mu_+(x)}{p^n} \\ &= \frac{1}{p^n} \sum_{\zeta \in \mu_{p^n}} \zeta^{-a} \int_{\mathbb{Z}_p} \zeta^x \mu_+ \\ &= \frac{1}{p^n} \sum_{\zeta \in \mu_{p^n}} \zeta^{-a} \log_p^+(\zeta - 1) \end{aligned}$$

Suppose that n is odd. [Lemma 2.1](#) together with [Remark 2.2](#) allows us to rewrite this sum as

$$\begin{aligned} \mu_+(a + p^n \mathbb{Z}_p) &= \frac{1}{p^n} \sum_{\zeta \in \mu_{p^n}} \zeta^{-a} p^{-(n+1)/2} \prod_{j=1}^{(n-1)/2} \Phi_{2j}(\zeta) \\ &= \frac{1}{p^{(3n+1)/2}} \sum_{\zeta \in \mu_{p^n}} \zeta^{-a} \prod_{j=1}^{(n-1)/2} \Phi_{2j}(\zeta). \end{aligned}$$

When n is even, $n + 1$ is odd and every ζ in the sum above is a p^{n+1} -st root of unity. On applying [Lemma 2.1](#) and [Remark 2.2](#) with n replaced by $n + 1$, we deduce that

$$\begin{aligned} \mu_+(a + p^n \mathbb{Z}_p) &= \frac{1}{p^n} \sum_{\zeta \in \mu_{p^n}} \zeta^{-a} p^{-(n+2)/2} \prod_{j=1}^{n/2} \Phi_{2j}(\zeta) \\ &= \frac{1}{p^{(3n+2)/2}} \sum_{\zeta \in \mu_{p^n}} \zeta^{-a} \prod_{j=1}^{n/2} \Phi_{2j}(\zeta). \end{aligned}$$

This finishes the proof. \square

Corollary 3.6. *Let a and n be as given in [Proposition 3.5](#). Then,*

$$\begin{aligned} \mu_+(a + p^n \mathbb{Z}_p) &= \frac{1}{p^{\lfloor (3n+2)/2 \rfloor}} \sum_{i \in R_{\lfloor n/2 \rfloor}^+} \sum_{\zeta \in \mu_{p^n}} \zeta^{i-a}, \\ \mu_-(a + p^n \mathbb{Z}_p) &= \frac{1}{p^{\lfloor (3n+1)/2 \rfloor + 1}} \sum_{i \in R_{\lfloor (n+1)/2 \rfloor}^-} \sum_{\zeta \in \mu_{p^n}} \zeta^{i-a}. \end{aligned}$$

Proof. This follows from combining [Proposition 3.5](#) with [Lemmas 3.1](#) and [3.2](#). \square

Let us now recast the sets S_n^\pm given in the introduction as follows:

$$\begin{aligned} S_n^+ &= \{a \in \mathbb{Z}_p : \exists b \in R_{\lfloor n/2 \rfloor}^+, a \equiv b \pmod{p^n}\}, \\ S_n^- &= \{a \in \mathbb{Z}_p : \exists b \in R_{\lfloor (n+1)/2 \rfloor}^-, a \equiv b \pmod{p^n}\}. \end{aligned}$$

Corollary 3.7. *Let a and n be as given in [Proposition 3.5](#). The values of $\mu_\pm(a + p^n \mathbb{Z}_p)$ are given by*

$$\begin{aligned} \mu_+(a + p^n \mathbb{Z}_p) &= \begin{cases} p^{-\lfloor (n+2)/2 \rfloor} & \text{if } a \in S_n^+, \\ 0 & \text{otherwise,} \end{cases} \\ \mu_-(a + p^n \mathbb{Z}_p) &= \begin{cases} p^{-\lfloor (n+3)/2 \rfloor} & \text{if } a \in S_n^-, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Proof. As before, we only prove this for $\mu_+(a + p^n\mathbb{Z}_p)$. Let i be any integer. We have

$$\sum_{\zeta \in \mu_{p^n}} \zeta^{i-a} = \begin{cases} p^n & i \equiv a \pmod{p^n}, \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, if $a \in S_n^+$, the sum in [Corollary 3.6](#) simplifies to

$$\frac{1}{p^{\lfloor (3n+2)/2 \rfloor}} \times p^n.$$

Otherwise, it is 0. Hence the result. \square

Remark 3.8. Since \log_p^\pm are both $o(\log_p)$, the distributions μ_\pm are uniquely determined by the values given in [Corollary 3.7](#).

4. Generalization for two-variable logarithms

In this section, we apply our result on the one-variable plus and minus logarithms to their two-variable counterparts defined by Loeffler in [\[9\]](#).

Definition 4.1. For $*, \circ \in \{+, -\}$, we define four two-variable logarithms by using \log_p^+ and \log_p^- :

$$\log_p^{*\circ}(T_1, T_2) := \log_p^*(T_1) \cdot \log_p^\circ(T_2).$$

For $\mathbf{a} = (a, b)$ and $\mathbf{n} = (n, m)$, we shall write $\mathbf{a} + p^n\mathbb{Z}_p$ for the open set $(a + p^n\mathbb{Z}_p, b + p^m\mathbb{Z}_p)$ in \mathbb{Z}_p^2 . Furthermore, we denote the characteristic function of $\mathbf{a} + p^n\mathbb{Z}_p$ on \mathbb{Z}_p^2 by $\chi_{\mathbf{a} + p^n\mathbb{Z}_p}$.

Proposition 4.2. Let $\mathbf{a} = (a, b)$ and $\mathbf{n} = (n, m)$. The characteristic function $\chi_{\mathbf{a} + p^n\mathbb{Z}_p}$ is given by

$$\chi_{\mathbf{a} + p^n\mathbb{Z}_p}(x, y) = \frac{1}{p^{n+m}} \left(\sum_{\zeta \in \mu_{p^n}} \zeta^{x-a} \right) \left(\sum_{\zeta \in \mu_{p^m}} \zeta^{y-b} \right).$$

Proof. This is a straightforward generalization of the analogous result in the 1-dimensional case. \square

Definition 4.3. For $\mu \in \mathcal{D}(\mathbb{Z}_p^2, \mathbb{Q}_p)$, the two-dimensional Amice transform is defined by the formula

$$A_\mu(T_1, T_2) = \int_{\mathbb{Z}_p^2} (1 + T_1)^x (1 + T_2)^y \mu(x, y).$$

Remark 4.4. In the notation of [\[9\]](#), the two-variable plus/minus logarithms are in fact contained $\mathcal{D}^{(1/2, 1/2)}(\mathbb{Z}_p^2, \mathbb{Q}_p)$ under the canonical quasi-factorisation of \mathbb{Z}_p^2 . Consequently, they are uniquely determined by their values evaluated on open sets of the form $\mathbf{a} + p^n\mathbb{Z}_p$ (see [Definition 5](#) and [Theorem 3](#) of [\[9\]](#)).

It is immediate from the definition that $\log_p^{*\circ}(z - 1, w - 1) = \int_{\mathbb{Z}_p^2} z^x w^y \mu_{*\circ}(x, y)$ if z and w are elements of \mathbb{C}_p with $|z|_p, |w|_p < 1$.

Proposition 4.5. Let $\mathbf{a} = (a, b)$ and $\mathbf{n} = (n, m)$, then

$$\mu_{*\circ}(\mathbf{a} + p^n\mathbb{Z}_p) = \mu_*(a + p^n\mathbb{Z}_p) \cdot \mu_\circ(b + p^m\mathbb{Z}_p).$$

Proof. We follow the same argument as in the one-dimensional case. [Proposition 4.2](#) gives

$$\mu_{*\circ}(\mathbf{a} + p^n\mathbb{Z}_p) = \frac{1}{p^{n+m}} \int_{\mathbb{Z}_p^2} \left(\sum_{\zeta \in \mu_{p^n}} \zeta^{x-a} \right) \left(\sum_{\zeta \in \mu_{p^m}} \zeta^{y-b} \right) \mu_{*\circ}(x, y).$$

Choose ζ_n and ζ_m to be some generators of μ_{p^n} and μ_{p^m} respectively. Then,

$$\begin{aligned}
 \mu_{*o}(\mathbf{a} + p^n \mathbb{Z}_p) &= \frac{1}{p^{n+m}} \int_{\mathbb{Z}_p^2} \sum_{\substack{0 \leq i < p^n \\ 0 \leq j < p^m}} (\zeta_n^i)^{x-a} (\zeta_m^j)^{y-b} \mu_{*o}(x, y) \\
 &= \frac{1}{p^{n+m}} \sum_{\substack{0 \leq i < p^n \\ 0 \leq j < p^m}} \zeta_n^{-ai} \zeta_m^{-bj} \int_{\mathbb{Z}_p^2} (\zeta_n^i)^x (\zeta_m^j)^y \mu_{*o}(x, y) \\
 &= \frac{1}{p^{n+m}} \sum_{\substack{0 \leq i < p^n \\ 0 \leq j < p^m}} \zeta_n^{-ai} \zeta_m^{-bj} \log_p^{*o}(\zeta_n^i - 1, \zeta_m^j - 1) \\
 &= \frac{1}{p^{n+m}} \sum_{\substack{0 \leq i < p^n \\ 0 \leq j < p^m}} \zeta_n^{-ai} \log_p^*(\zeta_n^i - 1) \zeta_m^{-bj} \log_p^o(\zeta_m^j - 1) \\
 &= \frac{1}{p^{m+n}} \left(\sum_{\zeta \in \mu_{p^n}} \zeta^{-a} \log_p^*(\zeta - 1) \right) \left(\sum_{\zeta \in \mu_{p^m}} \zeta^{-b} \log_p^o(\zeta - 1) \right).
 \end{aligned}$$

Then, as in the middle of the proof of Proposition 3.5, we may rewrite this as

$$\begin{aligned}
 \mu_{*o}(\mathbf{a} + p^n \mathbb{Z}_p) &= \frac{p^n p^m}{p^{m+n}} \mu_*(a + p^n \mathbb{Z}_p) \cdot \mu_o(b + p^m \mathbb{Z}_p) \\
 &= \mu_*(a + p^n \mathbb{Z}_p) \cdot \mu_o(b + p^m \mathbb{Z}_p),
 \end{aligned}$$

as required. \square

Corollary 4.6. *The values of μ_{*o} are given by*

$$\begin{aligned}
 \mu_{++}(\mathbf{a} + p^n \mathbb{Z}_p) &= \begin{cases} p^{-\lfloor (n+2)/2 \rfloor - \lfloor (m+2)/2 \rfloor} & \text{if } a \in S_n^+ \text{ and } b \in S_m^+, \\ 0 & \text{otherwise,} \end{cases} \\
 \mu_{+-}(\mathbf{a} + p^n \mathbb{Z}_p) &= \begin{cases} p^{-\lfloor (n+2)/2 \rfloor - \lfloor (m+3)/2 \rfloor} & \text{if } a \in S_n^+ \text{ and } b \in S_m^-, \\ 0 & \text{otherwise,} \end{cases} \\
 \mu_{-+}(\mathbf{a} + p^n \mathbb{Z}_p) &= \begin{cases} p^{-\lfloor (n+3)/2 \rfloor - \lfloor (m+2)/2 \rfloor} & \text{if } a \in S_n^- \text{ and } b \in S_m^+, \\ 0 & \text{otherwise,} \end{cases} \\
 \mu_{--}(\mathbf{a} + p^n \mathbb{Z}_p) &= \begin{cases} p^{\lfloor (n+3)/2 \rfloor - \lfloor (m+3)/2 \rfloor} & \text{if } a \in S_n^- \text{ and } b \in S_m^-, \\ 0 & \text{otherwise.} \end{cases}
 \end{aligned}$$

Proof. This follows directly from combining Proposition 4.5 with Corollary 3.7. \square

Acknowledgements

The authors would like to thank Daniel Delbourgo for answering our questions during the preparation of this article. The authors’ research was supported by the NSERC Discovery Grants Program 05710 and an ISM summer scholarship.

References

[1] Y. Amice, J. Vélu, Distributions p -adiques associées aux séries de Hecke, *Astérisque* 1975 (24–25) (1975) 119–131.
 [2] F. Castella, X. Wan, Λ -adic Gross-Zagier formula for supersingular primes, arXiv:1607.02019, 2016.
 [3] H. Darmon, A. Iovita, The anticyclotomic main conjecture for elliptic curves at supersingular primes, *J. Inst. Math. Jussieu* 7 (2) (2008) 291–325.
 [4] A. Iovita, R. Pollack, Iwasawa theory of elliptic curves at supersingular primes over \mathbb{Z}_p -extensions of number fields, *J. Reine Angew. Math.* 598 (2006) 71–103.
 [5] B. Du Kim, Signed-Selmer groups over the \mathbb{Z}_p^2 -extension of an imaginary quadratic field, *Can. J. Math.* 66 (4) (2014) 826–843.
 [6] B. Du Kim, Two-variable p -adic L -functions of modular forms for non-ordinary primes, *J. Number Theory* 144 (2014) 188–218.
 [7] S.-I. Kobayashi, Iwasawa theory for elliptic curves at supersingular primes, *Invent. Math.* 152 (1) (2003) 1–36.
 [8] N. Koblitz, p -Adic Numbers, p -Adic Analysis, and Zeta-Functions, *Graduate Texts in Mathematics*, vol. 58, Springer-Verlag, New York–Heidelberg, 1977.
 [9] D. Loeffler, p -Adic integration on ray class groups and non-ordinary p -adic L -functions, in: *Iwasawa Theory 2012*, in: *Contrib. Math. Comput. Sci.*, vol. 7, Springer, Heidelberg, Germany, 2014, pp. 357–378.
 [10] M. Longo, S. Vigni, Plus/minus Heegner points and Iwasawa theory of elliptic curves at supersingular primes, preprint, arXiv:1503.07812, 2015.
 [11] R. Pollack, On the p -adic L -function of a modular form at a supersingular prime, *Duke Math. J.* 118 (3) (2003) 523–558.
 [12] M.M. Visik, Nonarchimedean measures associated with Dirichlet series, *Mat. Sb. (N.S.)* 99(141) (2) (1976) 248–260, 296, MR 0412114.
 [13] X. Wan, Iwasawa main conjecture for supersingular elliptic curves, preprint, arXiv:1411.6352, 2014.