

1-19-2023

## A Training Curriculum for an mHealth Supported Peer Counseling Program to Promote Exclusive Breastfeeding in Rural India

Roopa M. Bellad

Niranjana S. Mahantshetti

Umesh S. Charantimath

Tony Ma

Yukiko Washio

*See next page for additional authors*

Follow this and additional works at: <https://jdc.jefferson.edu/pedsfp>



Part of the [Health Information Technology Commons](#), and the [Pediatrics Commons](#)

**[Let us know how access to this document benefits you](#)**

---

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Pediatrics Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

---

**Authors**

Roopa M. Bellad, Niranjana S. Mahantshetti, Umesh S. Charantimath, Tony Ma, Yukiko Washio, Vanessa L. Short, Katie Chang, Parth Lalakia, Frances J. Jaeger, Patricia J. Kelly, Geetanjali Mungarwadi, Chandrashekar C. Karadiguddi, Shivaprasad S. Goudar, and Richard J. Derman

BRIEF REPORT

Open Access



# A training curriculum for an mHealth supported peer counseling program to promote exclusive breastfeeding in rural India

Roopa M Bellad<sup>1</sup>, Niranjana S Mahantshetti<sup>1</sup>, Umesh S Charantimath<sup>1</sup>, Tony Ma<sup>2</sup>, Yukiko Washio<sup>3</sup>, Vanessa L Short<sup>4</sup>, Katie Chang<sup>2</sup>, Parth Lalakia<sup>4</sup>, Frances J Jaeger<sup>4</sup>, Patricia J Kelly<sup>4\*</sup>, Geetanjali Mungarwadi<sup>1</sup>, Chandrashekar C Karadiguddi<sup>1</sup>, Shivaprasad S Goudar<sup>1</sup> and Richard J Derman<sup>4</sup>

## Abstract

**Background** Despite strong evidence about the benefits of exclusive breastfeeding, that is the baby receiving only breast milk, no other foods or liquids, rates have remained relatively unchanged over the past two decades in low- and middle-income countries. One strategy for increasing exclusive breastfeeding is through community-based programs that use peer counselors for education and support. The use of mobile health applications is also gaining increasing applicability in these countries. Minimal information is available about training peer counselors in the use of mobile technologies to support exclusive breastfeeding. The present article describes our curriculum in the state of Karnataka, India for supporting new mothers to exclusively breastfeed using a mobile health application in rural India.

**Methods** Twenty-five women from the community surrounding the city of Belgavi, Karnataka, India were trained to be peer counselors and to use a mobile health application to conduct a structured curriculum to support new mothers in exclusive breastfeeding. The three-day interactive training, conducted in March 2018, was based on the WHO breastfeeding course, translated, and adapted to the local culture. The curriculum, which included information collected during a formative research process, consisted of eight visits, two during the antenatal period and continuing for six months postpartum. Twelve nursing and obstetric experts validated curriculum content. Pre-post-evaluation of the training focused on breastfeeding knowledge, self-efficacy, skills, and app usability.

**Results** We observed a significant increase in the mean scores for knowledge ( $P < 0.0001$ ) and skills ( $P = 0.0006$ ) from pre- to post-training. Age of the peer counselors and their own breastfeeding experience correlated significantly with the acquisition of knowledge and skills. The mobile health app showed high usability scores.

**Conclusions** The culturally adapted curriculum presented here, combined with an mHealth app, can be an important educational strategy for training rural women in the acquisition of exclusive breastfeeding knowledge and skills.

**Keywords** mHealth application, Breastfeeding, Exclusive breastfeeding, Training curriculum, Peer counseling

\*Correspondence:

Patricia J Kelly  
patricijaneKelly@gmail.com

<sup>1</sup> KLE Academy of Higher Education and Research, Jawaharlal Nehru Medical College, Belagavi, Karnataka, India

<sup>2</sup> Benten Technologies Inc, Manassas, Virginia, USA

<sup>3</sup> Substance Use, Gender and Applied Research, RTI International, Research Triangle Park, North Carolina, USA

<sup>4</sup> Thomas Jefferson University, Philadelphia, Pennsylvania, USA



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

Exclusive breastfeeding (EBF), the practice of a baby receiving only breast milk, no other foods or liquids, significantly decreases mortality and morbidity in children less than five years of age, resulting in improved child survival and health [1, 2]. Exclusive breastfeeding can reduce infant mortality by preventing diarrhea and acute respiratory infections [3]. Increasing the rate of EBF in the first six months of life to at least 50 % is one of the six World Health Organization (WHO) global nutrition targets for 2025 [4].

Global rates of optimal breastfeeding practices, especially EBF, have remained stagnant over the past two decades in India and other low-and middle-income countries (LMICs), with only one in three infants under six months being exclusively breastfed [3]. The most recent India National Family Health Survey found that for the first six months of life, 56.4 % of infants in Karnataka, India were exclusively breastfed [5]. Achieving EBF requires a multi-pronged approach that involves both healthcare providers and policymakers, along with community participation and support [6, 7].

One strategy for increasing EBF is the use of community-based programs that use peer counselors (PCs) to educate and support mothers. Such programs have shown to be effective in increasing the initiation and duration of breastfeeding in diverse populations and settings, including LMICs. Successful PC programs have been shown to address a myriad of health-related problems, including HIV prevention, immunizations, alcohol use, and depression in several LMICs [8–18]. Effective training of PCs is a crucial prerequisite for the success of such programs.

Mobile health (mHealth) applications (apps) use mobile devices such as smartphones or handheld mobile tablets to enhance teaching, collaboration, and provision of medical care [19]. Such mHealth apps are increasingly popular to reach rural populations in LMICs with strong internet access capability [20–22], and is commonly available in India, mHealth-based programs have the potential to provide scalable public health interventions [23].

There is evidence that mHealth-supported community programs can improve the behaviors of Indian mothers, including the uptake of health services among pregnant and breastfeeding women with HIV [24]. A recent meta-analysis of studies conducted in six countries suggested that mHealth may be associated with improved maternal breastfeeding attitude, knowledge, initiation, and EBF duration [25]. Two additional reviews found evidence of positive results on EBF and other neonatal and maternal outcomes but stressed the need for both strong research methods and personalized contact [26]. It remains

unknown, though, whether utilizing mHealth platforms to enhance PC efforts is effective at improving the infant feeding behaviors of mothers in India.

As part of a funded research project, *BEST4Baby*, we adapted the WHO's breastfeeding curriculum to the local culture and trained rural mothers with prior breastfeeding experiences to serve as breastfeeding peer counselors. The training prepared them to counsel and support antepartum and postpartum mothers around the topic of EBF and infant feeding. To maximize the utility of the PCs, we designed the delivery of training content via a mHealth app on a Samsung Android tablet. The present article describes our curriculum in the state of Karnataka, India for supporting new mothers to exclusively breastfeed and documents pre-post changes of the peer counselors training curriculum.

## Development of the training curriculum

Using the results of qualitative research that we conducted through focus group discussions, we adapted the World Health Organization's breastfeeding counseling course, and Haider and colleagues' breastfeeding PCs training module to the cultural setting of the southern Karnataka state in the Kannada language [8, 27, 28]. Local breastfeeding practices, misconception and dangers of the use of top feeding were incorporated into the curriculum since the results of our formative research described mixed practices about prelacteal and supplemental feeding, reflecting older, traditional views. Names and figures in the module were based on how women looked and spoke in local communities. From the formative research, the involvement of the mother-in-law was added, as well as specifics about cultural practices such as the use of *tim tim* (herbal drops) and *guti* (locally made gruel mixtures). Research staff were experienced in the content area and are designated as national breastfeeding trainers in India; additional guidance was sought from twelve nursing and obstetric experts to develop the curriculum.

## Training curriculum

A three-day PC training was conducted in the local language at the academic health center. Two days were dedicated to breastfeeding education, skills building, and the use of *Best4Baby* kit, such as a doll for positioning. The last day was dedicated to a half-day on the use of the mHealth app, with the remaining time for protocol training and skills assessment. Pedagogy across the three days included interactive lectures, demonstrations, brainstorming sessions, case-based learning, role-playing, and continued positive feedback to enhance self-efficacy specific to breastfeeding teaching and support.

Training materials included a branded kit with the following contents to support in-home breastfeeding counseling sessions with new mothers:

- Life-size newborn doll to demonstrate positioning;
- A skin-colored sock to prepare a breast model for demonstrating proper latching;
- A digital scale for weighing and assessing the growth of the infant;
- A nipple plunger to mitigate the problem of an inverted nipple;
- A Samsung Android tablet with the *BEST4Baby* app pre-loaded with wireless service and secured to allow for its sole use with the app.

Training content covered three key components: breastfeeding knowledge, counseling and communication skills, the use of the *BEST4Baby* app, and familiarity with overall PC responsibilities for each of the nine visits in the protocol for expectant or new mothers. Each session's learning objectives, learning strategies, and evaluation process were predefined. The curriculum was designed to educate PCs for delivering timely information related to the mother's stage of gestation (e.g., antepartum at 28 -32 weeks and 32-36 weeks) and the infant's age (e.g., postpartum at 1-3 days, 15 days, 1 month, 2, 4 and 6 months) during nine home visits to provide support to the first-time mothers to practice exclusive breastfeeding. Tables 1 and 2 provide specific information about the curriculum content.

The *BEST4Baby* app was designed to reinforce training by allowing PCs to use the device during and after the training sessions and in the field. A complete description of app content with screenshots is described in the work by Ma and colleagues [29]. The app's design included a step-by-step guide for each visit to cover relevant topics for that visit. Educational videos for the PCs to share with the mothers in the app included content on early initiation of breastfeeding, breastfeeding techniques, breastfeeding position and attachment, expression of breast milk, and myths of breastfeeding. The PCs were given the opportunity to use the app, including the process of logging in, scheduling visits, and practicing each visit. Time was allotted to practice the content of the training modules incorporated in the app.

## Methods

We describe here the pre- and post-evaluation of the initial training program for peer counselors in EBF and the use of the mHealth app conducted at the Research Unit of the Jawaharalal Nehru Medical College in

Belagavi, India. The Ethics Committee of the College approved the protocol for our work. All participants signed consent forms.

## Participants

Potential peer counselors were identified by staff from five primary health centers located in Belagavi, Karnataka, India and interviewed by research staff about their breastfeeding experience. Other inclusion criteria were (1) residing in the local community; (2) having breastfed within the past five years; (3) having at least 10 years of formal education; (4) having an available mobile phone; (5) being familiar with operating an Android phone; and (6) being able to read, write, and communicate in the local language.

## Assessment instruments

We evaluated participants in the breastfeeding training curriculum on knowledge, self-efficacy, skills in counseling, and usability of the mHealth application. These skill sets were evaluated employing a 30-item survey, adapted from the literature, to reflect curriculum content, validated in advance by twelve nursing and obstetric content experts who were members of the study's Breastfeeding Advisory Panel. This panel included Karnataka MCH clinicians and academics, a Karnataka State Ministry of Health official, Reproductive and Child Health Officers, District Health Officers, and MCH advocacy group, La Leche League, and Baby Friendly Initiative representatives [30, 31]. The survey contained 20 knowledge items (e.g., colostrum is the first breast milk; breastfeeding helps in mother and child bonding); three counseling items (e.g., mother has a complaint about something that you, as a peer counselor, know little about. how do you respond?); two self-efficacy items (e.g., how confident are you in your ability to address mother's concerns about breastfeeding?) and five observational skill items (e.g., is the infant correctly positioned?). Possible scores ranged from 0 to 20 on knowledge, 0-3 on counseling, 0-8 on self-efficacy, and 0-10 on skills subscales. At the end of the training, we also evaluated the app's usability using a modified System Usability Scale, a simple, widely available, validated 10-item survey based on a 5-point Likert scale and included two self-efficacy items [32]. Modification of this scale is commonly standardized by replacing "the system" with the name of the entity being evaluated, in this case, the app name. All instruments were translated into the local language of Karnataka, back-translated for accuracy of content, and pilot tested with local women.

**Table 1** Peer counsellor training content

| Session | Day 1   | Time   |
|---------|---|--------|
| 1       | Registration; consent form signing; pre-test  | 30 min |
| 2       | Introduction<br>Importance of breastfeeding and recommended practices<br>Benefits of exclusive breastfeeding for 6 months<br>Benefits for the mother<br>Benefits for the child<br>Benefits for the family<br>Recommended practices<br>How breastfeeding works   | 1 hour |
| 3       | Local breastfeeding situation and misconception   | 1 hour |
| 4       | Importance of Early initiation of EBF and pregnancy counselling   | 30 min |
| 5       | Demonstration of correct positioning and attachment<br>Helping mothers hold the infant appropriately, with proper positioning<br>Recommended rules of correct attachment<br>Results of poor attachment<br>How to produce sufficient milk<br>How mother will understand that the infant is getting enough milk | 30 min |
| 6       | Breastfeeding video and explanation   | 45 min |
| 7       | Expression of breast milk – Demonstration and practice  | 45 min |
|         | <b>Day 2</b>  |        |
| 8       | Review of day 1 training  | 1 hour |
| 9       | Care of pregnant and lactating mothers and family planning<br>Mother's food and rest during pregnancy<br>Iron tablets<br>Check-up<br>Delivery related information<br>Family planning  | 30 min |
| 10      | Counselling skills with demonstration   | 1 hour |
| 11      | Counselling in pregnancy and role play (by the participants)  | 1 hour |
| 12      | Counselling in breastfeeding mother and role play   | 1 hour |
| 13      | Common breastfeeding difficulties and role play   | 2 hour |
| 14      | Review and clarification  | 90 min |
| 15      | Peer counsellor responsibilities and practical training sessions  | 1 hour |
|         | <b>Day 3</b>  |        |
| 16      | App training with study protocol training   | 3 hour |
| 17      | Protocol review   | 3 hour |
| 18      | Post-test   | 30 min |

**Data collection**

Data were collected from the PCs on paper forms at the start and completion of the training. Each PC was given a unique identifier to match their pre- and post-surveys.

**Data analysis**

Data were entered into SAS 9.4 statistical program. [33] (SAS Institute, 2002-2012). In addition to descriptive statistics, matched pair T-tests were conducted to assess pre-and post- training changes.

**Results/evaluation of training**

Fifty-six potential PCs were identified by staff from five primary health centers; 25 were selected who met

the inclusion criteria. The 25 PCs had an average age of 29 years (SD 4.4; range 23-40), 88 % were married for an average of 11 years; 56 % had attended at least 11-16 years of school and 76 % had some form of formal work experience. All 25 PCs completed both pre and post-test assessments. Mean scores significantly improved from pre- to post-training on the breastfeeding knowledge ( $p < 0.0001$ ), counseling ( $p = 0.0006$ ), and skills ( $p = 0.0006$ ) modules. Scores on the self-efficacy items did not change significantly. After the half-day training, scores on the Peer Counselor Usability Scale for the app were found to be highly usable, with the 25 PCs reporting an average score of 87.5 (SD  $\pm$  8.2; range 72.5-100).

**Table 2** Peer counselor training curriculum

| Day Session          | Content  | Learning Objectives   | Learning Activity  | Evaluation   |
|----------------------|--|---|--|--|
| Day 1<br>Session 1-4 | Breastfeeding knowledge  | Importance of breastfeeding and recommended practices<br>Local breastfeeding situation and misconception<br>Importance of early initiation of EBF<br>Care of pregnant and lactating mothers and family planning | Brain storming<br>Case based Interactive lectures  | Pre- and post- questionnaire                                   |
| Day 1<br>Session 4-7 | Breastfeeding skills   | Demonstration of correct positioning and attachment<br>Assessment and observation of breastfeeding<br>Expression of breast milk   | Demonstration and practice with breastfeeding model and doll<br>Breastfeeding video<br>Assessment and observation of breastfeeding in real patients using the WHO Breastfeeding observation form                   | Pre- and post- questionnaire with LATCH <sup>a</sup> criteria  |
| Day 2<br>Session 8-9 | Counseling skills during visits<br>Breastfeeding mothers and mothers with breastfeeding problems | Understand counseling skills<br>Learning and listening<br>Confidence building and helping skills<br>Using non-verbal and verbal technics to encourage mother to talk  | Demonstration<br>Interactive lecture /Brain storming<br>case scenarios/role play of different scenarios<br>Practice sessions -counseling sessions of real mothers in the postnatal ward<br>Debriefing/ reflections | Pre- and post- Questionnaire                                   |
| Day 2<br>Session 10  | Peer counseling training   | Understand responsibilities of peer counselors at each visit  | Interactive lecture<br>Role plays  |  |
| Day 3<br>Session 11  | Use of Mhealth app   | Understand usage and application of mobile app<br>Study protocol training   | Interactive lecture<br>Demonstration and Practice  | Peer Counselor usability scale for breastfeeding education app |

<sup>a</sup> LATCH= "L" = how well the infant latches onto the breast; "A" = amount of audible swallowing noted; "T" = mother's nipple type; "C" = mother's level of comfort; "H" amount of help the mother needs to hold her infant to the breast.

## Conclusions

This study described and evaluated a new training curriculum for PCs to support EBF practices with effective counseling and a mHealth app. Results demonstrated that a brief 3-day training curriculum effectively trained regular mothers to become PCs with significant increases in the acquisition of PC's knowledge, counseling, communication skills, and their ability to use the mHealth app.

Despite not having substantial prior knowledge and training in breastfeeding counseling, PCs obtained their skills through focused training, interactive pedagogy, and support to perform their counseling tasks effectively. Limited literature is available describing training curricula for PCs to promote breastfeeding within the local context. A comprehensive review of the PC training protocols and breastfeeding peer support programs revealed a wide divergence of content, type, tools, and aids, as well as the duration of the training and type of curriculum used; in several studies, few details were provided regarding the training process [34].

Effective training and support of PCs is a crucial step in ensuring the success of community-based peer support programs and is one of the proven effective strategies for increasing breastfeeding initiation and duration [35–37].

Our work highlights the need for a standardized training program and the development of training tools/aids, as well as the utilization of a mHealth app to support the implementation of a PC breastfeeding program. Similar to this study, all training must be adapted to local needs, provide up-to-date scientific information, and ensure skills development.

Peer Counselor training programs cited in earlier studies have used different modules and curricula for training (see examples in [36]). A systematic review and meta-analyses examined the effectiveness of community-based peer counseling on breastfeeding practices and found that most of the studies conducted in both developed and developing countries, employed the WHO/UNICEF breastfeeding manual/course for training [36]. Our curriculum had three key components: improving breastfeeding knowledge, counseling and communication skills, the use of a specially designed mHealth app, and elucidation of PC responsibilities during each of the nine visits to expectant or recently delivering mothers. While there exist varying modules for expanding breastfeeding knowledge and skills, we believe this to be among the first curricula which incorporate a specific mHealth app to enhance the impact of PC training on breastfeeding practices within a nine-visit breastfeeding intervention that

provides “just-in-time” information to support mothers in their breastfeeding effort.

Limitations of this work include its one site, one time use. Future applications are planned, depending on funding availability. Strengths include the fact that, to our knowledge, a mHealth app for use by PCs to support lactation counseling has not been previously reported. Breastfeeding apps have been developed for use by mothers and support networks, including fathers [21, 22, 38]. The *BEST4Baby* mHealth app was specifically designed to provide a step-by-step guide for the PCs to conduct each of the nine visits and provide relevant content at each visit. Building the content for the mHealth app and training women with limited smartphone usage experience were among the greatest challenges in implementation.

Our work highlights the value of developing a structured, effective training curriculum for PCs to implement a community-based EBF initiative. Apart from facilitating the acquisition of knowledge and skills, the training also provided the PCs with the ability (self-efficacy) to use the mHealth app effectively to support the EBF program. The curriculum recognizes the importance of a mHealth app in implementing the program and ensuring adherence to the intervention protocol. The long-term impact of such PC training on EBF practices needs further study and ideally should be evaluated in diverse settings employing a larger group of PCs. The implementation fidelity of the training curriculum requires further validation.

#### Abbreviations

|         |                                 |
|---------|---------------------------------|
| EBF     | Exclusive breastfeeding         |
| PCs     | Peer counselors                 |
| WHO     | World Health Organization       |
| LMICs   | Low-and middle-income countries |
| mHealth | Mobile health                   |
| apps    | Applications                    |

#### Acknowledgements

Not applicable

#### Authors' contributions

RB, NM, UC, GM, CK, and SG performed the research. TM, YW, KC and PL developed content for the training. VS, FJ, PK, and RD wrote the manuscript and contributed essential analyses. All authors have read and approved the final manuscript.

#### Funding

U.S. National Institute of Health, Fogarty International Center, R21TW010609, Richard Derman, Principal Investigator

#### Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

#### Declarations

##### Ethics approval and consent to participate

The Ethics Committee of the Jawaharlal Nehru Medical College from Belagavi, India approved the conduct of our work. All participants provided signed, informed consent.

#### Consent for publication

No individual data are contained in the manuscript. All authors provided consent for publication

#### Competing interests

The authors declare that they have no financial and non-financial competing interests

Received: 29 July 2022 Accepted: 12 January 2023

Published online: 19 January 2023

#### References

- Horta BL, Victora CG. Long-term effects of breastfeeding: A systematic review. Geneva (Switzerland): World Health Organization; 2014. Available from: [https://apps.who.int/iris/bitstream/handle/10665/79198/9789241505307\\_eng.pdf;jsessionid=174639532695322456BF33A46598C0CE?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/79198/9789241505307_eng.pdf;jsessionid=174639532695322456BF33A46598C0CE?sequence=1) [Cited 2020 Dec 15]
- Victora CG, Bahl R, Barros AJ, Franca G, Horton S, Frasevec J, et al. Lancet Breastfeeding Series Group. Breastfeeding in the 21st century: Epidemiology, mechanisms, and lifelong effect. *Lancet*. 2016;387(10017):475–90.
- Chandhiok N, Singh K, Sahu D, Singh L, Pandey A. Changes in exclusive breastfeeding practices and its determinants in India, 1992–2006: Analysis of national survey data. *Int Breastfeed J*. 2015;10:34. <https://doi.org/10.1186/s13006-015-0059-0>.
- World Health Organization. Global nutrition targets 2025. Geneva (Switzerland): World Health Organization; 2014. (Policy brief series: WHO/NMH/NHD/14.2). Available from: <https://www.who.int/nutrition/global-target-2025/en/>. [Cited Dec 15]
- National Family Health Survey. KEY FINDINGS FROM NFHS-5; 2019-20. Available from: [http://rchiips.org/NFHS/NFHS-5\\_FCTS/NFHS-5%20State%20Factsheet%20Compendium\\_Phase-I.pdf](http://rchiips.org/NFHS/NFHS-5_FCTS/NFHS-5%20State%20Factsheet%20Compendium_Phase-I.pdf)
- Sinha B, Chowdhury R, Sankar MJ, Martines J, Taneja S, Mazumder S, et al. Interventions to improve breastfeeding outcomes: A systematic review and meta-analysis. *Acta Paediatr*. 2015;104(467):114–34.
- Sharma IK, Byrne A. Early initiation of breastfeeding: A systematic literature review of factors and barriers in south Asia. *Int Breastfeed J*. 2016;11:17. <https://doi.org/10.1186/s13006-016-0076-7>.
- Haider R, Ashworth A, Kabir I, Huttly S. Effect of community-based peer counsellors on exclusive breastfeeding practices in Dhaka, Bangladesh: a randomised controlled trial. *Lancet*. 2000;356(9242):1643–7.
- Kushwaha KP, Singh YD, Rathi AK, Bhatia V, Gupta Y. Importance of community support in breastfeeding promotion. *J Neonatology*. 2003;18:60–1.
- Ara G, Khanam M, Papri N, Nahar B, Haque MA, Kabir I, Dibley MJ. Peer counselling improves breastfeeding practices: a cluster randomized controlled trial in urban Bangladesh. *Matern Child Nutr*. 2018;14(3):e12605. <https://doi.org/10.1111/mcn.12605>.
- Anderson AK, Damio G, Young S, Chapman DJ, Pérez-Escamilla R. A randomized trial assessing the efficacy of peer counseling on exclusive breastfeeding in a predominantly latina low-income community. *Arch Pediatr Adolesc Med*. 2005;159(9):836–41.
- Nankunda J, Tumwine JK, Soltvedt A, Semiyaga N, Ndeezi G, Tylleskär T. Community based peer counsellors for support of exclusive breastfeeding: Experiences from rural Uganda. *Int Breastfeed J*. 2006;1:19. <https://doi.org/10.1186/1746-4358-1-19>.
- Durgappa A, Raghavendra B, Parimala B, et al. Effect of peer counseling on breastfeeding practices in the rural community: Randomized control trial. *Int J Health Sci*. 2012;82(8):8–19.
- Das A, Chatterjee R, Karthick M, Mahapatra T, Chaudhuri I. The influence of seasonality and community-based health worker provided counselling on exclusive breastfeeding - findings from a cross-sectional survey in India. *PLoS One*. 2016;11(8):e0161186. <https://doi.org/10.1371/journal.pone.0161186>.
- Das A, George B, Ranebennur V, Parthasarathy MR, Shreenivas GS, Todankar P, Shrivastav A, Reddy AK, Akolo C, Cassell M, Mane S, Tripathi D, Baishya J. Getting to the first 90: Incentivized peer mobilizers promote HIV testing services to men who have sex with men using social media in Mumbai. *India Glob Health Sci Pract*. 2019;7(3):469–77.



16. Gianfredi V, Moretti M, Lopalco PL. Countering vaccine hesitancy through immunization information systems, a narrative review. *Hum Vaccin Immunother*. 2019;15(11):2508–26.
17. Nadkarni A, Weobong B, Weiss HA, et al. Counselling for alcohol problems (CAP), a lay counsellor-delivered brief psychological treatment for harmful drinking in men, in primary care in India: A randomised controlled trial. *Lancet*. 2017;389(10065):186–95.
18. Chowdhary N, Anand A, Dimidjian S, Shinde S, Weobong B, Balaji M, et al. The healthy activity program lay counsellor delivered treatment for severe depression in India: Systematic development and randomised evaluation. *Br J Psychiatry*. 2016;208(4):381–8.
19. Garavand A, Samadbeik M, Nadri H, et al. Effective factors in adoption of mobile health applications between medical sciences students using the UTAUT model. *Methods Inf Med*. 2019;58(4–05):131–9.
20. Wu Q, Huang Y, van Velthoven MH, et al. The effectiveness of using a WeChat account to improve exclusive breastfeeding in Huzhu County Qinghai Province, China: Protocol for a randomized control trial. *BMC Public Health*. 2019;19:1603. <https://doi.org/10.1186/s12889-019-7676-2>.
21. Wang CJ, Chaovalit P, Pongnumkul S. A breastfeed-promoting mobile app intervention: Usability and usefulness study. *JMIR Mhealth Uhealth*. 2018;6(1):e27. <https://doi.org/10.2196/mhealth.8337>.
22. Wheaton N, Lenehan J, Amir LH. Evaluation of a breastfeeding app in rural Australia: Prospective cohort study. *J Hum Lact*. 2018;34(4):711–20. <https://doi.org/10.1177/0890334418794181>.
23. Bassi A, John O, Praveen D, Maulik PK, Panda R, Jha V. Current status and future directions of mHealth interventions for health system strengthening in India: Systematic review. *JMIR Mhealth Uhealth*. 2018;6(10):e11440. <https://doi.org/10.2196/11440>.
24. Suryavanshi N, Kadam A, Kanade S, Gupte N, Gupta A, Bollinger R, et al. Acceptability and feasibility of a behavioral and mobile health intervention (COMBIND) shown to increase uptake of prevention of mother to child transmission (PMTCT) care in India. *BMC Public Health*. 2020;20:752. <https://doi.org/10.1186/s12889-020-08706-5>. doi:10.2196/11440.
25. Lau Y, Htun TP, Tam WS, et al. Efficacy of e-technologies in improving breastfeeding outcomes among perinatal women: A meta-analysis. *Matern Child Nutr*. 2016;12(3):381–401. <https://doi.org/10.1111/mcn.12202>.
26. Almohanna AA, Win KT, Meedy S. Effectiveness of internet-based electronic technology interventions on breastfeeding outcomes: Systematic review. *J Med Internet Res*. 2020;22(5):e17361. <https://doi.org/10.2196/17361>.
27. Charantimath U, Bellad R, Majantashetti N, Washio Y, Derman R, Kelly PJ, et al. Facilitators and challenges to exclusive breastfeeding in Belagavi district, Karnataka, India. *PLoS One*. 2020;15(5):e0231755. <https://doi.org/10.1371/journal.pone.0231755>.
28. World Health Organization/UNICEF. Breastfeeding counseling: A training course. Geneva (Switzerland): WHO; 1993. Available from: [https://www.who.int/maternal\\_child\\_adolescent/documents/who\\_cdr\\_93\\_3/en/](https://www.who.int/maternal_child_adolescent/documents/who_cdr_93_3/en/). [Cited 2020 Dec 15]
29. Ma T, Chang K, Alyusuf A, Bajracharya E, Washio Y, Kelly PJ, et al. Design, development, and testing of BEST4Baby, an mHealth technology to support exclusive breastfeeding in India: Pilot study. *MIR Form Res*. 2022;6(9):e32795. <https://doi.org/10.2196/32795>.
30. Vijayalakshmi P, Susheela T, Mythili D. Knowledge, attitudes, and breast feeding practices of postnatal mothers: A cross sectional survey. *Int J Health Sci (Qassim)*. 2015;9(4):364–74.
31. Jensen D, Wallace S, Kelsay P. LATCH: A breastfeeding charting system and documentation tool. *J Obstet Gynecol Neonatal Nurs*. 1994;23(1):27–32.
32. This is correct. Can add DOI Klug B. An Overview of the system usability scale in library website and system usability testing. *Weave*. 2017;1(6). <https://doi.org/10.3998/weave.12535642.0001.602>.
33. SAS Institute. (2002–2012). SAS/STAT 9.4 user's guide. Cary, NC: SAS Institute Inc.
34. Chapman DJ, Morel K, Anderson AK, Damio G, Pérez-Escamilla R. Breastfeeding peer counseling: From efficacy through scale-up. *J Hum Lact*. 2010;26(3):314–26.
35. Haider R, Kabir I, Huttly SR, et al. Training peer counselors to promote and support exclusive breastfeeding in Bangladesh. *J Hum Lact*. 2002;18:7–12.
36. Shakya P, Kunieda MK, Koyama M, et al. Effectiveness of community-based peer support for mothers to improve their breastfeeding practices: A systematic review and meta-analysis. *PLoS One*. 2017;12(5):e0177434. <https://doi.org/10.1371/journal.pone.0177434>.
37. Mihrshahi S, Tait H, Haider R, et al. Characteristics and experiences of peer counsellors in urban Dhaka: A structured interview study. *Int Breastfeed J*. 2019;14:48. <https://doi.org/10.1186/s13006-019-0240-y>;14:48.y.
38. White BK, Burns SK, Giglia RC, et al. Designing evaluation plans for health promotion mHealth interventions: a case study of the Milk Man mobile app. *Health Promot J Austr*. 2016;27(3):198–203.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

