

Episodic memory encoding in middle age: effects of ageing and cognitive fatigue on brain activation

Citation for published version (APA):

Klaassen, E., Evers, L., De Groot, R., Backes, W., Veltman, D., & Jolles, J. (2012). *Episodic memory encoding in middle age: effects of ageing and cognitive fatigue on brain activation*. Poster session presented at Annual meeting for the international society for neuroimaging in psychiatry, Heidelberg, Germany.

Document status and date:

Published: 02/02/2012

Document Version:

Peer reviewed version

Document license:

CC BY-NC-ND

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

<https://www.ou.nl/taverne-agreement>

Take down policy

If you believe that this document breaches copyright please contact us at:

pure-support@ou.nl

providing details and we will investigate your claim.

Downloaded from <https://research.ou.nl/> on date: 15 Feb. 2023

Open Universiteit
www.ou.nl



Episodic memory encoding in middle age: effects of aging and cognitive fatigue on brain activation

E. Klaassen^A, E. A. T. Evers^B, R. H. M. de Groot^{AC}, W. H. Backes^{A,D}, D. Veltman^E, J. Jolles^F

^A Cognitive Neuropsychiatry and Clinical Neuroscience division, Faculty of Health, Medicine and Life Science, Maastricht University, The Netherlands
^B Centre for Functional Magnetic Resonance Imaging, Department of Radiology, University of California San Diego (UCSD), San Diego, USA
^C Centre for Learning Sciences and Technologies (CELSTEC), Open University, The Netherlands
^D Department of Radiology, Maastricht University Medical Centre, The Netherlands
^E Department of Psychiatry, VU University Medical Centre, The Netherlands
^F Centre for Brain and Learning, AZIRE Research Institute and Faculty of Psychology and Education, VU University Amsterdam, The Netherlands

Background

Little is known about age-related changes in episodic memory in middle age. Less still is known about age-related differences in the context of induced cognitive fatigue. Yet insight into changes present during middle age and the effects of factors encountered during daily life, such as fatigue, are important to our understanding of the process of cognitive aging and the ecological validity of this understanding.

Introduction

Healthy cognitive aging is thought to impact most heavily on episodic memory [1]. However, changes in episodic memory prior to the age of 60 are more controversial than changes in older adults [2]. Furthermore, cognitive decline already present in middle age may not yet manifest in behavior due to the action of neural compensation processes that preserve performance at the behavioral level. Therefore, fMRI can provide valuable insights into age-related changes present in middle age [3]. It is also important to determine the extent to which middle-aged adults must compensate for the effects of cognitive aging in order to maintain performance not just in the short-term, but following sustained, fatigue inducing task performance likely, for example, to be commonly encountered during the workday. Therefore, in the current study, each participant was tested twice: once following a fatiguing condition involving the sustained performance of cognitively demanding tasks and once following a less demanding baseline condition.

Methods

Participants: 14 young (25-35) and 15 middle-aged (50-60) male teachers

Procedure: Participants completed the fatigue manipulation before entering the scanner to perform the episodic memory task.

Manipulation: Baseline condition - 1.5 hrs of low demand tasks
 Fatigue condition - 1.5 hrs of cognitively demanding tasks

Subjective fatigue measure: Profile of Mood States fatigue subscale (before and after the fatigue manipulation, and after the scanning session)

Episodic memory task: Encoding - categorisation of 100 words into 4 semantic categories

Recognition - presentation of the 100 encoding phase words plus 100 new words. Participants indicated whether words were 'old', 'maybe old', 'maybe new', or 'new'.

fMRI analysis: SPM8 was used to examine 'successful encoding' using the subsequent memory paradigm [4]: encoding task items subsequently remembered on the recognition task minus subsequently forgotten items.

Objective

We examined differences in episodic memory encoding performance and brain activation between young and middle-aged adults at baseline and in the context of induced cognitive fatigue. Furthermore, we examined the effect of induced fatigue compared to baseline within each of these age groups.

We hypothesized that fMRI would show brain activation differences between the two age groups at baseline (specifically, increased brain activation in frontal regions in middle-aged adults reflecting neural compensation), and that induced fatigue would increase the level of demand experienced on the task (leading to an exacerbation of activation differences between the two age groups, or an exhaustion of cognitive resources in the middle-aged group).

Conclusions

1. At baseline, activation was greater, primarily in PFC regions, in middle-aged compared to young adults. This suggests increased exertion of top-down cognitive control in middle-aged adults during successful encoding.
2. In the fatigue condition, activation differences between the two age groups were no longer apparent. Activation in both age groups, but particularly the middle-aged group, decreased in comparison to baseline activation. Therefore, in a state of induced fatigue, middle-aged adults no longer showed greater exertion of cognitive control than young adults, and instead showed activation changes suggestive of an exhaustion of cognitive resources.

Results

Subjective ratings: Fatigue ratings (Figure 1) increased significantly following the fatiguing manipulation (time point 1), but did not differ between age groups.




Figure 1: Subjective ratings on the Fatigue subscale (Profile of Mood States).

Task performance: Accuracy (Table 1) and RT were not significantly affected by age or the fatigue manipulation.

Table 1: Episodic memory recognition task performance. Mean proportions (SD) of each judgment type.

Judgment (confidence)	Young		Middle age	
	Baseline	Fatigue	Baseline	Fatigue
Old words				
Recognition (high)	0.59 (0.16)	0.61 (0.12)	0.50 (0.22)	0.52 (0.21)
Recognition (low)	0.15 (0.11)	0.16 (0.10)	0.18 (0.11)	0.16 (0.06)
Forgotten (low & high)	0.25 (0.09)	0.21 (0.10)	0.31 (0.15)	0.31 (0.17)
New words				
Correct rejection (high)	0.43 (0.23)	0.39 (0.22)	0.41 (0.20)	0.42 (0.20)
Correct rejection (low)	0.27 (0.13)	0.27 (0.13)	0.35 (0.19)	0.34 (0.15)
False alarm (low & high)	0.28 (0.13)	0.33 (0.22)	0.23 (0.17)	0.22 (0.16)

Task-related brain activation: Successful encoding was associated with activation in the left VLPFC, ACC, left OFC, left temporal pole, left lateral temporal cortex, and left pallidum (at $p(\text{FWE}) < 0.05$).

Baseline age group differences: Increased activation was found in the middle-aged compared to young adults in the regions shown in Table 1 and Figure 2 (at $p_{\text{uncorrected}} < .001$, masked inclusively at $p_{\text{uncorrected}} < .05$, cluster threshold > 5)

Table 2: Regions showing greater activation in middle-aged than in young adults at baseline.

Region (BA)	MNI coordinates X Y Z	T-value	Cluster Size (voxels)
Left Dorsolateral PFC (9)	-30 27 45	3.45	6
Left Dorsomedial PFC (32)	-12 42 27	3.65	22
Left OFC (11)	-24 48 -12	3.90	16
Right OFC (11)	24 54 -12	3.59	6
Left Angular gyrus (7)	-39 -72 42	3.59	5




Figure 2: Regions showing greater activation in middle-aged than in young adults at baseline.

Fatigue condition age group differences: No differences.

Effects of fatigue within the young group: Activation decreased in the fatigue condition compared to baseline in the ACC.

Effects of fatigue within the middle-aged group: Activation decreased in the fatigue condition compared to baseline in the regions shown in Table 2 and Figure 3.

Table 3: Regions showing greater activation in the baseline than fatigue condition in middle-aged adults.

Region (BA)	MNI coordinates X Y Z	T-value	Cluster Size (voxels)
Left Dorsomedial PFC (32)	-15 15 45	3.93	53
Right Anterior cingulate (32)	12 36 18	3.57	8
Left hippocampal formation (20)	-27 -24 -12	4.12	51
Right Angular gyrus (39)	51 -51 33	3.78	46
Left Precuneus	-9 -66 33	3.72	67
Right Precuneus	6 -60 33	4.01	
Right Occipital	21 -69 27	3.81	16
Left Pallidum	-21 3 3	4.26	53




Figure 3: Regions showing greater activation in the baseline than fatigue condition in middle-aged adults.

References

- 1.Reuter-Lorenz, P.A. & Park, D.C., (2010). Human neuroscience and the aging mind: at old problems a new look. *J Gerontol B-Psychol*, 65, 405-415
- 2.Salthouse, T.A. (2010). Selective review of cognitive aging. *J Int Neuropsych Soc*, 16, 754-760
- 3.Park, D.C. & Reuter-Lorenz, P.A., (2009). The adaptive brain: aging and neurocognitive scaffolding. *Annu Rev Psychol*, 60, 173-196
- 4.Paller, K.A. & Wagner, A.D., (2002). Observing the transformation of experience into memory. *Trends Cogn Sci*, 6, 93-102