

SHORT REPORT

Open Access

# Establishment of a human pluripotent stem cell-derived MKX-tdTomato reporter system



Yuki Fujisawa<sup>†</sup>, Lu Ming<sup>†</sup>, Daisuke Yamada, Tomoka Takao and Takeshi Takarada<sup>\*</sup>

## Abstract

Tendon regeneration is difficult because detailed knowledge about tendon progenitor cells (TPCs), which produce tenocytes to repair tendon tissue, has not been revealed. Mohawk homeobox (MKX) is a marker of TPCs or tenocytes, but a human pluripotent stem cell (hPSC)-based reporter system that visualizes MKX<sup>+</sup> cells has not been developed. Here, we established an hPSC-derived MKX-tdTomato reporter cell line and tested the induction ratio of MKX-tdTomato<sup>+</sup> cells using our stepwise/xeno-free differentiation protocol. MKX-tdTomato<sup>+</sup> cells were generated with high efficiency and expressed tendon-specific markers, including *MKX*, *SCX*, *TNMD*, and *COL1A1*. Our MKX-tdTomato hPSC line would be a useful tool for studying the development or regeneration of tendon tissue.

**Keywords:** Tendon, Human pluripotent stem cells, Tenocytes, Tendon regeneration

## Introduction

Tendons are fibrous connective structures composed of collagen fibers that connect muscles to bones. Tendons are easily damaged by injury, overuse, or age-related degeneration, and tendinopathy is common and hard to recover from due to the poor regenerative potential of tendons [1, 2]. An effective treatment to induce tendon regeneration is still needed to improve patients' quality of life. In recent decades, cellular therapies have been proposed as a promising approach to overcome tendon defects [3]; cell types commonly used in tendon healing include mesenchymal stem cells, tendon stem/progenitor cells, induced pluripotent stem cells (iPSCs), and embryonic stem cells (ESCs) [4–8]. Although several tenogenic differentiation protocols using pluripotent stem cells (PSCs/ESCs) have been reported, none of them describe the expansion capacity of these differentiated cells.

During embryogenesis, paraxial mesoderm is considered to differentiate into several cell types, including skeletal muscle cells, chondrocytes, osteocytes, dermal fibroblasts, and tenocytes [9]. Tenocytes are specific fibroblast cells that constitute the tendon. Some researchers have recently reported the successful differentiation of pluripotent stem cells (PSCs)-derived tenocytes in vitro [7, 10] and their promising therapeutic applications, indicating important insights into tendon regeneration. In this study, we used an induction protocol for the efficient differentiation of the paraxial mesoderm into tenocytes derived from human pluripotent stem cells (hPSCs).

Mohawk homeobox (MKX) is an essential transcription factor that is persistently expressed during tendon development and plays a crucial role in tendon maturation and maintenance [11, 12]. Here, we established an MKX-tdTomato reporter hPSC line to generate tenocytes using our tenogenic differentiation protocol.

## Materials and methods

### Cell culture

Human ESC cell line SEES4 (donated by RIKEN BRC, Japan) was cultured and maintained using StemFit (AK02N, Ajinomoto). Before reaching subconfluency,

<sup>†</sup>Yuki Fujisawa and Lu Ming contributed equally to this work

<sup>\*</sup>Correspondence: takarada@okayama-u.ac.jp

Department of Regenerative Science, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama 700-8558, Japan



the cells were dissociated with TrypLE Select (Thermo Fisher)/0.25 mM EDTA and suspended in StemFit containing 10 μM Y-27632. The cells ( $1 \times 10^4$ ) were then suspended in StemFit containing 10 μM Y27632 and 8 μl


iMatrix511 (human laminin-511 E8 fragment, Nippi) and added to a 6 cm dish. Next day, the culture media were replaced with fresh StemFit without Y-27632. After that, the media were replaced every two days until the next passage (Table 1).

**Table 1** Oligos for PX459-MKX gRNA

Purpose	Name	Sequence
MKX 3'UTR knock-in	hMKX-CRPs2	caccGGCTAATAAGCATATGGCGT
	hMKX-CRPa2	aaacACGCCATATGCTTATTAGCC

**Establishment of an MKX-tdTomato reporter cell line**

Figure 1A depicts the targeting strategy for the knock-in of the IRRS-tdTomato-PGK-Neo cassette at the MKX 3' untranslated region (UTR) to achieve MKX and tdTomato coexpression. To construct the targeting vector



(pEXA2J2-hMKX HA-IRES-tdTomato-PGK-Neo), primers listed in Table 2, were used and IRES-tdTomato-PGK-Neo fragments were inserted into the pEXA2J2-hMKX homology arm (pEXA2J2-hMKX HA; artificially synthesized by Eurofins) using an In-Fusion HD cloning kit (Takara).

Guide RNAs (gRNA) were designed to target the protospacer adjacent motif (PAM) sequence-located MKX locus (CCAACGCCATATGCTTATTAGCC; the PAM sequence is indicated in bold font). The number of potential target sites in the human genome are 1 site (20 mer + PAM), 1 site (12 mer + PAM) and 389 sites (8 mer + PAM). gRNA oligos (Table 1) were designed and subcloned into the PX459 vector (Addgene, #62988) harboring a Cas9 expression cassette (PX459-MKX gRNA). To generate the MKX-tdTomato reporter line,  $1 \times 10^6$  SEES4 hESCs were electroporated with pEXA2J2-hMKX HA-IRES-tdTomato-PGK-Neo (1  $\mu$ g) and PX459-MKX gRNA (10  $\mu$ g). Selection with G418 (Life Technologies) was performed until stable colonies appeared; then, colonies were selected for expansion. To verify the precise integration of IRES-tdTomato-PGK-Neo cassette into the MKX 3' UTR, genomic DNA was amplified using the primers in the 5' and 3' direction for PCR genotyping. The primers used are listed in Table 3.

### Tenogenic differentiation of hPSCs

The hPSCs suspension ( $3 \times 10^4$  cells in 1 ml of StemFit (Ajinomoto) containing 10  $\mu$ M Y27632 (Wako), and 4  $\mu$ l of iMatrix511 (1:250 dilution)) was seeded onto a 3.5-cm

culture dish. The culture medium was replaced the next day with fresh StemFit without Y-27632. After culturing for 2 days, the cells were washed with PBS, and differentiation was induced by changing the culture medium at each time point. A chemically defined CDM2 medium was used as the basal culture medium supplemented with cytokines and chemicals to prepare each differentiation medium. The composition of the CDM2 basal medium was as follows: 50% IMDM (+ GlutaMAX; Gibco), 50% F12 (+ GlutaMAX; Gibco), 1 mg/ml polyvinyl alcohol (Sigma-Aldrich), 1% (vol/vol) chemically defined lipid concentrate (Gibco), 450  $\mu$ M monoethioglycerol (Sigma-Aldrich), 7  $\mu$ g/ml insulin (Sigma-Aldrich), 15  $\mu$ g/ml transferrin (Sigma-Aldrich), and 1% (vol/vol) penicillin-streptomycin (Gibco). On day 0, hPSCs were differentiated into the anterior primitive streak in the CDM2 medium supplemented with 30 ng/ml Activin A (R&D), 4  $\mu$ M CHIR99021 (GSK3 $\beta$  inhibitor; Axon Medchem), 20 ng/ml FGF2 (Wako), 100 nM PIK90 (PI3K inhibitor; Millipore), and 10  $\mu$ M Y-27632 for 24 h. For paraxial mesoderm (PM) differentiation, CDM2 medium containing 1  $\mu$ M A-83-01 (ALK4/5/7 inhibitor; Tocris), 3  $\mu$ M CHIR99021, 250 nM LDN-193189 (ALK2/3 inhibitor; ReproCELL), 20 ng/ml FGF2, and 10- $\mu$ M Y-27632 was added for 24 h. On day 2, cells were cultured in a differentiation medium supplemented with 1  $\mu$ M A-83-01, 250 nM LDN-193189, 1  $\mu$ M C59 (PORCN inhibitor; Cologen Technology), 500 nM PD0325901 (MEK inhibitor; Tocris), and 10  $\mu$ M Y-27632 for 24 h. On day 3, somite cells were differentiated into sclerotome (SCL) cells after

**Table 2** Oligos for pEXA2J2-3'MKX HA-IRES-tdTomato-PGK-Neo

Purpose	Name	Sequence
MKX 3'UTR Homology arm	F1_FW for MKX HA	CCTCAAATGCCAACGGTACCGAGCTCGGATCC
	F1_RV for MKX HA	TAATAAGCATATGGCGGTCTAGACTCGAGGCGG
IRES-tdTomato-PGKNeo (In-fusion)	15 bp + IresTomato_PX459	GTTCTCTCCCTCTGGTACCGAGCTCGGATCCGCCCC
	IresTomato + 15bp_PX459	TGGTGGTGGTATCCCGGTCTAGACTCGAGGCGGCCGC
pEXA2J2- 3'MKX HA (In-fusion)	Vec_FW for MKX HA	GCCATATGCTTATTAGCCTG
	Vec_RV for MKX HA	GTTGGCATTGAGGCATAGC

**Table 3** Genotyping primers for MKX-tdTomato reporter

Purpose	Name	Sequence
Knock-in check(WT: 2130 bp, KI: 3037 bp)	MKXreporter_OUT(F)	CCCTGACATTGTGGGAGGTC
	MKXreporter_OUT(R)	ACTGGCTGCACTATTGACCC
Cloning primers for 5' boundary	out-F(1)	TGCAATTACAGAAACCCA CCA
	ires-R(1)	CACACCGGCCTTATCCAAG
Cloning primers for 3' boundary	PGK-F(2)	CCAGACTGCCTTGGGAAAAG
	out-R(2)	ACCAGAGCTCAGGCTCCAAA

two days of culture in a CDM2 medium containing 5 nM SAG 21 K (SMO agonist, R&D), 1  $\mu$ M C59, and 10  $\mu$ M Y-27632. Subsequently, for syndetome (SYN)/tenocytes induction, cells were cultivated in the CDM2 medium supplemented with 20 ng/ml FGF8 (BioLegend) and 10  $\mu$ M Y-27632 for the first three days and in the CDM2 medium supplemented with 10 ng/ml TGF  $\beta$  3 (BioLegend), 10 ng/ml BMP7 (BioLegend), and 10  $\mu$ M Y-27632 for the next 18 days.

#### Immunocytochemistry

Cultured cells were washed with PBS, fixed with 4% paraformaldehyde (PFA) for 20 min at room temperature, and incubated with blocking solution (3% normal goat serum and 0.1% Triton X-100 in PBS) for 1 h at room temperature. Next, the cells were incubated with primary antibodies (1:200 dilution) at 4 °C overnight. The secondary antibodies (1:500 dilution) were subsequently added to the cells for 1 h at room temperature. After incubation, 0.1  $\mu$ g/ml DAPI (Thermo Fisher) in PBS was used to counterstain the nuclei. The samples were then observed using a BZ-X710 fluorescence microscope (Keyence). The antibodies used are listed in Table 4.

#### RNA extraction and quantitative reverse transcription-polymerase chain reaction

RNA was extracted using an RNeasy kit (Qiagen), and complementary DNA was synthesized using M-MLV reverse transcriptase (Thermo Fisher) and random primers (Thermo Fisher). The expression of specific genes was analyzed by qPCR using an AriaMX real-time PCR system (Agilent). The cycle parameters include denaturation at 95 °C for 30 s, annealing at 62 °C for 30 s, and elongation at 72 °C for 30 s. The mRNA expression levels of each gene were normalized to  $\beta$ -Actin (*ACTB*) and quantified using the  $2^{-\Delta\Delta C_t}$  method. The primer sequences are listed in Table 5.

#### Flow cytometry


Dissociated cells were suspended in 100  $\mu$ l of 2% FBS/PBS containing 10 ng/ml DAPI. Furthermore, tdTomato expression was detected and analyzed using a CytoFLEX S flow cytometer (Beckman Coulter) and FlowJo software (FlowJo LLC), respectively.

**Table 4** Antibodies used for immunocytochemistry

Name	Company	Catalog Number	Clone	Dilution
SOX2	Cell Signaling Technology	4900	L1D6A2	1:200
CDX2	Cell Signaling Technology	12306	D11D10	1:200
SOX9	MERK	AB5535		1:200
MKX	Atlas antibodies	A83377		1:200
anti-rabbit Alexa Fluor 488 Fab2	Cell Signaling Technology	4412		1:200
anti-rabbit Alexa Fluor 647 Fab2	Cell Signaling Technology	4414		1:200
anti-mouse Alexa Fluor 647 Fab2	Cell Signaling Technology	4410		1:200

**Table 5** qRT-PCR primers

Gene	Forward primer sequence	Reverse primer sequence
<i>ACTB</i>	AGAAAATCTGGCACACACC	AGAGGCGTACAGGGATAGCA
<i>CDX2</i>	GGGCTCTCTGAGAGGCAGGT	CCTTTGCTCTGCGTTCTG
<i>SOX9</i>	AAGCTCTGGAGACTTCTGAACGA	CGCCTTGAAGATGGCGTTGG
<i>MKX</i>	CTCGCAGATGACGCTAGTGC	TGGCTGTCGAACGGTATTCTT
<i>SCX</i>	GAGAAAGTTGAGCAAGGACCG	CCAGCTCAGGTCCAAGGTG
<i>TNMD</i>	TGGCCGGAGGTACCCAAAAA	AAGTAGATGCCAGTGATCCGTTT
<i>COL1A1</i>	GACTGGTGAGACCTGCGTGT	GCCGCCATACTCGAACTGGA
<i>tdTomato</i>	CTGTTCTGGGGCATGGCA	CGGCCATGTTGTTGCTCTCG
<i>ISL1</i>	AGATTATATCAGTTGTACGGGATCA	ACACAGCGGAAACACTCGAT
<i>GATA6</i>	CCCACAACACAACCTACAGC	GCGAGACTGACGCTATGTA
<i>OTX2</i>	TGCTAGAGCAGCCCTCACTC	TGGGTTTGGAGCAGTGAACCTTA




## Results

### Establishment of an MKX-tdTomato reporter hPSC line

To visualize MKX<sup>+</sup> cells at each step of tenogenic differentiation, we utilized PX459-MKX guide RNA to recombine the targeting vector harboring IRES-tdTomato-PGK-Neo cassettes to the 3' UTR region of *MKX* in SEES4 hESCs (Fig. 1A). After electroporation of these plasmids, cells were treated with G418, and single-cell cloning was performed to establish the MKX-tdTomato reporter hPSC line (Fig. 1B). Genomic DNA was isolated from each established clone, and PCR was performed

using primers that recognize the sequence of the 5' or 3' homology arm. As shown in Fig. 1C (Original image is Additional file 1: Fig. S1), SEES4 wild type (hereafter wild type) had wild type alleles (2130 bp), but only recombinant alleles (5167 bp) were amplified in an established clone. Although off-target genome editing in possible 389 target sites (8 mer + PAM) within human genome has not been assessed in the present study, these results indicate the successful establishment of the MKX-tdTomato reporter hPSC line.



### Differentiation of SCL from hPSCs

To test the differentiation capacity of our MKX-tdTomato reporter hPSCs, we performed stepwise differentiation to induce SCL that generates syndetome (SYN) or tendon progenitor cells (TPCs). During mesoderm development, pluripotent epiblast cells differentiate into the primitive streak and paraxial mesoderm (PM), subsequently producing somites. Somites are divided into two compartments, namely dermomyotome dorsally and SCL/TPCs ventrally [13, 14] (Fig. 2A). Here, we modified a previously reported protocol [13] to induce sclerotome from hPSCs. The successful transition from wild type or MKX-tdTomato reporter hPSCs to SCL was demonstrated by immunocytochemical analysis of pluripotency (SOX2,

day 0), PM (CDX2, day 2), and SCL markers (SOX9, day5) (Fig. 2B). Quantitative reverse transcription–polymerase chain reaction (qRT-PCR) of the mRNA expression of each marker revealed similar results (Fig. 2C). These data suggest that MKX-tdTomato reporter hPSCs maintained the differentiation capacity to generate SCL cells.

### Induction of MKX<sup>+</sup> tenocytes from SCL

FGF8 signaling is required for SYN differentiation in the early phase, and BMP and TGFβ signaling pathways are involved in the development and maintenance of tendons and ligaments [15–17]. To induce MKX<sup>+</sup> tenocytes, SCL cells were treated with FGF8/Y-27632 for three days and TGFβ3/BMP7/Y-27632 for 18 days (Fig. 3A). On day 26,

MKX-tdTomato reporter-derived cells expressed tdTomato and showed spindle-shaped morphologies that resembled tenocytes (Fig. 3B). Flow cytometry revealed that almost all cells became MKX-tdTomato<sup>+</sup> (Fig. 3C). As shown in Fig. 3D, the mRNA expression levels of syndetome- and tenocyte-specific marker genes (*MKX*, *SCX*, *TNMD*, and *COL1A1*) and *tdTomato* were significantly increased on day 26 after tenogenic induction. When each lineage marker expression was compared between Day 2 and Day 26, *CDX2* was downregulated at Day 26 but the expression of other lineage markers, including *OTX2* (neuroectoderm), *ISL1* (lateral plate mesoderm) and *GATA6* (definitive endoderm) was not changed (Additional file 1: Fig. S2). Importantly, the expression signature of *tdTomato* was similar to that of *MKX*, and tdTomato fluorescence was detected only at day 26 (Additional file 1: Fig. S3). Furthermore, immunocytochemical analysis showed that our protocol induced MKX and tdTomato coexpression in MKX reporter cells, confirming that the reporter system correctly visualized MKX<sup>+</sup> cells (Fig. 3E). These results demonstrate that our tenogenic induction protocol generated hPSC-derived MKX-tdTomato<sup>+</sup> tenocytes with high efficiency.

## Conclusions

Herein, we utilized a CRISPR/Cas9-mediated homologous recombination system to establish an hPSC-derived reporter hPSC line that allows us to visualize MKX<sup>+</sup> cells by tdTomato fluorescence. Additionally, our stepwise/xeno-free induction protocol generated MKX-tdTomato<sup>+</sup> tenocytes with high efficiency, which may promote further understanding of tenocyte development or provide novel insight into hPSC-based tendon regeneration.

## Abbreviations

TPCs: Tendon progenitor cells; MKX: Mohawk homeobox; hPSCs: Human pluripotent stem cells; MSCs: Mesenchymal stem cells; TSPCs: Tendon stem/progenitor cells; iPSCs: Induced pluripotent stem cells; ESCs: Embryonic stem cells; PSCs: Pluripotent stem cells; gRNA: Guide RNA; WT: Wild type; SCL: Sclerotome; SYN: Syndetome; PM: Paraxial mesoderm; SM: Somite; SOX: Sry-related HMG box; HMG: High-mobility group; TGF: Transforming growth factor; BMP: Bone morphogenetic protein; SCX: Scleraxis; TNMD: Tenomodulin; COL: Collagen.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13287-022-03203-5>.

**Additional file 1. Supplemental Fig. S1** Full-length blot related to Fig. 1c.

## Acknowledgements

The authors thank the National Research Institute for Child Health and Development (Tokyo, Japan) for providing the hESC line SEES4.

## Author contributions

YF and LM performed the experiments, analyzed the data, and wrote the manuscript. DY and ToT discussed the data and provided critical advice. TaT supervised the project and wrote the manuscript. All authors read and approved the final manuscript.

## Funding

This work was mainly supported by Projects for Grants-in-aid for Scientific Research from the Japan Society for the Promotion of Science (Grant No. 21H02643 to TaT) and AMED (Grant No. 18bm0704024h0001 to TaT). However, these funders played no role in the study design, data collection and analysis, publication decision, or manuscript preparation.

## Availability of data and materials

All data generated and/or analyzed during this study are included in this published article and its supplementary information.

## Declarations

### Ethics approval and consent to participate

The Ethics Committee of Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, approved the experimental protocols for studies of human subjects. Written informed consent was provided by each donor.

### Consent for publication

Not applicable.

### Competing interests

The authors declare no competing interests.

Received: 24 August 2022 Accepted: 27 October 2022

Published online: 12 November 2022

## References

- Sharma P, Maffulli N. Biology of tendon injury- healing, modeling and remodeling. *J Musculoskelet Neuronal Interact*. 2006;6(2):181–90.
- Docheva D, Muller SA, Majewski M, Evans CH. Biologics for tendon repair. *Adv Drug Deliv Rev*. 2015;84:222–39.
- Chen HS, Chen YL, Harn HJ, Lin JS, Lin SZ. Stem cell therapy for tendon injury. *Cell Transplant*. 2013;22(4):677–84.
- Conrad S, Weber K, Walliser U, Geburek F, Skutella T. Stem cell therapy for tendon regeneration: current status and future directions. *Adv Exp Med Biol*. 2019;1084:61–93.
- Chamberlain CS, Clements AEB, Kink JA, Choi U, Baer GS, Halanski MA, et al. Extracellular vesicle-educated macrophages promote early Achilles tendon healing. *Stem Cells*. 2019;37(5):652–62.
- Bi Y, Ehrichiou D, Kilts TM, Inkson CA, Embree MC, Sonoyama W, et al. Identification of tendon stem/progenitor cells and the role of the extracellular matrix in their niche. *Nat Med*. 2007;13(10):1219–27.
- Komura S, Satake T, Goto A, Aoki H, Shibata H, Ito K, et al. Induced pluripotent stem cell-derived tenocyte-like cells promote the regeneration of injured tendons in mice. *Sci Rep*. 2020;10(1):3992.
- Chen X, Song XH, Yin Z, Zou XH, Wang LL, Hu H, et al. Stepwise differentiation of human embryonic stem cells promotes tendon regeneration by secreting fetal tendon matrix and differentiation factors. *Stem Cells*. 2009;27(6):1276–87.
- Nakajima T, Ikeya M. Development of pluripotent stem cell-based human tenocytes. *Dev Growth Differ*. 2021;63(1):38–46.
- Nakajima T, Shibata M, Nishio M, Nagata S, Alev C, Sakurai H, et al. Modeling human somite development and fibrodysplasia ossificans progressiva with induced pluripotent stem cells. *Development*. 2018;145(16):165431.
- Ito Y, Toriuchi N, Yoshitaka T, Ueno-Kudoh H, Sato T, Yokoyama S, et al. The Mohawk homeobox gene is a critical regulator of tendon differentiation. *Proc Natl Acad Sci U S A*. 2010;107(23):10538–42.

12. Liu H, Zhang C, Zhu S, Lu P, Zhu T, Gong X, et al. Mohawk promotes the tenogenesis of mesenchymal stem cells through activation of the TGF-beta signaling pathway. *Stem Cells*. 2015;33(2):443–55.
13. Loh KM, Chen A, Koh PW, Deng TZ, Sinha R, Tsai JM, et al. Mapping the pairwise choices leading from pluripotency to human bone, heart, and other mesoderm cell types. *Cell*. 2016;166(2):451–67.
14. Christ B, Huang R, Scaal M. Formation and differentiation of the avian sclerotome. *Anat Embryol (Berl)*. 2004;208(5):333–50.
15. Brent AE, Schweitzer R, Tabin CJ. A somitic compartment of tendon progenitors. *Cell*. 2003;113:235–48.
16. Pryce BA, Watson SS, Murchison ND, Staverosky JA, Dunker N, Schweitzer R. Recruitment and maintenance of tendon progenitors by TGFbeta signaling are essential for tendon formation. *Development*. 2009;136(8):1351–61.
17. Schwarting T, Lechler P, Struewer J, Ambrock M, Frangen TM, Ruchholtz S, et al. Bone morphogenetic protein 7 (BMP-7) influences tendon-bone integration in vitro. *PLoS ONE*. 2015;10(2):e0116833.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

