

XVI MEETING OF PLANT MOLECULAR BIOLOGY



# ABSTRACT BOOK



## PGI1-MEDIATED VASCULAR PENTOSE PHOSPHATE PATHWAY ACTIVITY DETERMINES GROWTH, PHOTOSYNTHESIS AND METABOLISM THROUGH 2-C-METHYL-D-ERYTHRITOL 4-P PATHWAY ACTION IN ARABIDOPSIS

Ángela María Sánchez-López<sup>1</sup>, **Abdellatif Bahaji**<sup>1</sup>, Samuel Gámez-Arcas<sup>1</sup>, Nuria De Diego<sup>2</sup>, Edurne Baroja-Fernández<sup>1</sup>, Francisco José Muñoz<sup>1</sup>, Goizeder Almagro<sup>1</sup>, Karel Doležal<sup>3,4</sup>, Ondřej Novák<sup>3</sup>, José María Seguí-Simarro<sup>5</sup>, Mercedes Taberero-Mendoza<sup>5</sup>, Rafael Jorge León Morcillo<sup>6</sup> and Javier Pozueta-Romero<sup>1,6</sup>.

<sup>1</sup>Instituto de Agrobiotecnología (IdAB), CSIC-Gobierno de Navarra, Mutilva, Spain, <sup>2</sup>Centre of Region Haná for Biotechnological and Agricultural Research, Czech Advanced Technology and Research Institute, Olomouc, Czech Republic, <sup>3</sup>Department of Chemical Biology, Faculty of Science, Palacký University, Olomouc, Czech Republic, <sup>4</sup>Laboratory of Growth Regulators, Faculty of Science of Palacký University and Institute of Experimental Botany of the Czech Academy of Sciences, Olomouc, Czech Republic, <sup>5</sup>COMAV - Universitat Politècnica de València, Valencia, Spain, <sup>6</sup>Institute for Mediterranean and Subtropical Horticulture "La Mayora" (IHSM), CSIC-UMA, Málaga, Spain.

Corresponding author: Abdellatif Bahaji, [abdellatif.bahaji@csic.es](mailto:abdellatif.bahaji@csic.es)



Phosphoglucose isomerase is involved in the early steps of glycolysis and regeneration of glucose-6-phosphate pools in the pentose phosphate pathway (PPP). In Arabidopsis, plastidial phosphoglucose isomerase (PGI1) is an important determinant of growth, metabolism and photosynthesis, probably due to its involvement in the synthesis of 2-C-methyl-D-erythritol 4-P (MEP)-derived hormones in root tips and vascular tissues (Bahaji *et al.*, 2015; Bahaji *et al.*, 2018). To test this hypothesis, we conducted proteomic and metabolic characterization of *PGI1*-null *pgi1-2* plants. We also characterized *pgi1-2* plants ectopically expressing *PGI1* under the control of a root tip- and vascular tissue-specific promoter. Furthermore, we characterized *pfk4/pfk5* knockout plants impaired in the early steps of plastidial glycolysis, and *pgl3-1* plants with reduced activity of the plastidial PPP enzyme 6-phosphogluconolactonase 3. The overall data obtained in this work provide strong evidence that root tip and vascular PGI1-mediated plastidial PPP determines growth, development and photosynthesis through MEP pathway action.

\_63

Bahaji A, *et al* (2015) Plastidic phosphoglucose isomerase is an important determinant of starch accumulation in mesophyll cells, growth, photosynthetic capacity, and biosynthesis of plastidic cytokinins in Arabidopsis. *PLoS One* 10: e0119641.

Bahaji A, *et al* (2018) Plastidial phosphoglucose isomerase is an important determinant of seed yield through its involvement in gibberellin-mediated reproductive development and storage reserve biosynthesis in Arabidopsis. *Plant Cell* 30: 2082–2098.

This work was supported by the Ministerio de Ciencia e Innovación (MCIN) and Agencia Estatal de Investigación (AEI) / 10.13039/501100011033/ (grants BIO2016-78747-P, PID2019-104685GB-I00) and the Ministry of Education, Youth and Sport of the Czech Republic and ERDF project entitled "Plants as a tool for sustainable global development" (No. CZ.02.1.01/0.0/0.0/16\_019/0000827).

## THE MICROBIAL VOLATILE-RESPONSIVE REDOX-SENSITIVE CYS<sup>154</sup> RESIDUE OF THE CALVIN-BENSON ENZYME FRUCTOSE-1,6-BISPHOSPHATASE 1 IS AN IMPORTANT DETERMINANT OF PHOTOSYNTHETIC ACTIVITY IN ARABIDOPSIS

Samuel Gámez-Arcas<sup>1</sup>, Francisco José Muñoz<sup>1</sup>, Antonio J. Serrato<sup>2</sup>, Ángela María Sánchez-López<sup>1</sup>, Edurne Baroja-Fernández<sup>1</sup>, Abdellatif Bahaji<sup>1</sup>, Goizeder Almagro<sup>1</sup>, Jesús Leal-López<sup>3</sup>, Rafael Jorge León Morcillo<sup>3</sup>, **Javier Pozueta-Romero**<sup>1,3</sup>.

<sup>1</sup>Instituto de Agrobiotecnología (IdAB), CSIC-Gobierno de Navarra. Iruñako etorbidea 123, 31192 Mutiloabeti, Nafarroa, Spain, <sup>2</sup>Departamento de Bioquímica, Biología Molecular y Celular de Plantas, Estación Experimental del Zaidín, Consejo Superior de Investigaciones Científicas, C/Profesor Albareda 1, 18008, Granada, Spain, <sup>3</sup>Institute for Mediterranean and Subtropical Horticulture “La Mayora” (IHSM), CSIC-UMA, Campus de Teatinos, Avda. Louis Pasteur, 49, 29010 Málaga, Spain.

Corresponding author: Javier Pozueta-Romero, [Javier.pozueta@csic.es](mailto:Javier.pozueta@csic.es)



Microorganisms emit volatile compounds (VCs) that promote plant growth and photosynthesis as well as strong developmental and metabolic changes through multiple and complex regulatory mechanisms including global reduction of the thiol redox proteome (Gámez-Arcas *et al.* 2022). Recently, we found that microbial VC treatment promotes the reduction of the Cys<sub>154</sub> residue of the redox-regulated Calvin-Benson cycle enzyme fructose-1,6-bisphosphatase 1 (cFBP1) (Ameztoy *et al.* 2019). Although highly conserved throughout land plants and algae, this residue is not located in the proposed regulatory Trx redox or catalytic domains of cFBP1. To investigate the role played by Cys<sup>154</sup> in the activity of cFBP1 and the response of plants to microbial VCs, we produced and characterized recombinantly produced wild type (WT) cFBP1 (cFBP1wt) and a mutated form of cFBP1 in which the Cys<sub>154</sub> residue has been replaced by serine (cFBP1mut). We also produced and characterized *cfbp1* plants ectopically expressing cFBP1wt and cFBP1mut under the control of the cFBP1 promoter. In native gels, the electrophoretic mobilities of recombinantly produced cFBP1 and cFBP1mut were different. In addition, recombinant cFBP1mut had ca. 85% less activity than cFBP1wt. The ectopic expression of cFBP1wt, and to a lesser extent that of cFBP1mut, countered the reduced photosynthetic activity of cFBP1-lacking *cfbp1* plants, reverting it to the WT. Results presented in this work provide strong evidence that the Cys<sub>154</sub> residue of cFBP1 is an important determinant of photosynthetic activity in Arabidopsis.

### References:

Ameztoy K, Baslam M, Sánchez-López ÁM, Muñoz FJ, Bahaji A, Almagro G, García-Gómez P, Baroja-Fernández E, De Diego N, Humplík JF, Ugena L, Spíchal L, Dolezal K, Kaneko K, Mitsui T, Cejudo FJ, Pozueta-Romero J (2019) Plant responses to fungal volatiles involve global post-translational thiol redox proteome changes that affect photosynthesis. *Plant Cell Environ* 42: 2627–2644.

Gámez-Arcas S, Baroja-Fernández E, García-Gómez P, Muñoz FJ, Almagro G, Bahaji A, Sánchez-López ÁM, Pozueta-Romero J (2022) Action mechanisms of small microbial volatile compounds in plants. *J Exp Bot.* 73: 498–510.

### Acknowledgements & Funding:

This work was supported by the Ministerio de Ciencia e Innovación (MCIN) and Agencia Estatal de Investigación (AEI) / 10.13039/501100011033/ (grants BIO2016-78747-P, PID2019-104685GB-I00 and PGC2018-096851-B-C21).

## VASCULAR AND ROOT TIP GPT2 EXPRESSION MEDIATES THE *PGI1*-INDEPENDENT RESPONSE OF ARABIDOPSIS TO SMALL MICROBIAL VOLATILES

Samuel Gámez-Arcas<sup>1</sup>, **Francisco José Muñoz**<sup>1</sup>, Adriana Ricarte-Bermejo<sup>1</sup>, Ángela María Sánchez-López<sup>1</sup>, Marouane Baslam<sup>1,2</sup>, Edurne Baroja-Fernández<sup>1</sup>, Abdellatif Bahaji<sup>1</sup>, Goizeder Almagro<sup>1</sup>, Nuria De Diego<sup>3</sup>, Karel Doležal<sup>4,5</sup>, Ondřej Novák<sup>4</sup>, Jesús Leal-López<sup>6</sup>, Rafael Jorge León Morcillo<sup>6</sup>, Araceli G. Castillo<sup>6</sup> and Javier Pozueta-Romero<sup>1,6</sup>.

<sup>1</sup>Instituto de Agrobiotecnología (IdAB), CSIC-Gobierno de Navarra, Mutilva, Spain, <sup>2</sup>Laboratory of Biochemistry, Faculty of Agriculture, Niigata University, Niigata, Japan, <sup>3</sup>Centre of Region Haná for Biotechnological and Agricultural Research, Czech Advanced Technology and Research Institute, Olomouc, Czech Republic, <sup>4</sup>Department of Chemical Biology, Faculty of Science, Palacký University, Olomouc, Czech Republic, <sup>5</sup>Laboratory of Growth Regulators, Faculty of Science of Palacký University and Institute of Experimental Botany of the Czech Academy of Sciences, Olomouc, Czech Republic, <sup>6</sup>Institute for Mediterranean and Subtropical Horticulture "La Mayora" (IHSM), CSIC-UMA, Málaga, Spain. Corresponding author: Francisco José Muñoz, [francisco.munoz@csic.es](mailto:francisco.munoz@csic.es)



Microorganisms emit a plethora of volatile compounds (VCs) that promote plant growth and photosynthesis as well as strong developmental and metabolic changes. In Arabidopsis, the plastidial isoform of phosphoglucose isomerase PGI1 mediates photosynthesis, metabolism and development, probably due to its involvement in the synthesis of isoprenoid-derived signals in vascular tissues (Bahaji *et al.*, 2015; Bahaji *et al.*, 2018). Like in wild-type (WT) plants, microbial VCs promote growth and photosynthesis as well as starch and CK accumulation in PGI1-lacking *pgi1-2* plants (Sánchez-López *et al.* 2016). A striking alteration in the transcriptome of leaves of small fungal VC-treated plants involves strong up-regulation of levels of transcripts of GPT2 (At1g61800), a gene that codes for a plastidial G6P/Pi transporter. We hypothesized that the PGI1-independent response to microbial volatile emissions involves GPT2 action. To test this hypothesis, we characterized responses of *WT*, *GPT2-null gpt2-1*, *PGI1-null pgi1-2* and *pgi1-2gpt2-1* plants to small fungal VCs. In addition, we characterized responses of *pgi1-2gpt2-1* plants expressing GPT2 under the control of a vascular tissue- and root tip-specific promoter to small fungal VCs. Results presented in this work provide evidence that, under conditions in which PGI1 activity is reduced, long-distance action of GPT2 plays an important role in the response of plants to small VCs through mechanisms involving resetting of the photosynthesis-related proteome in leaves and complex GPT2 regulation.

Bahaji A, *et al* (2015) Plastidic phosphoglucose isomerase is an important determinant of starch accumulation in mesophyll cells, growth, photosynthetic capacity, and biosynthesis of plastidic cytokinins in Arabidopsis. *PLoS One* 10: e0119641.

Bahaji A, *et al* (2018) Plastidial phosphoglucose isomerase is an important determinant of seed yield through its involvement in gibberellin-mediated reproductive development and storage reserve biosynthesis in Arabidopsis. *Plant Cell* 30: 2082–2098.

Sánchez-López ÁM, *et al* (2016) Arabidopsis responds to *Alternaria alternata* volatiles by triggering plastid phosphoglucose isomerase-independent mechanisms. *Plant Physiol* 172: 1989–2001.

This work was supported by the Ministerio de Ciencia e Innovación (MCIN) and Agencia Estatal de Investigación (AEI) / 10.13039/501100011033/ (grants BIO2016-78747-P, PID2019-104685GB-I00 and PID2019-107657RB-C22) and the Ministry of Education, Youth and Sport of the Czech Republic and ERDF project entitled "Plants as a tool for sustainable global development" (No. CZ.02.1.01/0.0/0.0/16\_019/0000827).