




Effects of Feeding Legume Silage With Differing Tannin Levels on Lactating Dairy Cattle

U. C. Hymes Fecht
U.S. Department of Agriculture

G. A. Broderick
U.S. Department of Agriculture

R. E. Muck
U.S. Department of Agriculture

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium2/45>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Effects of feeding legume silage with differing tannin levels on lactating dairy cattle

U.C. Hymes Fecht, G.A. Broderick and R.E. Muck

United States Department of Agriculture, Agricultural Research Service, U.S. Dairy Forage Research Center, 1925 Linden Drive West Madison, Wisconsin 53706-1108 U.S.A. Email: uchymesfecht@wisc.edu

Keywords: condensed tannins, birdsfoot trefoil, N utilisation, milk production

Introduction Condensed tannins (CT) bind to plant proteins in the rumen, reducing protein degradation to ammonia and increasing milk production and milk protein (e.g. Waghorn, 1987). Previous research showed that the reduced soluble non-protein nitrogen (NPN) content of red clover (*Trifolium pratense*) silage (RCS) was related to its greater N efficiency relative to lucerne (*Medicago sativa*) silage (LS) (Broderick *et al.*, 2001). Commercial cultivars of birdsfoot trefoil (*Lotus corniculatus*; BFT) contain modest levels of CT which reduce NPN formation in silage (Albrecht & Muck, 1991). The objective was to compare silages made from BFT with RCS and LS for milk production and N efficiency in lactating dairy cows.

Materials and methods Twenty-five lactating Holstein cows (5 fitted with ruminal cannulae) were randomly assigned to incomplete 5x5 Latin squares to assess effects on milk production and N utilisation. Diets contained (DM basis) 50% of LS, RCS or one of 3 BFT lines that contained low (LTBFT), normal (NTBFT) or high (HTBFT) concentrations of CT. The HTBFT and LTBFT lines were developed from the NTBFT by selecting for high and low CT by Dr Nancy Ehlke (University of Minnesota, USA). The remainder of the ration consisted of maize silage, high moisture maize and soybean meal 48% CP.

Results Characteristics of the silages and animal performance on the silages are shown in Table 1. There were differences in CP among silages: LS and LTBFT were highest, NTBFT and HTBFT intermediate, and RCS lowest ($P<0.01$). The levels of NDF were higher in RCS and LS than in the BFT silages ($P<0.01$). There were no differences in DM intake or in milk composition due to silage source ($P>0.01$). However, yield of milk and FCM was higher on NTBFT and HTBFT than LTBFT, which was higher than that on LS or RCS ($P<0.01$). Fat yield was 0.19 kg/d higher on NTBFT than on LS, with the other 3 diets being intermediate ($P<0.01$). Protein yield on all 3 BFT diets, regardless of CT level, was higher than on LS and RCS, despite the fact that the BFT diets contained about 1% less CP ($P<0.01$). Milk urea nitrogen (MUN) was lower on NTBFT and HTBFT than on LTBFT, LS and RCS ($P<0.01$). Differences in milk yield may have been confounded by the BFT diets being lower in fibre (27% NDF) than the LS and RCS diets (29% NDF) ($P<0.01$). However, these results suggest CT concentration was directly related to improved utilisation of CP in BFT silages.

Table 1 Silage characteristics, intake, milk production and milk constituents

Item	LS	RCS	LTBFT	NTBFT	HTBFT	SE	P Value
Silage NDF (% DM)	35.3 ^b	42.7 ^a	32.4 ^c	32.4 ^c	32.2 ^c	0.4	<0.01
Silage CP (% DM %)	22.0 ^a	18.1 ^c	21.7 ^a	20.4 ^b	20.1 ^b	0.3	<0.01
DMI (kg/d)	24.4	25.6	25.2	23.3	24.5	0.7	0.41
Milk (kg/d)	30.2 ^c	31.1 ^c	32.9 ^b	34.6 ^a	34.3 ^a	0.5	<0.01
3.5% FCM (kg/d)	31.4 ^d	32.6 ^{cd}	33.8 ^{bc}	36.3 ^a	35.3 ^{ab}	0.6	<0.01
Fat (kg/d)	1.13 ^c	1.17 ^{bc}	1.20 ^{bc}	1.32 ^a	1.24 ^{ab}	0.03	<0.01
Protein (kg/d)	0.94 ^b	0.96 ^b	1.04 ^a	1.09 ^a	1.07 ^a	0.02	<0.01
MUN (mg/dl)	10.8 ^a	11.0 ^a	10.8 ^a	9.3 ^b	9.2 ^b	0.3	<0.01

^{a,b,c} Means within a row with unlike superscripts differ ($P<0.05$)

Conclusions The results indicate that the condensed tannins in BFT improved N utilisation in the dairy cow with no ill effects on milk production.

References

- Albrecht, K.A. & R.E. Muck (1991). Proteolysis in ensiled forage legumes that vary in tannin concentration. *Crop Science*, 31, 464-469.
- Broderick, G.A., R.P. Walgenbach & S. Maignan (2001). Production of lactating dairy cows fed alfalfa or red clover silage at equal dry matter or crude protein contents in the diet. *Journal of Dairy Science*, 84, 1728-1737.
- Waghorn, G.C., M.J. Ulyatt, A. John, & M.T. Fisher (1987). The effect of condensed tannins on the site of digestion of amino acids and other nutrients in sheep fed on *Lotus corniculatus* L. *British Journal of Nutrition*, 57, 115-126.