

2023

Assessment of Novel Semen Evaluation Technologies and Breed Comparisons in Yearling Beef Bulls

A. R. Hartman
Kansas State University, arhartma@k-state.edu

I. E. Batey
Kansas State University, iebatey@k-state.edu

D. M. Grieger
Kansas State University, dgrieger@ksu.edu

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Beef Science Commons](#)

Recommended Citation

Hartman, A. R.; Batey, I. E.; Grieger, D. M.; and Fike, K. E. (2023) "Assessment of Novel Semen Evaluation Technologies and Breed Comparisons in Yearling Beef Bulls," *Kansas Agricultural Experiment Station Research Reports*: Vol. 9: Iss. 1. <https://doi.org/10.4148/2378-5977.8416>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2023 Kansas State University Agricultural Experiment Station and Cooperative Extension Service. Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Assessment of Novel Semen Evaluation Technologies and Breed Comparisons in Yearling Beef Bulls

Abstract

Objective: The objectives were 1) to evaluate the iSperm, when conducting breeding soundness exams (BSE) on bulls by comparing sperm motility to a technician's assessment and 2) to evaluate correlations between sperm response to reactive oxygen species (ROS) and functional sperm measurements.

Study Description: Ejaculates were collected via electroejaculation from yearling bulls as part of a BSE. All BSE were conducted by one veterinarian and ejaculates were evaluated by a single technician. Additional sperm motility analysis was conducted with the iSperm analyzer. Ejaculates meeting minimum thresholds for passing a BSE were diluted and sent overnight for flow cytometry evaluation. Data were analyzed using Pearson's correlation coefficients in SAS.

Results: Both gross and progressive motilities were significantly ($r = 0.30; 0.38; P < 0.001$) correlated to the technician's assessment of progressive motility. Percentage of live spermatozoa with positive ROS status was correlated ($r = 0.53; P < 0.001$) with percentage progressive motility. Percentage of live spermatozoa with negative ROS status was moderately correlated with percentage spermatozoa exhibiting secondary abnormalities ($r = 0.33; P = 0.02$). Percentage live spermatozoa that had disrupted acrosomes was strongly correlated with percentage live spermatozoa with negative ROS ($r = 0.66; P < 0.001$) and moderately negatively correlated with percentage live spermatozoa with positive ROS ($r = -0.31; P = 0.04$). Percentage of live spermatozoa with positive ROS status was correlated ($r = 0.58; P < 0.001$) with percentage of spermatozoa with active mitochondrial membranes. Percentage of live spermatozoa with positive ROS status was strongly correlated ($r = 0.92; P < 0.001$) with percentage of live spermatozoa with intact acrosomes.

The Bottom Line: The iSperm can be used to produce semen assessments similar to those of a trained technician and may offer a useful tool for producers to perform on-farm semen analysis. Sperm health and function continue to be related to negative ROS status.

Keywords

sperm motility, morphology, concentration

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Cover Page Footnote

We greatly appreciate Fink Beef Genetics for providing the bulls for this project, Sharon Tucker for performing the semen evaluation, Blue Rapids Veterinary for performing the BSE, and the crew at Fort Keogh Livestock & Range Research Laboratory for performing the flow cytometry analysis.

Authors

A. R. Hartman, I. E. Batey, D. M. Grieger, and K. E. Fike

Assessment of Novel Semen Evaluation Technologies and Breed Comparisons in Yearling Beef Bulls

A.R. Hartman, I.E. Batey, D.M. Grieger, and K.E. Fike

Abstract

The objective of this study was to evaluate correlations of sperm quality assessments and breed comparisons as observed during yearling beef bull breeding soundness exams (BSE). Ejaculates were collected via electroejaculation from yearling Charolais ($n = 23$) and Angus ($n = 23$) bulls as part of BSE. One veterinarian conducted BSE, and one technician conducted sperm quality assessments. Additional sperm motility analysis was conducted with the iSperm. Ejaculates meeting minimum thresholds for passing a BSE were subjected to flow cytometry. Pearson's correlation coefficients were determined, and breed comparisons were made using GLIMMIX in SAS. The iSperm analyzer gross and progressive motilities were correlated ($r = 0.30; 0.38; P < 0.001$) with the progressive motility assessed by the technician. Neither iSperm ($P = 0.26$) nor visual assessment ($P = 0.66$) of sperm motility differed between breeds. Bull breed did not influence total percentage of viable cells ($P = 0.83$), percentage of viable cells with intact acrosomes ($P = 0.83$), or percentage of live sperm cells with positive reactive oxygen species (ROS) status ($P = 0.92$). Sperm from Charolais bulls ($31.1\% \pm 3.35$) tended ($P = 0.10$) to have greater percentage of positive mitochondrial energy potential as compared with Angus bulls ($17.6\% \pm 3.35$). Percentage of live spermatozoa with negative ROS status was moderately correlated with the percentage of spermatozoa exhibiting secondary abnormalities ($r = 0.33; P = 0.02$). Percentage of live spermatozoa with disrupted acrosomes was strongly correlated ($r = 0.66; P < 0.001$) with percentage of live spermatozoa with negative ROS. Percentage of live spermatozoa with positive ROS status was correlated ($r = 0.58; P < 0.001$) with percentage of spermatozoa with active mitochondrial membranes. Sperm motility data assessed by the technician and iSperm data are positively correlated, offering producers an on-farm evaluation tool. Though the bull breed had little influence on sperm quality assessments, negative ROS status appears to impair sperm health and function.

Introduction

There is currently no definitive test that evaluates a bull's fertility. Current semen evaluation techniques include evaluation of motility and morphology, and although these are insightful tools for bull fertility, they are not definitive fertility tests and are highly subjective. In recent years many new fertility markers have been identified that provide an objective analysis of spermatozoa and an insight into identifying sub-fertile bulls. The use of flow cytometry is a method of analysis that works through excitation and

emissions spectra and aids in the detection of fertility markers. Specific colors bind to sperm based on the functional status of the individual cells. In recent years development of assays that target components known to be essential to bull fertility has advanced our knowledge of sperm functional statuses (Bucher et al., 2019). Reactive oxygen species are endogenous, highly reactive, oxygen- and nitrogen-bearing molecules that can be found throughout the body (Krumova and Gonzalo, 2016). In semen, ROS affects spermatozoa characteristics, including mitochondrial membrane potential, acrosomal integrity, and structural abnormalities that can influence spermatozoa function. In bulls, the presence of ROS in semen has been shown to have a direct impact on the function of spermatozoa as well as a relationship to bull fertility (Kumaresan et al., 2017; Leite et al., 2022).

Use of on-farm technologies for semen evaluation in cattle is limited; however, a product called the iSperm offers hope that these technologies may become more accessible. The iSperm works through the camera on an iPad Mini and is a relatively easy-to-use, affordable, and portable semen analysis device. This device has been validated in equines and canines but not in the bovine (Moraes et al., 2019; Dini et al., 2019; Domain et al., 2022).

Bull breed influences bull fertility. Barth and Waldner (2002) found that Angus bulls were more likely to pass a BSE than Charolais bulls, and Brito et al. (2002) found differences in motility and ejaculate concentrations among *Bos taurus* bulls when compared to *Bos indicus* but proposed no suggested explanation for these differences. Others have shown that breed influences motility, morphology, concentration, and volume of ejaculates (Hartman, 2021). As the need for a better understanding of bull fertility grows, the influence of breed has been largely understudied.

Experimental Procedures

Ejaculates were collected via electroejaculation on one of three consecutive days from Angus and Charolais yearling bulls (403 ± 11 days of age; $n = 46$) as part of a BSE. One veterinarian conducted all BSE, and ejaculates were evaluated by one technician. Ejaculates were diluted in BoviFree to accomplish a 1:5 dilution based on manufacturer recommendations, and an additional sperm motility and concentration analysis was conducted with the iSperm analyzer. Ejaculates meeting minimum thresholds for passing a BSE were diluted to 70 million cells/mL using BoviFree and sent overnight for flow cytometry evaluation. Flow cytometry assays included acrosome and cell membrane integrity, mitochondrial energy potential, and oxidation status.

Data for analysis comparisons were assessed using Pearson's correlation coefficients in SAS. The GLIMMIX procedure of SAS with bull as experimental unit, bull breed as the main effect, and collection date as a random variable was used to assess potential differences in sperm quality variables between breeds.

Results and Discussion

Percentage of live spermatozoa with positive ROS status was correlated ($r = 0.53$; $P < 0.001$) with percentage progressive motility (Table 1). Percentage of live spermatozoa with negative ROS status was moderately correlated with percentage spermatozoa exhibiting secondary abnormalities ($r = 0.33$; $P = 0.02$) and tended to be lowly correlated ($r = 0.28$; $P = 0.06$) with percentage spermatozoa exhibiting primary abnor-

malities. Percentage of live spermatozoa that had disrupted acrosomes was strongly correlated ($r = 0.66$; $P < 0.001$) with percentage live spermatozoa with negative ROS and moderately negatively correlated ($r = -0.31$; $P = 0.04$) with percentage live spermatozoa with positive ROS. These results for the relationship between ROS and acrosome integrity are similar to those observed by Kumaresan et al. (2017). The percentage of live spermatozoa with positive ROS status was correlated ($r = 0.58$; $P < 0.001$) with the percentage of spermatozoa with active mitochondrial membranes. Leite et al. (2022) found that when there were increased levels of ROS and impaired mitochondrial membranes, this often resulted in lower fertility in bulls, which supports our findings of this relationship. Live spermatozoa with positive ROS were strongly correlated ($P < 0.001$) with live spermatozoa ($r = 0.94$) and live spermatozoa with intact acrosomes ($r = 0.92$). Bucher et al. (2019) found that evaluation of the viability, acrosomal status, and mitochondrial function of cryopreserved bovine sperm could be predictive of sperm functional status. Thus, our data confirm previous research showing detrimental effects of ROS on spermatozoa function.

Both gross and progressive motilities were significantly correlated ($r = 0.30$; 0.38 ; $P < 0.001$) to the technician's assessment of progressive motility. These results are similar to previous research comparing assessments by technicians to the iSperm when evaluating stallion semen (Morales et al., 2019; Dini et al., 2019). Our results are comparable to previous results when validating the iSperm for canine use (Domain et al., 2022).

Neither iSperm ($P = 0.26$) nor visual assessment ($P = 0.66$) of sperm motility differed among breeds (Table 2). Bull breed did not influence the total percentage of viable cells ($P = 0.83$) or viable cells with intact acrosomes ($P = 0.83$). When evaluating oxidation status by measuring reactive oxygen species, the bull breed did not influence ($P = 0.92$) the percentage of live sperm cells with positive reactive oxygen species status. There was a tendency ($P = 0.10$) for a greater percentage of sperm from Charolais bulls ($31.1\% \pm 3.35$) to have positive mitochondrial energy potential as compared with Angus bulls ($17.6\% \pm 3.35$). These results differ from those found by Barth and Waldner (2002), who saw differences in motility in relation to BSE between Angus and Charolais bulls. In our study, the bull breed appears to have little influence on sperm quality assessments among yearling bulls meeting the threshold requirements for passing BSE.

Implications

Technician and iSperm assessment of sperm motility data are positively correlated, offering producers an on-farm evaluation tool. Though the bull breed has little influence on sperm quality assessments, negative ROS status in sperm appears to impair sperm health and function.

Acknowledgments

We greatly appreciate Fink Beef Genetics for providing the bulls for this project, Sharon Tucker for performing the semen evaluation, Blue Rapids Veterinary for performing the BSE, and the crew at Fort Keogh Livestock & Range Research Laboratory for performing the flow cytometry analysis.

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. Persons using such products assume responsibility for their use in accordance with current label directions of the manufacturer.

References

- Barth, A. D., and C. L. Waldner. 2002. Factors affecting breeding soundness classification of beef bulls examined at the Western College of Veterinary Medicine. *Can. Vet. J.* 43(4):274-284.
- Brito, L. F. C., A. E. D. F. Silva, L. H. Rodrigues, F. V. Vieira, L. A. G. Deragon, and J. P. Kastelic. 2002. Effects of environmental factors, age and genotype on sperm production and semen quality in *Bos indicus* and *Bos taurus* AI bulls in Brazil. *Anim. Reprod. Sci.* 70(3-4):181-190. doi:10.1016/S0378-4320(02)00009-X.
- Bucher, K., E. Malama, M. Siuda, F. Janett, and H. Bollwein. 2019. Multicolor flow cytometric analysis of cryopreserved bovine sperm: a tool for the evaluation of bull fertility. *J. Dairy Sci.* 102(12):11652-11669.
- Dini, P., L. Troch, I. Lemahieu, P. Deblende, and P. Daels. 2019. Validation of a portable device (iSperm®) for the assessment of stallion sperm motility and concentration. *Repro. Dom. Anim.* 54(8):1113-1120.
- Domain, G., P. Banchi, H. Ali Hassan, A. Eilers, J. Lannoo, E. Wydooghe, W. Nizański, and A. Van Soom. 2022. Sperm gone smart: A portable device (iSperm®) to assess semen concentration and motility in dogs. *Animal* 12(5):652.
- Hartman, A. R. 2021. The influence of environmental and managerial factors on the characteristics of beef bull semen. Master's thesis, Kansas State University, Manhattan, KS.
- Krumova, K. and C. Gonzalo. 2016. Chapter 1: Overview of Reactive Oxygen Species. In: *Singlet Oxygen: Applications in Biosciences and Nanosciences* 1:1-21.
- Kumaresan, A., A. Johannisson, E. M. Al-Essawe, and J. M. Morrell. 2017. Sperm viability, reactive oxygen species, and DNA fragmentation index combined can discriminate between above-and below-average fertility bulls. *J. Dairy Sci.* 100(7):5824-5836.
- Leite, R. F., J. D. de Agostini Losano, G. K. V. Kawai, B. R. Rui, K. K. Nagai, V. C. Castiglioni, A. F. P. Siqueira, M.E.O.D.A. Assumpção, P. S. Baruselli, and M. Nichi. 2022. Sperm function and oxidative status: Effect on fertility in *Bos taurus* and *Bos indicus* bulls when semen is used for fixed-time artificial insemination. *Anim. Reprod. Sci.* 237:106922.
- Moraes, C. R., E. E. Runcan, B. Blawut, M. A. Coutinho da Silva. 2019. Technical Note: The use of iSperm technology for on-farm measurement of equine sperm motility and concentration. *Trans. Anim. Sci.* 3(4):1513–1520.

Table 1. Pearson's correlation coefficients of sperm attributes from ejaculates collected following breeding soundness exams

| Item | % Live negative ROS spermatozoa ¹ | % Live positive ROS spermatozoa ² |
|---|---|---|
| | r (P-value) | |
| % Primary abnormalities ³ | 0.28 (0.06) | -0.15 (0.33) |
| % Secondary abnormalities ⁴ | 0.33 (0.02) | -0.23 (0.12) |
| % Progressive motility ⁵ | -0.27 (0.10) | 0.53 (<0.001) |
| % Live with intact acrosome ⁶ | -0.16 (0.29) | 0.92 (<0.001) |
| % Live with disrupted acrosome ⁷ | 0.66 (<0.001) | -0.31 (0.04) |
| % Live ⁸ | -0.19 (0.22) | 0.94 (<0.001) |
| % Polarized ⁹ | 0.03 (0.84) | 0.58 (<0.001) |

¹Percentages of spermatozoa from ejaculate with an intact cell membrane that have negative reactive oxygen species (ROS).

²Percentages of spermatozoa from ejaculate with an intact cell membrane that have positive ROS.

³Percentage of spermatozoa from ejaculate exhibiting primary abnormalities.

⁴Percentage of spermatozoa from ejaculate exhibiting secondary abnormalities.

⁵Percentages of spermatozoa from ejaculate that are progressively motile.

⁶Percentage of spermatozoa from ejaculate with an intact cell membrane and acrosome.

⁷Percentage of spermatozoa from ejaculate with an intact cell membrane and disrupted acrosome.

⁸Percentage of spermatozoa from ejaculate with an intact cell membrane.

⁹Percentage of spermatozoa from ejaculate with polarized mitochondrial membranes.

Table 2. Sperm quality assessments using visual analysis and flow cytometry on ejaculates from Angus and Charolais breeds of yearling bulls meeting BSE threshold requirements

| Factor | Least squares mean ± Standard error of mean | | |
|--|--|---------------------|----------------------|
| | Angus n = 23 | Charolais n = 23 | P-value of factor |
| Bull age, days | 402.9 ± 2.36 | 403.3 ± 2.36 | 0.90 |
| Semen characteristic | | | |
| Technician progressive motility, ¹ % | 43.7% ± 1.69 | 47.39 ± 1.69 | 0.26 |
| iSperm progressive motility, ² % | 50.1 ± 2.26 | 47.8 ± 3.18 | 0.66 |
| iSperm gross motility, ² % | 71.6 ± 2.78 | 70.5 ± 2.78 | 0.82 |
| Cells live and viable, ³ % | 42.3 ± 3.95 | 43.6 ± 3.95 | 0.83 |
| Cells live with intact acrosome, ⁴ % | 41.5 ± 3.40 | 42.6 ± 3.40 | 0.83 |
| Cells viable with positive reactive oxygen species, ⁵ % | 29.1 ± 3.52 | 28.5 ± 3.52 | 0.92 |
| Active mitochondrial potential, ⁶ % | 17.6 ± 3.35 | 31.1 ± 3.35 | 0.10 |

¹Ejaculate gross motility was analyzed by a single veterinarian as a part of a breeding soundness exam.

²Progressive and gross motility of each ejaculate were analyzed using the iSperm software and manufacturer recommendations.

³Percentages of live and viable cells were determined by flow cytometry using the Invitrogen Live/Dead sperm viability kit.

⁴Percentages of live cells with intact acrosomes and sperm membrane integrity were determined by flow cytometry using the IMV Technologies acrosome and sperm membrane integrity assay.

⁵Percentages of viable cells with a positive reactive oxygen species were determined by flow cytometry using the IMV Technologies Easy Kit 3: Oxidation molecule D assay.

⁶Percentages of spermatozoa with active mitochondrial potential were determined by flow cytometry using IMV Technologies Easy Kit 2: Mitochondrial activity assay.