



UWS Academic Portal

The impact of job-demand-control-support on leptin and ghrelin as biomarkers of stress in emergency healthcare workers

Bouillon-Minois, Jean Baptiste; Outrey, Justin; Pereira, Bruno; Adeyemi, Oluwaseun John; Sapin, Vincent; Bouvier, Damien; Thivel, David; de Saint-Vincent, Sarah; Ugbolue, Ukadike Chris; Baker, Julien S.; Bagheri, Reza; Schmidt, Jeannot; Trousselard, Marion; Dutheil, Frédéric

Published in:
Nutrients

DOI:
[10.3390/nu14235009](https://doi.org/10.3390/nu14235009)

Published: 25/11/2022

Document Version
Publisher's PDF, also known as Version of record

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Bouillon-Minois, J. B., Outrey, J., Pereira, B., Adeyemi, O. J., Sapin, V., Bouvier, D., Thivel, D., de Saint-Vincent, S., Ugbolue, U. C., Baker, J. S., Bagheri, R., Schmidt, J., Trousselard, M., & Dutheil, F. (2022). The impact of job-demand-control-support on leptin and ghrelin as biomarkers of stress in emergency healthcare workers. *Nutrients*, 14(23), [5009]. <https://doi.org/10.3390/nu14235009>

General rights















Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Article

The Impact of Job-Demand-Control-Support on Leptin and Ghrelin as Biomarkers of Stress in Emergency Healthcare Workers

Jean-Baptiste Bouillon-Minois ^{1,*} , Justin Outrey ² , Bruno Pereira ³ , Oluwaseun John Adeyemi ⁴ , Vincent Sapin ⁵ , Damien Bouvier ⁵ , David Thivel ⁶ , Sarah de Saint-Vincent ⁷ , Ukadike Chris Ugbole ⁸ , Julien S. Baker ⁹ , Reza Bagheri ¹⁰ , Jeannot Schmidt ¹ , Marion Trousselard ¹¹  and Frédéric Dutheil ¹² 

- ¹ Emergency Department, CHU Clermont-Ferrand, Université Clermont Auvergne, CNRS, LaPSCo, Physiological and Psychosocial Stress, F-63000 Clermont-Ferrand, France
 - ² Emergency Department, CHU de Besançon, F-25000 Besançon, France
 - ³ Clinical Research and Innovation Direction, CHU Clermont-Ferrand, F-63000 Clermont-Ferrand, France
 - ⁴ Ronald O. Perelman Department of Emergency Medicine, NYU School of Medicine, New York University Langone Health, New York, NY 10016, USA
 - ⁵ Centre de Biologie, CHU Clermont-Ferrand, F-63003 Clermont-Ferrand, France
 - ⁶ Research Center in Human Nutrition, Laboratory AME2P, Université Clermont Auvergne, F-63120 Aubière, France
 - ⁷ Occupational and Environmental Medicine, Université Clermont Auvergne, F-63000 Clermont-Ferrand, France
 - ⁸ School of Health and Life Sciences, Institute for Clinical Exercise & Health Science, University of the West of Scotland, Glasgow G1 1XW, UK
 - ⁹ Centre for Health and Exercise Science Research, Department of Sport, Physical Education and Health, Hong Kong Baptist University, Kowloon Tong 999077, Hong Kong
 - ¹⁰ Department of Exercise Physiology, University of Isfahan, Isfahan 81746-73441, Iran
 - ¹¹ Neurophysiology of Stress, French Armed Forces Biomedical Research Institute, IRBA, F-91223 Brétigny-sur-Orge, France
 - ¹² CHU Clermont-Ferrand, Université Clermont Auvergne, CNRS, LaPSCo, Physiological and Psychosocial Stress, Occupational and Environmental Medicine, WittyFit, F-63000 Clermont-Ferrand, France
- * Correspondence: jbbouillon-minois@chu-clermontferrand.fr; Tel.: +33-4-73-75-19-99; Fax: +33-4-73-27-46-49



Citation: Bouillon-Minois, J.-B.; Outrey, J.; Pereira, B.; Adeyemi, O.J.; Sapin, V.; Bouvier, D.; Thivel, D.; de Saint-Vincent, S.; Ugbole, U.C.; Baker, J.S.; et al. The Impact of Job-Demand-Control-Support on Leptin and Ghrelin as Biomarkers of Stress in Emergency Healthcare Workers. *Nutrients* **2022**, *14*, 5009. <https://doi.org/10.3390/nu14235009>

Academic Editor: Carlos Diéguez González

Received: 20 October 2022
Accepted: 22 November 2022
Published: 25 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Despite the available literature on the consequences of night shiftwork on stress and food intake, its impact on leptin and ghrelin has never been studied. We previously demonstrated that leptin and ghrelin were biomarkers related to stress, and acute stress-induced a decrease in leptin levels and an increase in ghrelin levels. We performed a prospective observational study to assess the influence of night work, nutrition, and stress on the levels of ghrelin and leptin among emergency healthcare workers (HCWs). We took salivary samples at the beginning of a day shift and/or at the end of a night shift. We also monitored stress using the job demand-control-support model of Karasek. We recorded 24-h food intake during the day shift and the consecutive night shift and during night work and the day before. We included 161 emergency HCWs. Emergency HCWs had a tendency for decreased levels of leptin following the night shift compared to before the dayshift ($p = 0.067$). Furthermore, the main factors explaining the decrease in leptin levels were an increase in job-demand (coefficient -54.1 , 95 CI -99.0 to -0.92) and a decrease in job control (-24.9 , -49.5 to -0.29). Despite no significant changes in ghrelin levels between shifts, social support was the main factor explaining the increase in ghrelin (6.12, 0.74 to 11.5). Food intake (kcal) also had a negative impact on leptin levels, in addition to age. Ghrelin levels also decreased with body mass index, while age had the opposite effect. In conclusion, we confirmed that ghrelin and leptin as biomarkers of stress were directly linked to the job demand-control-support model of Karasek, when the main cofounders were considered.

Keywords: nutrients; work; well-being; quality of life; leptin; ghrelin

1. Introduction

Stress at work is a major public health concern. This is especially true among healthcare workers (HCWs) and even more concerning in the Emergency Departments (ED) due to shifting work, fatigue and lack of sleep [1], poor food intake [2], and life-threatening emergencies in the context of overcrowding and job demands [3,4]. Currently, the best scale to assess stress levels at work is the job-demand-control-support model (JDCS), a self-reported psychological questionnaire created and validated by Karasek in 1981 [5,6]. Night work, defined as “all work which is performed during a period of not less than seven consecutive hours, including the interval from midnight to 5 a.m.” [7], has many health consequences. These include disturbance of circadian rhythm [8], obesity, and cardiometabolic disorders [9–11] which are putative consequences of eating disorders [12] and stress [13]. Eating behavior can be influenced by sociodemographic criteria such as age and gender [14] and by occupational characteristics such as experience [15] and workload [16,17]. Night work also induces a conflict between socially determined diurnal mealtimes, eating habits, the biological rhythm of hunger, satiety, and metabolism [18,19]. The main hormones that control the physiology of appetite are ghrelin and leptin. Ghrelin leads the orexigenic role [20]. Increased ghrelin levels influence meal initiation and food-seeking behavior [21]. In response to a meal, levels vary quickly and intensely due to its short half-life [22,23]. Leptin, a peptide mainly produced in white adipose tissue, is commonly known as a satiety hormone, with low levels observed during meal initiation and high levels after the meal [24]. Its baseline blood levels are influenced by body mass index (BMI), mainly due to the white adipose tissue component [25]. Although the classical hormones to assess stress are cortisol, epinephrine, interleukin and melatonin, both ghrelin and leptin are influenced by stress and can be considered biomarkers of stress [26–31]. Indeed, we previously performed a meta-analysis on the impact of both physical and psychological stress on ghrelin and leptin levels, measured using both blood or saliva. We found that ghrelin has a short-term increase following acute stress with a higher and prolonged response among obese individuals [27]. Regarding leptin, we found a decrease following acute stress with a higher variation of leptin levels after stress among normal-weight individuals and women [26]. Emergency HCWs are a perfect example for studying the impact of night shift and stress on biomarkers of nutrition [32]. Indeed, HCWs work under stressful conditions such as overcrowding—lack of beds in hospitals [3], life-threatening emergencies [33], the wait for possible disasters [34] all having consequences on the biomarkers of stress [31,35]. Furthermore, the night shift has a bad influence on water consumption and food intake among emergency HCWs, in both quantity and quality [2]. To date, no studies have assessed the influence of night work and stress on the levels of nutrition biomarkers among emergency HCWs, nor in relation to their occupational characteristics.

Therefore, we performed this study to assess the influence of night work, stress, and sociodemographic characteristics on ghrelin and leptin levels among emergency HCWs.

2. Materials and Methods

2.1. Study Design

We performed a prospective nationwide observational study in five French hospitals—two university hospitals and three non-university hospitals. The main inclusion criterion was to work as an emergency HCW. Exclusion criteria were refusal to participate and pregnancy. This study was part of the SEEK protocol [2,32,34]. We obtained ethical approval from the French Ethics Committee South-East I with reference DC-2014-2151, and the protocol was registered on ClinicalTrials.gov, number NCT02401607. The study was performed during two shifts (day shift and night shift). During the day shift, participants started to work between 7:30 and 8:30 a.m. and finished between 6:30 and 7:30 p.m. For nightshift, they began between 6:30 and 7:30 p.m. and finished between 7:30 and 8:30 a.m. the following day. All participants volunteered and signed consent forms before answering questionnaires and providing samples. They performed the study for one dayshift and/or

one nightshift. Salivary sampling was collected at the beginning of a dayshift and at the end of a nightshift, i.e., between 7:00 a.m. and 8:30 a.m. for both conditions. Furthermore, food intake was monitored twice over 24 consecutive hours: (1) during a day shift (from 8:30 a.m. to 6:30 p.m.) + the night after (no work, 6:30 p.m. to 8.30 a.m.), and (2) during a rest day before (from 8:30 a.m. to 6:30 p.m.) and a night shift (6:30 p.m. to 8:30 a.m.) [2]. Participants also had to complete a questionnaire collecting their level of stress using the JDCS model of Karasek [6] and sociodemographic data (Figure 1).




Figure 1. Study design. Among five French Emergency Departments, we were able to recruit 161 emergency healthcare workers for a total of 185 salivary samples. Twenty-four were excluded because of a low quantity of saliva. Ghrelin and leptin levels were assessed from 161 samples.

2.2. Outcomes

Ghrelin and leptin levels were measured from saliva samples. We used saliva preferentially because its collection is non-invasive and results are acceptable [36]. Saliva samples were stored at -80°C at the Institute of Occupational Medicine of Clermont-Ferrand. Total ghrelin was assessed by ELISA using commercial kits (CEA991Hu Cloud Clone Corp[®] (CLOUD-CLONE CORP. (CCC, Los Angeles, CA, USA)) with a detection range from 12.35 to 1000 pg/mL, a sensitivity level of $<4.87\text{pg/mL}$, an intra-assay coefficient $<10\%$ and an inter-assay coefficient $<12\%$. Leptin was assessed using ALPCO, Salem, NH, USA/22-LEPHUU-e01/kit ultrasensitive with a range from 0.05 to 5 ng/mL, a sensitivity level at 0.01 ng/mL, an intra-assay coefficient at 7.2% and an inter-assay coefficient at 4.35%. Karasek's job-content

questionnaire (JCQ) is composed of three dimensions: psychological demands, decision latitude, and social support. Participants were assessed by the 26 items of the JCQ (nine for both decision latitude and psychological demand and eight for social support). Each participant was asked to answer using a 4-level Likert-type scale for each item, ranging from 1 (strongly disagree) to 4 (strongly agree). Decision latitude was calculated using the following formula: $4 \times Q4 + 4 \times (5 - Q6) + 4 \times Q8 + 2 \times (5 - Q2) + 2 \times Q5 + 2 \times Q7 + 2 \times Q1 + 2 \times Q3 + 2 \times Q9$. A score less than 71 reflects low decision latitude. Psychological demand was calculated with the following formula: $Q10 + Q11 + Q12 + (5 - Q13) + Q14 + Q15 + Q16 + Q17 + Q18$. A score below 20 reflected a low psychological demand. Social support was calculated with the following formula: $Q19 + Q20 + Q21 + Q22 + Q23 + Q24 + Q25 + Q26$. A score below 24 reflected low social support [6]. Participants were also asked to complete a 24-h dietary recall that was explained and detailed to them by a member of the investigation team. The participants were asked to indicate as precisely as possible all the details regarding the food ingested during each meal and in-between meals. The diaries were reviewed with the participants during a dedicated interview. We next used Nutrilog[®] (Nutrilog, version 3.2), a diet and nutrition software for health care professionals, to translate participants' answers on nutrient intake using a nutrition table Ciqua based on this software. We calculated total energy intake (kcal), carbohydrates, lipids, proteins (grams and % energy intake), and water consumption in milliliters (mL) [2]. Participants also had to complete a questionnaire about their sociodemographics (age, gender, marital status, kids at home, tobacco) and work situation (occupation, seniority).

2.3. Statistics

Continuous variables were expressed as mean and standard deviation (SD), or median [interquartile range] (IQR), and categorical variables were expressed as numbers (percentage—%). The assumption of Gaussian distribution was analyzed by the Shapiro-Wilk test. As data could be repeated for a part of participants' habits (measures for nightshift and dayshift), the linear mixed method was performed (i) to compare values between nightshift and dayshift (i.e., to assess the influence of night work on ghrelin and leptin levels) and then (ii) to assess the influence on ghrelin and leptin levels on stress (using job demand, job control, and social support), nutrition (food intake in kcal, carbohydrates, lipids and proteins in grams), water consumption and sociodemographic data (job, age, BMI). Time (nightshift versus dayshift) was considered as a fixed effect, whereas participants were considered as random effects in order to model between and within-subject variability. The normality of residuals was analyzed with the Shapiro-Wilk test and a visual plot. Accordingly, ghrelin and leptin levels were log-transformed. The results were expressed as coefficient regressions and 95 confidence intervals (95 CI). To increase readability, results were expressed using the following formulas: $1000 \times \log(\text{leptin})$ or $100 \times \log(\text{ghrelin})$. Statistical analyses were performed with Stata software (v17, College Station, TX, USA). Significance was set at the $p < 0.05$ level. Sensitivity analysis was conducted among emergency HCWs that engaged in the study on both the dayshift and nightshift. Both were compared for their main characteristics to evaluate the sample representativeness.

3. Results

3.1. Characteristics of the Population

We enrolled 161 emergency HCWs from five hospitals and obtained a total of 185 samples of saliva. Twenty-four samples were excluded due to the insufficiency of saliva. In total, we analyzed 161 saliva samples. Twenty-four (14.9%) performed the study twice (one during the day shift and one during the night shift) (Figure 1). The representativeness of the emergency HCWs was examined. We did not find any differences between the HCWs that performed the study twice versus those who performed only one shift. Fifty-seven HCWs were physicians, 79 were nurses, and 24 had other occupations (administrative, maid). Participants were 37.4 ± 10.4 years old, mainly female (57.5%, sex ratio = 1.37), with

a mean BMI of 23.2 ± 4.4 kg/m². Fifty-three (32.9%) were single, and 77 (58.3%) had no children. Median seniority as a HCWs was 6 [IQR 13] years, 3 [9] years in the ED (Table 1).

Table 1. Sociodemographic. SD = standard deviation, BMI = body mass index, kg = kilograms, kg/m² = kilogram per square meter, IQR = interquartile range, *n* = number, % = percentage. Results are expressed as mean \pm standard deviation or median [interquartile range] or number (percentage).

	Total <i>n</i> = 161	Physician <i>n</i> = 57	Paramedics <i>n</i> = 79	Other <i>n</i> = 24
Age, years, mean \pm SD	37.4 \pm 10.4	35.7 \pm 10.4	39.5 \pm 10.2	34.9 \pm 10.3
Sex, <i>n</i> (%) male	68 (42.5)	29 (50.9)	28 (35.4)	11 (45.8)
BMI, kg/m ² , mean \pm SD	23.2 \pm 4.4	22.9 \pm 4.3	23.2 \pm 3.6	23.7 \pm 4.4
Underweight, <i>n</i> (%)	10 (6.2)	3 (5.3)	6 (7.6)	1 (4.2)
Normal weight, <i>n</i> (%)	107 (66.5)	39 (68.4)	53 (67.1)	15 (62.5)
Overweight, <i>n</i> (%)	36 (22.4)	14 (24.6)	17 (21.5)	5 (20.8)
Obesity class 1, <i>n</i> (%)	4 (2.5)	0	1 (1.3)	3 (12.5)
Obesity class 2, <i>n</i> (%)	1 (0.6)	0	1 (1.3)	0
Obesity class 3, <i>n</i> (%)	3 (1.9)	1 (1.8)	1 (1.3)	0
Seniority, years, median [IQR]				
In the job	6 [13]	4 [8.5]	11 [13]	4.5 [8.5]
In the department	3 [9]	2 [4.5]	5 [9]	1 [6]
Physical activity, h/week, median [IQR]	2 [4]	2 [5]	2 [4]	2 [3]
Tea/coffee, cup/day, median [IQR]	3 [3]	4 [2]	6 [12]	5 [3.5]
Smoker, <i>n</i> (%)	57 (35.6)	17 (29.8)	31 (39.7)	9 (37.5)
Cig/day for smokers, median [IQR]	6 [9]	8 [5]	6 [12]	5 [3.5]
Family situation				
Married-engaged, <i>n</i> (%)	108 (67.1)	35 (61.4)	63 (79.7)	9 (37.5)
Single-divorced, <i>n</i> (%)	53 (32.9)	22 (38.6)	16 (20.2)	15 (62.5)
Kids at home, <i>n</i> (%)	55 (41.7)	16 (33.3)	35 (53.0)	4 (22.2)

3.2. Assessment of Ghrelin and Leptin Levels

We did not find any statistically significant change on the level of both ghrelin and leptin after the end of the night shift compared to the beginning of the day shift (Figure 2).

3.3. Impact of Stress, Night Shift, Sociodemographic, and Nutrition on Leptin and Ghrelin Levels

Regarding leptin levels (expressed as $1000 \times \log$ leptin)—we found a negative impact of job demand on leptin levels (Coefficient -54.1 per 10-unit; 95 CI -99.0 to -0.92) while job control increased leptin levels (24.9 per 10-unit; 0.29 to 49.5). Age increases parallelly with leptin level (1.93 per year; 0.26 to 3.61). We also found a negative tendency for night shift (-40.2 ; -83.2 to 2.82), BMI (-5.36 per kg/m²; -10.8 to 0.09) and food intake (-2.39 per 100 kcal/24 h; -4.83 to 0.58) while water consumption tended to increase leptin levels (2.94 per dL/24 h; -2.58 to 6.13). Regarding ghrelin (expressed as $100 \times \log$ ghrelin), we found a positive impact of social support (6.12 per unit; 0.74 to 11.5), and age (2.34 per year; 0.46 to 4.25) and a negative impact for BMI (-8.31 per kg/m²; -14.0 to -2.60) (Figure 3).




Figure 2. Median [interquartile range] salivary levels of leptin and ghrelin before the beginning of a day shift and at the end of a night shift. The results of leptin are expressed in nanograms per deciliter. Results of ghrelin are expressed in picogram per milliliter. Ng = nanogram, pg = picogram, dL = deciliter, mL = milliliter, NS = not significant, n = number of samples studied, SD = standard deviation, IQR = interquartile range.




Figure 3. Impact of stress assessed using Karasek jobdemand-control-support, night, sociodemographic status and nutrition on leptin and ghrelin salivary levels using a linear mixed model. Job demand refers to the psychological needs imposed by daily working activities, job control refers to the latitude of decision and is composed of two components: skill discretion and decision authority. Social support represents the support from colleagues and/or from the hierarchy. Results were expressed using coefficient regression (95% confidence interval). Results were considered significant for *p*-value < 0.05 and 95 CI not containing 0. BMI = body mass index, kg = kilogram, m² = square meter, g = gram, dL = deciliter.

4. Discussion

We demonstrated that both stress, nightshift, and food intake have an impact on nutrition biomarkers represented by ghrelin and leptin.

4.1. *The Impact of Night Shift and Stress*

We found a lower level of both ghrelin and leptin after the night shift compared to before the beginning of the day shift. Considering that these hormones act inversely, this result needs to be discussed. Ghrelin is considered the main orexigenic hormone. A high level of ghrelin promotes food intake, while a low level is observed at the end of a meal. Literature on the impact of night work on ghrelin levels is not strong and is controversial. Indeed, one study did not find any difference in 24 h ghrelin levels recorded between day and night shifts [37], while another study found a lower ghrelin level among night shift workers compared to dayshift workers [38]. However, all studies observed that ghrelin levels are unstable (in a range of minutes) to be considered using mean levels. Conversely, leptin is the main satiety hormone, with low levels observed during meal initiation and high levels after the meal. However, the impact of night work on leptin levels seems to be negative. Indeed, it seems that average 24 h leptin levels decreased up to 40% during the night shift compared to baseline in two studies that experimentally created night and day shifts [37,39]. Furthermore, this impact seems to increase for participants that have a non-synchronized schedule [40]. However, a further study noted different results, i.e., an increase in leptin levels among hospital nurses among night workers that worked exclusively during the night for at least one year [41]. Regarding the impact of stress on ghrelin, we did not find any significant impact on job demand and job control. However, it seems that social support increases ghrelin levels (coeff 6.12; 95 CI 0.74 to 11.5). Regarding leptin, job demand decreased leptin levels while job control increased levels. This confirms the influence of stress on leptin levels and, more widely, the impact of stress on biomarker responses. We previously published that both ghrelin and leptin, measured using both blood or saliva, are influenced by stress. More precisely, ghrelin increases very quickly once the stress (physical, psychological of both) is induced—within less than five minutes—and returns to baseline levels within 45 min with a higher and prolonged response among obese individuals. Leptin decreases more slowly, with a significant decrease after 60 min, with a higher variation of leptin levels after stress among normal-weight individuals and women [26,27].

4.2. *Impact of Food Intake and Sociodemographic Status*

We found a negative impact of kcal per day intake on leptin levels with no impact on ghrelin levels. This result is not logical when initially considered. Indeed, leptin is a satiety hormone, so we would expect that greater amounts of food would induce higher levels of leptin. Furthermore, we also found a negative impact of BMI (coef -0.005 ; 95 CI -0.011 to 0.0001). Indeed, overweight individuals have a higher circulating leptin level and leptin resistance due to the availability of highly palatable food that increases triglyceride production and, therefore, white adipose tissue production of leptin [42,43]. A possible explanation for this could be the negative impact of sleep deprivation. Indeed, several studies found lower leptin levels among healthy volunteers with acute sleep deprivation or chronic partial sleep deprivation [44–46]. Very interestingly, we did not find any impact of qualitative nutrients, i.e., carbohydrates, lipids, or proteins, on biomarker levels. Furthermore, water consumption seems to increase leptin levels, although we did not find an explanation for why in the literature. A possible answer is that participants declared drinking soft drinks, not water, a high-concentrated fructose drink, is linked to obesity [47]. Lastly, we found a higher level of leptin among older participants, which is consistent with the literature [48]. Regarding ghrelin, we did not find significant covariates. This could be explained by fluctuations in homeostasis. We only found a negative correlation between BMI and ghrelin level, which is consistent with the literature [49]. However, we found a positive impact of age on ghrelin levels, which is controversial in the literature [50–52].

No other data was significant regarding food intake, water consumption, stress, or quality of food.

4.3. Limitations

Our study has some limitations, none of which affects our results. First, we performed only one measurement, but both ghrelin and leptin have a circadian rhythm [53]. This rhythm can be disturbed by night-shift. Furthermore, the two sessions have different periods. The day-shift is four-hours shorter than the night-shift. However, every sample was collected and studied under the same conditions. Second, total ghrelin was assayed rather than its active, acylated form. Nevertheless, most of the physiology of ghrelin has been defined based on the assay of total ghrelin. Thirdly, even if we tried to monitor the time of breakfast, data were too imprecise to be used. Indeed, HCWs eat their breakfast before work during the day shift, but it is not consistent with those working on the night shift. Considering that ghrelin levels decrease quickly and intensely after the beginning of food intake and increase slowly until lunch, the putative results could be more significant. Some devices are now able to continuously monitor certain biomarkers. Advances in technology should soon facilitate the circadian rhythm of leptin and ghrelin in emergency HCWs, making it feasible to study the underlying mechanisms of stress regulation.

5. Conclusions

In conclusion, we did not find any difference between night and day regarding ghrelin and leptin levels. However, we found that ghrelin and leptin can be considered as biomarkers of stress directly linked to the job demand-control-support model of Karasek, in a study that considered the main cofounders. Understanding mechanisms of stress in a comprehensive study may help to build efficient health promotion and/or preventive strategies.

Author Contributions: Conceptualization, F.D. and J.S.; methodology, F.D. and B.P.; validation, All authors; formal analysis, J.-B.B.-M., S.d.S.-V. and F.D.; investigation, F.D. and J.S.; writing—original draft preparation, J.-B.B.-M.; writing—review and editing, J.O., O.J.A., V.S., D.B., D.T., U.C.U., J.S.B., R.B. and M.T.; supervision, F.D. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: The study was conducted in accordance with the Declaration of Helsinki, and approved by the French Ethics Committee South-East I with reference DC-2014-2151, and the protocol was registered on ClinicalTrials.gov number NCT02401607.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: Data are available by request to the corresponding author.

Acknowledgments: To all the Emergency HealthCare Workers that participated in this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Dutheil, F.; Danini, B.; Bagheri, R.; Fantini, M.L.; Pereira, B.; Moustafa, F.; Trousselard, M.; Navel, V. Effects of a Short Daytime Nap on the Cognitive Performance: A Systematic Review and Meta-Analysis. *Int. J. Environ. Res. Public Health* **2021**, *18*, 10212. [[CrossRef](#)]
2. Bouillon-Minois, J.-B.; Thivel, D.; Croizier, C.; Ajebo, É.; Cambier, S.; Boudet, G.; Adeyemi, O.J.; Ugbole, U.C.; Bagheri, R.; Vallet, G.T.; et al. The Negative Impact of Night Shifts on Diet in Emergency Healthcare Workers. *Nutrients* **2022**, *14*, 829. [[CrossRef](#)] [[PubMed](#)]
3. Bouillon-Minois, J.-B.; Raconnat, J.; Clinchamps, M.; Schmidt, J.; Dutheil, F. Emergency Department and Overcrowding During COVID-19 Outbreak; a Letter to Editor. *Arch. Acad. Emerg. Med.* **2021**, *9*, e28. [[CrossRef](#)] [[PubMed](#)]
4. Bouillon-Minois, J.-B.; Schmidt, J.; Dutheil, F. SARS-CoV-2 Pandemic and Emergency Medicine: The Worst Is yet to Come. *Am. J. Emerg. Med.* **2020**, *42*, 246–247. [[CrossRef](#)]
5. Karasek, R.A. Job Demands, Job Decision Latitude, and Mental Strain: Implications for Job Redesign. *Adm. Sci. Q* **1979**, *24*, 285–308. [[CrossRef](#)]

6. Karasek, R.A. *Job Decision Latitude, Job Design, and Coronary Heart Disease*; Machine pacing and occupational stress; Taylor and Francis Ltd.: London, UK, 1981.
7. Convention C171—Night Work Convention, 1990 (No. 171). Available online: https://www.ilo.org/dyn/normlex/en/f?p=NORMLEXPUB:12100:0::NO::P12100_INSTRUMENT_ID:312316 (accessed on 4 November 2021).
8. Boivin, D.B.; Boudreau, P. Impacts of Shift Work on Sleep and Circadian Rhythms. *Pathol. Biol.* **2014**, *62*, 292–301. [[CrossRef](#)]
9. Ganesan, S.; Magee, M.; Stone, J.E.; Mulhall, M.D.; Collins, A.; Howard, M.E.; Lockley, S.W.; Rajaratnam, S.M.W.; Sletten, T.L. The Impact of Shift Work on Sleep, Alertness and Performance in Healthcare Workers. *Sci. Rep.* **2019**, *9*, 4635. [[CrossRef](#)]
10. Kosmadopoulos, A.; Kervezee, L.; Boudreau, P.; Gonzales-Aste, F.; Vujovic, N.; Scheer, F.A.J.L.; Boivin, D.B. Effects of Shift Work on the Eating Behavior of Police Officers on Patrol. *Nutrients* **2020**, *12*, 999. [[CrossRef](#)]
11. Lim, Y.C.; Hoe, V.C.W.; Darus, A.; Bhoo-Pathy, N. Association between Night-Shift Work, Sleep Quality and Metabolic Syndrome. *Occup. Environ. Med.* **2018**, *75*, 716–723. [[CrossRef](#)] [[PubMed](#)]
12. Keski-Rahkonen, A.; Mustelin, L. Epidemiology of Eating Disorders in Europe: Prevalence, Incidence, Comorbidity, Course, Consequences, and Risk Factors. *Curr. Opin. Psychiatry* **2016**, *29*, 340–345. [[CrossRef](#)]
13. Duthel, F.; Chaplais, E.; Vilmant, A.; Courteix, D.; Duche, P.; Abergel, A.; Pfabigan, D.M.; Han, S.; Mobdillon, L.; Vallet, G.T.; et al. Stress Management in Obesity during a Thermal Spa Residential Programme (ObesiStress): Protocol for a Randomised Controlled Trial Study. *BMJ Open* **2019**, *9*, e027058. [[CrossRef](#)] [[PubMed](#)]
14. Abdella, H.M.; El Farssi, H.O.; Broom, D.R.; Hadden, D.A.; Dalton, C.F. Eating Behaviours and Food Cravings; Influence of Age, Sex, BMI and FTO Genotype. *Nutrients* **2019**, *11*, E377. [[CrossRef](#)]
15. Gifkins, J.; Johnston, A.; Loudoun, R. The Impact of Shift Work on Eating Patterns and Self-Care Strategies Utilised by Experienced and Inexperienced Nurses. *Chronobiol. Int.* **2018**, *35*, 811–820. [[CrossRef](#)]
16. Padilla, H.M.; Wilson, M.; Vandenberg, R.J.; Davis, M.; Clark, M.A. Health Behavior among Working Adults: Workload and Exhaustion Are Associated with Nutrition and Physical Activity Behaviors That Lead to Weight Gain. *J. Health Psychol.* **2021**, *26*, 892–904. [[CrossRef](#)]
17. Nishitani, N.; Sakakibara, H.; Akiyama, I. Eating Behavior Related to Obesity and Job Stress in Male Japanese Workers. *Nutrition* **2009**, *25*, 45–50. [[CrossRef](#)]
18. Palla, L.; Almoosawi, S. Diurnal Patterns of Energy Intake Derived via Principal Component Analysis and Their Relationship with Adiposity Measures in Adolescents: Results from the National Diet and Nutrition Survey RP (2008–2012). *Nutrients* **2019**, *11*, 422. [[CrossRef](#)]
19. Bonham, M.P.; Kaias, E.; Zimberg, I.; Leung, G.K.W.; Davis, R.; Sletten, T.L.; Windsor-Aubrey, H.; Huggins, C.E. Effect of Night Time Eating on Postprandial Triglyceride Metabolism in Healthy Adults: A Systematic Literature Review. *J. Biol. Rhythms* **2019**, *34*, 119–130. [[CrossRef](#)]
20. Bali, A.; Jaggi, A.S. An Integrative Review on Role and Mechanisms of Ghrelin in Stress, Anxiety and Depression. *Curr. Drug Targets* **2016**, *17*, 495–507. [[CrossRef](#)] [[PubMed](#)]
21. Hansson, C.; Shirazi, R.H.; Näslund, J.; Vogel, H.; Neuber, C.; Holm, G.; Anckarsäter, H.; Dickson, S.L.; Eriksson, E.; Skibicka, K.P. Ghrelin Influences Novelty Seeking Behavior in Rodents and Men. *PLoS ONE* **2012**, *7*, e50409. [[CrossRef](#)] [[PubMed](#)]
22. Cummings, D.E. Ghrelin and the Short- and Long-Term Regulation of Appetite and Body Weight. *Physiol. Behav.* **2006**, *89*, 71–84. [[CrossRef](#)] [[PubMed](#)]
23. Nagaya, N.; Kojima, M.; Uematsu, M.; Yamagishi, M.; Hosoda, H.; Oya, H.; Hayashi, Y.; Kangawa, K. Hemodynamic and Hormonal Effects of Human Ghrelin in Healthy Volunteers. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2001**, *280*, R1483–R1487. [[CrossRef](#)] [[PubMed](#)]
24. Münzberg, H.; Morrison, C.D. Structure, Production and Signaling of Leptin. *Metabolism* **2015**, *64*, 13–23. [[CrossRef](#)]
25. Gijón-Conde, T.; Graciani, A.; Guallar-Castillón, P.; Aguilera, M.T.; Rodríguez-Artalejo, F.; Banegas, J.R. Leptin Reference Values and Cutoffs for Identifying Cardiometabolic Abnormalities in the Spanish Population. *Rev. Esp. Cardiol.* **2015**, *68*, 672–679. [[CrossRef](#)]
26. Bouillon-Minois, J.-B.; Trousselard, M.; Thivel, D.; Benson, A.C.; Schmidt, J.; Moustafa, F.; Bouvier, D.; Duthel, F. Leptin as a Biomarker of Stress: A Systematic Review and Meta-Analysis. *Nutrients* **2021**, *13*, 3350. [[CrossRef](#)] [[PubMed](#)]
27. Bouillon-Minois, J.-B.; Trousselard, M.; Thivel, D.; Gordon, B.A.; Schmidt, J.; Moustafa, F.; Oris, C.; Duthel, F. Ghrelin as a Biomarker of Stress: A Systematic Review and Meta-Analysis. *Nutrients* **2021**, *13*, 784. [[CrossRef](#)] [[PubMed](#)]
28. Zefferino, R.; Di Gioia, S.; Conese, M. Molecular Links between Endocrine, Nervous and Immune System during Chronic Stress. *Brain Behav.* **2021**, *11*, e01960. [[CrossRef](#)]
29. Pulpulos, M.M.; Baeken, C.; De Raedt, R. Cortisol Response to Stress: The Role of Expectancy and Anticipatory Stress Regulation. *Horm. Behav.* **2020**, *117*, 104587. [[CrossRef](#)] [[PubMed](#)]
30. Wong, D.L.; Tai, T.C.; Wong-Faull, D.C.; Claycomb, R.; Meloni, E.G.; Myers, K.M.; Carlezon, W.A.; Kvetnansky, R. Epinephrine: A Short- and Long-Term Regulator of Stress and Development of Illness: A Potential New Role for Epinephrine in Stress. *Cell Mol. Neurobiol.* **2012**, *32*, 737–748. [[CrossRef](#)]
31. Duthel, F.; Trousselard, M.; Perrier, C.; Lac, G.; Chamoux, A.; Duclos, M.; Naughton, G.; Mnatzaganian, G.; Schmidt, J. Urinary Interleukin-8 Is a Biomarker of Stress in Emergency Physicians, Especially with Advancing Age—The JOBSTRESS* Randomized Trial. *PLoS ONE* **2013**, *8*, e71658. [[CrossRef](#)] [[PubMed](#)]

32. Bouillon-Minois, J.-B.; Trousselard, M.; Pereira, B.; Schmidt, J.; Clinchamps, M.; Thivel, D.; Ugbohue, U.C.; Moustafa, F.; Ocelli, C.; Vallet, G.; et al. Protocol of the Study on Emergency Health Care Workers' Responses Evaluated by Karasek Questionnaire: The SEEK-Study Protocol. *Int. J. Environ. Res. Public Health* **2021**, *18*, 4068. [[CrossRef](#)] [[PubMed](#)]
33. Durand, A.-C.; Bompard, C.; Sportiello, J.; Michelet, P.; Gentile, S. Stress and Burnout among Professionals Working in the Emergency Department in a French University Hospital: Prevalence and Associated Factors. *Work* **2019**, *63*, 57–67. [[CrossRef](#)] [[PubMed](#)]
34. Bouillon-Minois, J.-B.; Roux, V.; Pereira, B.; Flannery, M.; Pelissier, C.; Ocelli, C.; Schmidt, J.; Navel, V.; Dutheil, F. Stress among Emergency Health Care Workers on Nuclear or Radiation Disaster: A Preliminary State Survey. *Int. J. Environ. Res. Public Health* **2021**, *18*, 8384. [[CrossRef](#)]
35. Dutheil, F.; Boudet, G.; Perrier, C.; Lac, G.; Ouchchane, L.; Chamoux, A.; Duclos, M.; Schmidt, J. JOBSTRESS Study: Comparison of Heart Rate Variability in Emergency Physicians Working a 24-Hour Shift or a 14-Hour Night Shift—A Randomized Trial. *Int. J. Cardiol.* **2012**, *158*, 322–325. [[CrossRef](#)]
36. Aydin, S.; Halifeoglu, I.; Ozercan, I.H.; Erman, F.; Kilic, N.; Aydin, S.; Ilhan, N.; Ilhan, N.; Ozkan, Y.; Akpolat, N.; et al. A Comparison of Leptin and Ghrelin Levels in Plasma and Saliva of Young Healthy Subjects. *Peptides* **2005**, *26*, 647–652. [[CrossRef](#)] [[PubMed](#)]
37. McHill, A.W.; Melanson, E.L.; Higgins, J.; Connick, E.; Moehlman, T.M.; Stothard, E.R.; Wright, K.P. Impact of Circadian Misalignment on Energy Metabolism during Simulated Nightshift Work. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 17302–17307. [[CrossRef](#)] [[PubMed](#)]
38. Crispim, C.A.; Waterhouse, J.; Dâmaso, A.R.; Zimberg, I.Z.; Padilha, H.G.; Oyama, L.M.; Tufik, S.; de Mello, M.T. Hormonal Appetite Control Is Altered by Shift Work: A Preliminary Study. *Metabolism* **2011**, *60*, 1726–1735. [[CrossRef](#)]
39. Scheer, F.A.J.L.; Hilton, M.F.; Mantzoros, C.S.; Shea, S.A. Adverse Metabolic and Cardiovascular Consequences of Circadian Misalignment. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 4453–4458. [[CrossRef](#)]
40. Nguyen, J.; Wright, K.P. Influence of Weeks of Circadian Misalignment on Leptin Levels. *Nat. Sci. Sleep* **2010**, *2*, 9–18. [[CrossRef](#)] [[PubMed](#)]
41. Molzof, H.E.; Peterson, C.M.; Thomas, S.J.; Gloston, G.F.; Johnson, R.L.; Gamble, K.L. Nightshift Work and Nighttime Eating Are Associated with Higher Insulin and Leptin Levels in Hospital Nurses. *Front. Endocrinol.* **2022**, *13*, 876752. [[CrossRef](#)] [[PubMed](#)]
42. Izquierdo, A.G.; Crujeiras, A.B.; Casanueva, F.F.; Carreira, M.C. Leptin, Obesity, and Leptin Resistance: Where Are We 25 Years Later? *Nutrients* **2019**, *11*, 2704. [[CrossRef](#)] [[PubMed](#)]
43. Myers, M.G.; Leibel, R.L.; Seeley, R.J.; Schwartz, M.W. Obesity and Leptin Resistance: Distinguishing Cause from Effect. *Trends Endocrinol. Metab.* **2010**, *21*, 643–651. [[CrossRef](#)] [[PubMed](#)]
44. Taheri, S.; Lin, L.; Austin, D.; Young, T.; Mignot, E. Short Sleep Duration Is Associated with Reduced Leptin, Elevated Ghrelin, and Increased Body Mass Index. *PLoS Med.* **2004**, *1*, e62. [[CrossRef](#)]
45. Mullington, J.M.; Chan, J.L.; Van Dongen, H.P.A.; Szuba, M.P.; Samaras, J.; Price, N.J.; Meier-Ewert, H.K.; Dinges, D.F.; Mantzoros, C.S. Sleep Loss Reduces Diurnal Rhythm Amplitude of Leptin in Healthy Men. *J. Neuroendocrinol.* **2003**, *15*, 851–854. [[CrossRef](#)]
46. Spiegel, K.; Tasali, E.; Penev, P.; Van Cauter, E. Brief Communication: Sleep Curtailment in Healthy Young Men Is Associated with Decreased Leptin Levels, Elevated Ghrelin Levels, and Increased Hunger and Appetite. *Ann. Intern. Med.* **2004**, *141*, 846–850. [[CrossRef](#)] [[PubMed](#)]
47. Bray, G.A. Soft Drink Consumption and Obesity: It Is All about Fructose. *Curr. Opin. Lipidol.* **2010**, *21*, 51–57. [[CrossRef](#)]
48. Biercewicz, M.; Slusarz, R.; Kedziora-Kornatowska, K.; Filipiska, K.; Bielawski, K.; Ruskowska-Ciastek, B. Assessment of Leptin-to-Adiponectin Ratio in Prediction of Insulin Resistance and Nutrition Status in a Geriatric Female Population. *J. Physiol. Pharmacol.* **2020**, *71*, 35–46. [[CrossRef](#)]
49. Langenberg, C.; Bergstrom, J.; Laughlin, G.A.; Barrett-Connor, E. Ghrelin and the Metabolic Syndrome in Older Adults. *J. Clin. Endocrinol. Metab.* **2005**, *90*, 6448–6453. [[CrossRef](#)] [[PubMed](#)]
50. Rigamonti, A.E.; Pincelli, A.I.; Corrà, B.; Viarengo, R.; Bonomo, S.M.; Galimberti, D.; Scacchi, M.; Scarpini, E.; Cavagnini, F.; Müller, E.E. Plasma Ghrelin Concentrations in Elderly Subjects: Comparison with Anorexic and Obese Patients. *J. Endocrinol.* **2002**, *175*, R1–R5. [[CrossRef](#)] [[PubMed](#)]
51. Marchesini, G.; Bianchi, G.; Lucidi, P.; Villanova, N.; Zoli, M.; De Feo, P. Plasma Ghrelin Concentrations, Food Intake, and Anorexia in Liver Failure. *J. Clin. Endocrinol. Metab.* **2004**, *89*, 2136–2141. [[CrossRef](#)] [[PubMed](#)]
52. Bauer, J.M.; Wirth, R.; Troegner, J.; Erdmann, J.; Eberl, T.; Heppner, H.J.; Schusdziarra, V.; Sieber, C.C. Ghrelin, Anthropometry and Nutritional Assessment in Geriatric Hospital Patients. *Z Gerontol. Geriatr.* **2007**, *40*, 31–36. [[CrossRef](#)] [[PubMed](#)]
53. Al-Rawi, N.; Madkour, M.; Jahrami, H.; Salahat, D.; Alhasan, F.; BaHammam, A.; Faris, M.A.-I. Effect of Diurnal Intermittent Fasting during Ramadan on Ghrelin, Leptin, Melatonin, and Cortisol Levels among Overweight and Obese Subjects: A Prospective Observational Study. *PLoS ONE* **2020**, *15*, e0237922. [[CrossRef](#)] [[PubMed](#)]