

BMJ Open DE-PASS Best Evidence Statement (BEST): modifiable determinants of physical activity and sedentary behaviour in children and adolescents aged 5–19 years—a protocol for systematic review and meta-analysis

Mohammed Khudair ¹, Anna Marcuzzi ², Kwok Ng,^{3,4} Gavin Daniel Tempest,¹ František Bartoš,⁵ Ratko Peric ⁶, Maximilian Maier,⁷ Flavia Beccia,⁸ Stefania Boccia,⁸ Mirko Brandes ⁹, Greet Cardon,¹⁰ Angela Carlin,¹¹ Carolina Castagna,⁸ Helmi Chaabene,^{12,13} Anna Chalkley,^{14,15} Simone Ciaccioni,¹⁶ Joanna Cieślińska-Świder,¹⁷ Vilma Čingienė,¹⁸ Cristina Cortis,¹⁹ Chiara Corvino,⁸ Eco JC de Geus,^{20,21} Angela Di Baldassarre,²² Andrea Di Credico ²², Patrik Drid ²³, Rosa Ma Fernández Tarazaga,²⁴ Francesca Gallè,²⁵ Esther García Sánchez,²⁴ Mekdes Gebremariam,²⁶ Barbara Ghinassi,²² Marios Goudas,²⁷ Grainne Hayes,^{28,29} Samuel Honorio,³⁰ Pascal Izzicupo,²² Henriette Jahre,³¹ Judith Jelsma,²¹ Petra Juric,³² Athanasios Kolovelonis,²⁷ Atle Kongsvold,² Evangelia Kouidi ³³, Fiona Mansergh,³⁴ Bojan Masanovic,^{35,36} Teferi Mekonnen,³⁷ Paul Jarle Mork,² Marie Murphy,³⁸ Kelly O'Hara,³⁹ Ayse Ozbil Torun,⁴⁰ Federico Palumbo,¹⁶ Stevo Popovic,^{35,41} Olaf Prieske,⁴² Zrinka Puharic,^{43,44} José Carlos Ribeiro,⁴⁵ Penny Louise Sheena Rumbold,¹ Petru Sandu,⁴⁶ Maroje Sorić,^{32,47} Mette Stavnsbo ⁴⁸, Ioannis Sympas,²⁷ Hidde P van der Ploeg,²¹ Aurélie Van Hoye,²⁹ Sofia Vilela,^{49,50} Catherine Woods,²⁹ Kathrin Wunsch,⁵¹ Laura Capranica,¹⁶ Ciaran MacDonncha,²⁹ Fiona Chun Man Ling ¹ On behalf of DE-PASS

To cite: Khudair M, Marcuzzi A, Ng K, *et al*. DE-PASS Best Evidence Statement (BEST): modifiable determinants of physical activity and sedentary behaviour in children and adolescents aged 5–19 years—a protocol for systematic review and meta-analysis. *BMJ Open* 2022;**12**:e059202. doi:10.1136/bmjopen-2021-059202

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-059202>).

Received 16 November 2021
Accepted 15 July 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Mohammed Khudair;
mohammed2.khudair@northumbria.ac.uk

ABSTRACT

Introduction Physical activity among children and adolescents remains insufficient, despite the substantial efforts made by researchers and policymakers. Identifying and furthering our understanding of potential modifiable determinants of physical activity behaviour (PAB) and sedentary behaviour (SB) is crucial for the development of interventions that promote a shift from SB to PAB. The current protocol details the process through which a series of systematic literature reviews and meta-analyses (MAs) will be conducted to produce a best-evidence statement (BEST) and inform policymakers. The overall aim is to identify modifiable determinants that are associated with changes in PAB and SB in children and adolescents (aged 5–19 years) and to quantify their effect on, or association with, PAB/SB. **Methods and analysis** A search will be performed in MEDLINE, SportDiscus, Web of Science, PsychINFO and Cochrane Central Register of Controlled Trials. Randomised controlled trials (RCTs) and controlled trials (CTs) that investigate the effect of interventions

on PAB/SB and longitudinal studies that investigate the associations between modifiable determinants and PAB/SB at multiple time points will be sought. Risk of bias assessments will be performed using adapted versions of Cochrane's RoB V.2.0 and ROBINS-I tools for RCTs and CTs, respectively, and an adapted version of the National Institute of Health's tool for longitudinal studies. Data will be synthesised narratively and, where possible, MAs will be performed using frequentist and Bayesian statistics. Modifiable determinants will be discussed considering the settings in which they were investigated and the PAB/SB measurement methods used. **Ethics and dissemination** No ethical approval is needed as no primary data will be collected. The findings will be disseminated in peer-reviewed publications and academic conferences where possible. The BEST will also be shared with policy makers within the DE-PASS consortium in the first instance. **Systematic review registration** CRD42021282874.

**STRENGTHS AND LIMITATIONS OF THIS STUDY**

- ⇒ Modifiable determinants will be summarised and described within the settings in which they were investigated to contextualise how they interact with other determinants and subsequently affect physical activity and sedentary behaviour in children and adolescents.
- ⇒ The body of evidence from high-quality research will be summarised, accounting for differences in study designs, methodological quality and measurement methods of physical activity and sedentary behaviour of children and adolescents.
- ⇒ Bayesian meta-analysis will be used in addition to frequentist meta-analysis to allow for assessment of the plausibility of the results and provide more nuanced conclusions regarding the effectiveness of physical activity and sedentary behaviour interventions in children and adolescents.
- ⇒ Modifiable determinants reported in study designs which are not included in the current works may be overlooked and should be investigated in future reviews as they may provide insights into potentially effective interventions.
- ⇒ While our aim is to quantify the effect of modifiable determinants on physical activity and sedentary behaviour of children and adolescents, the analyses of most included studies might not permit the quantification, thus a narrative approach will be adopted.

INTRODUCTION

Physical inactivity among children and adolescents is a global public health issue. Four in five (81%) adolescents across the world do not meet the WHO's physical activity (PA) guidelines.^{1 2} Physical inactivity is a contributing factor to the high prevalence of cardiovascular, metabolic and bone health-related conditions.³ Reducing levels of physical inactivity from a young age has a positive impact on physical and mental health as children and adolescents transition into adulthood.⁴ It is therefore important to promote physical activity behaviour (PAB) and minimise sedentary behaviour (SB) as part of a healthy lifestyle in children and adolescents to mitigate the negative effects of physical inactivity.⁵ In the global action plan on PA 2018–2030, the WHO adopted a target to reduce physical inactivity worldwide by 15% by 2030.⁶ To achieve this target, evidence-based policies need to be created and adopted worldwide.⁷ Furthermore, the fact that PA guidelines are currently not met in a large proportion of young people points towards a lack of understanding and insufficient translation of the evidence behind what makes children and adolescents physically active into policy and public interventions.^{8 9} Therefore, a better understanding of the determinants of PAB/SB is a crucial first step in developing interventions that lead to a sustained increase in PAB and reduced SB and a foundation for PA policy development.^{10 11} In the current protocol, we refer to 'determinants' of PAB or SB as mechanisms that drive and explain behaviour adaptation in specific contexts.^{11 12} We focus on modifiable determinants, signifying those which are malleable and can be altered through interventions, and present opportunities to intervene from public health and policy perspectives.^{11 13} Using a rigorous methodology, our goal is to synthesise high-quality evidence on the effectiveness and association of key modifiable

determinants on PAB/SB and produce a Best Evidence Statement (BEST) which can inform future interventions. We also aim to identify the settings for interventions that are most readily translatable to policy.

The current evidence of the effectiveness of modifiable determinants on PAB/SB is fragmented due to considerable variations in the methodologies used and the methodological quality across the available studies, which has contributed to largely inconclusive findings in systematic literature reviews (SLRs) and meta-analyses (MAs).^{8–10 13–18} To limit the variations across studies and extract trustworthy evidence, it is important to identify high-quality studies. Factors that contribute to methodological quality include research design and PAB/SB measurement methods. A range of research designs have been applied in existing PA research (eg, cross-sectional, longitudinal, randomised controlled trials (RCTs) and controlled trials (CTs)). Potential causality between modifiable determinants and the outcome measures can be indicated by RCTs and CTs, and well-designed RCTs can minimise bias through randomisation and intention-to-treat analyses.^{10 19 20} However, challenges in randomisation of PAB/SB interventions have been recognised,²¹ therefore, CTs might be the next most credible alternative. While RCTs are regarded as the 'gold standard', high-quality longitudinal studies can provide indications of a causal relationship between modifiable determinants and the outcome measures by virtue of the repeated measurements over time.⁸ Furthermore, RCTs and CTs can be short-lasting and may not capture the prolonged exposures that can be explored in longitudinal follow-ups.¹⁰ Therefore, we consider RCTs, CTs and longitudinal studies to be among the highest quality of evidence appropriate to develop the BEST.

Methods for measurement of PAB/SB contribute to the disparities in the methodologies used between studies. Data obtained from self-report methods are generally considered to be less sensitive to change than data obtained via device-based methods due to recall errors, underestimation/overestimation or interpretation discrepancies.^{14 15 22 23} On the one hand, device-based measurements are deemed to be more sensitive to behaviour change and can detect cognitively salient behaviours, such as time spent in SB.²³ On the other hand, many studies rely on self-report measurements as they are less costly, logistically easier to implement and are more applicable in some domains of behaviour (eg, strength training) than device-based measurements.²³ Given that both device-based and self-report methods present strengths and weaknesses, we consider it methodologically appropriate to include both in BEST, provided that validity and reliability of the instruments are assessed and reported thoroughly in the included studies. However, as previous research has shown low levels of agreement between the two measurement methods, we will conduct separate analyses per method within SLRs and MAs.²⁴

Over the years, PAB/SB measurements have been used to assess different forms of PA, such as structured PA (eg,

physical education), leisure-time PA and active transport PA and different domains where sedentary time is spent, such as screen-based activities (eg, doing homework on computers), leisure-based activities (eg, sitting and reading) and transport-related activities (eg, sitting in a bus).¹⁵ Recently, there has been an increased emphasis on identifying the settings (or contexts) in which PAB/SB take place and the determinants at work within the settings, so that the settings of the most impactful, modifiable determinants can be targeted when translating research into policy.^{8 25} Answering the questions about what works for whom (children and/or adolescents), why (determinants and their interactions) and when/where (settings) is critical to advance our understanding of the implementation and possible effectiveness of interventions.²⁶ Therefore, to produce the BEST, we aim to investigate the modifiable determinants in their respective settings in SLRs and MAs so that our results can inform future interventions within settings that speak to policy makers.

The current protocol will be used to produce a series of SLRs and MAs aiming to investigate the effectiveness of modifiable determinants on PAB/SB in children and adolescents using high-quality evidence available. Investigating the modifiable determinants of PAB/SB in their respective settings will help contextualise their modifiability and effect. Therefore, to produce the BEST, it is important to ascertain methodological rigour which is set apart from previous efforts in understanding PAB/SB determinants in children and adolescents. By considering the settings of the modifiable determinants, our results can readily inform policy makers and future PA interventions.

Objectives

The overarching aim of the proposed SLRs and MAs is to identify modifiable determinants that are associated with changes in PAB and SB in children and adolescents (aged 5–19). Specific aims are:

- ▶ To investigate which modifiable determinants of PAB and SB have been targeted in interventions designed to promote PA in children and adolescents in RCTs and CTs.
- ▶ To investigate which modifiable determinants are associated with PAB and SB in children and adolescents in longitudinal studies.
- ▶ To investigate the strength of the association between such modifiable determinants and PAB/SB in children and adolescents.

METHODS AND ANALYSIS

The current protocol was registered in the International prospective register of systematic reviews (PROSPERO) on 12 October 2021 with the registration number: CRD42021282874. The reporting in the current protocol manuscript was guided by the Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols (PRISMA-P).²⁷

The modifiable determinants that have been targeted in all included studies will be listed and analysed narratively in SLRs. Meta-analytic methods will be applied to the data from intervention and longitudinal studies. Analyses will be performed for different categories of studies based on (1) methods for measurement of PAB/SB (eg, self-report, device-based) and (2) age (eg, children aged 5–12 years, adolescents aged 12–19 years) in a series of SLRs and MAs with varying focus. Study settings (eg, school, home, community) will also be identified.

Population

Studies targeting children and adolescents with and without disabilities aged 5–19 years will be included. According to the International Classification of Functioning, Disability and Health (ICF),²⁸ disability is an umbrella term for impairments, activity limitations and participation restrictions, denoting the negative aspects of the interaction between an individual and that individual's contextual factors. Studies that include children and/or adolescents with any reported ongoing diagnosed medical conditions known to affect PA participation and include patients under treatment on all levels of care will be excluded (eg, studies including patients with cancer or individuals with anterior cruciate ligament injury, or studies where the intervention takes place in a clinical setting). Studies that report data for ages exceeding the specified age range will be excluded, unless data for a subgroup within the eligible mean age can be extracted.

Types of studies

We will include studies examining modifiable PAB/SB determinants in RCTs, CTs and longitudinal studies. RCTs and CTs that investigate the effectiveness of interventions aiming to promote PA or reduce SB in children and adolescents, should include control groups or other intervention groups, that are matched to the experimental groups, and report preintervention and postintervention measurements of both outcome measures and modifiable determinants. Longitudinal studies should investigate the association between modifiable determinants of PA and PAB/SB in children and adolescents and report measurements of both the modifiable determinants and PAB/SB at least at two time-points. No control groups or comparisons will be required for the longitudinal studies. Length of follow-up or length of intervention in any of the study designs will not be restricted, data will be extracted if reported for participants within the specified age range (5–19 years).

Outcomes

The main outcome measures targeted in the current protocol are PAB and SB. PA is defined as any bodily movement produced by skeletal muscles that requires energy expenditure, thus including any modality of movement at any intensity.² As such, PAB encompasses behaviours of sedentary, light, moderate and vigorous intensity PA and SB includes any waking behaviour characterised by

an energy expenditure of 1.5 METs or lower while sitting, reclining or lying.^{2–29} Therefore, we will categorise PAB into light, moderate and vigorous intensity and SB-based types of activities reported in the included studies. Any of the two types of measurement methods for PAB/SB, including self-report methods (eg, questionnaires, diaries, recall), and device-based methods (eg, accelerometers, pedometers) will be included.²³ Moreover, we target studies which have reported modifiable determinants as secondary measures. Modifiable determinants will be identified based on the context of each study, where manipulation of the determinant is hypothesised to have an effect on PAB/SB. Where possible, we will explore the mediating effect of the modifiable determinants in the changes in PAB/SB by analysing the structural relationship between the modifiable determinants and PAB/SB.

Comparators

The main comparator will include PAB/SB measurement methods. The included studies will comprise those adopting self-report or device-based measures of PAB/SB or both as outcome measures. Self-report and device-based measures will be analysed separately. In studies where both device-based and self-report measures are reported, the data for both measurement methods will be extracted and analysed separately. In addition, to strengthen the BEST, results from the respective measurement methods will be compared with provide further indication of the strength of the evidence yielded from studies, depending on their measurement methods for PAB/SB. Classification of the settings in which the modifiable determinants were targeted will be identified once data have been extracted.

Search strategy

A search will be performed in MEDLINE (Ovid), PsycINFO (EBSCO), Web of Science, Sport Discus and Cochrane Central Register of Controlled Trials (CENTRAL). The piloted search strategy is presented in [table 1](#). The search strategy is built using the main outcome measures of (1) PAB and (2) SB, and synonyms of PAB/SB that are commonly used in PA research; (3) the targeted study designs (ie, RCTs, CTs and longitudinal studies) and related terms; (4) determinant and synonyms that are commonly used in PA research; (5) the targeted population, to identify children and adolescents and synonyms that are commonly used in PAB/SB research; and (6) measurement methods for PAB/SB such as accelerometer or pedometer for device-based methods and diary and activity recall for self-report methods.

For languages other than English, studies will be included if an English version is available, or if a translation can be obtained through members of the review team. We will include studies published from 2010—which was the year when the first global PA guidelines were published by WHO³⁰ and around the time previous SLRs with similar aims were published.^{31–32} Only peer-reviewed studies will be included and grey literature such as research reports, working papers, conference proceedings and theses will be excluded during the search and at the initial screening of the studies.

Study records

At the initial screening, records of grey literature and duplicates from the different databases will be excluded. The initial screening will be performed before the start of the blinded review process by one member of the review team. For this, EndNote x9³³—a reference management

Table 1 The search terms, Boolean commands and field indicators, presented for each domain

Domain	Search terms
Outcome: Physical activity behaviour*	('Physical activ**') OR (exercise) OR (sport*) OR (play) OR (exertion) OR (recreation) OR (training) OR ('motor activit**') OR ('physical performance') OR ('physical movement') OR ('physical effort') OR (exergaming)
OR	
Outcome: Sedentary behaviour*	(sedentar*) OR ('screen time') OR (gaming) OR ('computer use') OR (sitting) OR (inactiv*) OR ('seated posture') OR ((watch* or view*) N/2 (TV or television))
AND	
Target population*	(child*) OR (youth) OR (adolescen*) OR ('young people') OR ('school age**') OR (p?ediatric) OR (juvenile) OR (teen*)
AND	
Study design†	(RCT) OR ('control* trial**') OR (quasi) OR (longitudinal) OR (intervention*) OR (prospective) OR ('follow-up')
OR	
Determinants†	(determinant*) OR (antecedent*) OR (predictor*) OR (mediator*) OR (moderator*) OR (exposure*)
AND	
Measurement methods†	(acceleromet*) OR ('activity profile') OR (recall) OR (diary) OR ('activity monitor**') OR ('heart rate monitor**') OR ('direct observation') OR (actigraph*) OR ('activity track**') OR ('self report**') OR (survey) OR (pedomet*) OR (wearable*)
*Restricted search to title, abstract and keywords.	
†Search in entire study.	

software will be used. The same member of the review team will upload the resulting list to Covidence³⁴—an online tool for SLRs in which the blinded review process, including title and abstract screening, full-text screening, study selection, data extraction and risk of bias assessment, will be completed. Covidence allows the distribution of studies among several reviewers in a process based on the PRISMA flow diagram for SLRs.³⁵

Several workshops will be held before the commencement of the respective stages (ie, study screening, risk of bias assessments and data extraction) to ensure that all reviewers will be proficient in the procedures and to ensure agreement among them. As the review team consists of 31 members, an online communication tool—Slack³⁶—will be used to maintain communication among the members of the review team throughout the review process to respond to queries and provide updates on the process. A core group of the review team will guide and support the review team members throughout the review process.

Screening process

At title and abstract screening and full-text screening, each study will be screened by two blinded independent reviewers of the review team. Any conflicts between the independent reviewers will be resolved by a third reviewer, who is a member of the core group. An equal number of studies will be distributed among reviewers and random studies are selected by Covidence to be distributed to each reviewer. At the first stage, titles and abstracts will be assessed for eligibility using a prepiloted decision tree based on the inclusion/exclusion criteria expected to be found in either the title or abstract. The full-text version of the studies that remain after title and abstract screening will then be uploaded to Covidence. At the second stage, full texts will be assessed for eligibility using the full inclusion/exclusion criteria. Reasons for exclusion of studies at the full-text stage will be recorded. Following the full-text screening, the included studies will be checked by one reviewer to exclude any duplicate reporting, that is, reporting of the results from the same sample in multiple studies or studies that have been published more than once. For this purpose, study information will be compared between studies, such as authors, study locations and settings, intervention content and design, sample size, demographic information and ethical committee approval number.³⁷ If duplicate reporting is detected among included studies, the reviewers will attempt to identify the main study which was duplicated. If the main study cannot be identified, the study with the longest follow-up or highest number of measurement time points will be selected for inclusion.^{38 39}

Data extraction

A data extraction form will be created in Covidence and piloted ahead of the data extraction stage. The data extraction from each study will be completed by two

independent reviewers. If any information or data are missing, or if clarifications are needed, the corresponding author of the respective studies will be contacted. If a response is not provided before data extraction completes, or if the reporting remains incomplete, the study will be excluded. Following the independent data extraction, the two reviewers will perform a consensus procedure to resolve any conflicts and ascertain the correctness of the extracted data.

The data extracted will include the following items:

- ▶ Study/intervention description: study design, brief study intervention description, description of intervention design and content, description of control group activity and study setting.
- ▶ Sample information: sample size, sample age (including age by sex), sex (including grouping based on sex; % Male, % Female) and population type (disability/non-disability).
- ▶ Outcome measures and modifiable determinants: PAB/SB outcome measurement method type (eg, self-report, device-based) and instrument (eg, Acti-Graph, Youth Activity Profile, 7-day recall), length of device-based PAB/SB measurement (days), days of the week for device-based PAB/SB measurement (weekdays/weekend day), wear-time requirement for device-based PAB/SB measurement, unit of measure for PAB/SB, reported validity and reliability of PAB/SB measurements, modifiable determinant measurement instruments and their reported validity and reliability.
- ▶ Time frames: intervention length (weeks), intervention location (country), number of measurement time points and length of follow-up (weeks).
- ▶ Results data: PAB/SB outcome data (mean, measures of variance) and modifiable determinant data (mean, measures of variance).

Risk of bias

Different scales will be used for the assessment of risk of bias depending on the study design of each included study. For RCTs, a modified version of the Cochrane risk of bias tool for randomised trials (RoB V.2.0) will be used.⁴⁰ For CTs without randomisation, a modified version of Cochrane's Risk of Bias in Non-randomised Studies-of Interventions (ROBINS-I) will be used.⁴¹ The Cochrane tools, RoB V.2.0 and ROBINS-I, are modified to include an additional domain concerning the bias in measurement of the determinants. For longitudinal studies, an adapted version of the National Institutes of Health quality assessment tool will be used.⁴² The adaptation of the latter tool involves the exclusion/addition of items relevant to longitudinal studies, based on the tool used by Kontostoli *et al.*⁴³

The two independent reviewers who extract the data from the respective studies will perform the risk of bias assessment to ensure familiarity with the studies. The risk of bias assessment will be completed in forms created in Covidence with the respective risk of bias tools as

templates. Following the independent data extraction, the two reviewers will perform a consensus procedure to resolve any conflicts and ascertain the correctness of the assessment.

Data synthesis

Data extraction will yield a data file containing data for the included RCTs, CTs and longitudinal studies, and include populations with and without disabilities. A summary table will be created describing the overall characteristics of the included studies with information on the methods (ie, intervention description for intervention studies/exposure for longitudinal studies), settings, modifiable determinants, sample characteristics (ie, sample size, age) and outcomes (ie, outcome measures, measure type, number of measures, measurement time points). Results of the risk of bias assessment will be reported in a separate table.⁴⁴

Findings will be synthesised narratively to identify and list the modifiable determinants and the settings they were investigated in. Studies for disability and non-disability populations, and studies reporting PAB/SB measured using self-report and device-based methods will be discussed separately. The findings will be discussed considering the different settings and the quality of evidence included in the review.

Most data extracted from the included studies are expected to be continuous. Where possible, meta-analytic methods will be applied. MAs will be performed using both frequentist and Bayesian approaches to statistical inference in JASP statistics software.⁴⁵ MAs will be performed for intervention studies (RCTs and CTs) to investigate the effect of the interventions on PAB/SB and determinants and for longitudinal studies to investigate the strength of the association between identified modifiable determinants and PAB/SB. For studies including more than one experimental group or modifiable determinant, each will be included in the MAs.

Direct effect will be investigated in frequentist pairwise comparisons, for which the standardised mean difference (SMD) and the 95% CIs will be calculated. We expect the presence of heterogeneity among included studies in each MA due to the nature, settings or types of interventions. Therefore, the MAs will be conducted using random effects models. For intervention studies, the postintervention data will be used to calculate the between-group difference while controlling for baseline differences. For longitudinal studies, the within-group difference will be calculated as control groups are not expected to be included in longitudinal studies. For data interpretation, effect size values of $SMD < 0.50$ indicate small, $0.50 \leq SMD < 0.80$ indicate medium and $SMD \geq 0.80$ indicate large effects.⁴⁶ Heterogeneity will be identified using Cochrane's Q , which is based on a χ^2 test using the CI size in relation to the df. Heterogeneity will also be quantified by using I^2 , which represents the degree (in %) of methodological consistency across studies using the χ^2 statistic Q in relation to the df. For interpretation of heterogeneity, $I^2 < 25\%$ indicates low heterogeneity, $25\% < I^2 < 50\%$ indicates moderate heterogeneity and I^2

$> 75\%$ indicates high heterogeneity.⁴⁷ Benchmarks will be used to give an approximation for the level of heterogeneity—0%–40%: might not be important; 30%–60%: may represent moderate heterogeneity; 50%–90%: may represent substantial heterogeneity; and 75%–100%: considerable heterogeneity.⁴⁸ The level for statistical significance will be set to $\alpha < 0.05$.

The Bayesian approach to statistical inference will be applied for the MAs using random effects models. The primary benefits of using Bayesian meta-analysis in addition to frequentist meta-analysis include (1) the ability to include prior knowledge of the effect into a model, updating the existing knowledge as evidence accumulates; (2) the ability to make more nuanced conclusions that expand on a simple presence or absence of support for the hypotheses based on a p value and (3) the ability to assess the plausibility of the results and to make conclusions based on the probability that the results are within a given range.^{49 50} For the Bayesian meta-analysis, Gibbs sampling of the Markov Chain Monte Carlo algorithm will be used in JASP.⁴⁵ The probability for publication bias will also be calculated using the JASP extension Robust Bayesian Meta-analysis (RoBMA). We will apply RoBMA to conduct state-of-the-art publication bias-adjusted MA.^{51 52} The Bayesian framework will allow for Bayesian model averaging,⁴⁹ taking several plausible models into account and alleviating concerns about selecting the right model from the variety of adjustment methods available.⁵³ In addition, RoBMA has several other benefits—it allows researchers to (1) quantify evidence on a continuous scale, including for the null, (2) avoid accumulation bias and (3) ease estimation problems by using prior distributions. We will use the prior specifications⁵¹ and models with the modification of removing the fixed-effects models.

Additionally, the mediation effects of determinants on PAB/SB will be investigated using frequentist meta-analytical structural equation modelling (meta-SEM).⁵⁴ To conduct meta-SEM, the covariance structure of the mediation is required. If this information is not presented in a primary study, the authors will be contacted. We will conduct meta-SEM only when we can extract the required data.

ETHICS AND DISSEMINATION

The current protocol describes the process through which a series of SLRs and MAs will be performed, with the aim to identify modifiable determinants that are (in)effective in influencing PAB and SB in children and adolescents. The findings of the resultant studies will be disseminated in peer-reviewed publications and academic conferences, where possible. Modifiable determinants from studies with different study designs and measured using self-report or device-based methods will be reported separately in different publications. The BEST will also be shared with policymakers within the DE-PASS consortium in the first instance. As no primary data will be collected, no ethical approval is required.

Author affiliations

- ¹Department of Sport, Exercise and Rehabilitation, Faculty of Health and Life Sciences, Northumbria University, Newcastle upon Tyne, UK
- ²Department of Public Health and Nursing, Norwegian University of Science and Technology, Trondheim, Norway
- ³School of Educational Sciences and Psychology, University of Eastern Finland, Joensuu, Finland
- ⁴Department of Physical Education and Sport Sciences, University of Limerick, Limerick, Ireland
- ⁵Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands
- ⁶Exercise Physiology Laboratory, OrthoSport Banja Luka, Banja Luka, Bosnia and Herzegovina
- ⁷University College London, London, UK
- ⁸Department of Psychology, Università Cattolica del Sacro Cuore, Milano, Italy
- ⁹Department of Prevention and Evaluation, Leibniz Institute for Prevention Research and Epidemiology, Bremen, Germany
- ¹⁰Department of Movement and Sports Sciences, Ghent University, Ghent, Belgium
- ¹¹Centre for Exercise Medicine, Physical Activity and Health, Sport and Exercise Sciences Research Institute, University of Ulster, Jordanstown Campus, UK
- ¹²Department of Sports and Health Sciences, Faculty of Human Sciences, University of Potsdam, Potsdam, Germany
- ¹³High Institute of Sports and Physical Education of Kef, University of Jendouba, Jendouba, Tunisia
- ¹⁴Faculty of Health Studies, University of Bradford, Bradford, UK
- ¹⁵Center for Physically Active Learning, Faculty of Education, Arts and Sports, Western Norway University of Applied Sciences, Norway, Bergen, Norway
- ¹⁶Department of Movement, Human and Health Sciences, University of Rome "Foro Italico", Rome, Italy
- ¹⁷Department of Physiotherapy, The Jerzy Kukuczka Academy of Physical Education in Katowice, Katowice, Poland
- ¹⁸Institute of Leadership and Strategic Management, Faculty of Public Governance and Business, Mykolas Romeris University, Vilnius, Lithuania
- ¹⁹Department of Human Sciences, Society and Health, University of Cassino and Lazio Meridionale, Cassino, Italy
- ²⁰Department of Biological Psychology, Vrije Universiteit, Amsterdam, Netherlands
- ²¹Department of Public and Occupational Health, Amsterdam Public Health Institute, Amsterdam University Medical Centres, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
- ²²Department of Medicine and Aging Sciences, Gabriele d'Annunzio University of Chieti and Pescara, Chieti, Italy
- ²³Faculty of Sport and Physical Education, University of Novi Sad, Novi Sad, Serbia
- ²⁴Fundación para la Formación e Investigación Sanitarias de la Región de Murcia, Murcia, Spain
- ²⁵Department of Movement Sciences and Wellbeing, University of Naples-Parthenope, Napoli, Italy
- ²⁶Department of Community Medicine and Global Health, Institute of Health and Society, University of Oslo, Oslo, Norway
- ²⁷Department of Physical Education and Sport Science, University of Thessaly, Volos, Greece
- ²⁸Department of Sport and Early Childhood Studies, Technological University of the Shannon: Midlands Midwest, Moylish Campus, Limerick, Ireland
- ²⁹Physical Activity for Health Research Cluster, Health Research Institute, University of Limerick, Limerick, Ireland
- ³⁰Sports, Health and Exercise Research Unit (SHERU), Polytechnic Institute of Castelo Branco, Castelo Branco, Portugal
- ³¹Department of Physiotherapy, Oslo Metropolitan University, Oslo, Norway
- ³²Faculty of Kinesiology, University of Zagreb, Zagreb, Croatia
- ³³Laboratory of Sports Medicine, Department of Physical Education and Sports Science, Aristotle University of Thessaloniki, Thessaloniki, Greece
- ³⁴Health and Wellbeing Programme, Department of Health, Dublin, Ireland
- ³⁵Faculty for Sport and Physical Education, University of Montenegro, Podgorica, Montenegro
- ³⁶Montenegrin Sports Academy, Podgorica, Montenegro
- ³⁷Department of Nutrition, Institute of Basic Medical Sciences, Faculty of Medicine, University of Oslo, Oslo, Norway
- ³⁸Doctoral College, Ulster University, Newtownabbey, UK
- ³⁹Sport Science Department, University of Beira Interior, Covilha, Portugal
- ⁴⁰Department of Architecture and Built Environment, Northumbria University, Newcastle upon Tyne, UK

⁴¹Western Balkan Sport Innovation Lab, Podgorica, Montenegro

⁴²University of Applied Sciences for Sport and Management Potsdam, Potsdam, Germany

⁴³University of Applied Sciences Bjelovar, Bjelovar, Croatia

⁴⁴Faculty of dental medicine and health, Osijek, Croatia

⁴⁵Faculty of Sport, University of Porto, Porto, Portugal

⁴⁶National Institute of Public Health in Romania, Bucharest, Romania

⁴⁷Faculty of Sport, University of Ljubljana, Ljubljana, Slovenia

⁴⁸Department of Sport Science and Physical Education, Faculty of Health and Sport Sciences, University of Agder, Kristiansand, Norway

⁴⁹EPIUnit - Institute of Public Health, University of Porto, Porto, Portugal

⁵⁰Laboratory for Integrative and Translational Research in Population Health (ITR), Porto, Portugal

⁵¹Institute of Sports and Sports Science, Karlsruhe Institute of Technology, Karlsruhe, Germany

Correction notice This article has been corrected since it first published. Author name 'Laura Capranica' has been updated.

Twitter Mohammed Khudair @mo_Khudair, Gavin Daniel Tempest @gdtempest, Ratko Peric @RatkoPeric, Mirko Brandes @MirkoBrandes, Angela Carlin @AngelaCarlin7, Anna Chalkley @AnnaChalkley, Cristina Cortis @cortis_cristina and Fiona Chun Man Ling @drflingUK

Contributors The current systematic review protocol was produced by members of the COST Action CA19101 Determinants of Physical Activities in Settings (DE-PASS). The protocol was conceived and designed by CM, LC, FCML, MK, AM, GDT and KN. The methodology was planned and outlined by CM, FCML, MK, AM, GDT, KN, FB, RP and MM. The protocol was initially drafted by MK, AM, GDT and FCML. Subsequent drafts were reviewed by the included members of DE-PASS: FrantišekB, RP, MM, FlaviaB, SB, MB, GC, AC, CarolinaC, HC, AC, SC, JC-S, VČ, CristinaC, ChiaraC, EJdG, ADB, ADC, PD, RMFT, FG, EGS, MekdesG, BG, MariosG, GH, SH, PI, HJ, JJ, PJ, AthanasiosK, AtleK, EK, FM, BM, TM, PJM, MM, KOH, AOT, FP, SP, OP, ZP, JCR, PR, PS, MarojeS, MetteS, IS, HPvdP, AVH, SV, CW, KW, LC and CM. Revisions were made by MK, AM, GDT, KN and FCML. All authors have read and agreed to the submitted version of the manuscript.

Funding This article is based on work from COST Action CA19101 Determinants of Physical Activities in Settings (DE-PASS), supported by COST (European Cooperation in Science and Technology). The content of this article reflects only the authors' views and the European Community is not liable for any use that may be made of the information contained therein. COST (European Cooperation in Science and Technology) is a funding agency for research and innovation networks. Our Actions help connect research initiatives across Europe and enable scientists to grow their ideas by sharing them with their peers. This boosts their research, career and innovation. www.cost.eu.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Mohammed Khudair <http://orcid.org/0000-0002-8164-6406>

Anna Marcuzzi <http://orcid.org/0000-0002-5460-4788>

Ratko Peric <http://orcid.org/0000-0003-3345-6293>

Mirko Brandes <http://orcid.org/0000-0003-2926-4758>

Andrea Di Credico <http://orcid.org/0000-0002-8388-9305>

Patrik Drid <http://orcid.org/0000-0002-2075-6038>

Evangelia Kouidi <http://orcid.org/0000-0002-5023-2542>

Mette Stavnsbo <http://orcid.org/0000-0002-0920-964X>

Fiona Chun Man Ling <http://orcid.org/0000-0003-2357-3779>

REFERENCES

- 1 Guthold R, Stevens GA, Riley LM, *et al.* Global trends in insufficient physical activity among adolescents: a pooled analysis of 298 population-based surveys with 1.6 million participants. *Lancet Child Adolesc Health* 2020;4:23–35.
- 2 Bull FC, Al-Ansari SS, Biddle S, *et al.* World Health organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med* 2020;54:1451–62.
- 3 World Health Organization. *Global status report on noncommunicable diseases 2014*, 2014.
- 4 Biddle SJH, Ciaccioni S, Thomas G, *et al.* Physical activity and mental health in children and adolescents: an updated review of reviews and an analysis of causality. *Psychol Sport Exerc* 2019;42:146–55.
- 5 Telama R. Tracking of physical activity from childhood to adulthood: a review. *Obes Facts* 2009;2:187–95.
- 6 World Health Organization. *Global action plan on physical activity 2018–2030: more active people for a healthier world*, 2018.
- 7 Sallis JF, Bull F, Guthold R, *et al.* Progress in physical activity over the Olympic quadrennium. *Lancet* 2016;388:1325–36.
- 8 Brug J, van der Ploeg HP, Loyen A, *et al.* Determinants of diet and physical activity (DEDIPAC): a summary of findings. *Int J Behav Nutr Phys Act* 2017;14:1–24.
- 9 Puggina A, Aleksovska K, Buck C, *et al.* Policy determinants of physical activity across the life course: a 'DEDIPAC' umbrella systematic literature review. *Eur J Public Health* 2018;28:105–18.
- 10 Bauman AE, Reis RS, Sallis JF, *et al.* Correlates of physical activity: why are some people physically active and others not? *Lancet* 2012;380:258–71.
- 11 Keller C, Ferrer RA, King RB, *et al.* Future directions of the National Institutes of health science of behavior change program. *Transl Behav Med* 2021;11:1795–801.
- 12 HM Treasury. Magenta book: central government guidance on evaluation, 2020. Available: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/879438/HMT_Magenta_Book.pdf
- 13 Condello G, Puggina A, Aleksovska K, *et al.* Behavioral determinants of physical activity across the life course: a "DEterminants of Diet and Physical ACTivity" (DEDIPAC) umbrella systematic literature review. *Int J Behav Nutr Phys Act* 2017;14.
- 14 Aleksovska K, Puggina A, Giraldo L, *et al.* Biological determinants of physical activity across the life course: a "Determinants of Diet and Physical Activity" (DEDIPAC) umbrella systematic literature review. *Sports Med Open* 2019;5:2.
- 15 Cortis C, Puggina A, Pesce C, *et al.* Psychological determinants of physical activity across the life course: A "DEterminants of Diet and Physical ACTivity" (DEDIPAC) umbrella systematic literature review. *PLoS One* 2017;12:1–25.
- 16 Jaeschke L, Steinbrecher A, Luzak A, *et al.* Socio-cultural determinants of physical activity across the life course: a 'Determinants of Diet and Physical Activity' (DEDIPAC) umbrella systematic literature review. *Int J Behav Nutr Phys Act* 2017;14:1–15.
- 17 O'Donoghue G, Kennedy A, Puggina A. Socio-economic determinants of physical activity across the life course: A 'DEterminants of Diet and Physical ACTivity' (DEDIPAC) umbrella literature review. *Sport Med - Open* 2020;6:1–24.
- 18 Martin Ginis KA, van der Ploeg HP, Foster C, *et al.* Participation of people living with disabilities in physical activity: a global perspective. *Lancet* 2021;398:443–55.
- 19 Elferink-Gemser MT, te Wierike SC, Visscher C. Multidisciplinary longitudinal studies: A Perspective from the field of sports. In: Ericsson AK, Hoffman R, Kozbelt A, eds. *The Cambridge Handbook of expertise and expert performance*. Cambridge: Cambridge University Press, 2018: 271–90.
- 20 Lefebvre C, Glanville J, Briscoe S, *et al.* Chapter 4: Searching for and selecting studies. In: Higgins J, Thomas J, Chandler J, eds. *Cochrane Handbook for systematic reviews of interventions version 6.2*. Cochrane, 2021. www.training.cochrane.org/handbook
- 21 Dobbins M, Husson H, DeCorby K, *et al.* School-based physical activity programs for promoting physical activity and fitness in children and adolescents aged 6 to 18. *Cochrane Database Syst Rev* 2013:CD007651.
- 22 Van Der Horst K, Paw MJCA, Twisk JWR, *et al.* A brief review on correlates of physical activity and sedentariness in youth. *Med Sci Sports Exerc* 2007;39:1241–50.
- 23 Troiano RP, Gabriel KKP, Welk GJ, *et al.* Reported physical activity and sedentary behavior: why do you ask? *J Phys Act Health* 2012;9:S68–75.
- 24 Gorzelitz J, Peppard PE, Malecki K, *et al.* Predictors of discordance in self-report versus device-measured physical activity measurement. *Ann Epidemiol* 2018;28:427–31.
- 25 Newman L, Baum F, Javanparast S, *et al.* Addressing social determinants of health inequities through settings: a rapid review. *Health Promot Int* 2015;30:ii126–43.
- 26 Rycroft-Malone J, McCormack B, Hutchinson AM, *et al.* Realist synthesis: illustrating the method for implementation research. *Implement Sci* 2012;7:1–10.
- 27 Shamseer L, Moher D, Clarke M, *et al.* Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2015;349:g7647–25.
- 28 World Health Organization. *International classification of functioning disability and health: ICF*, 2001. <https://apps.who.int/iris/handle/10665/42407>
- 29 Tremblay MS, Aubert S, Barnes JD, *et al.* Sedentary behavior research network (SBRN) – terminology consensus project process and outcome. *Int J Behav Nutr Phys Act* 2017;14:1–17.
- 30 World Health Organization. *Global recommendations on physical activity for health*. Geneva: World Health Organisation, 2010.
- 31 Craggs C, Corder K, van Sluijs EMF, *et al.* Determinants of change in physical activity in children and adolescents: a systematic review. *Am J Prev Med* 2011;40:645–58.
- 32 Uijtendewilligen L, Nauta J, Singh AS, *et al.* Determinants of physical activity and sedentary behaviour in young people: a review and quality synthesis of prospective studies. *Br J Sports Med* 2011;45:896–905.
- 33 Clarivate. Endnote version x9, 2020. Available: <https://endnote.com/>
- 34 Veritas Health Innovation. Covidence systematic review software. Available: www.covidence.org
- 35 Page MJ, McKenzie JE, Bossuyt PM. PRISMA 2020 statement: updated guidelines for reporting systematic reviews. *BMJ* 2021;372.
- 36 Slack. Slack is where the future works. Available: www.slack.com
- 37 Higgins J, Deeks J. Selecting studies and collecting data. *Cochrane Handb Syst Rev Interv* 2008:151–85.
- 38 Wood JA. Methodology for dealing with duplicate study effects in a meta-analysis. *Organ Res Methods* 2008;11:79–95.
- 39 von Elm E, Poggio G, Walder B, *et al.* Different patterns of duplicate publication: an analysis of articles used in systematic reviews. *JAMA* 2004;291:974.
- 40 Higgins JPT, Altman DG, Gøtzsche PC, *et al.* The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:1–9.
- 41 Sterne JA, Hernán MA, Reeves BC, *et al.* ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;355:4–10.
- 42 National Institute of Health. Quality assessment tool for observational cohort and cross-sectional studies, 2014. Available: <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>
- 43 Kontostoli E, Biddle SJH, Jones AP. Age-related change in sedentary behavior during childhood and adolescence: A systematic review and meta-analysis. *Obes Rev* 2021;1–11.
- 44 Schünemann H, Higgins J, Vist G. Chapter 14: Completing 'Summary of findings' tables and grading the certainty of the evidence. In: Higgins J, Thomas J, Chandler J, eds. *Cochrane Handbook for systematic reviews of interventions version 6.2*. Cochrane, 2021. <http://www.training.cochrane.org/handbook>
- 45 Jasp Team. JASP (version 0.14.1), 2020
- 46 Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. New Jersey: Lawrence Erlbaum Associates Inc, 1988.
- 47 Higgins JPT, Thompson SG, Deeks JJ. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
- 48 Deeks JJ, Higgins JPT, Altman DG. Analysing data and undertaking meta-analyses. *Cochrane Handb Syst Rev Interv* 2019:241–84.
- 49 Hinne M, Gronau QF, van den Bergh D, *et al.* A conceptual introduction to Bayesian model averaging. *Adv Methods Pract Psychol Sci* 2020;3:200–15.
- 50 Gronau QF, Heck DW, Berkhout SW, *et al.* A primer on Bayesian Model-Averaged meta-analysis. *Adv Methods Pract Psychol Sci* 2021;4:251524592110312.
- 51 Bartoš F, Maier M, Wagenmakers E-J. No need to choose: robust Bayesian meta-analysis with competing publication bias adjustment methods. *Psyarxiv* 2021.
- 52 Maier M, Bartoš F, Wagenmakers E-J. Robust Bayesian meta-analysis: addressing publication bias with model-averaging. *Psychol Methods* 2022. doi:10.1037/met0000405. [Epub ahead of print: 19 May 2022].
- 53 Carter EC, Schönbrodt FD, Gervais WM, *et al.* Correcting for bias in psychology: a comparison of meta-analytic methods. *Adv Methods Pract Psychol Sci* 2019;2:115–44.
- 54 Jak S, Li H, Kolbe L, *et al.* Meta-analytic structural equation modeling made easy: a tutorial and web application for one-stage MASEM. *Res Synth Methods* 2021;12:590–606.