



Universiteit  
Leiden  
The Netherlands

## Observability of evaporating lava worlds

Zilinskas, M.; Buchem, C.P.A. van; Miguel, Y.; Louca, A.J.; Lupu, R.; Zieba, S.; Westrenen, W. van

### Citation

Zilinskas, M., Buchem, C. P. A. van, Miguel, Y., Louca, A. J., Lupu, R., Zieba, S., & Westrenen, W. van. (2022). Observability of evaporating lava worlds. *Bulletin Of The American Astronomical Society*, (5), 503.05. Retrieved from <https://hdl.handle.net/1887/3561556>

Version: Publisher's Version  
License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)  
Downloaded from: <https://hdl.handle.net/1887/3561556>

**Note:** To cite this publication please use the final published version (if applicable).

**Bulletin of the AAS • Vol. 54, Issue 5**

# **Observability of Evaporating Lava Worlds**

**Mantas Zilinskas<sup>1</sup> Christiaan van Buchem<sup>1</sup> Yamila Miguel<sup>1</sup>  
Amy Louca<sup>1</sup> Roxana Lupu<sup>2</sup> Sebastian Zieba<sup>3</sup> Wim van Westrenen<sup>4</sup>**

<sup>1</sup>Leiden University, <sup>2</sup>Eureka Scientific Inc, <sup>3</sup>Max-Planck-Institut für Astronomie,  
<sup>4</sup>Vrije Universiteit Amsterdam

**Published on:** Jun 20, 2022

**URL:** <https://baas.aas.org/pub/2022n5i503p05>

**License:** [Creative Commons Attribution 4.0 International License \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/)

Lava worlds belong to a class of short orbital period planets reaching surface temperatures high enough to melt their silicate crust. Theory predicts that the resulting lava oceans outgas their volatile components, attaining equilibrium with the overlying vapour. This creates a tenuous, silicate-rich atmosphere that may be confined to the permanent dayside of the planet. With the recently successful deployment of JWST it is now possible to characterise these worlds. We assess JWST observability of key spectral features by self-consistently modelling silicate atmospheres for all the currently confirmed targets having sufficient substellar temperatures. We use outgassed equilibrium chemistry and radiative transfer methods to compute temperature-pressure profiles, atmospheric chemical compositions and emission spectra. Our results indicate that SiO and SiO<sub>2</sub> infrared features are the best, unique identifiers of silicate atmospheres, detectable using the MIRI instrument of JWST. Detection of these two species in emission would allow for strong constraints on atmospheric thermal structure and possibly the composition of the melt. We also propose that certain species, e.g., TiO or MgO, may be directly tied to different classes of melts, possibly revealing surface and interior dynamics. Currently, there are nearly a dozen confirmed lava planets ideal for characterisation using JWST, but only two of these have been accepted for the initial General Observers program.