



Arbovirus circulation, epidemiology and spatiotemporal distribution in Uganda

Timothy Byaruhanga , John T. Kayiwa , Annet M. Nankya , Irene J. Ataliba , C. Patrick McClure , Jonathan K. Ball , Julius J. Lutwama

PII: S2772-7076(23)00013-9
DOI: <https://doi.org/10.1016/j.ijregi.2023.01.013>
Reference: IJREGI 215

To appear in: *IJID Regions*

Received date: 1 December 2022
Revised date: 25 January 2023
Accepted date: 26 January 2023

Please cite this article as: Timothy Byaruhanga , John T. Kayiwa , Annet M. Nankya , Irene J. Ataliba , C. Patrick McClure , Jonathan K. Ball , Julius J. Lutwama , Arbovirus circulation, epidemiology and spatiotemporal distribution in Uganda, *IJID Regions* (2023), doi: <https://doi.org/10.1016/j.ijregi.2023.01.013>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Ltd on behalf of International Society for Infectious Diseases.
This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Arbovirus circulation, epidemiology and spatiotemporal distribution in Uganda

Timothy Byaruhanga ^{1,2}, John T. Kayiwa ², Annet M. Nankya ², Irene J. Ataliba ², C. Patrick McClure ¹, Jonathan K. Ball ¹ and Julius J. Lutwama ².

¹ University of Nottingham School of Life Sciences, Wolfson Centre for Global Virus Research, Nottingham, UK

² Uganda Virus Research Institute, Department of Arbovirology, Emerging and Re-emerging infectious diseases.

Timothy Byaruhanga; tssekandi@gmail.com

Highlights

- The IgM seroprevalence of arboviruses in Uganda was estimated at 9.5%
- Yellow fever virus was the most prevalent arbovirus
- Spatiotemporal analysis identified four districts with high arbovirus risk
- There is a high burden of arboviral disease among the 5-14 years age group
- There's a need for mass YFV vaccination and continuous arbovirus surveillance

Abstract

Background: Arboviruses are endemic in Uganda; however, little is known about their epidemiology, seasonality, and spatiotemporal distribution. This study sought to provide information on arbovirus outbreaks from acute clinical presentations.

Methods: A retrospective analysis of IgM and confirmatory Plaque Reduction Neutralization Test (PRNT) and results for arbovirus diagnosis of samples collected from 2016 to 2019 was carried out. Demographic data were used to determine the epidemiology and spatiotemporal distribution of arboviruses using the SaTScan and SPSS software.

Results: Arbovirus activity peaked consistently during March-May rainy seasons. The overall arbovirus seroprevalence was 9.5% (137/1441). Of the 137 IgM positives, 72 (52.6%) were

confirmed by PRNT, of which the central region (53/72; 73.6%) and YFV (20/72; 27.8%) had the highest prevalence. The 5-14 age group were four times more likely to be infected with an arbovirus $p=0.003$, 4.1 (1.3- 12.3 CI). Significant arboviral activity was observed among indoor ($p=0.003$) and outdoor ($p=0.05$) patients. Spatiotemporal analysis indicated arboviral activity in 23 districts with five distinct clusters in 6 districts. Masaka, in the Central region, was the most affected among the districts, with a significant YFV cluster ($p<0.001$) from March to May 2016.

Interpretation: This study shows that arbovirus activity peak during the March-May rainy season and highlights the need for YFV mass vaccination to reduce the clinical burden of arboviruses transmitted within the region.

Introduction

Arboviruses are arthropod-borne viruses transmitted by mosquitoes, biting midges, and ticks; grouped within viral families such as Flaviviridae, Rhabdoviridae, Reoviridae, Togaviridae, Nairoviridae, Peribunyaviridae, and Phenuiviridae.

The global burden of arboviruses is significant, with up to 700,000 deaths annually attributed to arbovirus-related diseases [1]. Among mosquito-borne arboviruses, Dengue virus (DENV) has a massive impact on public health, with over 390 million cases reported annually, mainly affecting Asia and Latin America [2]. Yellow Fever Virus (YFV), which is mainly a problem in Africa [3] and South America [4], causes over 200,000 global cases and 30,000 deaths annually [5]. In Uganda, earlier serological studies by Henderson *et al.* reported a 16.8% prevalence for West Nile Virus (WNV), YFV, and Zika virus (ZIKV) [6]. The 2010 YFV outbreak in northern Uganda had a case fatality rate of over 24%, with a 7.5% seroprevalence amongst the population [7]. Among alphaviruses, the immunoglobulin G (IgG)

seroprevalence of Chikungunya virus (CHIKV) and O'nyong'nyong virus (ONNV) is estimated at 31 – 38.6% in Uganda [6]. Although IgM serology provides evidence of recent transmission, the method is plagued by cross-reactivity of antibodies with envelope proteins of related arboviruses [8] in arbovirus endemic regions. Cross-neutralisation is common in regions where more than one arbovirus circulates or in areas where vaccination is deployed [9]. Therefore, the laboratory testing algorithm for arboviruses at the Uganda Virus Research Institute (UVRI) involves initial serological detection of IgM antibodies to YFV, WNV, DENV 1-4, ZIKV, and CHIKV, followed by the plaque reduction neutralisation test (PRNT) confirmation and, where possible, RT-PCR for the detection of viral nucleic acids [10].

Despite continuous surveillance, there is little information about the epidemiology, spatiotemporal distribution, and seasonality of mosquito-borne arboviruses in Uganda. The *Aedes* species (in particular, *Aedes africanus* Theobald, for Uganda) a known competent virus transmitter of YFV, ZIKV, and CHIKV [11], has a significant abundance and distribution within Uganda [12]. However, the co-circulation of these viruses within the country is not known. Therefore, in addition to defining the epidemiology, spatiotemporal distribution and seasonality, this study sought to provide evidence supporting the co-circulation of YFV, ZIKV, and CHIKV during outbreaks.

Methods

Study population, sentinel sites and period

Demographic and laboratory data were selected from a cohort of patients enrolled at hospital-based sentinel sites to monitor arboviruses across Uganda [13]. In regions where no sentinel sites exist, samples were obtained through physician laboratory requests for YFV or any other haemorrhagic fever testing. The data were collected from 2016 to 2019 from patients two months and older with a recorded temperature of $\geq 37.5^{\circ}\text{C}$ or a history of fever not spanning more than seven days at the time of presentation.

Laboratory detection of arboviruses

Laboratory analyses were carried out and interpreted as described by Kayiwa *et al.* [14]. Briefly, immunoglobulin M (IgM) antibodies for YFV and ZIKV were detected using the CDC IgM antibody capture- enzyme-linked immunosorbent assay (CDC-MAC-ELISA) [15]. Diagnosis of WNV, DENV, and CHIKV was made using the InBios West Nile Virus Detect IgM Capture ELISA, DENV Detect IgM Capture ELISA, and the CHIKV Detect IgM Capture, respectively (InBios international, inc., Seattle, WA, USA). Furthermore, IgM-positive samples were confirmed by carrying out the PRNT test and interpreted using the algorithm described by Lindsey *et al.*, 2018 [8].

Univariate analysis was done using SPSS v.26. Spatial and Spatio-temporal data analyses were done using the SaTScan software package (version 10) [16]. A retrospective analysis of positive cases defined by purely spatial and space-time models was made relative to the district population projections predicted by the Uganda Bureau of Statistics from 2016 to 2019 [17]. A Poisson distribution was assumed to determine the number of expected positive cases within the population of a given district. Risk factors for arboviral infection were assessed by categorising patients into three groups, i.e. Outdoor workers group (comprising farmers, fishmongers, security personnel, butchery operators, builders, soldiers, and business persons), Indoor workers (patients with office-related jobs, housewives, student, teachers, children, bar and hotel maids, tailors, and waiters) and healthcare workers (nurses, veterinary officers, clinical and medical officers, and laboratory technicians). Participants provided written consent and assent where necessary.

Results

Description of the study population characteristics

Demographic and laboratory data were available for 1900 anonymised samples collected from the Northern, Central, Western, and Eastern regions (Figure 1a). Of the 1900 samples, complete data were available for 1,441 results, of which 137 (9.5%) had a positive IgM result for an arbovirus. Of the 137 positives, 72 (52.6%) were confirmed positive by the PRNT. Among the PRNT confirmed cases, 53 (73.6%) were from the Central, 13 (18.1%) from the Western, and six (8.3%) from the Eastern and Northern regions (Figure 1b). Flaviviruses accounted for 55.6% of the total PRNT confirmed cases, whilst the rest (44.4%) were confirmed as alphavirus infections. YFV had the highest prevalence of the total positives with 20 cases (27.8%), while the PRNT test confirmed no DENV.

Seasonality of arboviruses

The East African region has two main rainy seasons from March-May and October to December [18]. Arbovirus activity occurred throughout the study period, with higher peaks consistently observed during March-May than in the October – December rainy season (Figure 2). The observed flavivirus peak in March-May 2016 (Figure 2a) is consistent with the YFV outbreak in Central and Western regions (Figure 2b) [19] that continued until June 2016. In addition, three previously unregistered YFV outbreaks occurred, with the first occurring in the Central region in September-October 2017 (Masaka district) and the second in April 2018 (Wakiso district), while the third spread across the Central, North, and West regions from March-September 2019. Generally, alphaviruses circulated throughout the year without a specific pattern (Figure 2a). Only one CHIKV and seven (7) ONNV -positives were confirmed by PRNT whilst 24 cases were unspecific alphaviruses.

Risk factors for Arbovirus infections

On univariate analysis, no risk was observed due to gender among the patients (Table 1).

Among the 1-4-year-old age group, no arbovirus IgM antibodies were observed. However, the risk of an arbovirus infection, especially flaviviruses, was significant in the 5-14 age group $F(1-1439) = 8.2, p = 0.004$. In addition, patients in the 5-14 age group were four times more likely to be infected with an arbovirus (OR=4.1 (1.3- 12.3), $p = 0.003$). Patients within the 55-64 age group were 1.4 times more likely to test positive for an arbovirus infection. Despite IgM antibody and PRNT confirmation of arboviruses within the 15-34, 35-54 and over the 65 years, these patients had a lower risk of arbovirus infection.

Patients within the Central and Western regions had a significantly high likelihood of arboviral infection $F(1, 1439) = 6.0, p = 0.014$ and $F(1, 1439) = 7.5, p = 0.006$, respectively, than patients from the Northern and Western regions. Furthermore, patients from the Central region were 1.9 times 1.9 (1.1- 3.1, CI 95%), $p = 0.006$ more likely to have an arboviral infection than other regions.

Of the 72 PRNT confirmed positives, 44 (61.1%) were patients identified as outdoor workers ($P = 0.05$), and 13 (18.1%) were indoor patients ($p = 0.003$). In addition, indoor patients were 1.8 times (1.2- 2.9, CI 95%) more likely to have an arboviral disease than any other occupation category. Furthermore, the univariate analysis by linear regression showed that the type of occupation was significantly associated with risk of infection (Table 1).

Spatial and spatiotemporal analysis of Arboviruses

Spatial and spatiotemporal analyses were used to understand arbovirus transmission patterns by analysing districts with more than expected arbovirus incidences. Arboviral incidences were analysed by applying a cluster restriction of three cases for areas with high rates and a maximum spatial cluster size set at a radius of 50km. Upon spatial analysis, 23 districts with arboviral activity and five distinct clusters in six districts were recorded (Figure 3a). Of the five clusters, cluster A, in the Buikwe district (geographical Coordinates 0.301 N, 33.012 E)

comprised 15 cases observed within an average population of 194,530. Two clusters, B and C, overlapped Masaka (0.341 S, 31.736 E) and Sembalule districts (0.066 S, 31.483 E), with an average radius of 41.47 km and consisted of 16 cases in an average population of 142,091. Three spatial clusters (D- F) were observed in the South-western region of Uganda, comprising districts, *i.e.* Ntungamo (0.881 S, 30.25 E), Rubanda (1.183 S, 29.85 E) and Rukungiri (0.783 S, 29.33E), respectively. These clusters covered an average radius of 45.4 km comprising four cases in an average population of 23,481.

The spatial-temporal (space-time) scan for clusters applies a cylinder whose circular base covers the geographical area analysed whilst the height measures the time frame of infection activity in the given area [20]. Overall, 6 clusters were detected, of which only two were associated with high rates (Figure 3ii). Clusters A to D were associated with low rates and occurred in Buikwe (A), Mukono (B- coordinates, 0.481 N, 32.771 E), Kampala (C- coordinates 0.363 N, 32.61 E), and Wakiso (D- 0.167 N, 32.5 E) districts. In contrast, clusters E and F were associated with high rates and were observed in Masaka and Sembabule districts, respectively. The cluster in Masaka (cluster E) was significant ($p=0.024$), with three cases giving an annual incidence of 21.2 cases per 100,000 from March to May 2016. The second cluster occurred from September to November 2017 and overlapped between Masaka and Sembabule districts, with five cases and an annual arboviral incidence of 7.5 cases per 100,000 ($p=0.059$).

Spatial and Spatio-temporal distribution of YFV in Uganda

The arboviral activity was further interrogated to analyse the role of each viral species relative to districts and time. In Uganda, the disease threshold for a YFV alert is actioned when there is one suspected or confirmed case in the country

(<http://library.health.go.ug/publications/disease-surveillance-outbreaks/case-definitions-and-epidemic-thresholds-integrated>). For this reason, whilst setting up the data analysis

parameters for YFV activity, we considered a cluster to contain at least two PRNT confirmed cases. As a result, between March 2016 and August 2019, twenty (20) YFV confirmed cases were detected in 10 districts (Figure 3iii). Furthermore, within the ten districts, YFV was more significant in Masaka district (C), with 6 cases giving an incidence of 4.1 per 100,000 ($p < 0.001$). Of interest, YFV activity was detected in all but the Eastern region. Space-time analysis of YFV activity in Uganda showed a significant cluster Masaka from March to May 2016, with three cases and an incidence of 22 cases per 100,000 ($p < 0.001$).

Spatial and Spatio-temporal distribution of alphaviruses in Uganda

Uganda has neither alert nor action level to track alphavirus outbreaks. Therefore, the spatial and space-time distribution of alphaviruses in Uganda was analysed by restricting high rate clusters to at least four cases. Spatial analysis revealed 16 districts with 32 cases between May 2016 and March 2019. Low rates of alphavirus activity were detected within the Wakiso, Kampala and Hoima (1.43 N, 31-35 E) districts (Figure 3iv). On space-time analysis, the cluster observed in Wakiso district occurred between August and December 2017, while the Hoima district cluster occurred from January to May 2018.

Discussion

This study describes the epidemiological pattern of arboviruses, highlighting the role of YFV in morbidity among patients in Uganda. In addition, this study describes the seasonality of arboviruses, defines the patient demographics during the acute phase of illness, and determines the country's high-risk areas.

The general prevalence of arboviruses from 2016 to 2019 was defined based on the PRNT results, highlighting potential outbreaks. Several previous studies have analysed arbovirus seroprevalence based on IgG studies [21], and a few more have attempted to analyse recent arboviral infections using IgM results [22]. However, unlike these investigations, this study

provides more accurate epidemiological and public health information on the nature of the outbreaks using PRNT confirmed results.

It is unclear why the Central region had a higher prevalence of arbovirus infection; however, one possibility is that Lake Victoria may play a significant role in arboviral prevalence in the region. Secondly, the trend of increasing temperatures and flooding [23] observed in Uganda, together with probable house infestation with mosquitoes, may promote the survival of arboviral vectors, increase risk of arbovirus transmission and, in turn, explain the endemicity of arboviruses within the region. Such an effect would not be unprecedented; a study in Brazil identified that environmental factors such as temperature played a vital role in the distribution of arboviruses in high-risk areas [24]. Furthermore, the factors that favoured the dominance of YFV and its expansion from the central and western regions into the northern region by 2019 are unknown, despite sharing a common vector with ZIKV, DENV, and CHIKV.

High arbovirus infection levels among outdoor and indoor workers may be associated with the vectors' feeding and resting behaviour patterns [25]. In addition, climate change has affected farmers' agricultural patterns [26], which would probably impact the length and period of exposure to arboviral vectors whilst on the fields. In our study, the arboviral incidence among health workers cannot be concluded as a work-related occurrence. However, before the 1970s, frequent arbovirus infection of laboratory personnel involved in studies manipulating arboviruses was common, with over 2700 cases and 107 fatalities [27]. Patients aged 1-4 years reported no arbovirus infection, contrasting with a Puerto Rico report that recorded CHIKV infections in infants below one year [28]. Our findings agree that arboviral infections have been documented in patients five years and older [29].

Spatial and spatiotemporal analysis mapped the arbovirus distribution and time pattern of YFV. The spatiotemporal distribution of WNV, ZIKV, and DENV were not analysed due to insufficient data to support statistical analysis. Furthermore, the arbovirus species' definitive spatial and spatiotemporal distribution could not be fully defined due to the existence of unspecified alpha and flaviviruses. The diagnosis of unspecific alpha and flaviviruses highlights the importance of early detection by applying assays such as molecular detection by PCR. As a result of few PRNT confirmed CHIKV infections, this study could not confirm CHIKV co-circulation with YFV. However, spatiotemporal analysis using presumptive IgM results identified a possible CHIKV outbreak in Buikwe district co-occurring with the YFV outbreak in Masaka and Rukungiri districts in February-May 2016. Simultaneous circulation of CHIKV and YFV has been reported elsewhere [30], yet the notifiable YFV outbreak overshadows CHIKV activities.

YFV cases were detected throughout the study period (2016-2019), of which 80% were observed within the central region. The reduction in YFV cases from 2017 to 2019 is attributed to mass vaccination [31] although new YFV activity was detected in northern Uganda. It is important to note that according to our study, no previous mass vaccinations had been conducted in northern Ugandan areas with new YFV activity.

Our study is limited to patients presenting with acute febrile illnesses at sentinel health facilities or physician requests, and thus unable to detect most asymptomatic and non-sentinel site-associated cases within the country. Furthermore, the study does not account for population movements which may facilitate the introduction of competent and virus infested vectors into new regions/niches. In addition, a study characterising arbovirus in indoor and outdoor vectors and the species involved, needs to be carried out to support the high arboviral incidence.

Conclusion

Arbovirus infections, mainly YFV, occur during the rainy seasons in Uganda. There is a need to reduce the transmission of arboviruses among outdoor and indoor workers through mass vaccination for preventable arboviral disease. Early mass vaccination of the 5-14 age group will significantly lower YFV cases in the country. In addition, because the Northern and Eastern regions have limited sentinel sites, active capture of arboviral outbreaks in the regions is dependent on incidence reporting by referral hospitals. Thus, vital information on the epidemiology of outbreaks may be missed out.

Conflict of Interest

All authors declare no conflict of interest

Sources of Funding

Laboratory tests were funded by the CDC Fort Collins grant to the Department of Arbovirology, Emerging and Re-emerging Infectious Diseases.

Ethical approval statement

Ethical approval was waived through the existing approval granted by the UVRI research ethics committee and the Uganda National Council of Science and Technology to the Department of Arbovirology.

References

- [1] H. Ketkar, D. Herman, P. Wang, Genetic determinants of the re-emergence of arboviral diseases, *Viruses* 11(2) (2019) 150.
- [2] A. Wilder-Smith, D.J. Gubler, S.C. Weaver, T.P. Monath, D.L. Heymann, T.W. Scott, Epidemic arboviral diseases: priorities for research and public health, *The Lancet infectious diseases* 17(3) (2017) e101-e106.
- [3] M.U. Kraemer, N.R. Faria, R.C. Reiner Jr, N. Golding, B. Nikolay, S. Stasse, M.A. Johansson, H. Salje, O. Faye, G.W. Wint, Spread of yellow fever virus outbreak in Angola and the Democratic Republic of the Congo 2015–16: a modelling study, *The Lancet infectious diseases* 17(3) (2017) 330-338.

- [4] P.F. Vasconcelos, J.E. Bryant, A.P.T. Da Rosa, R.B. Tesh, S.G. Rodrigues, A.D. Barrett, Genetic divergence and dispersal of yellow fever virus, Brazil, *Emerging infectious diseases* 10(9) (2004) 1578.
- [5] J.-P. Mutebi, A.D. Barrett, The epidemiology of yellow fever in Africa, *Microbes and infection* 4(14) (2002) 1459-1468.
- [6] B.E. Henderson, G.B. Kirya, L. Hewitt, Serological survey for arboviruses in Uganda, 1967-69, *Bulletin of the World Health Organization* 42(5) (1970) 797.
- [7] J.F. Wamala, M. Malimbo, C.L. Okot, A.D. Atai-Omoruto, E. Tenywa, J.R. Miller, S. Balinandi, T. Shoemaker, C. Oyoo, E.O. Omony, Epidemiological and laboratory characterization of a yellow fever outbreak in northern Uganda, October 2010–January 2011, *International Journal of Infectious Diseases* 16(7) (2012) e536-e542.
- [8] N.P. Lindsey, J.E. Staples, K. Powell, I.B. Rabe, M. Fischer, A.M. Powers, O.I. Kosoy, E.C. Mossel, J.L. Munoz-Jordan, M. Beltran, Ability to serologically confirm recent Zika virus infection in areas with varying past incidence of dengue virus infection in the United States and US territories in 2016, *Journal of clinical microbiology* 56(1) (2018) e01115-17.
- [9] I.B. Rabe, J.E. Staples, J. Villanueva, K.B. Hummel, J.A. Johnson, L. Rose, S. Hills, A. Wasley, M. Fischer, A.M. Powers, Interim guidance for interpretation of Zika virus antibody test results, *Morbidity and mortality weekly report* 65(21) (2016) 543-546.
- [10] J.T. Kayiwa, A.M. Nankya, I. Ataliba, C.A. Nassuna, I.E. Omara, J.W. Koehler, J.M. Dye, E.C. Mossel, J.J. Lutwama, Dengue fever and chikungunya virus infections: identification in travelers in Uganda–2017, *Tropical Diseases, Travel Medicine and Vaccines* 5(1) (2019) 1-5.
- [11] D. Diallo, G. Fall, C.T. Diagne, A. Gaye, Y. Ba, I. Dia, O. Faye, M. Diallo, Concurrent amplification of Zika, chikungunya, and yellow fever virus in a sylvatic focus of arboviruses in Southeastern Senegal, 2015, *BMC microbiology* 20(1) (2020) 1-11.
- [12] J.-P. Mutebi, M. Crabtree, R. Kading, A. Powers, J. Lutwama, B. Miller, Mosquitoes of western Uganda, *Journal of medical entomology* 49(6) (2014) 1289-1306.
- [13] J.J. Lutwama, Uganda Arboviruses Surveillance Activities, *African Journal of Pharmacology and Therapeutics* 1(4) (2012).
- [14] J.T. Kayiwa, A.M. Nankya, I.J. Ataliba, E.C. Mossel, M.B. Crabtree, J.J. Lutwama, Confirmation of Zika virus infection through hospital-based sentinel surveillance of acute febrile illness in Uganda, 2014–2017, *Journal of General Virology* 99(9) (2018) 1248-1252.
- [15] D.A. Martin, D.A. Muth, T. Brown, A.J. Johnson, N. Karabatsos, J.T. Roehrig, Standardization of immunoglobulin M capture enzyme-linked immunosorbent assays for routine diagnosis of arboviral infections, *Journal of clinical microbiology* 38(5) (2000) 1823-1826.
- [16] M. Kulldorff, A spatial scan statistic, *Communications in Statistics-Theory and methods* 26(6) (1997) 1481-1496.
- [17] S. UBOS, Uganda Bureau of Statistics, The National Population and Housing Census (2014).
- [18] D. Conway, E. Allison, R. Felstead, M. Goulden, Rainfall variability in East Africa: implications for natural resources management and livelihoods, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 363(1826) (2005) 49-54.
- [19] L. Kwagonza, B. Masiira, H. Kyobe-Bosa, D. Kadobera, E.B. Atuheire, B. Lubwama, A. Kagirita, E. Katushabe, J.T. Kayiwa, J.J. Lutwama, Outbreak of yellow fever in central and southwestern Uganda, February–May 2016, *BMC infectious diseases* 18(1) (2018) 1-9.
- [20] M. Kulldorff, W.F. Athas, E.J. Feurer, B.A. Miller, C.R. Key, Evaluating cluster alarms: a space-time scan statistic and brain cancer in Los Alamos, New Mexico, *American journal of public health* 88(9) (1998) 1377-1380.
- [21] Y. Ushijima, H. Abe, G. Nguema Ondo, R. Bikangui, M. Massinga Loembé, V.R. Zadeh, J.G. Essimengane, A.V. Mbouna, E.B. Bache, S.T. Agnandji, Surveillance of the major pathogenic arboviruses of public health concern in Gabon, Central Africa:

- increased risk of West Nile virus and dengue virus infections, *BMC Infectious Diseases* 21(1) (2021) 1-11.
- [22] J.-P. Carrera, Z.M. Cucunubá, K. Neira, B. Lambert, Y. Pittí, J. Liscano, J.L. Garzón, D. Beltran, L. Collado-Mariscal, L. Saenz, Endemic and Epidemic Human Alphavirus Infections in Eastern Panama: An Analysis of Population-Based Cross-Sectional Surveys, *The American journal of tropical medicine and hygiene* 103(6) (2020) 2429.
- [23] R. Twinomuhangi, H. Sseviiri, C. Mulinde, P.I. Mukwaya, A. Nimusiima, A.M. Kato, Perceptions and vulnerability to climate change among the urban poor in Kampala City, Uganda, *Regional Environmental Change* 21(2) (2021) 1-13.
- [24] C. Lorenz, T.S. Azevedo, F. Virginio, B.S. Aguiar, F. Chiaravalloti-Neto, L. Suesdek, Impact of environmental factors on neglected emerging arboviral diseases, *PLoS neglected tropical diseases* 11(9) (2017) e0005959.
- [25] H. Delatte, A. Desvars, A. Bouétard, S. Bord, G. Gimonneau, G. Vourc'h, D. Fontenille, Blood-feeding behavior of *Aedes albopictus*, a vector of Chikungunya on La Réunion, *Vector-Borne and Zoonotic Diseases* 10(3) (2010) 249-258.
- [26] J.S. Okonya, K. Syndikus, J. Kroschel, Farmers' perception of and coping strategies to climate change: Evidence from six agro-ecological zones of Uganda, *Journal of agricultural science* 5(8) (2013) 252.
- [27] R. Hanson, S. Sulkin, E. Buescher, W.M. Hammon, R. McKinney, T. Work, Arbovirus Infections of Laboratory Workers: Extent of problem emphasizes the need for more effective measures to reduce hazards, *Science* 158(3806) (1967) 1283-1286.
- [28] C.H. Hsu, F. Cruz-Lopez, D. Vargas Torres, J. Perez-Padilla, O.D. Lorenzi, A. Rivera, J.E. Staples, E. Lugo, J. Munoz-Jordan, M. Fischer, Risk factors for hospitalization of patients with chikungunya virus infection at sentinel hospitals in Puerto Rico, *PLoS neglected tropical diseases* 13(1) (2019) e0007084.
- [29] M.L.B. Ferreira, C.A.A. de Brito, R.F. de Oliveira França, Á.J.P. Moreira, M.Í. de Morais Machado, R. da Paz Melo, R. Medialdea-Carrera, S.D. Mesquita, M.L. Santos, R. Mehta, Neurological disease in adults with Zika and chikungunya virus infection in Northeast Brazil: a prospective observational study, *The Lancet Neurology* 19(10) (2020) 826-839.
- [30] S. Makiala-Mandanda, S. Ahuka-Mundeke, J.L. Abbate, E. Pukuta-Simbu, J. Nsio-Mbeta, N. Berthet, E.M. Leroy, P. Becquart, J.-J. Muyembe-Tamfum, Identification of dengue and chikungunya cases among suspected cases of yellow fever in the Democratic Republic of the Congo, *Vector-Borne and Zoonotic Diseases* 18(7) (2018) 364-370.
- [31] W.H. Organization, Weekly Bulletin on Outbreak and other Emergencies: Week 21: 20-26 May 2019, *Weekly Bulletin on Outbreaks and other Emergencies* (2019) 1-27.




Figure 1 Sample collection and arboviruses distribution by region: **A.** Extract of the map of Uganda showing sample collection by region, further divided into districts. The orange represents the north, green for the west, red for the central, blue for the east, and grey shows areas with no sample collection. The number of samples collected from each region is represented as a percentage of the total collection. **B.** Regions with arboviruses confirmed by PRNT. The map divisions represent districts and are coloured to represent geographical regions. The uncoloured districts represent areas where no arbovirus has been confirmed. The maps are not drawn to scale.




Figure 2. Seasonality of arboviruses in Uganda: **A.** Number of arbovirus positives by month in Uganda from February 2016 to September 2019. The alphaviruses (red) and flaviviruses (blue) incidences peaked between March and May every year. The March-May 2016 and 2019 peaks coincide with the YFV outbreak in Masaka and Koboko districts. **B.** PRNT confirmed YFV (red) and other arboviruses (blue) observed in Uganda from 2016 to 2019. Four distinct YFV outbreaks were observed, with the first occurring in the central region March-May 2016 in the central and western regions. The second was in September-October 2017 (Masaka district) and the third in April 2018 (Wakiso district), while the fourth spread across the central, north, and west regions from March to September 2019.




Figure 3. Spatio-temporal distribution of arboviruses in Uganda in clock-wise direction. I. Spatial analysis of arboviruses showing their distribution in Uganda. Arboviral incidences were observed in 23 districts (black and red dots). Red circles represent clusters in districts from which more than two arboviruses were detected, indicated as loci A- F. Districts A-F are: A. Buikwe, B. Masaka, C. Sembabule, D. Ntungamo, E. Rubanda, and F. Rukungiri. Abbreviation: DRC- Democratic Republic of Congo. **II.** Space-time scan of arbovirus activity in Uganda. The blue, black and red spots show areas where arbovirus activity was observed. Two clusters E (Masaka) and F (Sembabule) had high arboviral rates and occurred from March-May 2016 in Masaka district. Cluster F overlapped with E but occurred from September to February 2017. The blue dots represent districts with low arboviral rates. A- Buikwe, B- Mukono, C- Kampala and D – Wakiso. **III.** Spatial Analysis of YFV activity. The blue, black and red spots show areas where YFV activity was observed. Cluster C (in Masaka) represent significant YFV activity, whilst clusters A (Kampala) and B (Wakiso) showed low activity. All but the eastern region showed YFV activity. **IV.** Spatial distribution of alphaviruses May 2016 - March 2019. The black, grey and red spots show the 16 locations where alphaviruses were detected. The red spot shows the Hoima district cluster with a high rate of alphavirus activity, whilst the blue spots represent areas of low virus rates in Kampala and Wakiso.

Table 1. Epidemiological characteristics of arboviruses. Abbreviation: HCW- Healthcare workers.

	Total positive	Total negative	Univariate	CHI Square test: pValue, OR (95% CI)
	{n= 72 (%)}	{n= 1369, (%)}		
Prevalence (n= 1441)	4.9			
GENDER				
Female	39 (54.2)	670 (48.9)		
Male	33 (45.8)	699 (51.1)		
AGE GROUP				
0-4 yrs	Nil	83 (6.1)		
5-14 yrs	3 (4.2)	231 (16.9)	F(1,1439)= 8.2, p=0.004	4.1 (1.3- 12.3)
15-34 yrs	37 (51.4)	584 (42.7)		0.08 (0.6- 1.0)
35-54 yrs	24 (33.3)	322 (23.5)		0.7 (0.5- 1.0)
55-64 yrs	3 (4.2)	81 (5.9)		1.4 (0.5- 4.4)
65+ yrs	5 (6.9)	68 (4.9)		
REGION				
Central region	53 (73.6)	809 (59.1)	F(1, 1439)= 6.0, p=0.014	0.8 (0.6-0.93)
Western region	13 (18.1)	459 (33.5)	F(1, 1439)= 7.5, p=0.006	p0.006, 1.9 (1.1- 3.1)
Northern region	4 (5.5)	69 (5)		0.9 (0.3-2.4)
Eastern region	2 (2.8)	32 (2.4)		0.8 (0.2-3.1)
OCCUPATION				
Outdoors	44 (61.1)	674 (49.2)	F(1,1439)= 3.9, p=0.05	p0.05, 0.8(07- 1.0)
Indoors	13 (18.1)	526 (38.4)	F(1,1439)= 8.9, p=0.003	p0.003,1.8 (1.2- 2.9)
HCW	3 (4.2)	28 (2)		1.6 (0.2- 11.4)
Other	12 (16.7)	141 (10.3)		0.6 (0.4- 1.1)