

2022

## Does high-velocity resistance exercise elicit greater physical function benefits than traditional resistance exercise in older adults? A systematic review and network meta-analysis of 79 trials

Pedro Lopez

*Edith Cowan University*, p.lopezdacruz@ecu.edu.au

Anderson Rech

Maria Petropoulou

Robert Newton

*Edith Cowan University*, r.newton@ecu.edu.au

Dennis Taaffe

*Edith Cowan University*, d.taaffe@ecu.edu.au

*See next page for additional authors*

Follow this and additional works at: <https://ro.ecu.edu.au/ecuworks2022-2026>



Part of the [Sports Sciences Commons](#)

---

[10.1093/gerona/glac230](https://doi.org/10.1093/gerona/glac230)

Lopez, P., Rech, A., Petropoulou, M., Newton, R. U., Taaffe, D. R., Galvão, D. A., . . . Radaelli, R. (2022). Does high-velocity resistance exercise elicit greater physical function benefits than traditional resistance exercise in older adults? A systematic review and network meta-analysis of 79 trials. *The Journals of Gerontology: Series A*.

Advance online publication. <https://doi.org/10.1093/gerona/glac230>

This Journal Article is posted at Research Online.

<https://ro.ecu.edu.au/ecuworks2022-2026/1471>

---

**Authors**

Pedro Lopez, Anderson Rech, Maria Petropoulou, Robert Newton, Dennis Taaffe, Daniel A. Galvão, Douglas P. J. Turella, Sandro R. Freitas, and Régis Radaelli

## Review

# Does High-Velocity Resistance Exercise Elicit Greater Physical Function Benefits Than Traditional Resistance Exercise in Older Adults? A Systematic Review and Network Meta-Analysis of 79 Trials

Pedro Lopez, MSc,<sup>1,2,\*</sup> Anderson Rech, PhD,<sup>3</sup> Maria Petropoulou, PhD,<sup>4</sup> Robert U. Newton, PhD, DSc,<sup>1,2</sup> Dennis R. Taaffe, PhD, DSc,<sup>1,2</sup> Daniel A. Galvão, PhD,<sup>1,2</sup> Douglas J. P. Turella, BSc,<sup>3</sup> Sandro R. Freitas, PhD,<sup>5</sup> and Régis Radaelli, PhD<sup>5,6</sup>

<sup>1</sup>Exercise Medicine Research Institute, Edith Cowan University, Joondalup, Western Australia, Australia. <sup>2</sup>School of Medical and Health Sciences, Edith Cowan University, Joondalup, Western Australia, Australia. <sup>3</sup>Curso de Educação Física, Universidade de Caxias do Sul, Caxias do Sul, Rio Grande do Sul, Brazil. <sup>4</sup>Institute of Medical Biometry and Statistics, Faculty of Medicine and Medical Center, University of Freiburg, Freiburg, Germany. <sup>5</sup>Neuromuscular Research Lab, Faculty of Human Kinetics, University of Lisbon, Cruz Quebrada Dafundo, Portugal. <sup>6</sup>CIPER, Faculty of Human Kinetics, University of Lisboa, Cruz Quebrada, Dafundo, Portugal.

\*Address correspondence to: Pedro Lopez, MSc, Exercise Medicine Research Institute, Edith Cowan University, 270 Joondalup Drive, Joondalup, WA 6027, Australia. E-mail: [p.lopezdacruz@ecu.edu.au](mailto:p.lopezdacruz@ecu.edu.au)

Received: August 21, 2022; Editorial Decision Date: November 6, 2022

**Decision Editor:** Lewis A. Lipsitz, MD, FGSA

## Abstract

**Background:** A systematic review and network meta-analysis was undertaken to examine the effectiveness of different modes of resistance exercise velocity in fast walking speed, timed-up and go, 5-times sit-to-stand, 30-second sit-to-stand, and 6-minute walking tests in older adults. **Methods:** CINAHL, Embase, LILACS, PubMed, Scielo, SPORTDiscus, and Web of Science databases were searched up to February 2022. Eligible randomized trials examined the effects of supervised high-velocity or traditional resistance exercise in older adults (ie, ≥60 years). The primary outcome for this review was physical function measured by fast walking speed, timed-up and go, 5-times sit-to-stand, 30-second sit-to-stand, and 6-minute walking tests, while maximal muscle power and muscle strength were secondary. A random-effects network meta-analysis was undertaken to examine the effects of different resistance exercise interventions.

**Results:** Eighty articles describing 79 trials ( $n = 3\,575$ ) were included. High-velocity resistance exercise was the most effective for improving fast walking speed (standardized mean difference [SMD]  $-0.44$ , 95% confidence interval [CI]:  $0.00$  to  $0.87$ ), timed-up and go (SMD  $-0.76$ , 95% CI:  $-1.05$  to  $-0.47$ ), and 5-times sit-to-stand (SMD  $-0.74$ , 95% CI:  $-1.20$  to  $-0.27$ ), while traditional resistance exercise was the most effective for 30-second sit-to-stand (SMD  $1.01$ , 95% CI:  $0.68$  to  $1.34$ ) and 6-minute walking (SMD  $0.68$ , 95% CI:  $0.34$  to  $1.03$ ).

**Conclusion:** Our study provides evidence that resistance exercise velocity effects are specific in older adults, as evidenced by physical function test dependence. We suggest that prescriptions based on the velocity of contraction should be individualized to address the specific functional needs of participants.

**Keywords:** Muscle power, Physical function, Strength training

Aging, even in the absence of chronic disease, is associated with several biological changes in the neural and musculoskeletal systems (1,2). While reductions of ~1.0% per year are observed in muscle mass (3–5), declines in muscle strength and muscle power (ie, the

product of force production by the speed of movement) are greater and faster (6,7), affecting the ability to undertake activities of daily living in older adults (8,9). In addition, older adults presenting with lower levels of muscle strength and power are at increased risk of

physical disability (10), falling (11,12), and frailty (13), which ultimately lead to dependent living and higher risk of hospitalizations and all-cause mortality (14,15). Therefore, strategies to preserve muscle strength and power, and physical function are critical in older adults, as established by the World Health Organization's concept of healthy aging (16).

Resistance exercise is a well-established intervention to improve physical function in older adults (17–20). As previously demonstrated, various resistance exercise modes can improve muscle strength and power (19–21), which partially accounts for gains in physical function (22). Recently, exercise guidelines have advocated high-velocity resistance exercise (defined as resistance exercise attempted to move the load as fast as possible during the concentric phase) as the most effective resistance exercise mode for improving physical function in older adults (23,24). This potential superiority of high-velocity compared to traditional resistance exercise (defined as resistance exercise prescribed at a set cadence of  $\geq 2$  seconds during the concentric phase or not attempting to move the load as fast as possible) relies on superior neural rather than muscle fiber adaptations (25) and the greater relevance of muscle power production to the performance of daily living activities (8,9,26–28). However, conflicting results in physical function are derived from previous randomized trials (29–31) and pairwise meta-analyses (32–34). For example, previous pairwise meta-analyses from Steib et al. (32) and Tschopp et al. (33), involving a small number of studies, indicate an advantage in favor of high-velocity compared to traditional resistance exercise when several physical function tests are combined (32,33). In contrast, despite similar results, de Rosa Orssatto et al. (34) and Balachandran et al. (35) suggest that the evidence is inconclusive to support such superiority. This may be related to several factors, such as (a) studies involving participants with varying health status (eg, healthy, frail, and mobility-limited participants), (b) studies undertaking resistance exercise combined with aerobic or balance exercise interventions, and (c) analyses combining several physical function tests which would result in a composite measure. These issues likely affect comparisons between these 2 resistance exercise modes precluding individualization in older adults. Therefore, although previous exercise guidelines recommend high-velocity resistance exercise for greater improvements in physical function (23,24), a more comprehensive assessment of the literature is necessary to better target specific exercise prescriptions to older adults based on individual clinical needs.

The utilization of only direct comparisons between high-velocity and traditional resistance exercises may also represent a limitation in previous meta-analyses (32–34). For instance, the inclusion of studies comparing high-velocity or traditional resistance exercise versus control condition would improve precision and comprehensiveness in assessing the effects of different velocities of resistance exercise on physical function in older adults. Network meta-analysis, comparing multiple interventions simultaneously by combining direct and indirect evidence within a network of randomized trials, is considered very effective in providing comparative effectiveness and supporting clinical decision-making (36). As a result, for this systematic review and network meta-analysis we aimed to determine which resistance exercise velocity mode, high-velocity or traditional, is most effective for improving walking speed, lower-limb mobility, power, strength and endurance, cardiorespiratory fitness, and walking endurance by using a range of tests such as fast walking speed, timed-up and go, 5-times sit-to-stand, 30-second sit-to-stand, and 6-minute walking tests in older adults. These tests are commonly used in exercise trials

to assess physical function in this age group and are strongly associated with a variety of clinical endpoints such as mortality, cardiovascular disease and death, and physical disability (37–41). In addition, we also examined the moderating effects of physical health status on these outcomes.

## Methods

### Study Eligibility Criteria

To be included, published randomized trials compared traditional resistance exercise versus high-velocity resistance exercise versus control, traditional resistance exercise versus high-velocity resistance exercise, traditional resistance exercise versus control, or high-velocity resistance exercise versus control in untrained older adults (ie,  $\geq 60$  years). High-velocity resistance exercise was defined as resistance exercise that attempted to move the load as fast as possible during the concentric phase, while traditional resistance exercise was defined as resistance exercise prescribed at a set cadence (eg,  $\geq 2$  seconds) or not attempting to move the load as fast as possible. These programs were required to be delivered under supervision involving physically healthy or impaired older participants (eg, self-reported physical disability, mobility-limited, sarcopenia (42), and frailty). The exclusion criteria were: (a) studies including older adults exclusively with acute or chronic cardiovascular or metabolic conditions (eg, osteoarthritis, type II diabetes, cancer, chronic hemodialysis, heart failure, hospitalized, osteoporosis, and fibromyalgia); (b) studies involving within-participant design; (c) interventions involving any other type of exercise component (eg, aerobic exercise and balance exercise); (d) interventions combining hybrid traditional and high-velocity resistance exercise simultaneously; (e) interventions involving nutrition components (eg, protein supplementation and caloric restriction); (f) studies with interventions lasting less than 4 weeks; (g) not reporting specific outcomes included in this review; and (h) studies written in a language other than English, Portuguese, or Spanish.

### Data Searches and Sources

A systematic search was conducted by a researcher (P.L.) using CINAHL, Embase, LILACS, PubMed, Scielo, SPORTDiscus, and Web of Science databases from inception to February 2022. The search strategy is presented in the [Supplementary Appendix 1](#). All procedures undertaken in the present study were reported in accordance with the Cochrane Back Review Group (CRBG) (43), the Implementing Prisma in Exercise, Rehabilitation, Sport medicine and Sports science (PERSiST) (44), and the Preferred Reporting Items for Systematic Reviews and Meta-Analyses for Network Meta-analyses (PRISMA-NMA) statement (45) with registration at the international prospective register of systematic reviewers (PROSPERO identifier: CRD42022297254).

### Study Selection Process and Data Extraction

In the search strategy, titles and abstracts were first independently evaluated following the eligibility criteria. Eligibility was assessed independently in duplicate (P.L. and D.J.P.T.), with differences resolved by consensus. When abstracts did not provide sufficient information, they were selected for full-text evaluation. Full-text articles meeting criteria were retrieved and read independently by both reviewers and assessed for inclusion in the study. In addition, a manual search of references in selected studies was performed to detect studies potentially eligible for inclusion.

One reviewer (P.L.) extracted data using a standardized form, while a second reviewer (D.J.P.T.) checked the information. Relevant information extracted from studies included publication information (ie, authors and year of publication), demographic and clinical characteristics such as age, body mass index (BMI), and health status (eg, self-reported physical disability, mobility-limited, sarcopenia, and frailty), experimental design and sample size, prescription characteristics such as delivery, modality, frequency, volume, intensity, and concentric and eccentric time of contraction when available.

## End Points

The primary outcome for this review was objectively assessed physical function measured by: (a) fast walking speed (a measure of gait speed), (b) timed-up and go test (ie, time in seconds to rise from the chair, walk a distance, turn, walk back to the chair, and sit down; a measure of lower-limb function and mobility), (c) 5-times sit-to-stand test (ie, time in seconds to rise as fast as possible from the chair 5 times; a measure of lower-limb power and strength), (d) 30-second sit-to-stand test (ie, number of times that a participant can rise to a full stand from a seated position within 30 seconds; a measure of lower-limb strength and endurance), and (e) 6-minute walking test (ie, distance covered in meters when participants were instructed to walk as far as they could during 6 minutes; a measure of cardiorespiratory fitness and walking endurance). Secondary outcomes included maximal muscle power measured by a multijoint lower-limb exercise or task test (eg, leg press exercise, stair climbing, countermovement jump, and sit-to-stand test) and expressed as  $W$  or  $W \cdot kg^{-1}$ , and maximal muscle strength measured by the leg press 1-repetition maximum test (1-RM). Information was extracted from baseline versus immediate postintervention assessment. For the outcomes assessed, when articles did not include dispersion values of change such as standard deviation ( $SD$ ), standard errors, or 95% confidence intervals (95% CI), the  $SD$  of the change was calculated assuming a correlation of  $r = 0.5$  between the baseline and postintervention assessment measures by the square root of (46) ( $(SD_{Baseline}^2 + SD_{Post-intervention}^2) - (2 \times r \times SD_{Baseline} \times SD_{Post-intervention})$ ).

## Study Risk of Bias Assessment and Certainty of Evidence (GRADE)

The risk of bias was evaluated according to the Cochrane risk-of-bias tool 2.0 (RoB 2) (47), with each assessment focused on the outcome level. The study quality assessment for all included studies was performed independently by 2 reviewers (P.L. and D.J.P.T.), with disagreements resolved by consensus. The certainty of the evidence for the network of interventions was assessed using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach for network meta-analysis (48). The several domains considered for downgrading the certainty of evidence were study limitations (ie, based on the risk of bias assessment), indirectness (ie, based on transitivity assumption), inconsistency (ie, based on inconsistency assumption), imprecision (ie, uncertainty around the pairwise estimate with confidence intervals crossing the null value or including values favoring different treatments), and publication bias (ie, based on comparison-adjusted funnel plot and the Egger's test) (48).

## Data Synthesis and Analysis

Continuous outcome data in both pairwise and network meta-analyses (NMA) were summarized as standardized mean differences (SMD) and their 95% confidence intervals (95% CI). Pairwise comparisons between high-velocity resistance exercise, traditional

resistance exercise, and control conditions were conducted in R (R Development Core Team, Vienna, Austria) using the package "meta" (49). The frequentist graph theoretical was performed following the current PRISMA guideline for NMA (50,51), and it was conducted using the R package "netmeta" (52). A random-effects model was undertaken as studies differ both clinically and methodologically (ie, between-study variability). The between-study variability (ie, heterogeneity) of the intervention effects within each intervention comparison was assessed by  $I^2$ , and the magnitude of the between-study variance ( $\tau^2$ ) and estimated using the generalized DerSimonian and Laird estimator and the Q-profile approach. For each NMA, we assessed a priori the transitivity and consistency assumptions. Statistical assessment of the transitivity assumption implies that the distribution of potential treatment effect modifiers is balanced across the available direct comparisons (53). We used the percentage of male participants, average age, BMI, health status, and overall risk of bias as potential intervention effect modifiers, with values reported for each study. We evaluated each network for inconsistency using the random-effects design-by-treatment interaction model (54) and locally by splitting the direct and indirect evidence (55).

Intervention effects were ranked according to  $p$  scores, measuring the extent of certainty that an intervention is better than the other (56). Comparisons were made when more than 1 study was included for each comparator and were considered statistically significant when the 95% CI did not include the value of zero effect. According to Cohen (57), SMD values of 0.0 to  $\leq 0.5$  indicate small; 0.51 to 0.79, medium; and  $\geq 0.8$ , large effects. Extreme-study effects (ie, outliers) were explored with the forward search algorithm (58) using the R package "NMAoutlier" (59). When obvious outliers were detected, these were excluded in a sensitivity analysis to assess the robustness of the results. NMA with subsets and sensitivity analyses were conducted for the primary outcome with consideration for potential intervention effect modifiers, health status, and overall risk of bias. When the number of studies was more than 10, a comparison-adjusted funnel plot was drawn to assess publication bias and small-study effects.

## Results

Following the deletion of duplicates and records marked as ineligible, a total of 2 375 potential records were identified. Of these, 1 333 records were excluded based on titles and abstracts due to their irrelevance to the research question, and 14 reports were not available for full-text assessment, resulting in 1 028 records deemed eligible for full-text review. A total of 955 reports were excluded due to reasons described in Supplementary Figure 1. After including 7 additional studies (30,60–65) via reference lists, a total of 80 articles (29–31,60–136) describing 79 randomized trials were included in the present systematic review and network meta-analysis (fast walking speed,  $n = 12$ ; timed-up and go,  $n = 40$ ; 5-times sit-to-stand test,  $n = 18$ ; 30-second sit-to-stand test,  $n = 24$ ; 6-minute walking test,  $n = 15$ ; leg press muscle power [expressed in  $W$ ],  $n = 10$ ; leg press muscle power [expressed in  $W \cdot kg^{-1}$ ],  $n = 4$ ; stair climbing muscle power,  $n = 3$ ; sit-to-stand muscle power,  $n = 3$ ; countermovement jump muscle power,  $n = 3$ ; and leg press muscle strength,  $n = 24$ ).

## Study, Participant, and Intervention Characteristics

A total of 3 575 participants (women,  $n = 2 491$ ; men,  $n = 836$ ; not reported,  $n = 248$ ) with a median age of 70.2 years (interquartile range [IQR] = 67.3 to 72.6 yrs) and BMI of  $27.6 \text{ kg} \cdot \text{m}^{-2}$  (IQR = 25.6 to  $28.6 \text{ kg} \cdot \text{m}^{-2}$ ) were included. Most studies included physically

healthy participants (55 out of 79 studies, 69.2%), followed by participants with a self-reported disability or mobility limitations (8 out of 79 studies, 10.1%), sarcopenia (6 out of 79 studies, 7.6%), and frailty (5 out of 79 studies, 6.3%). Five studies included both physically healthy and participants with some physical disability (6.3%).

From the 80 articles included, a total of 101 interventions were analyzed. While 31 interventions examined high-velocity resistance exercise (30.7%), 70 examined traditional resistance exercise (69.3%). Regarding exercise prescription characteristics, the median intervention duration of studies evaluating high-velocity resistance exercise was 12 weeks (IQR: 12–14 weeks), with a median of 36 sessions (IQR: 24–36 sessions). Information about high-velocity resistance exercise volume was reported for 29 out of the 31 interventions (93.5%) with a median of 42 sets per week (IQR: 32–61 sets per week), while peak intensity was reported for 25 out of the 31 interventions (80.6%) with a median of 70% of 1-RM (IQR: 50%–75% of 1-RM). For studies examining traditional resistance exercise, the median duration was 12 weeks (IQR: 12–16 weeks) and 36 sessions (IQR: 24–38 sessions). Information about volume was reported for 60 out of the 70 interventions (85.7%) with a median of 50 sets per week (IQR: 36–72 sets per week), while intensity was reported for 50 out of the 70 interventions with a median of 75% of 1-RM (IQR: 70%–80% of 1-RM). Sixty-seven out of the 79 studies included were randomized controlled trials. Studies were designed to compare resistance exercise programs versus nonactive controls (53 out of 67 studies, 79.1%), stretching control (5 out of 67 studies, 7.5%), health education control (5 out of 67 studies, 7.5%), walking control (2 out of 67 studies, 3.0%), and nutrition placebo control (2 out of 67 studies, 3.0%). The characteristics of the individual studies are presented in [Supplementary Table 1](#).


### Risk of Bias Assessment

A total of 57 studies examined the effects of resistance exercise velocity modes on physical function, while 20 and 24 studies examined muscle power and muscle strength, respectively. For the primary outcome of this review, 70.2% of the studies presented a *high risk of bias* (40 out of 57 studies), and 26.3% of the studies presented *some concerns* in physical function assessment (15 out of 57 studies). For the secondary outcomes, 70.0% of the studies presented a *high risk of bias* (14 out of 20 studies) in muscle power assessment, while 79.2% of the studies presented a *high risk of bias* (19 out of 24 studies) in muscle strength assessment. The individual risk of bias assessment is presented in [Supplementary Tables 2–4](#).

### Analysis of the Outcomes

#### Physical function outcomes

All pairwise comparison results estimated in the meta-analysis model are reported in [Supplementary Tables 5–9](#). The network geometry of studies examining physical function outcomes is presented in [Figure 1](#). High-velocity resistance exercise was the most effective for improving fast walking speed ( $p$  score = 92.8%) and reducing the time to perform the timed-up and go ( $p$  score = 89.5%) and the 5-times sit-to-stand test ( $p$  score = 82.1%) compared to controls ([Table 1](#)). Traditional resistance exercise was the most effective intervention for improving 30-second sit-to-stand ( $p$  score = 85.1%) and 6-minute walking test ( $p$  score = 79.1%; [Table 1](#)). Statistically significant differences were not observed between traditional and high-velocity resistance exercise for physical function outcomes ( $p$  = .239–.837).



**Figure 1.** Network geometry of studies examining (A) fast walking speed, (B) timed-up and go, (C) 5-times sit-to-stand, (D) 30-second sit-to-stand, and (E) 6-minute walking.  $k$ , number of comparisons;  $n$ , sample size.

Subgroup analyses for physical health status are presented in [Figures 2](#) and [3](#). High-velocity resistance exercise resulted in significant improvements in fast walking speed (SMD = 1.28, 95% CI: 0.74 to 1.82), timed-up and go (SMD = -0.88, 95% CI: -1.21 to -0.54), 30-second sit-to-stand (SMD = 0.93, 95% CI: 0.43 to 1.43), and 6-minute walking test (SMD = 0.74, 95% CI: 0.11 to 1.38) for those physically healthy but not 5-times sit-to-stand where significant effects were only observed in physically impaired participants (SMD = -1.26, 95% CI: -2.20 to -0.32). In traditional resistance exercise, significant effects were observed regardless of physical health status on timed-up and go (SMD = -0.74 to -0.62), 5-times sit-to-stand (SMD = -1.09 to -0.52), and 30-second sit-to-stand (SMD = 0.77 to 1.10). For fast walking speed and 6-minute walking test, traditional resistance exercise resulted in significant effects for mixed physically healthy and physically impaired participants (SMD = 0.47, 95% CI: 0.10 to 0.84) and physically healthy participants (SMD = 0.76, 95% CI: 0.36 to 1.16), respectively.

The heterogeneity  $I^2$  ranged from 61.1% to 69.9% in NMA for the outcomes of physical function. The global test for inconsistency (ie, design-by-treatment interaction random-effects model) indicated inconsistency for fast walking speed ( $Q$  = 18.2,  $p$  < .001). The locally side-split analyses for fast walking speed indicated a source of inconsistency between direct and indirect evidence for high-velocity resistance exercise versus control ( $p$  = .049; [Supplementary Table 10](#)). Visual assessment of comparison-adjusted funnel plots suggested no evidence of small-study effects (Egger test,  $p$  = .137–.743), except 5-times sit-to-stand outcome (Egger test,  $p$  = .025; [Supplementary Figure 2](#)). There was no evidence of publication bias after omitting the outlier (Coelho-Junior & Uchida (64)) from 5-times sit-to-stand ( $p$  = .114). Outliers were detected with the forward search algorithm for fast walking speed (Ortega & Cuartas, 2020 (127)), 5-times sit-to-stand (Coelho-Junior & Uchida, 2021 (64)), 30-second sit-to-stand, and 6-minute walking test (Filho et al., 2022 (65); [Supplementary Figure 3](#)). After omitting outliers, sensitivity analyses results were not different from the primary analyses for 5-times sit-to-stand and 6-minute walking test. Sensitivity analysis for fast walking speed provided traditional resistance exercise (SMD = 0.27, 95% CI: 0.01 to 0.52,  $p$  = .042;  $p$  score = 76.6%) as the best intervention modality, while heterogeneity reduced from 63.8% in the primary analysis to 40.2% ([Supplementary Table 11](#)). For 30-second sit-to-stand, we found high-velocity resistance exercise (SMD = 1.08, 95% CI: 0.66 to 1.49,  $p$  < .001;  $p$  score = 85.8%) as the best intervention modality

**Table 1.** Network Meta-Analysis Results for Physical Function Outcomes

Comparisons	<i>k</i>	SMD	95% CI	<i>p</i> -Value	<i>I</i> <sup>2</sup>	<i>p</i> -Score	Certainty
<b>Fast walking speed</b>							
HVRE vs Control groups	5	0.44	0.00 to 0.87	.048	63.8%	HVRE: 92.8%	⊕⊕⊕⊕
TRE vs Control groups	9	0.16	-0.16 to 0.48	.329		TRE: 47.8%	Very low <sup>*,†,‡</sup>
HVRE vs TRE	3	0.28	-0.19 to 0.74	.239			
<b>Timed-up and go</b>							
HVRE vs Control groups	10	-0.76	-1.05 to -0.47	<.001	61.1%	HVRE: 89.5%	⊕⊕⊕⊕
TRE vs Control groups	32	-0.64	-0.83 to -0.45	<.001		TRE: 60.6%	Low <sup>*,‡</sup>
HVRE vs TRE	11	-0.11	-0.41 to 0.17	.422			
<b>Five-times sit-to-stand</b>							
HVRE vs Control groups	4	-0.74	-1.20 to -0.27	.002	63.1%	HVRE: 82.1%	⊕⊕⊕⊕
TRE vs Control groups	14	-0.66	-0.96 to -0.34	<.001		TRE: 67.9%	Very low <sup>*,‡,§</sup>
HVRE vs TRE	6	-0.08	-0.52 to 0.36	.716			
<b>30-Second sit-to-stand</b>							
HVRE vs Control groups	7	0.89	0.43 to 1.34	<.001	76.3%	TRE: 85.1%	⊕⊕⊕⊕
TRE vs Control groups	18	1.01	0.68 to 1.34	<.001		HVRE: 64.9%	Low <sup>*,‡</sup>
HVRE vs TRE	7	-0.12	-0.58 to 0.33	.597			
<b>6-Minute walking</b>							
HVRE vs Control groups	4	0.63	0.07 to 1.18	.027	69.9%	TRE: 79.1%	⊕⊕⊕⊕
TRE vs Control groups	14	0.68	0.34 to 1.03	<.001		HVRE: 70.3%	Low <sup>*,‡</sup>
HVRE vs TRE	5	-0.06	-0.59 to 0.48	.837			


Notes: 95% CI = 95% confidence interval; HVRE = high-velocity resistance exercise; *k* = number of comparisons; TRE = traditional resistance exercise; SMD = standardized mean difference.

\*Certainty of evidence downgraded due to study limitations, with studies most studies presenting with high risk in the risk of bias assessment.


†Certainty of evidence downgraded due to inconsistency, with a source of inconsistency between direct and indirect evidence identified.

‡Certainty of evidence downgraded due to imprecision, with confidence intervals from interventions crossing null values or including values favoring both interventions tested.

§Certainty of evidence downgraded due to publication bias, with visual assessment of comparison-adjusted funnel plots suggesting evidence of small-study effects.



**Figure 2.** Network meta-analysis on (A) fast walking speed, (B) 5-times sit-to-stand, (C) 30-second sit-to-stand, and (D) 6-minute walking test based on health status. White squares indicate studies involving physically impaired participants; gray squares indicate studies involving mixed participants; black squares indicate studies involving physically healthy participants. CI = confidence interval; HVRE = high-velocity resistance exercise; TRE = traditional resistance exercise.



**Figure 3.** Network meta-analysis on timed-up and go based on health status. White squares indicate studies involving physically impaired participants; gray squares indicate studies involving mixed participants; black squares indicate studies involving physically healthy participants. CI = confidence interval; HVRE = high-velocity resistance exercise; TRE = traditional resistance exercise.

with a small reduction of heterogeneity (Supplementary Table 11). Regarding the certainty of the evidence, the ranking of interventions for improving physical function outcome was graded *low* for timed-up and go, 30-second sit-to-stand, and 6-minute walking tests, while a *very low* grade was observed for fast walking speed and 5-times sit-to-stand (Table 1).

**Muscle power and muscle strength outcomes**

Pairwise meta-analysis models for muscle power outcomes and muscle strength are presented in Supplementary Tables 12–17. The network geometry of studies examining muscle power and muscle strength outcomes is presented in Supplementary Figure 4. High-velocity resistance exercise was the most effective for improving

leg press muscle power (expressed in W) compared to traditional resistance exercise and control condition ( $p$  score = 99.9%; Table 2). For leg press muscle strength, traditional resistance exercise was the most effective intervention compared to the control ( $p$  score = 86.6%; Table 2). There was a statistical difference between traditional and high-velocity resistance exercises for leg press muscle power ( $p = .003$ ) but not for leg press muscle strength ( $p = .538$ ). NMA were not undertaken for leg press muscle power (expressed in  $W \cdot kg^{-1}$ ), stair climbing, sit-to-stand, and countermovement jump muscle power, given the small number of studies for each comparator ( $\leq 1$ ).

Subset analyses for physical health status are presented in Supplementary Figures 5 and 6. Both traditional and high-velocity resistance exercise resulted in significant improvements in leg press muscle power (expressed in W) for those physically healthy (SMD = 0.61 to 0.99) and mixed physically healthy and physically impaired participants (SMD = 0.78 to 1.03), but not for those physically impaired. Improvements in leg press muscle strength were only observed in physically healthy participants following high-velocity resistance exercise (SMD = 1.12, 95% CI: 0.75 to 1.50), while traditional resistance exercise provided significant results for leg press muscle strength regardless of physical health status (SMD = 0.95 to 1.22).

The heterogeneity  $I^2$  was 49.6% and 58.6% for leg press muscle power and muscle strength, respectively. The global test for inconsistency was not significant for leg press muscle power ( $Q = 4.59$ ,  $p = .205$ ) and muscle strength ( $Q = 1.72$ ,  $p = .633$ ), and differences between direct and indirect evidence were not observed in locally side-split analyses (Supplementary Table 18). Visual assessment of comparison-adjusted funnel plots suggested no evidence of small-study effects for leg press muscle power (Egger test,  $p = .970$ ) but did for muscle strength (Egger test,  $p < .001$ ; Supplementary Figure 7). Outliers were detected with the forward search algorithm for leg press muscle power (Fielding et al. (29)) and muscle strength (Filho et al. (65); Supplementary Figure 8). After omitting outliers, sensitivity analyses results were not different from the primary analyses (Supplementary Table 19). Heterogeneity was reduced from 49.6% in the primary analysis to 0% in leg press muscle power, while a small reduction was observed in leg press muscle strength (Supplementary Table 19). Based on the GRADE approach, the certainty of evidence was considered *moderate* for leg press muscle power and *very low* for leg press muscle strength (Table 2).

**Table 2.** Network Meta-Analysis Results for Muscle Strength and Muscle Power Outcomes

Comparisons	<i>k</i>	SMD	95% CI	<i>p</i> -Value	$I^2$	<i>p</i> -Score	Certainty
<b>Leg press muscle power</b>							
HVRE vs Control groups	4	0.90	0.49 to 1.30	<.001	49.6%	HVRE: 99.9%	⊕⊕⊕⊕
TRE vs Control groups	5	0.35	-0.04 to 0.73	.080		TRE: 48.1%	Moderate*
HVRE vs TRE	7	0.55	0.19 to 0.92	.003			
<b>Leg press 1-RM</b>							
HVRE vs Control groups	8	1.10	0.75 to 1.46	<.001	58.6%	TRE: 86.6%	⊕⊕⊕⊕
TRE vs Control groups	23	1.21	0.95 to 1.46	<.001		HVRE: 63.4%	Very low* <sup>†</sup> ‡
HVRE vs TRE	11	-0.10	-0.44 to 0.23	.538			

Notes: 1-RM = 1-repetition maximum; 95% CI = 95% confidence interval; HVRE = high-velocity resistance exercise; *k* = number of comparisons; SMD = standardized mean difference; TRE = traditional resistance exercise; W = Watts.

\*Certainty of evidence downgraded due to imprecision, with confidence intervals from interventions crossing null values or including values favoring both interventions tested.

<sup>†</sup>Certainty of evidence downgraded due to study limitations, with studies most studies presenting with high risk in the risk of bias assessment.

<sup>‡</sup>Certainty of evidence downgraded due to publication bias, with visual assessment of comparison-adjusted funnel plots suggesting evidence of small-study effects.



## Discussion

In the present systematic review with network meta-analysis, we investigated the effects of high-velocity and traditional resistance exercise on physical function, muscle power, and strength outcomes in physically healthy and impaired older adults. Although both resistance exercise velocities effectively improved performance-based physical function outcomes in these participants, we found that the neuromuscular changes derived are velocity specific, as evidenced by physical function test dependence. While high-velocity resistance exercise provided better results for fast walking speed, 5-times sit-to-stand, timed-up and go, and muscle power outcomes, traditional resistance exercise resulted in greater effects on 30-second sit-to-stand, 6-minute walking, and muscle strength. These findings indicate that resistance exercise prescription based on the velocity of contraction should be individualized in older adults to optimize improvements in the specific physical function outcomes desired.

It is well recognized that resistance exercise is an efficient intervention to improve physical function in older adults (17–20,23,24). However, there is a lack of information on the most effective resistance exercise mode. In the current study, we found that resistance exercise velocity effects are specific and depend on the characteristics the test is quantifying. Attempting to achieve a high execution velocity resulted in greater improvements in physical function tests with a time component, while endurance tasks, where participants had to work longer, derived superior benefits from traditional low-to-moderate-velocity movement. This difference may be related to the specific neuromuscular adaptations promoted by fast or slow controlled contractions within resistance exercise (137). Therefore, we suggest that resistance exercise velocity should be individualized to specifically address deficits in older adults' neuromuscular performance per the concept of *one-size-does-not-fit-all*.

Older adults with a higher risk of disabilities, such as those suffering from frailty or mobility limitations, are vulnerable to stressors and adverse outcomes (13). Given reductions in a physical reserve capacity, interventions targeting basic tasks of daily living such as walking, change of direction, balance, and standing up from a chair are critical in these individuals. Therefore, older adults who are physically impaired could especially benefit from targeted high-velocity resistance exercise. This may provide a greater safety margin before the threshold for disabilities is reached and assist in reducing the risk of falls and hospitalizations (28,138,139). Subsequently, goals such as improving muscular endurance could be prioritized by introducing traditional resistance exercises or even different exercise components such as aerobic and balance exercises within a multi-modal exercise program (23,24).

In regards to physically healthy older adults, this group can be targeted based on initial assessment, preventing physical disability and maintaining healthy aging (16). For example, participants who perform poorly on 30-second sit-to-stand or 6-minute walking tests can be targeted with traditional resistance exercise, the most effective intervention for these outcomes. Alternatively, hybrid programs comprising both velocity modes can be used to develop different components of physical function simultaneously (140), although further investigations are required to elucidate the response to this model, including within an overall periodized program. Therefore, individualizing the resistance exercise program using high-velocity or traditional resistance exercises can target participants at different stages, such as reversing physical disability or preserving physical function before the threshold for disabilities.

The strengths of the present study are: (a) the inclusion of 79 resistance exercise randomized trials with 3 575 older adults; (b) studies examining resistance exercise modes without the interference of other exercise components such as aerobic or balance interventions; (c) a network meta-analysis involving simultaneous comparison among high-velocity and traditional resistance exercise and control conditions; (d) investigation of the most effective resistance exercise velocity mode for different physical function tests; and (e) subgroup analyses providing specific information for physically healthy or impaired older adults. However, there are also limitations worthy of comment. First, the majority of studies had a high risk of bias. Several concerns regarding the *randomization process*, *measurement of outcomes*, and *selection of reported results* were identified. These affected the precision, magnitude, and certainty of the evidence. Second, we observed a high heterogeneity across analyses on physical function. This may be a result of different settings or characteristics of studies included, in addition to the lack of or poor reporting of resistance exercise characteristics such as volume and intensity in the studies included. Third, participants in the high-velocity resistance exercise group likely had their performance affected by accumulated fatigue throughout sessions, sets and repetitions, reducing their ability to recruit all motor units at optimal discharge rates and maintain appropriate velocity (141). As a result, this may have reduced our ability to observe even greater differences between resistance exercise velocities. Fourth, we did not include other commonly used physical function tests, such as the usual walking speed or the 400-m walk. Given the availability of studies assessing performance measures or sensitivity to change following resistance exercise, this systematic review and meta-analysis were planned to focus on those with a higher number of studies available (eg, 6-minute walking test vs 400-m walk test) and higher sensitivity to change following resistance exercise programs (eg, fast walking speed and timed-up and go vs usual walking speed) in older adults (37,142). Finally, only a few studies were conducted on patients who were physically impaired, and these may have limited our findings for this group.

In conclusion, our study provides evidence that resistance exercise effects on physical function are velocity specific, as evidenced by physical function test dependence in older adults. While high-velocity resistance exercise promoted greater improvements in physical function tests with a time component, traditional resistance exercise was the most effective intervention for improving performance where participants had to work longer. These results are of clinical importance as they indicate that resistance exercise prescription based on the velocity of contraction should be individualized and specific to target the relative deficits of participants' and their needs within the resistance exercise program. Moreover, older adults will often present with a range of deficits in physical function and consequently, both high-velocity and traditional resistance exercise may be required to enhance multiple domains of physical function.

## Supplementary Material

Supplementary data are available at *The Journals of Gerontology, Series A: Biological Sciences and Medical Sciences* online.

## Funding

P.L. is supported by the National Health and Medical Research Council (NHMRC) Centre of Research Excellence (CRE) in Prostate Cancer Survivorship Scholarship. D.A.G. and R.U.N. are funded by a NHMRC CRE

in Prostate Cancer Survivorship. The results of the study are presented clearly, honestly, without fabrication, falsification, or inappropriate data manipulation.

Sponsors were not involved in the study design, analysis or interpretation of data, manuscript writing and decision to submit the manuscript for publication.

No financial support was received to conduct the present study, or for the preparation or publication of this manuscript.

## Conflict of Interest

None declared.

## Author Contributions

P.L. had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis; Conception and design: P.L., A.R., and R.R.; Acquisition, analysis, or interpretation of data: P.L., A.R., M.P., R.U.N., D.R.T., D.A.G., D.J.P.T., S.R.F., and R.R.; Drafting of the manuscript: P.L., A.R., M.P., R.U.N., D.R.T., D.A.G., D.J.P.T., S.R.F., and R.R.; Critical revision of the manuscript for important intellectual content: P.L., A.R., M.P., R.U.N., D.R.T., D.A.G., D.J.P.T., S.R.F., and R.R.; Statistical analysis: P.L. and M.P.

## Data Availability

Data used in the present study such as data extraction templates, forms, and analysis will be made available upon request to the corresponding author.

## References

- Rosenberg IH. Sarcopenia: origins and clinical relevance. *J Nutr*. 1997;127(5 Suppl):990s–1s. doi:10.1093/jn/127.5.990S
- Klass M, Baudry S, Duchateau J. Voluntary activation during maximal contraction with advancing age: a brief review. *Eur J Appl Physiol*. 2007;100(5):543–551. doi:10.1007/s00421-006-0205-x
- Lexell J, Taylor CC, Sjöström M. What is the cause of the ageing atrophy? Total number, size and proportion of different fiber types studied in whole vastus lateralis muscle from 15- to 83-year-old men. *J Neurol Sci*. 1988;84(2–3):275–294. doi:10.1016/0022-510x(88)90132-3
- Frontera WR, Hughes VA, Fielding RA, Fiatarone MA, Evans WJ, Roubenoff R. Aging of skeletal muscle: a 12-yr longitudinal study. *J Appl Physiol*. 2000;88(4):1321–1326. doi:10.1152/jappl.2000.88.4.1321
- Goodpaster BH, Park SW, Harris TB, et al. The loss of skeletal muscle strength, mass, and quality in older adults: the health, aging and body composition study. *J Gerontol A Biol Sci Med Sci*. 2006;61(10):1059–1064. doi:10.1093/gerona/61.10.1059
- Lynch NA, Metter EJ, Lindle RS, et al. Muscle quality. I. Age-associated differences between arm and leg muscle groups. *J Appl Physiol*. 1999;86(1):188–194. doi:10.1152/jappl.1999.86.1.188
- Delmonico MJ, Harris TB, Visser M, et al.; Health, Aging, and Body. Longitudinal study of muscle strength, quality, and adipose tissue infiltration. *Am J Clin Nutr*. 2009;90(6):1579–1585. doi:10.3945/ajcn.2009.28047
- Bean JF, Kiely DK, Herman S, et al. The relationship between leg power and physical performance in mobility-limited older people. *J Am Geriatr Soc*. 2002;50(3):461–467. doi:10.1046/j.1532-5415.2002.50111.x
- Foldvari M, Clark M, Laviolette LC, et al. Association of muscle power with functional status in community-dwelling elderly women. *J Gerontol A Biol Sci Med Sci*. 2000;55(4):M192–M199. doi:10.1093/gerona/55.4.m192
- Seeman TE, Merkin SS, Crimmins EM, Karlamangla AS. Disability trends among older Americans: National Health And Nutrition Examination Surveys, 1988–1994 and 1999–2004. *Am J Public Health*. 2010;100(1):100–107. doi:10.2105/AJPH.2008.157388
- Tinetti ME, Speechley M, Ginter SF. Risk factors for falls among elderly persons living in the community. *N Engl J Med*. 1988;319(26):1701–1707. doi:10.1056/NEJM198812293192604
- Campbell AJ, Borrie MJ, Spears GF, Jackson SL, Brown JS, Fitzgerald JL. Circumstances and consequences of falls experienced by a community population 70 years and over during a prospective study. *Age Ageing*. 1990;19(2):136–141. doi:10.1093/ageing/19.2.136
- Fried LP, Tangen CM, Walston J, et al.; Cardiovascular Health Study Collaborative Research Group. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*. 2001;56(3):M146–M156. doi:10.1093/gerona/56.3.m146
- Cawthon PM, Fox KM, Gandra SR, et al.; Health, Aging and Body Composition Study. Do muscle mass, muscle density, strength, and physical function similarly influence risk of hospitalization in older adults? *J Am Geriatr Soc*. 2009;57(8):1411–1419. doi:10.1111/j.1532-5415.2009.02366.x
- Hanlon P, Nicholl BI, Jani BD, Lee D, McQueenie R, Mair FS. Frailty and pre-frailty in middle-aged and older adults and its association with multimorbidity and mortality: a prospective analysis of 493 737 UK Biobank participants. *Lancet Public Health*. 2018;3(7):e323–e332. doi:10.1016/S2468-2667(18)30091-4
- Beard JR, Officer A, de Carvalho IA, et al. The World report on ageing and health: a policy framework for healthy ageing. *Lancet*. 2016;387(10033):2145–2154. doi:10.1016/S0140-6736(15)00516-4
- Fiatarone MA, Marks EC, Ryan ND, Meredith CN, Lipsitz LA, Evans WJ. High-intensity strength training in nonagenarians. Effects on skeletal muscle. *JAMA*. 1990;263(22):3029–3034.
- American College of Sports M. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc*. 2009;41(3):687–708. doi:10.1249/MSS.0b013e3181915670
- Borde R, Hortobágyi T, Granacher U. Dose–response relationships of resistance training in healthy old adults: a systematic review and meta-analysis. *Sports Med*. 2015;45(12):1693–1720. doi:10.1007/s40279-015-0385-9
- Lopez P, Pinto RS, Radaelli R, et al. Benefits of resistance training in physically frail elderly: a systematic review. *Ageing Clin Exp Res*. 2018;30(8):889–899. doi:10.1007/s40520-017-0863-z
- Straight CR, Lindheimer JB, Brady AO, Dishman RK, Evans EM. Effects of resistance training on lower-extremity muscle power in middle-aged and older adults: a systematic review and meta-analysis of randomized controlled trials. *Sports Med*. 2016;46(3):353–364. doi:10.1007/s40279-015-0418-4
- Buchner DM, Larson EB, Wagner EH, Koepsell TD, de Lateur BJ. Evidence for a non-linear relationship between leg strength and gait speed. *Age Ageing*. 1996;25(5):386–391. doi:10.1093/ageing/25.5.386
- Fragala MS, Cadore EL, Dorgo S, et al. Resistance training for older adults: position statement from the national strength and conditioning association. *J Strength Cond Res*. 2019;33(8):2019–2052. doi:10.1519/JSC.0000000000003230
- Izquierdo M, Merchant RA, Morley JE, et al. International Exercise Recommendations in Older Adults (ICFSR): expert consensus guidelines. *J Nutr Health Aging*. 2021;25(7):824–853. doi:10.1007/s12603-021-1665-8
- Clafin DR, Larkin LM, Cederna PS, et al. Effects of high- and low-velocity resistance training on the contractile properties of skeletal muscle fibers from young and older humans. *J Appl Physiol*. 2011;111(4):1021–1030. doi:10.1152/japplphysiol.01119.2010
- Bassey EJ, Fiatarone MA, O'Neill EF, Kelly M, Evans WJ, Lipsitz LA. Leg extensor power and functional performance in very old men and women. *Clin Sci*. 1992;82(3):321–327. doi:10.1042/cs0820321
- Byrne C, Faure C, Keene DJ, Lamb SE. Ageing, muscle power and physical function: a systematic review and implications for pragmatic training interventions. *Sports Med*. 2016;46(9):1311–1332. doi:10.1007/s40279-016-0489-x
- Simpkins C, Yang F. Muscle power is more important than strength in preventing falls in community-dwelling older adults. *J Biomech*. 2022;134:111018. doi:10.1016/j.jbiomech.2022.111018
- Fielding RA, LeBrasseur NK, Cuoco A, Bean J, Mizer K, Fiatarone Singh MA. High-velocity resistance training increases skeletal

- muscle peak power in older women. *J Am Geriatr Soc.* 2002;50(4):655–662. doi:10.1046/j.1532-5415.2002.50159.x
30. Henwood TR, Taaffe DR. Short-term resistance training and the older adult: the effect of varied programmes for the enhancement of muscle strength and functional performance. *Clin Physiol Funct Imaging.* 2006;26(5):305–313. doi:10.1111/j.1475-097X.2006.00695.x
  31. Reid KF, Callahan DM, Carabello RJ, Phillips EM, Frontera WR, Fielding RA. Lower extremity power training in elderly subjects with mobility limitations: a randomized controlled trial. *Aging Clin Exp Res.* 2008;20(4):337–343. doi:10.1007/BF03324865
  32. Steib S, Schoene D, Pfeifer K. Dose–response relationship of resistance training in older adults: a meta-analysis. *Med Sci Sports Exerc.* 2010;42(5):902–914. doi:10.1249/MSS.0b013e3181c34465
  33. Tschopp M, Sattelmayer MK, Hilfiker R. Is power training or conventional resistance training better for function in elderly persons? A meta-analysis. *Age Ageing.* 2011;40(5):549–556. doi:10.1093/ageing/afr005
  34. da Rosa Orssatto LB, de la Rocha Freitas C, Shield AJ, Silveira Pinto R, Trajano GS. Effects of resistance training concentric velocity on older adults' functional capacity: a systematic review and meta-analysis of randomised trials. *Exp Gerontol.* 2019;127:110731. doi:10.1016/j.exger.2019.110731
  35. Balachandran AT, Steele J, Angielczyk D, et al. Comparison of power training vs traditional strength training on physical function in older adults: a systematic review and meta-analysis. *JAMA Netw Open.* 2022;5(5):e2211623. doi:10.1001/jamanetworkopen.2022.11623
  36. Rouse B, Chaimani A, Li T. Network meta-analysis: an introduction for clinicians. *Intern Emerg Med.* 2017;12(1):103–111. doi:10.1007/s11739-016-1583-7
  37. Sabia S, Dumurgier J, Tavernier B, Head J, Tzourio C, Elbaz A. Change in fast walking speed preceding death: results from a prospective longitudinal cohort study. *J Gerontol A Biol Sci Med Sci.* 2014;69(3):354–362. doi:10.1093/gerona/glt114
  38. Chun S, Shin DW, Han K, et al. The Timed Up and Go test and the ageing heart: findings from a national health screening of 1,084,875 community-dwelling older adults. *Eur J Prev Cardiol.* 2021;28(2):213–219. doi:10.1177/2047487319882118
  39. Karlsen T, Nauman J, Dalen H, Langhammer A, Wisløff U. The combined association of skeletal muscle strength and physical activity on mortality in older women: the HUNT2 study. *Mayo Clin Proc.* 2017;92(5):710–718. doi:10.1016/j.mayocp.2017.01.023
  40. Tveter AT, Dagfinrud H, Moseng T, Holm I. Health-related physical fitness measures: reference values and reference equations for use in clinical practice. *Arch Phys Med Rehabil.* 2014;95(7):1366–1373. doi:10.1016/j.apmr.2014.02.016
  41. Yazdanyar A, Aziz MM, Enright PL, et al. Association between 6-minute walk test and all-cause mortality, coronary heart disease-specific mortality, and incident coronary heart disease. *J Aging Health.* 2014;26(4):583–599. doi:10.1177/0898264314525665
  42. Cruz-Jentoft AJ, Bahat G, Bauer J, et al.; Writing Group for the European Working Group on Sarcopenia in Older People 2 (EWGSP2), and the Extended Group for EWGSP2. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing.* 2019;48(1):16–31. doi:10.1093/ageing/afy169
  43. Furlan AD, Pennick V, Bombardier C, van Tulder M; Editorial Board CBRG. 2009 updated method guidelines for systematic reviews in the Cochrane Back Review Group. *Spine.* 2009;34(18):1929–1941. doi:10.1097/BRS.0b013e3181b1c99f
  44. Arden CL, Büttner F, Andrade R, et al. Implementing the 27 PRISMA 2020 Statement items for systematic reviews in the sport and exercise medicine, musculoskeletal rehabilitation and sports science fields: the PERSiST (implementing Prisma in Exercise, Rehabilitation, Sport medicine and Sports science) guidance. *Br J Sports Med.* 2022;56(4):175–195. doi:10.1136/bjsports-2021-103987
  45. Hutton B, Salanti G, Caldwell DM, et al. The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Ann Intern Med.* 2015;162(11):777–784. doi:10.7326/M14-2385
  46. Lefebvre C, Manheimer E, Glanville J, Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 (updated March 2011)*. The Cochrane Collaboration, 2011. Available from [www.cochrane-handbook.org](http://www.cochrane-handbook.org).
  47. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *BMJ.* 2019;366:l4898. doi:10.1136/bmj.l4898
  48. Salanti G, Del Giovane C, Chaimani A, Caldwell DM, Higgins JP. Evaluating the quality of evidence from a network meta-analysis. *PLoS One.* 2014;9(7):e99682. doi:10.1371/journal.pone.0099682
  49. Balduzzi S, Rucker G, Schwarzer G. How to perform a meta-analysis with R: a practical tutorial. *Evid Based Ment Health.* 2019;22(4):153–160. doi:10.1136/ebmental-2019-300117
  50. Rucker G. Network meta-analysis, electrical networks and graph theory. *Res Synth Methods.* 2012;3(4):312–324. doi:10.1002/jrsm.1058
  51. Shim SR, Kim SJ, Lee J, Rucker G. Network meta-analysis: application and practice using R software. *Epidemiol Health.* 2019;41:e2019013. doi:10.4178/epih.e2019013
  52. Rucker G, Krahn U, König J, et al. *netmeta: Network Meta-Analysis using Frequentist Methods. R package version 2.1-1*. 2022.
  53. Jansen JP, Naci H. Is network meta-analysis as valid as standard pairwise meta-analysis? It all depends on the distribution of effect modifiers. *BMC Med.* 2013;11:159. doi:10.1186/1741-7015-11-159
  54. Dias S, Welton NJ, Caldwell DM, Ades AE. Checking consistency in mixed treatment comparison meta-analysis. *Stat Med.* 2010;29(7–8):932–944. doi:10.1002/sim.3767
  55. König J, Krahn U, Binder H. Visualizing the flow of evidence in network meta-analysis and characterizing mixed treatment comparisons. *Stat Med.* 2013;32(30):5414–5429. doi:10.1002/sim.6001
  56. Rucker G, Schwarzer G. Ranking treatments in frequentist network meta-analysis works without resampling methods. *BMC Med Res Methodol.* 2015;15:58. doi:10.1186/s12874-015-0060-8
  57. Cohen J. Statistical power analysis. *Curr Dir Psychol Sci.* 1992;1(3):98–101. doi:10.1111/1467-8721.ep10768783
  58. Petropoulou M, Salanti G, Rucker G, Schwarzer G, Moustaki I, Mavridis D. A forward search algorithm for detecting extreme study effects in network meta-analysis. *Stat Med.* 2021;40(25):5642–5656. doi:10.1002/sim.9145
  59. Petropoulou M, Schwarzer G, Panos A, Mavridis D. *NMAoutlier: Detecting Outliers in Network Meta-Analysis. R package version 0.1.18*. 2021.
  60. Charette SL, McEvoy L, Pyka G, et al. Muscle hypertrophy response to resistance training in older women. *J Appl Physiol.* 1991;70(5):1912–1916. doi:10.1152/jappl.1991.70.5.1912
  61. Tiggemann CL, Dias CP, Radaelli R, et al. Effect of traditional resistance and power training using rated perceived exertion for enhancement of muscle strength, power, and functional performance. *Age.* 2016;38(2):42. doi:10.1007/s11357-016-9904-3
  62. Fernández-Lezaun E, Schumann M, Mäkinen T, Kyröläinen H, Walker S. Effects of resistance training frequency on cardiorespiratory fitness in older men and women during intervention and follow-up. *Exp Gerontol.* 2017;95:44–53. doi:10.1016/j.exger.2017.05.012
  63. Lopez P, Crosby BJ, Robetti BP, et al. Effects of an 8-week resistance training intervention on plantar flexor muscle quality and functional capacity in older women: a randomised controlled trial. *Exp Gerontol.* 2020;138:111003. doi:10.1016/j.exger.2020.111003
  64. Coelho-Júnior HJ, Uchida MC. Effects of low-speed and high-speed resistance training programs on frailty status, physical performance, cognitive function, and blood pressure in prefrail and frail older adults. *Front Med.* 2021;8:702436. doi:10.3389/fmed.2021.702436
  65. Filho MM, Venturini G, Moreira OC, et al. Effects of different types of resistance training and detraining on functional capacity, muscle strength, and power in older women: a randomized controlled study. *J Strength Cond Res.* 2022;36(4):984–990. doi:10.1519/jsc.0000000000004195
  66. Judge JO, Whipple RH, Wolfson LI. Effects of resistive and balance exercises on isokinetic strength in older persons. *J Am Geriatr Soc.* 1994;42(9):937–946. doi:10.1111/j.1532-5415.1994.tb06584.x

67. McCartney N, Hicks AL, Martin J, Webber CE. Long-term resistance training in the elderly: effects on dynamic strength, exercise capacity, muscle, and bone. *J Gerontol A Biol Sci Med Sci*. 1995;50(2):B97–B104. doi:10.1093/gerona/50a.2.b97
68. Sipilä S, Multanen J, Kallinen M, Era P, Suominen H. Effects of strength and endurance training on isometric muscle strength and walking speed in elderly women. *Acta Physiol Scand*. 1996;156(4):457–464. doi:10.1046/j.1365-201X.1996.461177000.x
69. Taaffe DR, Duret C, Wheeler S, Marcus R. Once-weekly resistance exercise improves muscle strength and neuromuscular performance in older adults. *J Am Geriatr Soc*. 1999;47(10):1208–1214. doi:10.1111/j.1532-5415.1999.tb05201.x
70. Haykowsky M, Humen D, Teo K, et al. Effects of 16 weeks of resistance training on left ventricular morphology and systolic function in healthy men >60 years of age. *Am J Cardiol*. 2000;85(8):1002–1006. doi:10.1016/s0002-9149(99)00918-2
71. Rhodes EC, Martin AD, Taunton JE, Donnelly M, Warren J, Elliot J. Effects of one year of resistance training on the relation between muscular strength and bone density in elderly women. *Br J Sports Med*. 2000;34(1):18–22. doi:10.1136/bjsm.34.1.18
72. Fatouros IG, Taxildaris K, Tokmakidis SP, et al. The effects of strength training, cardiovascular training and their combination on flexibility of inactive older adults. *Int J Sports Med*. 2002;23(2):112–119. doi:10.1055/s-2002-20130
73. Miszko TA, Cress ME, Slade JM, Covey CJ, Agrawal SK, Doerr CE. Effect of strength and power training on physical function in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci*. 2003;58(2):171–175. doi:10.1093/gerona/58.2.m171
74. Bean JF, Herman S, Kiely DK, et al. Increased Velocity Exercise Specific to Task (InVEST) training: a pilot study exploring effects on leg power, balance, and mobility in community-dwelling older women. *J Am Geriatr Soc*. 2004;52(5):799–804. doi:10.1111/j.1532-5415.2004.52222.x
75. Boshuizen HC, Stemmerik L, Westhoff MH, Hopman-Rock M. The effects of physical therapists' guidance on improvement in a strength-training program for the frail elderly. *J Aging Phys Act*. 2005;13(1):5–22. doi:10.1123/japa.13.1.5
76. de Vos NJ, Singh NA, Ross DA, Stavrinou TM, Orr R, Fiatarone Singh MA. Optimal load for increasing muscle power during explosive resistance training in older adults. *J Gerontol A Biol Sci Med Sci*. 2005;60(5):638–647. doi:10.1093/gerona/60.5.638
77. de Vreede PL, Samson MM, van Meeteren NL, Duursma SA, Verhaar HJ. Functional-task exercise versus resistance strength exercise to improve daily function in older women: a randomized, controlled trial. *J Am Geriatr Soc*. 2005;53(1):2–10. doi:10.1111/j.1532-5415.2005.53003.x
78. Sousa N, Sampaio J. Effects of progressive strength training on the performance of the Functional Reach Test and the Timed Get-Up-and-Go Test in an elderly population from the rural north of Portugal. *Am J Hum Biol*. 2005;17(6):746–751. doi:10.1002/ajhb.20446
79. Bottaro M, Machado SN, Nogueira W, Scales R, Veloso J. Effect of high versus low-velocity resistance training on muscular fitness and functional performance in older men. *Eur J Appl Physiol*. 2007;99(3):257–264. doi:10.1007/s00421-006-0343-1
80. Fahlman M, Morgan A, McNeven N, Topp R, Boardley D. Combination training and resistance training as effective interventions to improve functioning in elders. *J Aging Phys Act*. 2007;15(2):195–205. doi:10.1123/japa.15.2.195
81. Caserotti P, Aagaard P, Larsen JB, Puggaard L. Explosive heavy-resistance training in old and very old adults: changes in rapid muscle force, strength and power. *Scand J Med Sci Sports*. 2008;18(6):773–782. doi:10.1111/j.1600-0838.2007.00732.x
82. Bean JF, Kiely DK, LaRose S, O'Neill E, Goldstein R, Frontera WR. Increased velocity exercise specific to task training versus the National Institute on Aging's strength training program: changes in limb power and mobility. *J Gerontol A Biol Sci Med Sci*. 2009;64(9):983–991. doi:10.1093/gerona/glp056
83. Marsh AP, Miller ME, Rejeski WJ, Hutton SL, Kritchevsky SB. Lower extremity muscle function after strength or power training in older adults. *J Aging Phys Act*. 2009;17(4):416–443. doi:10.1123/japa.17.4.416
84. Kalapotharakos VI, Diamantopoulos K, Tokmakidis SP. Effects of resistance training and detraining on muscle strength and functional performance of older adults aged 80 to 88 years. *Aging Clin Exp Res*. 2010;22(2):134–140. doi:10.1007/BF03324786
85. Lustosa LP, Silva JP, Coelho FM, Pereira DS, Parentoni AN, Pereira LS. Impact of resistance exercise program on functional capacity and muscular strength of knee extensor in pre-frail community-dwelling older women: a randomized crossover trial. *Rev Bras Fisioter*. 2011;15(4):318–324.
86. Correa CS, LaRoche DP, Cadore EL, et al. 3 Different types of strength training in older women. *Int J Sports Med*. 2012;33(12):962–969. doi:10.1055/s-0032-1312648
87. Pereira A, Izquierdo M, Silva AJ, et al. Effects of high-speed power training on functional capacity and muscle performance in older women. *Exp Gerontol*. 2012;47(3):250–255. doi:10.1016/j.exger.2011.12.010
88. Raj IS, Bird SR, Westfold BA, Shield AJ. Effects of eccentrically biased versus conventional weight training in older adults. *Med Sci Sports Exerc*. 2012;44(6):1167–1176. doi:10.1249/MSS.0b013e3182442ecd
89. Sayers SP, Gibson K. Effects of high-speed power training on muscle performance and braking speed in older adults. *J Aging Res*. 2012;2012:426278. doi:10.1155/2012/426278
90. Wallerstein LF, Tricoli V, Barroso R, et al. Effects of strength and power training on neuromuscular variables in older adults. *J Aging Phys Act*. 2012;20(2):171–185. doi:10.1123/japa.20.2.171
91. Campos ALP, Del Ponte LS, Cavalli AS, Afonso MR, Schild JFG, Reichert FF. Efeitos do treinamento concorrente sobre aspectos da saúde de idosos. *Rev Bras Cineantropometria Desempenho Hum*. 2013;15(4):437–447. doi:10.1590/S1413-35552011000400010
92. Solberg PA, Kvamme NH, Raastad T, et al. Effects of different types of exercise on muscle mass, strength, function and well-being in elderly. *Eur J Sport Sci*. 2013;13(1):112–125. doi:10.1080/17461391.2011.617391
93. Yu W, An C, Kang H. Effects of resistance exercise using thera-band on balance of elderly adults: a randomized controlled trial. *J Phys Ther Sci*. 2013;25(11):1471–1473. doi:10.1589/jpts.25.1471
94. Andersen TR, Schmidt JF, Nielsen JJ, et al. Effect of football or strength training on functional ability and physical performance in untrained old men. *Scand J Med Sci Sports*. 2014;24(suppl 1):76–85. doi:10.1111/sms.12245
95. Balachandran A, Krawczyk SN, Potiaumpai M, Signorile JF. High-speed circuit training vs hypertrophy training to improve physical function in sarcopenic obese adults: a randomized controlled trial. *Exp Gerontol*. 2014;60:64–71. doi:10.1016/j.exger.2014.09.016
96. Beltran Valls MR, Dimauro I, Brunelli A, et al. Explosive type of moderate-resistance training induces functional, cardiovascular, and molecular adaptations in the elderly. *Age*. 2014;36(2):759–772. doi:10.1007/s11357-013-9584-1
97. Fragala MS, Fukuda DH, Stout JR, et al. Muscle quality index improves with resistance exercise training in older adults. *Exp Gerontol*. 2014;53:1–6. doi:10.1016/j.exger.2014.01.027
98. Lopes PB, Pereira G, de Souza DM, Rodacki ALF. Comparison between strength and power training on elderly force-generating ability. *Top Geriatr Rehabil*. 2014;30(4):264–269. doi:10.1097/TGR.0000000000000033
99. Pamukoff DN, Haakonssen EC, Zaccaria JA, Madigan ML, Miller ME, Marsh AP. The effects of strength and power training on single-step balance recovery in older adults: a preliminary study. *Clin Interv Aging*. 2014;9:697–704. doi:10.2147/CIA.S59310
100. Pinto RS, Correa CS, Radaelli R, Cadore EL, Brown LE, Bottaro M. Short-term strength training improves muscle quality and functional capacity of elderly women. *Age*. 2014;36(1):365–372. doi:10.1007/s11357-013-9567-2
101. Ramirez-Campillo R, Castillo A, de la Fuente CI, et al. High-speed resistance training is more effective than low-speed resistance training to increase functional capacity and muscle performance in older women. *Exp Gerontol*. 2014;58:51–57. doi:10.1016/j.exger.2014.07.001
102. Uematsu A, Tsuchiya K, Kadono N, et al. A behavioral mechanism of how increases in leg strength improve old adults' gait speed. *PLoS One*. 2014;9(10):e110350. doi:10.1371/journal.pone.0110350

103. Emerson NS, Stout JR, Fukuda DH, et al. Resistance training improves capacity to delay neuromuscular fatigue in older adults. *Arch Gerontol Geriatr*. 2015;61(1):27–32. doi:10.1016/j.archger.2015.04.002
104. Hvid LG, Strotmeyer ES, Skjoldt M, Magnussen LV, Andersen M, Caserotti P. Voluntary muscle activation improves with power training and is associated with changes in gait speed in mobility-limited older adults - a randomized controlled trial. *Exp Gerontol*. 2016;80:51–56. doi:10.1016/j.exger.2016.03.018
105. Lopes PB, Pereira G, Lodovico A, Bento PC, Rodacki AL. Strength and power training effects on lower limb force, functional capacity, and static and dynamic balance in older female adults. *Rejuvenation Res*. 2016;19(5):385–393. doi:10.1089/rej.2015.1764
106. Park J. Effects of 24-week resistance exercise training on carotid peak systolic and end diastolic flow velocity in healthy older adults. *J Phys Ther Sci*. 2016;28(10):2793–2797. doi:10.1589/jpts.28.2793
107. Sayers SP, Gibson K, Bryan Mann J. Improvement in functional performance with high-speed power training in older adults is optimized in those with the highest training velocity. *Eur J Appl Physiol*. 2016;116(11–12):2327–2336. doi:10.1007/s00421-016-3484-x
108. Edholm P, Strandberg E, Kadi F. Lower limb explosive strength capacity in elderly women: effects of resistance training and healthy diet. *J Appl Physiol*. 2017;123(1):190–196. doi:10.1152/jappphysiol.00924.2016
109. Liao CD, Tsao JY, Lin LF, et al. Effects of elastic resistance exercise on body composition and physical capacity in older women with sarcopenic obesity: a CONSORT-compliant prospective randomized controlled trial. *Medicine*. 2017;96(23):e1115. doi:10.1097/MD.00000000000007115
110. Ramírez-Campillo R, Martínez C, de La Fuente CI, et al. High-speed resistance training in older women: the role of supervision. *J Aging Phys Act*. 2017;25(1):1–9. doi:10.1123/japa.2015-0122
111. Vale RGS, Ferrão MLD, Nunes RAM, Silva JB, Nodari RJ, Dantas EHM. Muscle strength, GH and IGF-1 in older women submitted to land and aquatic resistance training. *Rev Bras Med Esporte*. 2017;23:274–279. doi:10.1590/1517-869220172304163788
112. Mazini Filho ML, Aidar FJ, Gama de Matos D, et al. Circuit strength training improves muscle strength, functional performance and anthropometric indicators in sedentary elderly women. *J Sports Med Phys Fitness*. 2018;58(7–8):1029–1036. doi:10.23736/s0022-4707.17.06903-1
113. Gargallo P, Colado JC, Juevas A, et al. The effect of moderate- versus high-intensity resistance training on systemic redox state and DNA damage in healthy older women. *Biol Res Nurs*. 2018;20(2):205–217. doi:10.1177/1099800417753877
114. Ramirez-Campillo R, Alvarez C, Garcia-Hermoso A, et al. High-speed resistance training in elderly women: effects of cluster training sets on functional performance and quality of life. *Exp Gerontol*. 2018;110:216–222. doi:10.1016/j.exger.2018.06.014
115. Saeterbakken AH, Bårdstu HB, Brudeseth A, Andersen V. Effects of strength training on muscle properties, physical function, and physical activity among frail older people: a pilot study. *J Aging Res*. 2018;2018:8916274. doi:10.1155/2018/8916274
116. Englund DA, Price LL, Grosicki GJ, et al. Progressive resistance training improves torque capacity and strength in mobility-limited older adults. *J Gerontol A Biol Sci Med Sci*. 2019;74(8):1316–1321. doi:10.1093/gerona/gly199
117. Lee SR, Jo E, Khamoui AV. Chronic fish oil consumption with resistance training improves grip strength, physical function, and blood pressure in community-dwelling older adults. *Sports*. 2019;7(7):1–10. doi:10.3390/sports7070167
118. Mafi F, Biglari S, Ghardashi Afousi A, Gaeini AA. Improvement in skeletal muscle strength and plasma levels of follistatin and myostatin induced by an 8-week resistance training and epicatechin supplementation in sarcopenic older adults. *J Aging Phys Act*. 2019;27(3):384–391. doi:10.1123/japa.2017-0389
119. Pirauá ALT, Cavalcante BR, de Oliveira VMA, et al. Effect of 24-week strength training on unstable surfaces on mobility, balance, and concern about falling in older adults. *Scand J Med Sci Sports*. 2019;29(11):1805–1812. doi:10.1111/sms.13510
120. Rashidi E, Hosseini Kakhak SAR, Askari R. The effect of 8 weeks resistance training with low load and high load on testosterone, insulin-like growth factor-1, insulin-like growth factor binding protein-3 levels, and functional adaptations in older women. *Iran J Ageing*. 2019;14(3):356–367. doi:10.32598/sija.13.10.470
121. Richardson DL, Duncan MJ, Jimenez A, Juris PM, Clarke ND. Effects of movement velocity and training frequency of resistance exercise on functional performance in older adults: a randomised controlled trial. *Eur J Sport Sci*. 2019;19(2):234–246. doi:10.1080/17461391.2018.1497709
122. Urzi F, Marusic U, Ličen S, Buzan E. Effects of elastic resistance training on functional performance and myokines in older women—a randomized controlled trial. *J Am Med Dir Assoc*. 2019;20(7):830–834.e2. doi:10.1016/j.jamda.2019.01.151
123. Vezzoli A, Mrakic-Sposta S, Montorsi M, et al. Moderate intensity resistive training reduces oxidative stress and improves muscle mass and function in older individuals. *Antioxidants*. 2019;8(10):1–17. doi:10.3390/antiox8100431
124. Baker BS, Weitzel KJ, Roysse LA, et al. Efficacy of an 8-week resistance training program in older adults: a randomized controlled trial. *J Aging Phys Act*. 2020;29(1):121–129. doi:10.1123/japa.2020-0078
125. Cavalcante BR, de Souza MF, Falck RS, et al. Effects of resistance exercise with instability on cognitive function (REI Study): a proof-of-concept randomized controlled trial in older adults with cognitive complaints. *J Alzheimers Dis*. 2020;77(1):227–239. doi:10.3233/JAD-200349
126. Choi HM, Hurr C, Kim S. Effects of elastic band exercise on functional fitness and blood pressure response in the healthy elderly. *Int J Environ Res Public Health*. 2020;17(19). doi:10.3390/ijerph17197144
127. Ortega JAF, Cuartas LAH. Efectos de la velocidad de entrenamiento en fuerza sobre diversas manifestaciones de la fuerza en mujeres adultas mayores. *Retos*. 2020;38:325–332.
128. Jaque C, Véliz P, Ramírez-Campillo R, Moran J, Gentil P, Cancino J. High-speed bodyweight resistance training improves functional performance through maximal velocity in older females. *J Aging Phys Act*. 2020;29(4):659–669. doi:10.1123/japa.2020-0129
129. Van Roie E, Walker S, Van Driessche S, Delabastita T, Vanwanseele B, Delecluse C. An age-adapted plyometric exercise program improves dynamic strength, jump performance and functional capacity in older men either similarly or more than traditional resistance training. *PLoS One*. 2020;15(8):e0237921. doi:10.1371/journal.pone.0237921
130. Lai X, Bo L, Zhu H, et al. Effects of lower limb resistance exercise on muscle strength, physical fitness, and metabolism in pre-frail elderly patients: a randomized controlled trial. *BMC Geriatr*. 2021;21(1):447. doi:10.1186/s12877-021-02386-5
131. Lee YH, Lee PH, Lin LF, Liao CD, Liou TH, Huang SW. Effects of progressive elastic band resistance exercise for aged osteosarcopenic adiposity women. *Exp Gerontol*. 2021;147:111272. doi:10.1016/j.exger.2021.111272
132. Lee SY, Goh A, Tan K, et al. Effectiveness of a community-delivered pneumatic machine resistance training programme (Gym Tonic) for older adults at neighbourhood senior centres - a randomized controlled trial. *Eur Rev Aging Phys Act*. 2021;18(1):21. doi:10.1186/s11556-021-00273-x
133. Osuka Y, Kojima N, Sasai H, et al. Effects of exercise and/or  $\beta$ -hydroxy- $\beta$ -methylbutyrate supplementation on muscle mass, muscle strength, and physical performance in older women with low muscle mass: a randomized, double-blind, placebo-controlled trial. *Am J Clin Nutr*. 2021;114(4):1371–1385. doi:10.1093/ajcn/nqab176
134. Seo MW, Jung SW, Kim SW, Lee JM, Jung HC, Song JK. Effects of 16 weeks of resistance training on muscle quality and muscle growth factors in older adult women with sarcopenia: a randomized controlled trial. *Int J Environ Res Public Health*. 2021;18(13):1–13. doi:10.3390/ijerph18136762
135. Vieira IP, Lobo PCB, Fisher J, Ramirez-Campillo R, Pimentel GD, Gentil P. Effects of high-speed versus traditional resistance training in older adults. *Sports Health*. 2022;14(2):283–291. doi:10.1177/19417381211015211
136. Griffen C, Duncan M, Hattersley J, Weickert MO, Dallaway A, Renshaw D. Effects of resistance exercise and whey protein supplementation on skeletal muscle strength, mass, physical function, and hormonal and inflammatory biomarkers in healthy active older men: a randomised,

- double-blind, placebo-controlled trial. *Exp Gerontol.* 2022;158:111651. doi:10.1016/j.exger.2021.111651
137. Mc Dermott EJ, Balshaw TG, Brooke-Wavell K, Maden-Wilkinson TM, Folland JP. Fast and ballistic contractions involve greater neuromuscular power production in older adults during resistance exercise. *Eur J Appl Physiol.* 2022. Epub 2022/04/17. doi:10.1007/s00421-022-04947-x
138. Struble-Fitzsimmons D, Pinto Zipp G, DeLuca D, Zhang N. Exploring the relationship between timed up and go test times and fall history in an inpatient geriatric psychiatry unit: a retrospective case-control study. *J Geriatr Phys Ther.* 2020;43(3):E25–E30. doi:10.1519/JPT.0000000000000225
139. Graham JE, Fisher SR, Bergés IM, Kuo YF, Ostir GV. Walking speed threshold for classifying walking independence in hospitalized older adults. *Phys Ther.* 2010;90(11):1591–1597. doi:10.2522/ptj.20100018
140. Holviala J, Kraemer WJ, Sillanpää E, et al. Effects of strength, endurance and combined training on muscle strength, walking speed and dynamic balance in aging men. *Eur J Appl Physiol.* 2012;112(4):1335–1347. doi:10.1007/s00421-011-2089-7
141. Kent-Braun JA. Skeletal muscle fatigue in old age: whose advantage? *Exerc Sport Sci Rev.* 2009;37(1):3–9. doi:10.1097/JES.0b013e318190ea2e
142. Hortobágyi T, Lesinski M, Gäbler M, VanSwearingen JM, Malatesta D, Granacher U. Effects of three types of exercise interventions on healthy old adults' gait speed: a systematic review and meta-analysis. *Sports Med.* 2015;45(12):1627–1643. doi:10.1007/s40279-015-0371-2