



UNIVERSITY OF LEEDS

This is a repository copy of *Eating Disorders In weight-related Therapy (EDIT) Collaboration: Rationale and study design*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/196639/>

Version: Accepted Version

Article:

Lister, NB, Baur, LA, Paxton, SJ et al. (20 more authors) (2023) Eating Disorders In weight-related Therapy (EDIT) Collaboration: Rationale and study design. *Nutrition Research Reviews*. pp. 1-30. ISSN 0954-4224

<https://doi.org/10.1017/S0954422423000045>

© The Author(s) 2023. This article has been published in a revised form in <https://doi.org/10.1017/S0954422423000045>. This version is free to view and download for private research and study only. Not for re-distribution or re-use.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Eating Disorders In weight-related Therapy (EDIT) Collaboration: Rationale and study design

Natalie B. Lister^{1,2}, Louise A. Baur^{1,2}, Susan J. Paxton⁴, Sarah P. Garnett^{1,3}, Amy L Ahern⁵, Denise Wilfley⁶, Sarah Maguire⁷, Amanda Sainsbury⁸, Katharine Steinbeck^{1,9}, Caroline Braet¹⁰, Andrew Hill¹¹, Dasha Nicholls^{12,13}, Rebecca A. Jones⁵, Genevieve Dammary⁷, Alicia Grunseit¹⁴, Kelly Cooper¹⁵, Theodore K Kyle¹⁶, Faith N. Heeren¹⁷, Kylie E Hunter¹⁸, Caitlin M McMaster¹, Brittany J Johnson¹⁹, Anna Lene Seidler¹⁸, Hiba Jebeile^{1,2}

¹ The University of Sydney, Children's Hospital Westmead Clinical School, Westmead, Australia 2145

² Charles Perkins Centre, The University of Sydney, Sydney, Australia 2145

³ Kids Research, Sydney Children's Hospital Network, Westmead, Australia 2145

⁴ School of Psychology and Public Health, La Trobe University, Melbourne, Victoria, Australia

⁵ MRC Epidemiology Unit, University of Cambridge, Cambridge, CB2 0QQ, UK

⁶ Washington University in St. Louis, St. Louis, USA

⁷ InsideOut Institute for Eating Disorders, Boden Collaboration for Obesity, Nutrition and Eating Disorders, Charles Perkins Centre, The University of Sydney, Sydney, New South Wales, Australia

⁸ The University of Western Australia, School of Human Sciences, Crawley, WA, Australia

⁹ The Academic Department of Adolescent Medicine, The Children's Hospital at Westmead, Westmead New South Wales Australia 2145

¹⁰ Department of Developmental, Personality and Social Psychology, Ghent University, Henri Dunantlaan 2, 9000 Ghent, Belgium

¹¹ Leeds Institute of Health Sciences, University of Leeds, United Kingdom

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI.

10.1017/S0954422423000045

Nutrition Research Reviews is published by Cambridge University Press on behalf of The Nutrition Society

¹² Division of Psychiatry, Imperial College London, 2nd Floor, Commonwealth Building, Du Cane Road, London, W12 0NN, UK

¹³NIHR ARC Northwest London, UK

¹⁴ The Children's Hospital at Westmead, Nutrition and Dietetics, Weight Management Services, Westmead, NSW 2145, Australia

¹⁵ Weight Issues Network, Australia

¹⁶ ConscienHealth, Pittsburgh, Pennsylvania, USA

¹⁷ Department of Health Outcomes and Biomedical Informatics, University of Florida College of Medicine, Gainesville, Florida, USA

¹⁸ National Health and Medical Research Council Clinical Trials Centre, The University of Sydney, Sydney, New South Wales, Australia

¹⁹Caring Futures Institute, College of Nursing and Health Sciences, Flinders University, Adelaide, South Australia, Australia, 5042

Corresponding author: Dr Natalie Lister, The University of Sydney, Children's Hospital Westmead Clinical School, Westmead, NSW, Australia. **Email:**

Natalie.Lister@sydney.edu.au

Shortened title: EDIT Collaboration: Rationale and study

Funding

The EDIT collaboration is funded by the Australian National Health and Medical Research Council Ideas Grant (GTN2002310). NBL is a recipient of a National Health and Medical Research Council Peter Doherty Early Career Fellowship (GTN1145748). ALA and RAJ are supported by the Medical Research Council (MRC) (Grant MC_UU_00006/6). AS is funded by a Senior Research Fellowship (#1135897) from the NHMRC of Australia. DN is supported by the National Institute for Health Research (NIHR) under the Applied Health Research (ARC) programme for Northwest London. The views expressed in this publication are those of the author(s) and not necessarily those of the National Health Service, the NIHR or the Department of Health in England. ALS is funded by an NHMRC Emerging Leadership Investigator Grant (#2009432). DW is supported by the Scott Rudolph University Endowed Professorship at Washington University in St. Louis School of Medicine. HJ is supported by the Sydney Medical School Foundation (University of Sydney).

Funders had no role in the design, analysis or writing of this article.

Conflict of Interest

AS owns 50% of the shares in Zuman International, which receives royalties for books AS has written and payments for presentations. AS additionally reports receiving presentation fees and travel reimbursements from Eli Lilly and Co, the Pharmacy Guild of Australia, Novo Nordisk, the Dietitians Association of Australia, Shoalhaven Family Medical Centres, the Pharmaceutical Society of Australia, and Metagenics, and serving on the Nestlé Health Science Optifast VLCD advisory board from 2016 to 2018. ALA is Principal Investigator on two publicly funded trials where the intervention is provided by WW (formerly Weight Watchers) at no cost and is a member of the WW Scientific Advisory Board. KS has received in kind support as meals from 'Lite and Easy' for a clinical trial of weight stigma in young women in the last 5 years. AJH reports receiving payment for advice given to Slimming World (UK). TTK reports professional fees from Emerald Lake Safety, Gelesis, Johnson & Johnson, Novo Nordisk, and Nutrisystem. FAH reports personal fees from Novo Nordisk, outside the submitted work.

Abstract

The cornerstone of obesity treatment is behavioural weight management, resulting in significant improvements in cardio-metabolic and psychosocial health. However, there is ongoing concern that dietary interventions used for weight management may precipitate the development of eating disorders. Systematic reviews demonstrate that, while for most participants medically supervised obesity treatment improves risk scores related to eating disorders, a subset of people who undergo obesity treatment may have poor outcomes for eating disorders. This review summarises the background and rationale for the formation of the Eating Disorders In weight-related Therapy (EDIT) Collaboration. The EDIT Collaboration will explore the complex risk factor interactions that precede changes to eating disorder risk following weight management. In this review, we also outline the program of work and design of studies for the EDIT Collaboration, including expected knowledge gains. The EDIT studies explore risk factors and the interactions between them using individual level data from international weight management trials. Combining all available data on eating disorder risk from weight management trials will allow sufficient sample size to interrogate our hypothesis: that individuals undertaking weight management interventions will vary in their eating disorder risk profile, based on personal characteristics and intervention strategies available to them. The collaboration includes the integration of health consumers in project development and translation. An important knowledge gain from this project is a comprehensive understanding of the impact of weight management interventions on eating disorder risk.

Keywords: Eating disorders, obesity, individual participant data, meta-analysis

Introduction

Obesity and eating disorders are both complicated by serious, short- and long-term health problems ^(1; 2; 3). The prevalence of both is increasing ^(4; 5), with some data suggesting rates of combined obesity and eating disorders are increasing faster than the prevalence of either obesity or eating disorders alone ⁽⁶⁾. For example, between 1995 to 2015 in a community sample of Australian adults, prevalence of obesity alone increased 1.7-fold, and binge eating episodes increased 3.5-fold, while prevalence of combined obesity and recurrent binge eating episodes increased 5.7-fold ⁽⁷⁾. Co-existence of obesity and a range of eating disorders is seldom acknowledged ⁽⁸⁾, with these conditions commonly stereotyped as existing on opposite ends of an eating disordered spectrum. A key distinction is that obesity is defined by a physical metric, whereas eating disorders are defined by well characterised cognitive and behavioural phenotypes ⁽⁹⁾. Obesity and eating disorders share many risk factors (e.g. weight concern, dieting) ⁽⁸⁾ and treatment approaches (e.g. health professional support, self-monitoring, goal setting, and normalising eating patterns) ⁽¹⁰⁾. Of concern, the focus on weight loss during obesity treatment may lead to under-diagnosis of eating disorders in people with obesity ⁽⁶⁾. While only a small proportion of individuals may develop or have exacerbated eating disorder symptoms during weight management ⁽¹¹⁾, the potential burden for this risk is high. Nevertheless, there is a paucity of research and limited treatment pathways for those affected by *both* obesity and eating disorders. Additionally eating disorders may develop over several years ⁽¹²⁾ and weight management interventions may only be a single experience in the eating disorder development pathway. However, this is a unique point of engagement with health services, where risk can be identified and addressed, thus representing an important research and practice gap.

The aim of this review is to provide the background, rationale and study designs for the Eating Disorders In weight-related Therapy (EDIT) Collaboration. We describe considerations for the nuances of eating disorder development, including risk factors from observational studies, and describe how such risk factors may be influenced by behavioural weight management. The EDIT Collaboration will bring together individual participant data (IPD) from relevant trials to understand how individual characteristics and components of weight management interventions may contribute to eating disorder risk. We hypothesise that individuals undertaking weight management interventions will vary in their eating disorder risk profile, based on personal characteristics and intervention strategies available to them.

Further, we propose individual characteristics can be identified and intervention strategies can be adapted to reduce eating disorder risk.

State of the literature: Obesity, weight management and eating disorders

Eating disorder prevalence and complications in people with obesity

Worldwide, in 2016, 39% of men and 40% of women were affected by overweight, while 11% of men and 15% of women were affected by obesity⁽¹³⁾; and prevalence is predicted to rise by the year 2030⁽¹⁴⁾. For children and adolescents (5-19 years), the prevalence of obesity was 5.6% in girls and 7.8% for boys⁽⁴⁾. Complications of obesity include type 2 diabetes, non-alcoholic fatty liver disease, cardiovascular disease, sleep apnoea and depression^(15; 16). There is growing evidence that prevalence of obesity is higher in populations with culturally diverse and lower socioeconomic backgrounds^(13; 17).

Eating disorders include anorexia nervosa, atypical anorexia nervosa, bulimia nervosa, binge eating disorder, and several other categories of feeding and eating disorders⁽¹⁸⁾. Eating disorders are severe mental and physical health conditions with a long duration⁽¹⁹⁾ and high morbidity^(2; 20; 21). Despite misconceptions that eating disorders are diseases of individuals with lower body weight, the prevalence of eating disorders is higher in both men and women with obesity compared to their healthy weight peers⁽²²⁾. In a survey of 12,337 adults in the US, the lifetime prevalence, i.e., proportion of people that had any eating disorder at any point in their life, was 2.2% in men and 4.9% in women⁽²²⁾. However, men and women with obesity had a higher prevalence of eating disorders compared to the general population, at 3.8% and 7.6%, respectively⁽²²⁾. Similarly in adolescents, the prevalence of eating disorders is associated with higher body mass index (BMI). For example, a study of 3,043 Canadian adolescents found 9.3% of male adolescents and 20.2% of female adolescents with obesity had a sub- or full- threshold eating disorder compared to respectively 2.1% and 8.4% of adolescents with a BMI in the normal range⁽²³⁾. Data from 5,191 Australian adolescents show those with overweight or obesity were more likely to experience an eating disorder⁽²⁴⁾. Further, several studies have identified an increase in disordered eating behaviours over time in community samples^(7; 25), particularly among individuals with overweight and obesity. It has been reported that approximately one in four adolescents with obesity engages in binge eating behaviours or experiences loss of control with eating^(26; 27). Binge eating and a loss of control with eating are also associated with

weight gain and symptoms of the metabolic syndrome and are important drivers of continuing weight gain^(2; 28). Moreover, emerging evidence suggests sociocultural factors such as food insecurity and childhood adversity influence both eating disorders^(29; 30) and obesity^(31; 32), and both eating disorder symptoms and weight gain have been exacerbated by the COVID-19 pandemic lockdowns^(33; 34; 35).

Compared to the general population, individuals with an eating disorder have an elevated risk of premature mortality, with a German study showing that the highest mortality risk (standardised mortality ratio) associated with anorexia nervosa was 5.35, compared to bulimia nervosa (1.49) and binge eating disorder (1.50)⁽³⁶⁾. Indeed, complications of anorexia nervosa are equally severe for individuals with weight within or above the normal range (i.e. atypical anorexia nervosa compared to anorexia nervosa)⁽³⁷⁾. People with binge eating disorder experience a high prevalence of both psychiatric comorbidities (e.g. mood, anxiety and substance use disorders)^(38; 39) and physical comorbidities (e.g. type 2 diabetes, hypertension and chronic pain)⁽⁴⁰⁾. These comorbidities are also associated with obesity^(41; 42), however higher weight only partly explains the association with binge eating disorder⁽⁴²⁾. Almost 30% of adults with binge eating disorder also report a history of childhood obesity⁽⁴³⁾. Thus, identifying eating disorders and reducing risk during weight management has potential to reduce or prevent a range of physical and psychological complications.

While there may be a growing recognition that people with obesity are at increased risk of developing eating disorders^(6; 44), there has been limited progress in the identification, prevention and treatment of eating disorders in the context of weight management.

Behavioural weight management

Multicomponent behavioural interventions are first-line treatment for adolescents and adults affected by obesity^(45; 46; 47). These interventions typically recommend a combination of diet physical activity and behavioural modifications. A 2018 systematic review of adult behavioural weight management interventions of at least 12 months in duration showed that they were likely to produce significantly more weight loss compared to standard care (mean difference in weight change [MD], -2.39 kg [95% CI -2.86 to -1.93]; 67 studies; n=22,065). Eligible studies included participants recruited from primary care or a health care system, and intervention groups experienced less weight regain during the follow-up periods (beyond 12-18 months, pooled MD compared to control, -1.59 kg [95% CI, -2.38 to -0.79]; n=1408)⁽⁴⁸⁾.

Further, the risk of developing diabetes over 1 to 9 years was substantially reduced (pooled risk ratio, 0.67 [95% CI, 0.51 to 0.89]; 9 trials; n=3140). Clinical practice guidelines for adolescents with overweight or obesity recommend a family-based approach to multicomponent behavioural interventions that address dietary, sedentary and sleep behaviours⁽⁴⁷⁾. A 2017 Cochrane review⁽⁴⁹⁾ found multicomponent behavioural interventions for adolescents (aged 12-17 years) with overweight or obesity resulted in a mean change in body weight of -3.67 kg (95% CI -5.21 to -2.13; 20 trials; n=1993) and BMI of -1.18 kg/m² (95% CI -1.67 to -0.69; 28 trials; n=2774). These effects were maintained at 24-months follow-up. A 2012 systematic review reported significant improvements in low-density lipoprotein cholesterol (-0.30 mmol/L, 95% CI -0.45 to -0.15), triglycerides (-0.15 mmol/L, 95% CI -0.24 to -0.07), fasting insulin (-55.1 pmol/L, 95% CI -71.2 to -39.1) and blood pressure up to one year from baseline following lifestyle interventions for children and adolescents. Hence, multicomponent intensive behavioural weight management interventions can effectively reduce body weight and cardiometabolic risk in both adolescents and adults.

An emerging area of research and practice are dietary interventions with the potential to induce greater weight loss and improve cardiometabolic complications of obesity^(50; 51; 52). These interventions may include very low energy diets (<800 kcal/d), very low carbohydrate diets (<50 g carbohydrate/d) or intermittent energy restriction; and require both medical and dietetic supervision^(53; 54; 55). Such interventions are recommended for adolescents and adults with obesity and associated complications, or with severe obesity^(47; 56). These restrictive approaches play an important role in effectively managing weight and cardiometabolic risk, particularly as an obesity management approach prior to bariatric surgery or when pharmacological and surgical approaches are not available or contraindicated. However, the effect of these interventions on eating disorder risk is unclear.

Weight management and eating disorder risk

There is concern that dietary interventions, the cornerstone of behavioural weight management, may promote disordered eating and worsen psychological health^(57; 58) in some individuals. This is informed by longitudinal data showing that dieting is an important step within eating disorder development^(59; 60). However, some of these studies have poorly characterised the population sampled and the definition of “dieting”. Nevertheless, energy restriction may trigger binge eating in some people and it is thought that dietary and weight

monitoring may trigger a preoccupation with food, weight and shape. Data from intervention studies including dietary components are described below.

Evidence from systematic reviews

Systematic reviews have examined the association between behavioural weight management and the change in eating disorder risk. A 2017 systematic review examining weight management interventions for adults identified five randomised controlled trials (RCTs), all of which reported beneficial outcomes for eating disorder symptoms, including a reduction in binge eating ⁽⁶¹⁾. Similarly, systematic reviews of pre-post studies and RCTs conducted in children and adolescents found no change or a small reduction in eating disorder symptoms, including binge eating and loss of control, following behavioural weight management ^(11; 62; 63; 64). Other related eating disorder risk factors, including depression, anxiety, and low body image and self-esteem have also been improved following weight management in both adults and adolescents ^(65; 66; 67; 68; 69). The effect of restrictive dietary approaches, including low or very low energy diets, on binge eating has been examined in a systematic review of 10 studies including 805 adults ⁽⁵⁸⁾. In participants with pre-treatment binge eating disorder, studies reported a reduction in binge eating. The evidence was mixed in studies with participants with sub-clinical or no binge eating symptoms prior to treatment, with some studies showing a reduction in symptoms and others showing no change or an increase. Importantly, two studies included in this review reported an increase in binge eating or the onset of binge eating disorder in 10 to 15% of participants ⁽⁵⁸⁾. To our knowledge, the evidence of restrictive dietary approaches on eating disorder risk in adolescents is yet to be synthesised. These interventions are likely to include delivery features very different from behavioural weight management interventions included in the reviews in adults above, and the implications of this are unknown.

In summary, previous evidence suggests that eating disorder risk is reduced for most participants following professionally supervised behavioural weight management; however, individual studies have reported a small subset of participants who develop an eating disorder during the intervention or in the years following intervention ^(70; 71; 72; 73; 74; 75). For example, scores increased to above a clinical cut-point in seven children in one study and three of 56 participants followed up at 6-years had developed binge eating disorder in another study ^{(71;}

⁷²⁾. The mechanisms by which behavioural weight management may increase or decrease eating disorder risk at the individual level are not clear.

Individual variation in responses to weight management interventions

Few studies have investigated individuals' characteristics for associations with eating disorder outcomes following behavioural weight management interventions. For some individuals within the general population, dietary restraint (a proxy marker of dieting behaviours) is associated with the development of symptoms of binge eating disorder and bulimia nervosa ^(76; 77). In contrast, in others it may be an important behaviour that enables improvements in weight management and cardiometabolic health ⁽⁷⁸⁾. This divergence in response may be in part explained by the difficulty in distinguishing between flexible (i.e. gradual reduction, foods are limited in quantity rather than eliminated) and rigid restraint (extreme, all or nothing mentality) ⁽⁷⁸⁾. Psychosocial predictors thought to play a role in the development of eating disorders include poor self-esteem, depression, anxiety, bulimic behaviours (i.e. binge eating with compensatory behaviours), body dissatisfaction, and drive for thinness ^(60; 79). It is possible that the interactions or clustering of such individual characteristics with dietary restraint may be important for eating disorder development. For example, Stice's Dual Pathway model hypothesises that pressure for thinness increases risk for body dissatisfaction, which in turn increases the risk for dietary restriction and/or negative affect, thereby increasing the risk for subsequent onset of binge eating-related disorders ⁽⁸⁰⁾. However, etiological models including these risk factors do not consistently predict onset of eating disorders ⁽¹²⁾, suggesting varied individual responses. Importantly, research identifying eating disorder risk factors has been conducted in predominantly healthy weight populations; risk factors specific to individuals with obesity, for the full spectrum of eating disorder diagnoses are needed ^(44; 81; 82).

Beyond individual characteristics, there may be components of behavioural weight management interventions that influence eating disorder risk. Behavioural weight management interventions typically include a combination of intervention strategies (e.g., related to diet, movement, eating behaviour), delivered through various approaches (e.g., in terms of mode of delivery, session frequency/duration). Some commonly used strategies within weight management interventions are considered disordered behaviours in the context of eating disorder development, or may be components of etiological models of eating

disorders. Two examples are the restriction of energy intake and self-monitoring of weight⁽⁸³⁾. In behavioural weight management interventions, the prescription of restriction of energy intake (i.e., reduced food intake), is thought to equate to dietary restriction, while a focus on monitoring of weight, is likened to pressure for thinness and weight preoccupation. Hence these two components of behavioural weight management are thought to tie into the Dual Pathway sequence to promote disordered eating behaviours⁽⁴⁴⁾. This is a contentious theory, with intervention studies of restricted energy intake for up to two years in individuals without obesity (such as the CALERIE trials^(84; 85)), demonstrating increases in dietary restraint without increases in binge eating or eating disorders⁽⁸⁶⁾. Indeed, this leads to the conceptualisation of dietary restraint as a successful self regulation strategy. Schaumburg et al⁽⁷⁸⁾ posit that when this self regulation fails, and is followed by a period of disinhibition, eating disorder risk is increased⁽⁷⁸⁾.

Moreover, intervention components such as self-monitoring improve weight-related outcomes and long term weight maintenance for most people⁽⁸⁷⁾. Large scale behavioural programs, such as the Diabetes Prevention Program (DPP) and Look AHEAD, demonstrate that self-monitoring of weight and dietary intake are positively related to weight loss^(88; 89; 90), however these factors are also associated with disordered eating in community samples⁽⁸³⁾. Furthermore, factors such as increased self-esteem and self-efficacy around healthy eating are thought to be protective against disordered eating⁽⁹¹⁾, suggesting that changes in individual characteristics may mediate changes in eating disorder risk during interventions. Indeed, eating disorder treatment programs may also include components of regular weighing and the development of healthy eating behaviours. This highlights the need to understand the complexities of intervention components of weight management interventions for eating disorder risk. There are also many components of behavioural weight management interventions which may be protective against disordered eating, including regular contact and support from healthcare professionals, and strategies such as promotion of self-efficacy, realistic goal-setting, supported problem-solving, and strategies for stimulus control or social support⁽⁹²⁾. Thus, a complex interaction between individual characteristics and intervention components is likely to influence eating disorder risk during weight management. To inform models of care for clinical practice it is vital to identify and understand components that may increase or decrease eating disorder risk for different individuals who participate in professionally supervised behavioural weight management programs. Addressing this is the central goal of the EDIT Collaboration.

Stakeholder engagement in obesity and eating disorder research

To identify individual or intervention-level attributes that influence eating disorder risk in the context of weight management, it is essential to bring together stakeholders including those with lived experience of obesity and eating disorders. Engagement of consumers in research development and dissemination is an important process for evidence-based medicine ^(93; 94; 95).

The engagement of consumers in obesity and eating disorder research must be sensitive to the stigma and potential harms for those affected by both conditions. Data on the lived experience of people with combined obesity and eating disorders are limited. However, qualitative studies in people with lived experience of obesity frequently identify the harm caused by obesity stigma and weight bias ^(96; 97). A 2017 systematic review of cross-sectional studies reported that more frequent weight stigma experiences were associated with poorer physiological and psychological health ⁽⁹⁸⁾. Indeed, using the term obesity is an important consideration for researchers, with studies reporting mixed responses to the term from people with higher weights ^(99; 100). Many people with obesity prefer neutral terms, such as ‘weight’ be used in clinical care. However one study found all weight related terms elicit negative emotions ⁽⁹⁹⁾. On the other hand, obesity is a defined medical condition by a number of international health professional organisations, including the World Health Organization ⁽¹⁰¹⁾. Systematic reviews and international guidelines make suggestions and recommendations for reducing weight stigma and bias, such as the Joint international consensus statement for ending stigma of obesity ^(102; 103; 104; 105). The EDIT Collaboration will use person-first language for scientific discourse and language used within the collaboration will be reviewed and adjusted as required.

Summary and rationale for the EDIT Collaboration

Behavioural weight management interventions form the first-line treatment approach for obesity ⁽⁴⁵⁾. Clinical trials and systematic reviews show that, in addition to improved weight and cardiometabolic health, adolescents and adults with obesity who participate in supervised weight management interventions overall have improved eating behaviours, and psychological outcomes ^(11; 67; 69; 106). However, there is ongoing concern that these interventions may promote disordered eating and worsen psychological health in some individuals ⁽⁵⁷⁾. Indeed, some studies report worsening eating disorder outcomes in some individual participants ⁽¹¹⁾. This suggests that those who have poorer outcomes following

weight management interventions are not captured when studies report *aggregate* risk scores. Due to the required large sample sizes and need for individual-level analysis (rather than pooling summary scores), the important research questions of whether weight management increases or decreases eating disorder risk for an individual is difficult if not impossible to answer using a single trial, qualitative methods, or standard aggregate data meta-analyses. Further, weight management interventions are complex and often poorly described. Research investigating whether certain intervention types or components of interventions may either increase or decrease eating disorder risk at an individual level is needed. It is likely that complex interactions between individual characteristics and intervention components influence eating disorder risk responses during weight management. Interventional evidence that addresses clinically supervised behavioural weight management for people with obesity should be examined to address these concerns.

Research program

The EDIT Collaboration will bring together clinicians, researchers, biostatisticians and individuals with lived experience from around the world to improve treatment for people affected by obesity and eating disorders. The EDIT Collaboration aims to: (1) To understand which participants experience a change in eating disorder risk, or related symptoms, during and following weight management interventions ; (2) understand which intervention components may contribute to eating disorder risk; (3) identify predictive pathways for increased or decreased eating disorder risk during weight management; and (4) develop resources and recommendations to reduce eating disorder development during obesity treatment. To achieve these aims, the EDIT Collaboration will conduct five related studies (Figure 1). Detailed methodologies will be published separately but a brief overview is provided below.

Scientific and stakeholder engagement

The work of the EDIT Collaboration is guided by Scientific and Stakeholder Advisory panels with international representation. The Scientific Advisory Panel includes experienced researchers and clinicians from the fields of obesity and eating disorders or those working across both conditions. The panel is responsible for overall program oversight and will provide strategic advice relating to the scientific rigour of each included study, contribute to

protocol development, scientific publications, guide project output and the translation of project outcomes. The Stakeholder Advisory Panel comprises consumers with a lived experience of eating disorders, obesity or both conditions. The Stakeholder Advisory Panel will provide strategic advice, contribute to protocol development, and guide project outputs and translation from the viewpoint of the end consumers of health services.

Eligible trials

Systematic searches of electronic databases and trial registries ⁽¹⁰⁷⁾ are being conducted to identify trials that meet our inclusion criteria: **1)** randomised controlled trial of behavioural weight management intervention; **2)** for adolescents and/or adults with obesity; **3)** report at least one measure of eating disorder symptoms or behaviours at baseline and post-intervention or follow-up using a validated self-report questionnaire (e.g., Eating Disorder Examination Questionnaire, Binge Eating Scale) and/or clinical assessment or diagnostic interview (e.g., Eating Disorder Examination). The protocol for this review is registered with PROSPERO (CRD42021265340), accessible from https://www.crd.york.ac.uk/prospERO/display_record.php?ID=CRD42021265340.

Representatives from each identified trial will be invited to join the collaboration and share IPD, i.e. line-by-line data for each individual participant data. The corresponding authors of identified trials are invited to join the EDIT Collaboration via email. If, after two attempts, no response has been received, other authors on the paper or listed on a registration record will be emailed. Finally, we will attempt to contact trialists via telephone, using our networks, via their institutions or at conferences. If trialists are unable to be contacted after multiple attempts, the trial will be excluded, since our analyses are not possible using published summary data alone. Trial representatives will have the opportunity to provide input to all major stages of the project including protocol development, analysis, results interpretation and translation.

Study 1: Consultation

We will identify individual participant characteristics and intervention strategies which may contribute to an increase or decrease in eating disorder risk during weight management interventions through broad stakeholder consultation. Using an online survey, we will canvass diverse opinions on possible causes of eating disorder development in weight

management interventions. Participants will be asked to rate the relevance of individual characteristics (e.g. body dissatisfaction, history of self-directed dieting, disinhibition related to eating) and intervention strategies (e.g. dietary monitoring, dietary behaviour change strategies, informed by a psychological framework or theory) and identify any not listed in the survey. Individual characteristics and intervention strategies listed in the survey are informed by relevant literature, expert and consumer consultation (Scientific and Stakeholder Advisory Panels). The outcomes of this survey will be used to inform the analyses in the studies below.

Studies 2 and 3: Individual participant data meta-analyses

Meta-analysis of IPD is considered the "gold standard" ⁽¹⁰⁸⁾ for meta-analysis, in part due to the opportunity to explore differences in treatment effects across subgroups. These subgroups might include subsets of participants, such as those with higher eating disorder risk at baseline, or subsets of studies, such as those with particular intervention strategies ⁽¹⁰⁹⁾.

We will collate all available data from all eligible studies to examine the individual risk of eating disorder development during weight management trials. IPD from collaborating trials will be collated into a central database. Trials will be identified through systematic searches, investigator networks and study branding (editcollaboration.com). The specific variables to be included in the analysis will be informed by stakeholder consultation (Study 1) and data availability, whereby any suggested predictor will be considered. We will conduct two IPD meta-analyses, with detailed methodology published elsewhere *a priori*. Study 2 aims to identify baseline participant risk factors which predict an increase or decrease in eating disorder risk, or related symptoms, during and following a weight management intervention. This will be a pre-post IPD meta-analysis. Study 3 aims to determine whether there are baseline participant risk factors which predict change in eating disorder risk, or related symptoms, if they receive any behavioural weight management intervention compared with no intervention (i.e., no treatment controls). This will be an IPD meta-analysis maintaining randomisation.

Study 4: Intervention deconstruction

Interventions included in the EDIT Collaboration will be deconstructed into their discrete components to improve understanding of what they involve. Intervention components will

include the intervention strategies to drive weight management (e.g. dietary or behaviour change strategies) and the features of how such interventions are delivered. Using a systematic coding framework developed for this study, we will then compare and synthesise the components of interventions targeting adolescent and adult populations. This project will allow for future quantitative analysis of intervention components and individual participant eating disorder risk.

Study 5: Predictive modelling

Data from Studies 2, 3 and 4 will be combined to identify any interactions between individual characteristics and intervention strategies which may increase or decrease the risk of eating disorders during weight management. Detailed methodology informed by the findings of studies 1-4 will be published separately.

Translation plan and recommendations

Knowledge gains (see **Box 1**) will inform the translation plan and recommendations. Workshops with the Stakeholder Advisory Panel (which includes stakeholders who are consumers with lived experience of obesity and eating disorders), Scientific Advisory Panel, and trial representatives will inform our translation action plan. A working group will be formed to *develop plans* for five key areas for translation: i) models of care for obesity management that consider eating disorder risk; ii) health professional education (e.g. training webinars, scientific publications, conferences, recommendations for screening and monitoring protocols); iii) community dissemination (e.g. website, newsletter, community seminars, decision aids for informing treatment consent); iv) identification of strategies to support further implementation; and v) policy briefing documents summarising key evidence that emerges from the research.

Strengths and Limitations

The strengths of this research are first the use of robust statistical methods to quantitatively examine individual-level and intervention-level eating disorder risk during behavioural weight management interventions. We will use statistical methods recommended by the Cochrane Collaboration^(108; 110), led by a team of biostatisticians with previous IPD meta-

analyses experience ^(111; 112). Secondly, this project incorporates consumers' views and broader stakeholder engagement to set research priorities and to translate the outcomes of the study. Thirdly, adolescents trials are included as an important life stage when the trajectories of both obesity and eating disorders become firmly established.

However, the studies outlined in this current research plan have limitations. This program will not report on qualitative experiences of participants who have undertaken weight management interventions. The IPD relies on retrospective analysis of data from clinical trials. Thus, there is an inherent risk that adverse eating disorder outcomes are not captured due to missing data, higher participant attrition among those at risk, or insufficient follow up of the included studies. However, IPD allow us to include data from excluded participants (e.g., outliers), more variables and timepoints from data sets that may not be included in a traditional aggregate data meta-analysis, thus increasing power to conduct subgroup analyses and detect adverse events. Further, where data are available, we will include in our analysis known psychosocial predictors of eating disorder development (e.g., self-esteem, depression, anxiety, bulimic behaviours, body dissatisfaction, and drive for thinness) to identify potential changes in an individual's risk profile that may precede changes in global eating disorder risk. Future research addressing qualitative experiences and a prospective data analysis will be important to complement the current research plan.

There is also the possibility that the clinical trials /interventions eligible for the EDIT Collaboration (i.e. including a validated comprehensive measure eating disorder risk) are not representative of broader weight management interventions and findings will not be generalisable to all weight management. Moreover, the withdrawal of interventions and support may also influence eating disorder risk, and long-term data may not capture this changing risk profile. All included trials are providing a weight management intervention, thus, whether eating disorder risk would differ for those not referred or enrolled in an intervention will not be determined. Instead, this project will identify adolescents and adults presenting for weight management: i) for whom weight management will likely improve physical and mental health; ii) for whom behavioural weight management is not recommended, and; iii) whether intervention components can improve outcomes for different individuals.

Summary and conclusion

The EDIT Collaboration will combine IPD meta-analysis and intervention coding to quantitatively explore the underlying pathways that increase or decrease eating disorder risk during behavioural weight management interventions. By understanding how individual participant characteristics may interact with intervention components to influence eating disorder risk, we have the potential to create an innovative toolbox for clinicians to build the safest interventions for each individual. Future combined research between obesity and eating disorder fields has the potential to lead to a tailored precision therapeutic response, improving both obesity and eating disorders care.

Acknowledgements

The authors would like to thank members of the TOPCHILD (Transforming Obesity Prevention for CHILDren) Collaboration for project support and feedback on study design.

Authorship

NBL, HJ, LAB, SJP, ALS conceived the idea for the study. NBL, HJ, LAB, SJP, SPG, ALA, DW, SM, AS, KS, BJJ, ALS developed the rationale and research question. NBL wrote the first draft of the manuscript. HJ, LAB, SJP, SPG, ALA, DW, SM, AS, KS, CB, AJH, DN, RAJ, GD, AMG, KC, TKK, FN, FQ, KEH, CM, BJJ, ALS provided critical review and feedback at each stage of the process. All authors critically revised the manuscript for intellectual content, and agreed and approved the final manuscript. NBL is the guarantor of the review.

References

1. US Preventive Services Task Force (2018) Behavioral weight loss interventions to prevent obesity-related morbidity and mortality in adults: US Preventive Services Task Force recommendation statement. *JAMA* **320**, 1163-1171.
2. Klump KL, Bulik CM, Kaye WH *et al.* (2009) Academy for eating disorders position paper: Eating disorders are serious mental illnesses. *International Journal of Eating Disorders* **42**, 97-103.
3. Steinbeck KS, Lister NB, Gow ML *et al.* (2018) Treatment of adolescent obesity. *Nature Reviews Endocrinology*.
4. NCD Risk Factor Collaboration (2017) Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128· 9 million children, adolescents, and adults. *Lancet*.
5. Galmiche M, Déchelotte P, Lambert G *et al.* (2019) Prevalence of eating disorders over the 2000–2018 period: a systematic literature review. *The American Journal of Clinical Nutrition* **109**, 1402-1413.
6. Da Luz FQ, Hay P, Touyz S *et al.* (2018) Obesity with Comorbid Eating Disorders: Associated Health Risks and Treatment Approaches. *Nutrients* **10**, 829.
7. da Luz FQ, Sainsbury A, Mannan H *et al.* (2017) Prevalence of obesity and comorbid eating disorder behaviors in South Australia from 1995 to 2015. *International Journal of Obesity* **41**, 1148-1153.
8. Haines J, Kleinman KP, Rifas-Shiman SL *et al.* (2010) Examination of Shared Risk and Protective Factors for Overweight and Disordered Eating Among Adolescents. *Archives of Pediatrics & Adolescent Medicine* **164**, 336-343.
9. Academy for Eating Disorders (AED) Nine more Truths about Eating Disorders: Weight and Weight Stigma. <https://www.aedweb.org/publications/nine-truths-weight> (2022)
10. Lister NB, Baur LA, Paxton SJ *et al.* (2021) Contextualising Eating Disorder Concerns for Paediatric Obesity Treatment. *Current Obesity Reports* **10**, 322-331.
11. Jebeile H, Gow ML, Baur LA *et al.* (2019) Treatment of obesity, with a dietary component, and eating disorder risk in children and adolescents: A systematic review with meta-analysis. *Obesity Reviews* **20**, 1287-1298.

12. Stice E, Van Ryzin MJ (2019) A prospective test of the temporal sequencing of risk factor emergence in the dual pathway model of eating disorders. *Journal of abnormal psychology* **128**, 119.
13. World Health Organization (2020) Overweight and obesity. In *Health at a Glance: Asia/Pacific 2020: Measuring Progress Towards Universal Health Coverage*. Paris: OECD Publishing.
14. Lobstein T, Brinsden H (2019) *Atlas of Childhood obesity*. World Obesity Federation.
15. Bray GA, Kim KK, Wilding JPH *et al.* (2017) Obesity: a chronic relapsing progressive disease process. A position statement of the World Obesity Federation. *Obesity Reviews* **18**, 715-723.
16. Pereira-Miranda E, Costa PRF, Queiroz VAO *et al.* (2017) Overweight and Obesity Associated with Higher Depression Prevalence in Adults: A Systematic Review and Meta-Analysis. *Journal of the American College of Nutrition* **36**, 223-233.
17. . Bann D, Johnson W, Li L *et al.* (2018) Socioeconomic inequalities in childhood and adolescent body-mass index, weight, and height from 1953 to 2015: an analysis of four longitudinal, observational, British birth cohort studies. *The Lancet Public Health* **3**, e194-e203.
18. American Psychiatric Association (2013) *Diagnostic and statistical manual of mental disorders (DSM-5®)*: American Psychiatric Pub.
19. . Schmidt U, Adan R, Böhm I *et al.* (2016) Eating disorders: the big issue. *The Lancet Psychiatry* **3**, 313-315.
20. Arcelus J, Mitchell AJ, Wales J *et al.* (2011) Mortality Rates in Patients With Anorexia Nervosa and Other Eating Disorders: A Meta-analysis of 36 Studies. *Archives of General Psychiatry* **68**, 724-731.
21. Ágh T, Kovács G, Supina D *et al.* (2016) A systematic review of the health-related quality of life and economic burdens of anorexia nervosa, bulimia nervosa, and binge eating disorder. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity* **21**, 353-364.
22. Duncan AE, Ziobrowski HN, Nicol G (2017) The Prevalence of Past 12-Month and Lifetime DSM-IV Eating Disorders by BMI Category in US Men and Women. *European Eating Disorders Review* **25**, 165-171.
23. Flament MF, Henderson K, Buchholz A *et al.* (2015) Weight Status and DSM-5 Diagnoses of Eating Disorders in Adolescents From the Community. *Journal of the American Academy of Child & Adolescent Psychiatry* **54**, 403-411.e402.

24. Mitchison D, Mond J, Bussey K *et al.* (2020) DSM-5 full syndrome, other specified, and unspecified eating disorders in Australian adolescents: prevalence and clinical significance. *Psychological Medicine* **50**, 981-990.
25. Darby A, Hay P, Mond J *et al.* (2009) The rising prevalence of comorbid obesity and eating disorder behaviors from 1995 to 2005. *International Journal of Eating Disorders* **42**, 104-108.
- 26 He J, Cai Z, Fan X (2017) Prevalence of binge and loss of control eating among children and adolescents with overweight and obesity: An exploratory meta-analysis. *International Journal of Eating Disorders* **50**, 91-103.
27. Goossens L, Braet C, Decaluwé V (2007) Loss of control over eating in obese youngsters. *Behaviour Research and Therapy* **45**, 1-9.
28. Tanofsky-Kraff M, Shomaker LB, Stern EA *et al.* (2012) Children's binge eating and development of metabolic syndrome. *International journal of obesity* **36**, 956-962.
29. Huryk KM, Drury CR, Loeb KL (2021) Diseases of affluence? A systematic review of the literature on socioeconomic diversity in eating disorders. *Eating Behaviors* **43**, 101548.
30. Rasmusson G, Lydecker JA, Coffino JA *et al.* (2019) Household food insecurity is associated with binge-eating disorder and obesity. *International Journal of Eating Disorders* **52**, 28-35.
31. Schroeder K, Schuler BR, Kobulsky JM *et al.* (2021) The association between adverse childhood experiences and childhood obesity: A systematic review. *Obesity Reviews* **22**, e13204.
32. St. Pierre C, Ver Ploeg M, Dietz WH *et al.* (2022) Food Insecurity and Childhood Obesity: A Systematic Review. *Pediatrics* **150**.
33. Sideli L, Lo Coco G, Bonfanti RC *et al.* (2021) Effects of COVID-19 lockdown on eating disorders and obesity: A systematic review and meta-analysis. *European Eating Disorders Review* **29**, 826-841.
34. Giel KE, Schurr M, Zipfel S *et al.* (2021) Eating behaviour and symptom trajectories in patients with a history of binge eating disorder during COVID-19 pandemic. *European Eating Disorders Review* **29**, 657-662.
35. Phillipou A, Meyer D, Neill E *et al.* (2020) Eating and exercise behaviors in eating disorders and the general population during the COVID-19 pandemic in Australia: Initial results from the COLLATE project. *International Journal of Eating Disorders* **53**, 1158-1165.

36. Fichter MM, Quadflieg N (2016) Mortality in eating disorders - results of a large prospective clinical longitudinal study. *International Journal of Eating Disorders* **49**, 391-401.
37. Walsh BT, Hagan KE, Lockwood C A systematic review comparing atypical anorexia nervosa and anorexia nervosa. *International Journal of Eating Disorders* **n/a**.
38. Araujo DMR, Santos GFdS, Nardi AE (2010) Binge eating disorder and depression: A systematic review. *The World Journal of Biological Psychiatry* **11**, 199-207.
39. Goossens L, Braet C, Van Vlierberghe L *et al.* (2009) Loss of control over eating in overweight youngsters: The role of anxiety, depression and emotional eating. *European Eating Disorders Review* **17**, 68-78.
40. Mitchell JE (2016) Medical comorbidity and medical complications associated with binge-eating disorder. *International Journal of Eating Disorders* **49**, 319-323.
41. McCuen-Wurst C, Ruggieri M, Allison KC (2018) Disordered eating and obesity: associations between binge-eating disorder, night-eating syndrome, and weight-related comorbidities. *Ann N Y Acad Sci* **1411**, 96-105.
42. Thornton LM, Watson HJ, Jangmo A *et al.* (2017) Binge-eating disorder in the Swedish national registers: Somatic comorbidity. *International Journal of Eating Disorders* **50**, 58-65.
43. Jacobi C, Hayward C, de Zwaan M *et al.* (2004) Coming to terms with risk factors for eating disorders: application of risk terminology and suggestions for a general taxonomy. *Psychological bulletin* **130**, 19.
44. Jebeile H, Lister N, Baur L *et al.* (2021) Eating disorder risk in adolescents with obesity. *Obesity reviews* **22**, e13173.
45. National Clinical Guideline Centre (UK) (2014) Identification, assessment and management of overweight and obesity in children, young people and adults. (NICE Clinical Guideline 189). www.nice.org.uk/guidance/cg189: London: National Institute for Health and Care Excellence (UK).
46. Semlitsch T, Stigler FL, Jeitler K *et al.* (2019) Management of overweight and obesity in primary care—A systematic overview of international evidence-based guidelines. *Obesity Reviews* **20**, 1218-1230.
47. Alman KL, Lister NB, Garnett SP *et al.* (2020) Dietetic management of obesity and severe obesity in children and adolescents: A scoping review of guidelines. *Obesity Reviews* **n/a**.
48. LeBlanc EL, Patnode CD, Webber EM *et al.* (2018) U.S. Preventive Services Task Force Evidence Syntheses, formerly Systematic Evidence Reviews. In *Behavioral and*

Pharmacotherapy Weight Loss Interventions to Prevent Obesity-Related Morbidity and Mortality in Adults: An Updated Systematic Review for the US Preventive Services Task Force. Rockville (MD): Agency for Healthcare Research and Quality (US).

49. Al-Khudairy L, Loveman E, Colquitt JL *et al.* (2017) Diet, physical activity and behavioural interventions for the treatment of overweight or obese adolescents aged 12 to 17 years. *The Cochrane Library*.

50. Mulholland Y, Nicokavoura E, Broom J *et al.* (2012) Very-low-energy diets and morbidity: a systematic review of longer-term evidence. *The British journal of nutrition* **108**, 832-851.

51. Asher RC, Burrows TL, Collins CE (2013) Very low-energy diets for weight loss in adults: A review. *Nutrition & Dietetics* **70**, 101-112.

52. Parretti HM, Jebb SA, Johns DJ *et al.* (2016) Clinical effectiveness of very-low-energy diets in the management of weight loss: a systematic review and meta-analysis of randomized controlled trials. *Obesity Reviews* **17**, 225-234.

53. Hoare JK, Jebeile H, Garnett SP *et al.* (2021) Novel dietary interventions for adolescents with obesity: A narrative review. *Pediatric Obesity* **16**, e12798.

54. Andela S, Burrows TL, Baur LA *et al.* (2019) Efficacy of very low-energy diet programs for weight loss: A systematic review with meta-analysis of intervention studies in children and adolescents with obesity. *Obesity reviews : an official journal of the International Association for the Study of Obesity* **20**, 871-882.

55. Gow ML, Garnett SP, Baur LA *et al.* (2016) The Effectiveness of Different Diet Strategies to Reduce Type 2 Diabetes Risk in Youth. *Nutrients* **8**, 486.

56. Jensen MD, Ryan DH, Apovian CM *et al.* (2014) 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Circulation* **129**, S102-138.

57. Academy for Eating Disorders (AED) (2019) Research Risk/Benefit Ethics Position Statement. https://higherlogicdownload.s3.amazonaws.com/AEDWEB/05656ea0-59c9-4dd4-b832-07a3fea58f4c/UploadedImages/Advocate/Research_Risk-Benefit_Ethics_Position_Statement_2_28_19_-_FINAL.pdf

58. da Luz FQ, Hay P, Gibson AA *et al.* (2015) Does severe dietary energy restriction increase binge eating in overweight or obese individuals? A systematic review. *Obesity Reviews* **16**, 652-665.

59. Neumark-Sztainer D, Wall M, Guo J *et al.* (2006) Obesity, Disordered Eating, and Eating Disorders in a Longitudinal Study of Adolescents: How Do Dieters Fare 5 Years Later? *Journal of the American Dietetic Association* **106**, 559-568.
60. Stice E, Marti CN, Durant S (2011) Risk factors for onset of eating disorders: Evidence of multiple risk pathways from an 8-year prospective study. *Behaviour Research and Therapy* **49**, 622-627.
61. Peckmezian T, Hay P (2017) A systematic review and narrative synthesis of interventions for uncomplicated obesity: weight loss, well-being and impact on eating disorders. *Journal of Eating Disorders* **5**, 15.
62. Moustafa AF, Quigley KM, Wadden TA *et al.* (2021) A systematic review of binge eating, loss of control eating, and weight loss in children and adolescents. *Obesity* **29**, 1259-1271.
63. Butryn ML, Wadden TA (2005) Treatment of overweight in children and adolescents: does dieting increase the risk of eating disorders? *International Journal of Eating Disorders* **37**, 285-293.
64. De Giuseppe R, Di Napoli I, Porri D *et al.* (2019) Pediatric obesity and eating disorders symptoms: the role of the multidisciplinary treatment. A systematic review. *Frontiers in pediatrics* **7**, 123.
65. Fabricatore AN, Wadden TA, Higginbotham AJ *et al.* (2011) Intentional weight loss and changes in symptoms of depression: a systematic review and meta-analysis. *International Journal of Obesity* **35**, 1363-1376.
66. Gow ML, Tee MS, Garnett SP *et al.* (2020) Pediatric obesity treatment, self-esteem, and body image: A systematic review with meta-analysis. *Pediatric Obesity*.
67. Jebeile H, Gow ML, Baur LA *et al.* (2019) Association of Pediatric Obesity Treatment, Including a Dietary Component, With Change in Depression and Anxiety: A Systematic Review and Meta-analysis. *JAMA pediatrics* **173**, e192841-e192841.
68. Murray M, Dordevic AL, Bonham MP (2017) Systematic review and meta-analysis: the impact of multicomponent weight management interventions on self-esteem in overweight and obese adolescents. *Journal of pediatric psychology* **42**, 379-394.
69. Jones RA, Lawlor ER, Birch JM *et al.* (2021) The impact of adult behavioural weight management interventions on mental health: A systematic review and meta-analysis. *Obesity Reviews* **22**, e13150.
70. Epstein LH, Valoski A, Wing RR *et al.* (1994) 10-year outcomes of behavioural family-based treatment for childhood obesity. *Health Psychology* **13**, 373-383.

71. Goossens L, Braet C, Verbeken S *et al.* (2011) Long-term outcome of pediatric eating pathology and predictors for the onset of loss of control over eating following weight-loss treatment. *International journal of eating disorders* **44**, 397-405.
72. Epstein LH, Paluch RA, Saelens BE *et al.* (2001) Changes in eating disorder symptoms with pediatric obesity treatment. *The Journal of pediatrics* **139**, 58-65.
73. Wadden TA, Foster GD, Sarwer DB *et al.* (2004) Dieting and the development of eating disorders in obese women: results of a randomized controlled trial. *Am J Clin Nutr* **80**, 560-568.
74. Raymond NC, de Zwaan M, Mitchell JE *et al.* (2002) Effect of a very low calorie diet on the diagnostic category of individuals with binge eating disorder. *Int J Eat Disord* **31**, 49-56.
75. Telch CF, Agras WS (1993) The effects of a very low calorie diet on binge eating. *Behavior Therapy* **24**, 177-193.
76. Stice E (2001) A prospective test of the dual-pathway model of bulimic pathology: mediating effects of dieting and negative affect. *Journal of abnormal psychology* **110**, 124.
77. Decaluwé V, Braet C (2005) The cognitive behavioural model for eating disorders: A direct evaluation in children and adolescents with obesity. *Eating Behaviors* **6**, 211-220.
78. Schaumberg K, Anderson D, Anderson L *et al.* (2016) Dietary restraint: what's the harm? A review of the relationship between dietary restraint, weight trajectory and the development of eating pathology. *Clinical Obesity* **6**, 89-100.
79. Russo J, Brennan L, Walkley J *et al.* (2011) Psychosocial Predictors of Eating Disorder Risk in Overweight and Obese Treatment-Seeking Adolescents. *Behaviour Change* **28**, 111-127.
80. Wonderlich S, Mitchell J, Steiger H (2005) *Annual Review of Eating Disorders 2006*: Radcliffe Publishing.
81. Hilbert A, Pike KM, Goldschmidt AB *et al.* (2014) Risk factors across the eating disorders. *Psychiatry research* **220**, 500-506.
82. Striegel-Moore RH, Fairburn CG, Wilfley DE *et al.* (2005) Toward an understanding of risk factors for binge-eating disorder in black and white women: a community-based case-control study. *Psychol Med* **35**, 907-917.
83. Pacanowski CR, Linde JA, Neumark-Sztainer D (2015) Self-Weighing: Helpful or Harmful for Psychological Well-Being? A Review of the Literature. *Current Obesity Reports* **4**, 65-72.

84. Williamson DA, Martin CK, Anton SD *et al.* (2008) Is caloric restriction associated with development of eating-disorder symptoms? Results from the CALERIE trial. *Health Psychol* **27**, S32-42.
85. Dorling JL, van Vliet S, Huffman KM *et al.* (2021) Effects of caloric restriction on human physiological, psychological, and behavioral outcomes: highlights from CALERIE phase 2. *Nutr Rev* **79**, 98-113.
86. Stewart TM, Martin CK, Williamson DA (2022) The Complicated Relationship between Dieting, Dietary Restraint, Caloric Restriction, and Eating Disorders: Is a Shift in Public Health Messaging Warranted? *Int J Environ Res Public Health* **19**.
87. Burke LE, Wang J, Sevick MA (2011) Self-Monitoring in Weight Loss: A Systematic Review of the Literature. *Journal of the American Dietetic Association* **111**, 92-102.
88. Knowler WC, Fowler SE, Hamman RF *et al.* (2009) 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* **374**, 1677-1686.
89. Apolzan JW, Venditti EM, Edelstein SL *et al.* (2019) Long-Term Weight Loss With Metformin or Lifestyle Intervention in the Diabetes Prevention Program Outcomes Study. *Annals of internal medicine* **170**, 682-690.
90. The Look AHEAD Research Group (2014) Eight-year weight losses with an intensive lifestyle intervention: The look AHEAD study. *Obesity* **22**, 5-13.
91. Neumark-Sztainer DR, Wall MM, Haines JI *et al.* (2007) Shared risk and protective factors for overweight and disordered eating in adolescents. *American journal of preventive medicine* **33**, 359-369. e353.
92. Hayes JF, Fitzsimmons-Craft EE, Karam AM *et al.* (2018) Disordered Eating Attitudes and Behaviors in Youth with Overweight and Obesity: Implications for Treatment. *Current Obesity Reports* **7**, 235-246.
93. Qaseem A, Forland F, Macbeth F *et al.* (2012) Guidelines International Network: toward international standards for clinical practice guidelines. *Annals of internal medicine* **156**, 525-531.
94. Schünemann HJ, Fretheim A, Oxman AD (2006) Improving the use of research evidence in guideline development: 1. Guidelines for guidelines. *Health Research Policy and Systems* **4**, 1-6.
95. Jarrett L, Unit PI (2004) A report on a study to evaluate patient/carer membership of the first NICE Guideline Development Groups. *National Institute for Clinical Excellence (NICE), National Health Service (NHS)*.

96. Pudney EV, Himmelstein MS, Puhl RM *et al.* (2020) Distressed or not distressed? A mixed methods examination of reactions to weight stigma and implications for emotional wellbeing and internalized weight bias. *Social Science & Medicine* **249**, 112854.
97. Hollett KB, Carter JC (2021) Separating binge-eating disorder stigma and weight stigma: A vignette study. *International Journal of Eating Disorders* **54**, 755-763.
98. Wu Y-K, Berry DC (2018) Impact of weight stigma on physiological and psychological health outcomes for overweight and obese adults: A systematic review. *Journal of Advanced Nursing* **74**, 1030-1042.
99. Brown A, Flint SW (2021) Preferences and emotional response to weight-related terminology used by healthcare professionals to describe body weight in people living with overweight and obesity. *Clinical Obesity* **11**, e12470.
100. Puhl RM (2020) What words should we use to talk about weight? A systematic review of quantitative and qualitative studies examining preferences for weight-related terminology. *Obesity Reviews* **21**, e13008.
101. World Health Organization (2022) Obesity. https://www.who.int/health-topics/obesity#tab=tab_1 (accessed 3rd October 2022 2022)
102. Pont SJ, Puhl R, Cook SR *et al.* (2017) Stigma Experienced by Children and Adolescents With Obesity. *Pediatrics* **140**, e20173034.
103. Eisenberg D, Noria S, Grover B *et al.* (2019) ASMBS position statement on weight bias and stigma. *Surgery for obesity and related diseases* **15**, 814-821.
104. Hill B, Bergmeier H, Incollingo Rodriguez AC *et al.* (2021) Weight stigma and obesity-related policies: A systematic review of the state of the literature. *Obesity Reviews* **22**, e13333.
105. Rubino F, Puhl RM, Cummings DE *et al.* (2020) Joint international consensus statement for ending stigma of obesity. *Nat Med* **26**, 485-497.
106. Braet C, Tanghe A, Decaluwé V *et al.* (2004) Inpatient treatment for children with obesity: weight loss, psychological well-being, and eating behavior. *J Pediatr Psychol* **29**, 519-529.
107. Hunter KE, Webster AC, Page MJ *et al.* (2022) Searching clinical trials registers: guide for systematic reviewers. *BMJ* **377**, e068791.
108. Stewart GB, Altman DG, Askie LM *et al.* (2012) Statistical analysis of individual participant data meta-analyses: a comparison of methods and recommendations for practice. *PloS one* **7**.

109. Sutton AJ, Kendrick D, Coupland CA (2008) Meta-analysis of individual-and aggregate-level data. *Statistics in medicine* **27**, 651-669.
110. Stewart L, Tierney J, Clarke M (2011) Chapter 18: reviews of individual patient data. *Cochrane handbook for systematic reviews of interventions version 5*, 18.11-18.19.
111. Askie LM, Espinoza D, Martin A *et al.* (2020) Interventions commenced by early infancy to prevent childhood obesity-The EPOCH Collaboration: An individual participant data prospective meta-analysis of four randomized controlled trials. *Pediatr Obes* **15**, e12618.
112. Seidler AL, Hunter KE, Baur L *et al.* (2022) Examining the sustainability of effects of early childhood obesity prevention interventions: Follow-up of the EPOCH individual participant data prospective meta-analysis. *Pediatr Obes* **17**, e12919.




Figure 1: Research plan for the Eating Disorders In weight-related Therapy (EDIT) Collaboration. IPD MA, individual participant data meta-analysis; RCTs, Randomised control trials.

List of Tables

Box 1: EDIT Collaboration knowledge gains

Expected knowledge gains

- Diverse community perspectives on the role of individual characteristics and intervention strategies that may increase or decrease eating disorder risk
- Comprehensive understanding of how weight management interventions impact eating disorder risk at an individual level
- Identify participant level predictors of change in eating disorder risk in the context of weight management
- Understand intervention strategies used during weight management interventions and how they may contribute to eating disorder risk
- Find interactions between individual characteristics and intervention components that contribute to an individuals increase or decrease in eating disorder risk
- Development of an international database of weight management interventions measuring eating disorder risk
- Development of recommendations for tailored models of care for obesity treatment