



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Serum Interleukin (IL)-23 and IL-17 Profile in Inflammatory Bowel Disease (IBD) Patients Could Differentiate between Severe and Non-Severe Disease

Citation for published version:

Lucaciu, LA, Ilie, M, Vesa, C, Seicean, R, Din, S, Iuga, CA & Seicean, A 2021, 'Serum Interleukin (IL)-23 and IL-17 Profile in Inflammatory Bowel Disease (IBD) Patients Could Differentiate between Severe and Non-Severe Disease', *Journal of personalized medicine*, vol. 11, no. 11, pp. 1130.
<https://doi.org/10.3390/jpm11111130>

Digital Object Identifier (DOI):

[10.3390/jpm11111130](https://doi.org/10.3390/jpm11111130)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of personalized medicine

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.




Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Article

Serum Interleukin (IL)-23 and IL-17 Profile in Inflammatory Bowel Disease (IBD) Patients Could Differentiate between Severe and Non-Severe Disease

Laura A. Lucaciu ^{1,†} , Maria Ilieș ^{2,†} , Ștefan C. Vesa ^{3,*}, Radu Seicean ⁴, Shahida Din ⁵, Cristina Adela Iuga ^{2,6}  and Andrada Seicean ^{1,7}

- ¹ Department of Gastroenterology and Hepatology, “Iuliu Hațieganu” University of Medicine and Pharmacy, Victor Babeș Street No. 8, 400000 Cluj-Napoca, Romania; lauraa.lucaciu@gmail.com (L.A.L.); andradaseicean@gmail.com (A.S.)
 - ² Department of Proteomics and Metabolomics, MedFuture-Research Centre for Advanced Medicine, “Iuliu Hațieganu” University of Medicine and Pharmacy, Victor Babeș Street No. 8, 400000 Cluj-Napoca, Romania; ilies.maria@umfcluj.ro (M.I.); iugac@umfcluj.ro (C.A.I.)
 - ³ Department of Pharmacology, Toxicology and Clinical Pharmacology, “Iuliu Hațieganu” University of Medicine and Pharmacy, Victor Babeș Street No. 8, 400000 Cluj-Napoca, Romania
 - ⁴ Department of General Surgery, First Surgical Clinic, “Iuliu Hațieganu” University of Medicine and Pharmacy, Victor Babeș Street No. 8, 400000 Cluj-Napoca, Romania; rseicean@yahoo.com
 - ⁵ Edinburgh IBD Unit, Western General Hospital, Edinburgh EH4 2XU, UK; sdin@exseed.ed.ac.uk
 - ⁶ Department of Pharmaceutical Analysis, Faculty of Pharmacy, “Iuliu Hațieganu” University of Medicine and Pharmacy, Victor Babeș Street No. 8, 400000 Cluj-Napoca, Romania
 - ⁷ “Prof. Dr. Octavian Fodor” Regional Institute of Gastroenterology and Hepatology, Croitorilor Street No. 19-21, 400162 Cluj-Napoca, Romania
- * Correspondence: stefan.vesa@umfcluj.ro
† These authors contributed equally to this work. Shared first authorship.



Citation: Lucaciu, L.A.; Ilieș, M.; Vesa, Ș.C.; Seicean, R.; Din, S.; Iuga, C.A.; Seicean, A. Serum Interleukin (IL)-23 and IL-17 Profile in Inflammatory Bowel Disease (IBD) Patients Could Differentiate between Severe and Non-Severe Disease. *J. Pers. Med.* **2021**, *11*, 1130. <https://doi.org/10.3390/jpm11111130>

Academic Editors: Michael Freeley and Kenneth P.H. Pritzker

Received: 16 September 2021
Accepted: 28 October 2021
Published: 2 November 2021

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Interleukin (IL)-17 and IL-23 are crucial for mediating gut mucosal inflammation in inflammatory bowel disease (IBD), which has led to new therapeutic strategies. We assessed the relevancy of IL-17 and IL-23 serum levels as potential biomarkers towards severe IBD discrimination and disease-related complications. Sixty-two patients diagnosed with Crohn’s disease (CD) and ulcerative colitis (UC) were included. Serum IL-17 and IL-23 were measured by sandwich enzyme-linked immunosorbent assays (ELISA). IL-23 and fecal calprotectin (FCal) were significantly higher in severe CD ($p < 0.001$) and UC ($p < 0.001$ and $p = 0.001$, respectively), compared to mild or moderate. Elevated C-reactive protein (CRP) was correlated with severe disease only in CD ($p = 0.008$), whereas for UC, disease severity was associated with increased IL-17 values ($p < 0.001$). Diagnostic role of IL-23 was superior to FCal in discriminating between severe and mild to moderate CD ($p < 0.001$). IL-23 levels were also significantly higher in CD patients with intestinal complications ($p = 0.04$). Both IL-17 and IL-23 correlate with IBD severity, and IL-23 might be a promising novel biomarker for severe CD. Identifying the dominant IL pathway involved in IBD severity could serve as guidance for clinical decision-making on biologic therapy.

Keywords: Crohn’s disease; ulcerative colitis; IBD severity; interleukin-17; interleukin-23; biomarker

1. Introduction

Ulcerative colitis (UC) and Crohn’s disease (CD) are chronic disorders of the gastrointestinal tract with a complex etiology that involves immune, genetic, and environmental factors. Disease progression and severity are highly variable across individuals with inflammatory bowel disease (IBD), which suggests that distinct cytokine pathways may be responsible for the heterogeneity of clinical outcomes [1].

Interleukin (IL)-23/T helper (Th)-17 cytokine pathway was found to have a key role in driving gut inflammation and the development of several other chronic inflammatory

diseases, such as psoriasis, rheumatoid arthritis, and multiple sclerosis [2]. Genome-wide association studies have linked IL-23 to IBD susceptibility via polymorphisms in the IL-23 receptor (IL-23-R) gene that enhances the activity of the IL-17/IL-23 pathway [3].

IL-23 levels were found to be positively correlated with other inflammatory biomarkers such as C-reactive protein (CRP) and rheumatoid factor in rheumatoid arthritis (RA) patients [4,5]. The association between disease activity score (DAS) and rheumatoid arthritis severity scale (RASS) suggested that IL-23 could be used as a biomarker to reflect RA activity [4]. Similarly, in ulcerative colitis patients, IL-23 serum levels were associated with disease severity and duration, leading to it being a possible disease diagnostic marker [6].

Furthermore, new therapeutic agents blocking components of this pathway have been developed; currently only ustekinumab is approved for the treatment of both CD and UC, but other IL-23p19 antagonists are in phase II or III development programs [7]. In contrast, IL-17A inhibition via secukinumab has been shown to worsen the outcomes in Crohn's disease, despite its success in psoriasis [8].

This has driven further research efforts into assessing whether molecular-based stratification with biomarkers according to underlying disease mechanisms could reveal which subgroup of patients would benefit from which therapies. This is particularly important for those with an unfavorable clinical course or complex disease when treatment should be tailored accordingly. Clinical trials aim to incorporate biomarker panels into their design, which are subsequently used in post-hoc analysis, to investigate their relationship with drug response [9]. Different cytokine profiles have been studied as potential biomarkers for response to therapy [10], but their role in predicting disease course is yet to be addressed.

In this study we postulated that quantifying the IL-17 and IL-23 serum levels in patients with IBD could differentiate subgroups of patients with severe and non-severe disease, hence providing a useful clinical tool for patients' stratification. We have tested this premise against biomarkers frequently used in clinical practice to assess disease activity, such as C-reactive protein (CRP), erythrocyte sedimentation rate (ESR), albumin, and fecal calprotectin (FCal).

2. Materials and Methods

2.1. Patients and Study Protocol

This was a pilot case-control prospective study conducted between October 2016 and December 2017. Sixty-two consecutive patients who had CD or UC, were either hospitalized for active disease or attended the day bed unit at the Regional Institute of Gastroenterology and Hepatology Cluj-Napoca, Romania, were recruited.

Eligible participants had a confirmed IBD diagnosis, based on Lennard-Jones criteria [11], serum and fecal samples collected at admission, underwent a colonoscopy or imaging study (CT scan/MRI) in the 6 months prior to admission, and agreed to participate in this study. Baseline demographic data, disease characteristics, phenotype, and the type and duration of IBD treatment (aminosalicylates, corticosteroids, immunosuppressive, and biologic agents) were recorded. Disease location, extension, and behavior were categorized according to the Montreal classification for both CD and UC. Disease activity scores were calculated using Crohn's Disease Activity Index (CDAI) and the Mayo score, respectively [12]. The endoscopic activity was assessed by calculating the Simplified Endoscopic Activity Score for Crohn's Disease (SES-CD) and Mayo endoscopic subscore [12].

Blood samples for inflammatory biomarkers (CRP, ESR, albumin), and fecal samples for FCal measurement were collected during admission as part of hospital protocol. Serum samples for cytokine analysis were aliquoted and stored at -80°C .

Cut-offs for severity thresholds for mild, moderate, or severe disease were established by international consensus, according to European Crohn's and Colitis Organization (ECCO) and American College of Gastroenterology (ACG) guidelines [12]. We classified severe disease patients as having stricturing and/or fistulizing CD, extensive UC, intestinal complications (abscess, fistula, strictures), previous IBD-related surgery, frequent relapses and need for corticosteroids, and >3 courses since diagnosis (Table S1).

The control group consisted of 15 age- and sex-matched subjects referred to our center for outpatient colonoscopy. They were selected from outpatients who did not have a history of autoimmune diseases, had a macroscopically normal colon, and negative fecal and serum inflammatory biomarkers (FCal, CRP, ESR), where available.

Patients that had a suspected or confirmed diagnosis of indeterminate colitis, infectious colitis or malignancy, patients that were pregnant at admission, or those who expressed their refusal to participate were excluded from the study.

2.2. Serum Cytokines Analysis

Serum IL-17 and IL-23 levels were determined using sandwich enzyme-linked immunosorbent assays (ELISA) (human IL-17 Quantikine ELISA kit D1700, human IL-23 Quantikine ELISA kit D2300B, R&D Systems, Minneapolis, MN, USA). Individual serum samples were prepared following manufacturer's instructions and using duplicate measurements. CLARIOstar microplate reader (BMG Labtech, Ortenberg, Germany) was used for reading absorbance. Raw data were acquired using Mars Data Analysis Software (BMG Labtech, Ortenberg, Germany). For quantification (pg/mL), a 4-Parameter fit based calibration curve was generated using known concentrations of the protein standard following the protocol. Final concentrations were calculated as the mean of the two measurements. Other biomarkers (CRP, ESR, albumin and FCal) were measured at the clinic following standard procedures.

2.3. Statistical Analysis

Statistical analysis was performed using MedCalc Statistical Software version 19.0.7 (MedCalc Software bvba, Ostend, Belgium; <https://www.medcalc.org>; 2019). The mean difference between groups for IL-23 was 1433 pg/mL. For a type 1 (alpha) error of 0.05 and a type 2 (beta) error of 0.01 we calculated a sample size of 15 patients per group. Continuous data were tested for normality of distribution using the Shapiro–Wilk test and characterized by mean and standard deviation, or the median and the 25th and the 75th percentiles. Qualitative data were expressed as absolute and relative frequency. Differences between groups were verified with the Mann–Whitney or the chi-square test, as appropriate. Areas under the curve (AUC) were calculated for discriminating between mild/moderate and severe disease. A p value of < 0.05 was considered statistically significant.

2.4. Ethical Considerations

This study was performed in accordance with the WMA Declaration of Helsinki, and written informed consent was obtained from each patient before being included in the study. This study was approved by the Ethical Commission of the “Prof. Dr. Octavian Fodor” Regional Institute of Gastroenterology and Hepatology, Cluj-Napoca, Romania (decision number 16265).

3. Results

3.1. Characteristics of the Study Participants

Table 1 summarizes the demographic and disease-related characteristics of subsets of patients with severe and mild or moderate CD and UC, respectively. Based on the criteria described in the method section (Table S1), 16 patients with CD and 14 UC patients were classified as having mild-to-moderate disease, whereas 15 CD and 17 UC patients were included in the severe disease group. Sixteen patients had anti-TNF treatment, adalimumab or infliximab (Table 1). Eight patients (5 with CD and 3 with UC) had been in remission by the time of inclusion in this study; therefore, they fell in the mild or moderate disease severity category. Younger patients had severe disease in both the CD ($p = 0.01$) and UC ($p = 0.02$) groups. Intestinal complications were more frequent in patients with severe CD ($p = 0.01$). The other variables were not significantly associated with the severity of CD or UC. There were no age or gender-related statistically significant differences between healthy participants and CD or UC patients.

Table 1. Baseline characteristics of the study participants.

	Mild or Moderate CD (n = 16)	Severe CD (n = 15)	p Value	Mild or Moderate UC (n = 14)	Severe UC (n = 17)	p Value	Controls (n = 15)
Age (Years), Mean ± SD	40.62 ± 10.54	31.8 ± 9.01	0.01	52.14 ± 18.79	37.47 ± 13.08	0.01	30.8 ± 6.12
Gender (F), n (%)	11 (68.75)	7 (46.67)	0.3	3 (21.43)	4 (23.53)	1	6 (5.3)
CDAI/Mayo Score, Median (IQR)	105(80.5–147.75)	368 (294–434)	<0.001	6 (2.25–7)	10 (10–12)	<0.001	
Age at Diagnosis, n, % (Montreal Classification, CD)							
A1 (n, %): <16	0 (0)	2 (13.33)	1				
A2 (n, %): 17–40	13 (81.25)	13 (86.67)					
A3 (n, %): >40	3 (18.75)	0 (0)					
Location, n, % (Montreal Classification, CD)							
L1: Ileum	10 (62.5)	5 (33.33)	0.2				
L2: Colon	2 (12.5)	2 (13.33)					
L3: Ileocolonic	4 (25)	8 (53.33)					
Location, n, % (Montreal Classification, UC)							
E1 (Distal Colitis)	-	-		1	1		
E2 (Left-Sided Colitis)	-	-		10	13		
E3 (Pancolitis)	-	-		3	3		
Behaviour, n, % (Montreal Classification, CD)							
B1: Non-Structuring, Non-Penetrating	11 (68.75)	0 (0)	<0.001				
B2: Structuring	2 (12.5)	6 (40)					
B3: Penetrating	2 (12.5)	7 (46.67)					
Mayo Endoscopic Score, Median, (IQR)				1 (0.5–1)	3 (2–3)		
SES-CD, Median (IQR)	5 (4–7)	13.5 (12–17.5)					
Complications, n, %, CD and UC							
Intestinal Complications, n (%)	With	6 (37.5)	13 (86.67)	0.01	0 (0)	2 (11.76)	
	Without	10 (62.5)	2 (13.3)		14 (100)	15 (88.2)	
Fistula, n (%)		0 (0)	1 (5.88)	0.08	-	-	-
Stenosis, n (%)		2 (12.5)	7 (46.7)	0.05	-	1 (5.9%)	1
Abscess, n (%)		0 (0)	2 (13.33)	0.2	-	-	-
Extraintestinal Complications, n (%)		3 (18.75)	2 (13.33)	1	-	1 (5.9)	1
Anti-TNF Therapy, n (%)		5 (31.25)	4 (26.67)	1	3 (21.43)	4 (25)	1

CD, Crohn’s disease. SD, standard deviation. UC, ulcerative colitis. SD, standard deviation. IQR, interquartile range.

3.2. Standard Inflammatory Biomarker Levels in IBD Patients by Disease Severity vs. Healthy Participants

Comparisons among groups between standard inflammatory biomarkers levels are shown in Figure 1 and Supplementary Table S2. FCal levels were significantly elevated in patients with severe CD and UC ($p < 0.001$) vs. mild or moderate disease groups and controls, with higher median values in UC patients. Higher CRP levels were associated with severe disease only in CD patients ($p = 0.008$). Lower albumin levels were more frequently observed in patients with severe UC ($p = 0.008$). ESR levels comparison between different groups of IBD severity did not yield significant results ($p = 0.1$).




Figure 1. Standard inflammatory biomarker distributions in IBD patients according to disease severity: (A) FCal levels in mild-moderate vs. severe CD and mild-moderate vs. severe UC groups. (B) CRP serum levels in mild-moderate vs. severe CD and mild-moderate vs. severe UC groups. (C) Albumin serum levels in mild-moderate vs. severe CD and mild-moderate vs. severe UC groups. (D) ESR serum levels in mild-moderate vs. severe CD and mild-moderate vs. severe UC groups. Biomarker levels in controls are also shown.

3.3. IL-17 and IL-23 Quantitative Serum Levels in IBD Patients vs. Healthy Controls

IL-17 levels were significantly higher in CD patients than in controls, with a median level of 949 (IQR 400.8–2242.9) pg/mL vs. 484.8 (IQR 139–560.5) pg/mL, $p = 0.002$. IL-23 levels were significantly higher in CD patients than in controls, with a median level of 937.4 (IQR 773.3–1477.4) pg/mL vs. 371.5 (IQR 362.8–380.1) pg/mL, $p < 0.001$.

In UC patients, IL-17 levels were significantly higher than in the control group, with a median level of 1167 (IQR 642.6–2405.7) pg/mL vs. 484.8 (IQR 139–560.5) pg/mL, $p < 0.001$. IL-23 levels were significantly higher in UC patients than in controls: 1365.1 (751.7–1512) pg/mL vs. 371.5 (362.8; 380.1) pg/mL, $p < 0.001$ (Figure 2 and Table S2).




Figure 2. IL-17 and IL-23 quantitative serum level distribution in IBD patients and the control group: (A) IL-17 serum levels in UC patients, CD patients, and the control group. (B) IL-23 serum levels in UC patients, CD patients, and the control group.

3.4. IL-17 and IL-23 Quantitative Serum Levels Associated with IBD Severity vs. Healthy Controls

IL-23 serum levels were significantly elevated in patients with severe CD and UC ($p < 0.001$), whereas IL-17 levels were significantly higher only in UC patients with severe disease ($p < 0.001$). In patients with severe CD, serum IL-17 levels were not significantly higher than in those with mild or moderate disease ($p = 0.1$) (Figure 3 and Table S2).




Figure 3. IL-17 and IL-23 quantitative serum level distribution in IBD patients according to disease severity: (A) IL-17 serum levels in mild-moderate vs. severe CD groups and mild-moderate vs. severe UC groups. (B) IL-23 serum levels in mild-moderate vs. severe CD groups and mild-moderate vs. severe UC groups.

3.5. Discriminating between Severe and Mild or Moderate IBD according to Biomarker Cut-Off Value

We calculated the performance of cut-off values for the biomarkers that discriminated between mild/moderate and severe disease (Table 2) using ROC (Receiver Operating Characteristic) Curve. Cut-off values for each biomarker were established where sensitivity and specificity were maximal. A significantly higher area under the ROC curve (AUC) of IL-23 for discriminating patients with severe CD was found, as compared to the AUC of CRP, FCal, or albumin ($p = 0.003$; $p = 0.04$; $p = 0.01$, respectively) (Table 2). For the identification of severe UC cases, the IL-23's AUC was only slightly better than FCal ($p = 0.09$), but significantly higher than the AUC of IL-17 or albumin ($p = 0.02$; $p = 0.01$, respectively). The specificity of IL-23 in discriminating severe UC cases was found to be higher compared to the FCal. The assessment of IL-17 and IL-23 levels according to biologic therapy has not yielded significant results ($p = 0.274$ and $p = 0.184$, respectively).

Table 2. AUC values for assessing disease severity.

Variable	AUC (CI 95%)	Cut-off	Sensitivity (CI 95%)	Specificity (CI 95%)	p Value	
CD	Albumin	0.733 (0.545–0.875)	3.7 g/dL	93.75 (69.8–99.8)	53.33 (26.6–78.7)	0.010
	CRP	0.781 (0.597–0.909)	1.26 mg/L	73.33 (44.9–92.2)	87.50 (61.7–98.4)	0.001
	FCal	0.877 (0.709–0.967)	50 µg/g	100 (78.2–100)	62.5 (35.4–84.8)	<0.001
	IL23	1 (0.888–1.000)	937.4 pg/mL	100 (78.2–100.0)	100 (79.4–100.0)	<0.001
	IL-17	0.667 (0.475–0.825)	>1536.15 pg/mL	53.33 (26.6–78.7)	87.50 (61.7–98.4)	0.010
UC	Albumin	0.773 (0.588–0.903)	3.7 g/dL	92.86 (66.1–99.8)	47.06 (23.0–72.2)	<0.001
	FCal	0.857 (0.685–0.956)	400 µg/g	94.12 (71.3–99.9)	71.43 (41.9–91.6)	<0.001
	IL-17	0.803 (0.621–0.923)	1512.05 pg/mL	64.71 (38.3–85.8)	100 (76.8–100)	<0.001
	IL-23	0.979 (0.851–1)	946.11 pg/mL	94.12 (71.3–99.9)	100 (76.8–100)	<0.001
	CRP	0.645 (0.453–0.808)	>0.89 mg/L	52.94 (27.8–77.0)	85.71 (57.2–98.2)	0.010

AUC, Area Under the Curve. CI, Confidence Interval. CD, Crohn’s disease. UC, Ulcerative Colitis. FCal, Faecal Calprotectin.

3.6. Assessment of CD Disease Complications according to Standard Inflammatory Biomarkers vs. IL-17 and IL-23 Serum Levels

We compared the median biomarker levels in CD patients with and without intestinal complications such as enteral fistulae, abscesses, or need for surgery (Table 3). IL-23 and, to a lesser degree, IL-17 had significantly higher levels in patients with intestinal complications ($p = 0.04$ and $p = 0.05$, respectively). There was a positive point-biserial (Pearson’s) correlation between both IL-17 and IL-23 and the presence of intestinal complications ($r_{pb} = 0.085$ and 0.243); this was statistically significant for IL-23 ($p = 0.04$) but not for IL-17 ($p = 0.51$).

Table 3. IL-17, IL-23, and standard inflammatory biomarker levels in patients with or without CD-related complications.

Biomarkers’ Serum Levels (Median, IQR)	Intestinal Complications		p Value
	Yes (n = 19)	No (n = 12)	
CRP	1.26 (0.36–3.80)	0.44 (0.33–6.44)	0.60
Albumin	4 (3.7; 4.2)	3.9 (3.65; 4.37)	0.90
Calprotectin	400 (150–850)	135 (50–400)	0.20
ESR	52 (21–82)	27 (13–90)	0.40
IL-17	1449.05 (708.50–2531.60)	638.77 (242.93–1322.96)	0.05
IL-23	1417.01 (799.22–1486.13)	797.06 (744.14–912.63)	0.04

IQR, Interquartile range. CRP, C-reactive protein. ESR, Erythrocyte sedimentation rate. IL, Interleukin.

4. Discussion

In order to achieve the goal of precision medicine, a clinical tool towards IBD patient stratification is urgently needed. So far, a molecular-based biomarker that could differentiate between subgroups of patients with severe and non-severe IBD is absent. We have therefore shown that serum levels of IL-17 and IL-23 could become a valuable biomarker for assessing disease severity subtypes in both CD and UC, by testing these molecules against biomarkers frequently used in clinical practice such as CRP, ESR, albumin, and FCal.

The results of our study highlight that serum IL-23 levels showed superior diagnostic potential to that of dedicated inflammatory biomarkers for IBD, such as FCal, CRP, and albumin. In contrast, higher IL-17 serum levels only demonstrated a modest diagnostic performance for both severe UC and CD. In addition, IL-17 and IL-23 serum levels were significantly more elevated in IBD patients compared to the control group.

It is known that cytokines amplify and sustain the inflammatory response in IBD; hence, elevated concentration in biological fluids is expected. This also indicates the pathway activation that drives inflammation and may lead to disease progression [13]. IL-23 is released by monocytes/macrophages/dendritic cells in response to bacterial stimulation, and subsequent interaction with its heterodimeric IL-23 receptor (IL-23R) leads

to production of inflammatory mediators (IL-17, IL-22, granulocyte-macrophage colony-stimulating factor (GM-CSF), TNF- α) through JAK-STAT pathway activation [14,15]. IL-17 acts as a proinflammatory cytokine produced by Th-17 cells [16] that has divergent functions, contributing to chronic inflammation via recruitment and activation of inflammatory mediators, and host defense against extracellular bacterial and fungal infections [17].

Elevated levels of IL-17 and IL-23 in IBD patients have previously been reported separately [6,18–21]. Fujino et al. [18] were the first to demonstrate a positive correlation between elevated levels of IL-17 in the inflamed mucosa and active lesions of IBD patients, as well as elevated serum levels of IL-17 and disease activity in both CD and UC. Similarly, Jiang et al. reported increased expression of Th-17 cytokines (IL-17, IL-21, and IL-22) in the intestinal mucosa in active IBD patients [19].

Studies assessing IL-23 expression in human IBD have also shown increased IL-23 production by lamina propria macrophages from CD patients but not in UC [20]. These findings are supported by studies in patients with concomitant inflammatory conditions, such as arthritis and sacroileitis [21], which reported higher IL-23 levels in CD rather than in those with UC. In contrast, Mirsattari et al. [6] showed that higher IL-23 levels correlate with both duration and severity of the disease (based on a modified Mayo score) but only in UC patients. These findings confirm the important role of IL-23 in IBD pathogenesis, making it not only an important therapeutic target but also a potential biomarker for IBD severity and prognosis.

Understanding and addressing disease prognosis in IBD is the basis for developing a precision medicine approach [22]. Stratifying patients in a timely manner and individualizing therapy is a continuous challenge in clinical practice. Many clinical parameters, such as serological markers, disease location, disease behavior, age and lifestyle, have been found to be associated with disease severity [23]. However, most of these predictors derive from observational studies and are not specific enough to alter early disease management. Therefore, the need for reliable biomarkers is paramount.

We obtained a high AUC for IL-23 in detecting CD patients prone to more severe disease. Furthermore, this was superior to the standard inflammation markers used in the clinical practice (CRP, FCal, albumin), therefore acting as potential noninvasive biomarker for identifying a complex disease phenotype in patients with CD.

While previous studies assessing the expression of IL-17 and IL-23 in IBD patients have classified patients according to “disease severity”, they have primarily relied on disease activity scores that have been symptom based at a point in time; however, patients may have severe disease, warranting aggressive therapies even if their disease presentation is not severe. For example, patients with extensive steroid-dependent ulcerative colitis or Crohn’s disease refractory to immunosuppressive therapies, with mild symptoms but on high doses of corticosteroids, can be considered [24]. We have therefore taken into consideration a more comprehensive approach when defining disease severity, beyond merely clinical presentation, such as disease burden and structural damage.

In CD patients IL-23 and IL-17 were significantly higher in patients who developed intestinal complications, such as fistulae, abscesses, and need for surgery at a certain point in time. Moreover, the point-biserial correlation coefficient showed a statistically significant correlation between IL-23 and CD patients with intestinal complications. It was not possible to comparatively assess complications for UC patients due to the small number of patients that had disease-related complications. Several biomarkers that are well established in clinical practice have been used to predict a complicated disease course in IBD. FCal has a crucial role in IBD diagnosis and disease activity assessment, prediction of relapse, response to therapy [25], disease extension, and severity [26]. Our findings have demonstrated inferior diagnostic value to that of IL-23 in discriminating a more severe CD outcome; in the UC cohort, however, the sensitivities for IL-23 and FCal were comparable, but with higher specificity for IL-23.

CRP is one of the most reliable and widely used, in conjunction with FCal, to assess IBD flares in clinical practice. Previous studies have shown that a high CRP is indicative of

severe disease complications (perforations, abscesses, and fistula) in CD [25,26], whereas in UC it is correlated with disease activity and severe symptoms but not with histologic inflammation [27]. In our study, CRP was inferior to IL-23 in accurately assessing disease severity in both CD and UC groups.

The ESR is a non-specific measure of systemic inflammation, and it can be used to monitor the acute-phase response of disease after 24 h [28]. We have shown that elevated ESR levels had no diagnostic value for a more severe disease course and was not correlated with the occurrence of intestinal complications in CD patients.

Serum albumin had the lowest specificity for diagnosing severe disease when compared to IL-17 and IL-23, amongst the other inflammatory biomarkers. Previous studies have shown that hypoalbuminemia in CD patients could predict poor postoperative outcomes [29], whereas in those with acute severe UC it is correlated with treatment failure, higher colectomy rate [30], and non-response to anti-TNF therapy [31]. In our CD patients, however, albumin levels had no significant association with disease-related complications.

The predictive potential of quantifying serum cytokines in IBD patients has previously been explored in the development of primary non-response in patients treated with anti-TNF α [32] and vedolizumab [33]. Furthermore, a transcriptomic analysis of the colonic mucosa in patients refractory to anti-TNF therapy showed that IL-23 gene expression is regulated upon anti-TNF blockade [34]; therefore, anti-TNF therapy is already effective in suppressing IL-23. It has been postulated that non-responders to anti-TNF might have less to gain from further IL-23 targeting. This is supported by the landmark UNITI-1, UNITI-2, and UNITI-IM studies that have shown better response rates in TNF-naive than in TNF experienced populations (54–58% vs 34%, respectively) [35]. This suggests that stratifying patients with different degrees of IBD severity, based on IL-17 and IL-23 serum profiling, could eventually be useful in guiding the choice of biologic therapy. This would entail, however, longitudinal comparisons between groups undergoing different targeted therapies.

This is the first study to explore the use of IL-17 and IL-23 in stratifying IBD patients by disease severity, in comparison with standard inflammatory tests used in clinical practice. Our results highlight that IL-23 was particularly efficient in differentiating IBD patients with a severe disease phenotype and that its performance surpassed that of faecal calprotectin. By also demonstrating a significant association with CD intestinal complications, we believe that IL-23 could become a reliable tool for stratifying CD patients since diagnosis. In comparison with the data already published, our findings have taken a step forward towards implementing these potential biomarkers in clinical practice.

So far, our results are based only on a small number of patient samples and a single time-point analysis. Serum cytokines and standard biomarker serum levels could not be included for comparison to the histological findings in the study as they were not assessed at the same time point. However, long term follow-up data for the IBD and UC patients are the focus of our future research, and the current findings will be applied with respect to disease progression in a future study.

5. Conclusions

Serum IL-23 levels were higher in both CD and UC patients and more effective than FCal in identifying the group with the highest disease severity. A significant correlation with intestinal complications was demonstrated in CD patients; therefore, IL-23 could become a useful biomarker in precision medicine. IL-17 was more elevated in UC patients with severe disease rather than CD but yielded lower diagnostic accuracy for disease severity when compared to other biomarkers. Further larger studies are necessary to establish whether the assessment of these cytokines in unresponsive patients to anti-TNF therapy could guide the choice of second-line biologics.

Supplementary Materials: The following are available online at <https://www.mdpi.com/article/10.3390/jpm11111130/s1>, Table S1: Classification of study participants according to disease sever-

ity, Table S2: Comparison between groups according to IL-17, IL-23, and standard inflammatory biomarkers levels.

Author Contributions: Conceptualization, L.A.L.; Data curation, L.A.L., Ş.C.V. and S.D.; Formal analysis, L.A.L., M.I. and Ş.C.V.; Funding acquisition, L.A.L.; Investigation, L.A.L.; Methodology, L.A.L.; Project administration, L.A.L. and A.S.; Resources, L.A.L.; Software, L.A.L., M.I. and Ş.C.V.; Supervision, Ş.C.V., R.S., C.A.I. and A.S.; Writing—original draft, L.A.L.; Writing—review and editing, L.A.L., M.I., Ş.C.V., R.S., S.D. and C.A.I. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the research grant no. 36/13/01/2017 awarded by “Iuliu Hațieganu” University of Medicine and Pharmacy Cluj-Napoca, Romania to Laura A Lucaciu.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Ethics Committee of “Prof. Dr. Octavian Fodor” Regional Institute of Gastroenterology and Hepatology (decision number 16265).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Marafini, I.; Sedda, S.; Dinallo, V.; Monteleone, G. Inflammatory cytokines: From discoveries to therapies in IBD. *Expert Opin. Biol. Ther.* **2019**, *19*, 1207–1217. [[CrossRef](#)]
2. Neurath, M.F. IL-23 in inflammatory bowel diseases and colon cancer. *Cytokine Growth Factor Rev.* **2019**, *45*, 1–8. [[CrossRef](#)]
3. Duerr, R.H.; Taylor, K.D.; Brant, S.R.; Rioux, J.D.; Silverberg, M.S.; Daly, M.J.; Steinhart, A.H.; Abraham, C.; Regueiro, M.; Griffiths, A.; et al. A genome-wide association study identifies IL23R as an inflammatory bowel disease gene. *Science* **2006**, *314*, 1461–1463. [[CrossRef](#)] [[PubMed](#)]
4. Alsheikh, M.M.; El-Shafey, A.M.; Gawish, H.H.; El-Desoky, E.T. Serum interleukin-23 level in rheumatoid arthritis patients: Relation to disease activity and severity. *Egypt. Rheumatol.* **2019**, *41*, 99–103. [[CrossRef](#)]
5. Abdo, A.I.K.; Tye, G.J. Interleukin 23 and autoimmune diseases: Current and possible future therapies. *Inflamm. Res.* **2020**, *69*, 463–480. [[CrossRef](#)] [[PubMed](#)]
6. Mirsattari, D.; Seyyedmajidi, M.; Zojaji, H.; Haghazali, M.; Orimi, P.G.; Shoushtarizadeh, T. The relation between the level of interleukin-23 with duration and severity of ulcerative colitis. *Gastroenterol. Hepatol.* **2012**, *5*, 49–53.
7. Sabino, J.; Verstockt, B.; Vermeire, S.; Ferrante, M. New biologics and small molecules in inflammatory bowel disease: An update. *Therap. Adv. Gastroenterol.* **2019**, *12*, 1756284819853208. [[CrossRef](#)]
8. Hueber, W.; Sands, B.E.; Lewitzky, S.; Vandemeulebroecke, M.; Reinisch, W.; Higgins, P.D.R.; Wehkamp, J.; Feagan, B.G.; Yao, M.D.; Karczewski, M.; et al. Secukinumab, a human anti-IL-17A monoclonal antibody, for moderate to severe Crohn’s disease: Unexpected results of a randomised, double-blind placebo-controlled trial. *Gut* **2012**, *61*, 1693–1700. [[CrossRef](#)]
9. Boyapati, R.K.; Kalla, R.; Satsangi, J.; Ho, G. Biomarkers in search of precision medicine in IBD. *Am. J. Gastroenterol.* **2016**, *111*, 1682–1690. [[CrossRef](#)] [[PubMed](#)]
10. Yarur, A.J.; Jain, A.; Quintero, M.A.; Czyl, F.; Deshpande, A.R.; Kerman, D.H.; Abreu, M.T. Inflammatory Cytokine Profile in Crohn’s Disease Nonresponders to Optimal Antitumor Necrosis Factor Therapy. *J. Clin. Gastroenterol.* **2019**, *53*, 210–215. [[CrossRef](#)] [[PubMed](#)]
11. Lennard-Jones, J.E. Classification of Inflammatory Bowel Disease. *Scand. J. Gastroenterol.* **2009**, *24*, 2–6. [[CrossRef](#)]
12. Sturm, A.; Maaser, C.; Calabrese, E.; Annesse, V.; Fiorino, G.; Kucharzik, T.; Vavricka, S.R.; Verstockt, B.; van Rheenen, P.; Tolan, D.; et al. ECCO-ESGAR Guideline for Diagnostic Assessment in IBD Part 2: IBD scores and general principles and technical aspects. *J. Crohns Colitis* **2019**, *13*, 273–284. [[CrossRef](#)] [[PubMed](#)]
13. Park, J.H.; Peyrin-Biroulet, L.; Eisenhut, M.; Shin, J.I. IBD immunopathogenesis: A comprehensive review of inflammatory molecules. *Autoimmun. Rev.* **2017**, *16*, 416–426. [[CrossRef](#)]
14. Moschen, A.R.; Tilg, H.; Raine, T. IL-12, IL-23 and IL-17 in IBD: Immunobiology and therapeutic targeting. *Nat. Rev. Gastroenterol. Hepatol.* **2019**, *16*, 185–196. [[CrossRef](#)]
15. Argollo, M.; Fiorino, G.; Hindryckx, P.; Peyrin-Biroulet, L.; Danese, S. Novel therapeutic targets for inflammatory bowel disease. *J. Autoimmun.* **2017**, *85*, 103–116. [[CrossRef](#)]
16. Abraham, C.; Dulai, P.S.; Vermeire, S.; Sandborn, W.J. Lessons Learned From Trials Targeting Cytokine Pathways in Patients with Inflammatory Bowel Diseases. *Gastroenterology* **2017**, *152*, 374–388. [[CrossRef](#)]
17. Omidian, Z.; Ahmed, R.; Giwa, A.; Donner, T.; Hamad, A.R.A. IL-17 and limits of success. *Cell. Immunol.* **2019**, *339*, 33–40. [[CrossRef](#)] [[PubMed](#)]
18. Fujino, S.; Andoh, A.; Bamba, S.; Ogawa, A.; Hata, K.; Araki, Y.; Bamba, T.; Fujiyama, Y. Increased expression of interleukin 17 in inflammatory bowel disease. *Gut* **2003**, *52*, 65–70. [[CrossRef](#)]

19. Jiang, W.; Su, J.; Zhang, X.; Cheng, X.; Zhou, J.; Shi, R.; Zhang, H. Elevated levels of Th17 cells and Th17-related cytokines are associated with disease activity in patients with inflammatory bowel disease. *Inflamm. Res.* **2014**, *63*, 943–950. [[CrossRef](#)] [[PubMed](#)]
20. Fuss, I.J.; Becker, C.; Yang, Z.; Groden, C.; Hornung, R.L.; Heller, F.; Neurath, M.F.; Strober, W.; Mannon, P.J. Both IL-12p70 and IL-23 are synthesized during active Crohn's disease and are down-regulated by treatment with anti-IL-12 p40 monoclonal antibody. *Inflamm. Bowel Dis.* **2006**, *12*, 9–15. [[CrossRef](#)] [[PubMed](#)]
21. Gheita, T.A.; El Gazzar, I.I.; El-Fishawy, H.S.; Aboul-Ezz, M.A.; Kenawy, S.A. Involvement of IL-23 in enteropathic arthritis patients with inflammatory bowel disease: Preliminary results. *Clin. Rheumatol.* **2014**, *33*, 713–717. [[CrossRef](#)]
22. Verstockt, B.; Noor, N.M.; Marigorta, U.M.; Pavlidis, P.; Deepak, P.; Ungaro, R.C. Results of the Seventh Scientific Workshop of ECCO: Precision medicine in IBD—Disease outcome and response to therapy. *J. Crohns Colitis* **2021**, *17*, jjab050.
23. Sandborn, W.J. Crohn's disease evaluation and treatment: Clinical decision tool. *Gastroenterology* **2014**, *147*, 702–705. [[CrossRef](#)] [[PubMed](#)]
24. Peyrin-Biroulet, L.; Panés, L.; Sandborn, W.J.; Vermeire, S.; Danese, S.; Feagan, B.G.; Colombel, J.F.; Hanauer, S.B.; Rycroft, B. Defining Disease Severity in Inflammatory Bowel Diseases: Current and Future Directions. *Clin. Gastroenterol. Hepatol.* **2016**, *14*, 348–354. [[CrossRef](#)] [[PubMed](#)]
25. Soubieres, A.A.; Poullis, A. Emerging Biomarkers for the Diagnosis and Monitoring of Inflammatory Bowel Diseases. *Inflamm. Bowel Dis.* **2016**, *22*, 2016–2022. [[CrossRef](#)]
26. Kawashima, K.; Ishihara, S.; Yuki, T.; Fukuba, N.; Oshima, N.; Kazumori, H.; Sonoyama, H.; Yamashita, N.; Tada, Y.; Kusunoki, R.; et al. Fecal calprotectin level correlated with both endoscopic severity and disease extent in ulcerative colitis. *BMC Gastroenterol.* **2016**, *16*, 47. [[CrossRef](#)] [[PubMed](#)]
27. Rogler, G.; Biedermann, L. Clinical Utility of Biomarkers in IBD. *Curr. Gastroenterol. Rep.* **2015**, *17*, 26. [[CrossRef](#)] [[PubMed](#)]
28. Lichtenstein, G.R.; McGovern, D.P.B. Using Markers in IBD to Predict Disease and Treatment Outcomes: Rationale and a Review of Current Status. *Am. J. Gastroenterol. Suppl.* **2016**, *3*, 17–26. [[CrossRef](#)]
29. Nguyen, G.C.; Du, L.; Chong, R.Y.; Jackson, T.D. Hypoalbuminaemia and Postoperative Outcomes in Inflammatory Bowel Disease: The NSQIP Surgical Cohort. *J. Crohns Colitis* **2019**, *13*, 1433–1438. [[CrossRef](#)]
30. Kumar, S.; Ghoshal, U.C.; Aggarwal, R.; Saraswat, V.A.; Choudhuri, G. Severe ulcerative colitis: Prospective study of parameters determining outcome. *J. Gastroenterol. Hepatol.* **2004**, *19*, 1247–1252. [[CrossRef](#)]
31. Morita, Y.; Bamba, S.; Takahashi, K.; Imaeda, H.; Nishida, A.; Inatomi, O.; Sasaki, M.; Tsujikawa, T.; Sugimoto, M.; Andoh, A. Prediction of clinical and endoscopic responses to anti-tumor necrosis factor- α antibodies in ulcerative colitis. *Scand. J. Gastroenterol.* **2016**, *51*, 934–941. [[CrossRef](#)] [[PubMed](#)]
32. Billiet, T.; Cleynen, I.; Ballet, V.; Claes, K.; Princen, F.; Singh, S.; Ferrante, M.; Van Assche, G.; Gils, A.; Vermeire, S. Evolution of cytokines and inflammatory biomarkers during infliximab induction therapy and the impact of inflammatory burden on primary response in patients with Crohn's disease. *Scand. J. Gastroenterol.* **2017**, *52*, 1086–1092. [[CrossRef](#)] [[PubMed](#)]
33. Bertani, L.; Baglietto, L.; Antonioli, L.; Fornai, M.; Tapete, G.; Albano, E.; Ceccarelli, L.; Mumolo, M.G.; Pellegrini, C.; Lucenteforte, E.; et al. Assessment of serum cytokines predicts clinical and endoscopic outcomes to vedolizumab in ulcerative colitis patients. *Br. J. Clin. Pharmacol.* **2020**, *86*, 1296–1305. [[CrossRef](#)] [[PubMed](#)]
34. Leal, R.F.; Planell, N.; Kajekar, R.; Lozano, J.J.; Ordas, I.; Dotti, I.; Esteller, M.; Masamunt, M.C.; Parmar, H.; Ricart, E.; et al. Identification of inflammatory mediators in patients with Crohn's disease unresponsive to anti-TNF α therapy. *Gut* **2015**, *64*, 233–242. [[CrossRef](#)]
35. Sandborn, W.; Feagan, B.G.; Gasink, C.; Jacobstein, D.; Gao, L.-L.; Johanns, J.; Sands, B.E.; Hanauer, S.B.; Targan, S.; Ghosh, S.; et al. 768 A Phase 3 Randomized, Multicenter, Double-Blind, Placebo-Controlled Study of Ustekinumab Maintenance Therapy in Moderate—Severe Crohn's Disease Patients: Results From IM-UNITI. *Gastroenterology* **2016**, *150*, S157–S158. [[CrossRef](#)]