



OPEN ACCESS

Survey of liver pathologists to assess attitudes towards digital pathology and artificial intelligence

Clare McGenity ^{1,2}, Rebecca Randell,^{3,4} Christopher Bellamy,⁵ Alastair Burt,⁶ Alyn Cratchley,² Robert Goldin,⁷ Stefan G Hubscher,⁸ Desley A H Neil ^{8,9}, Alberto Quaglia,¹⁰ Dina Tiniakos,^{11,12} Judy Wyatt,² Darren Treanor^{1,2,13,14}

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/jcp-2022-208614>).

For numbered affiliations see end of article.

Correspondence to

Dr Clare McGenity, University of Leeds, Leeds, UK; medcmcga@leeds.ac.uk

Received 6 October 2022
Accepted 24 November 2022

ABSTRACT

Aims A survey of members of the UK Liver Pathology Group (UKLPG) was conducted, comprising consultant histopathologists from across the UK who report liver specimens and participate in the UK National Liver Pathology External Quality Assurance scheme. The aim of this study was to understand attitudes and priorities of liver pathologists towards digital pathology and artificial intelligence (AI).

Methods The survey was distributed to all full consultant members of the UKLPG via email. This comprised 50 questions, with 48 multiple choice questions and 2 free-text questions at the end, covering a range of topics and concepts pertaining to the use of digital pathology and AI in liver disease.

Results Forty-two consultant histopathologists completed the survey, representing 36% of fully registered members of the UKLPG (42/116). Questions examining digital pathology showed respondents agreed with the utility of digital pathology for primary diagnosis 83% (34/41), second opinions 90% (37/41), research 85% (35/41) and training and education 95% (39/41). Fatty liver diseases were an area of demand for AI tools with 80% in agreement (33/41), followed by neoplastic liver diseases with 59% in agreement (24/41). Participants were concerned about AI development without pathologist involvement 73% (30/41), however, 63% (26/41) disagreed when asked whether AI would replace pathologists.

Conclusions This study outlines current interest, priorities for research and concerns around digital pathology and AI for liver pathologists. The majority of UK liver pathologists are in favour of the application of digital pathology and AI in clinical practice, research and education.

INTRODUCTION

Liver diseases are a major cause of global morbidity and mortality. Approximately two million deaths worldwide and over 150 000 across Europe are due to liver disease each year.^{1 2} Preventable deaths due to liver conditions amounted to 26 265 in England alone from 2015 to 2017.³ In liver disease, the histopathologist's tissue assessment frequently plays an important role in determining diagnosis, prognosis and response to treatment for many conditions, especially where there is uncertainty in the patient's clinical history and presentation.^{4 5} However, overall demand for pathology services is rising, the content of pathology reports is

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Interest in the uses and potential of digital pathology and artificial intelligence (AI) has grown in recent years. These technologies will influence the diagnosis and management of liver diseases.

WHAT THIS STUDY ADDS

⇒ This is the first study to examine the views of UK liver pathologists around digital pathology and AI and to highlight research priorities within liver pathology.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ The authors anticipate that this will influence the direction of research for digital pathology and AI applied to liver disease, by highlighting the demand and concerns from the UK liver pathology community.

increasingly complex and histopathology laboratories face ongoing workforce challenges.^{6 7}

Digital pathology is the digitisation of glass slides to make high-resolution images available for viewing on a computer screen.⁸ This is achieved through whole slide imaging, involving the use of scanners to capture an entire glass slide.⁸ Work is currently ongoing to deploy digital pathology to hospitals across the UK.⁹ Image analysis techniques ranging from conventional computerised morphology tools through to artificial intelligence (AI) have been used in pathology research for some years, although recent AI developments in deep learning have inspired researchers further.^{10 11} The increasing availability of digital pathology is bringing efforts to develop and clinically deploy AI tools closer to reality. In the face of the workforce and workflow challenges in pathology laboratories, the application of digital pathology and AI as tools to assist the histopathologist in reporting of liver specimens has generated substantial enthusiasm.^{12–17} However, understanding and targeting areas where most benefit can be achieved is important in addressing clinical demands and avoiding research waste.

The UK Liver Pathology Group (UKLPG) is a professional association that arose from the collaboration of several groups with liver pathology interests.¹⁸ They held their first meeting as the UKLPG in 2016 and its membership comprises consultant histopathologists (full members) from across the



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY. Published by BMJ.

To cite: McGenity C, Randell R, Bellamy C, et al. *J Clin Pathol* Epub ahead of print: [please include Day Month Year]. doi:10.1136/jcp-2022-208614

UK who report liver specimens in their clinical practice.¹⁸ The UKLPG is also responsible for running the UK liver pathology External Quality Assurance (EQA) scheme, and therefore, encompasses the vast majority of histopathologists who report liver cases from across the country. An academic subgroup of the UKLPG membership formed in 2020 to collaborate and focus on the development of digital pathology and AI for use in liver disease. Acknowledging the growing interest in this area, the group aimed to understand current attitudes and research priorities of UKLPG members for digital pathology and AI, and to report this to the wider research community.

METHODS

Study design

A cross-sectional survey was developed by three members of the UKLPG (CM, JW and DTr) from February 2021 to April 2021. Broader consultation across a series of meetings with the digital pathology and AI academic subgroup of the UKLPG involved exploring experiences and opinions of group members, considering publications in this area and identifying areas of clinical utility for these technologies to inform the survey content. Further discussions with a qualitative researcher (RR) took place in helping to inform the design and content of the survey. The rationale for canvassing opinion from the UK liver pathology community and the full survey were reviewed and approved by the UKLPG committee prior to distribution. The authors are not aware of any studies attempting to gather opinion from the UK liver pathology community on the subject of digital pathology and AI prior to this exercise. The survey comprised a total of 50 questions, divided into 11 sections of multiple choice questions (MCQs) and 2 free text questions at the end to allow open responses. For MCQs addressing topics on digital pathology and AI, the answers were selected from a five point grading system from 'strongly disagree' to 'strongly agree' and there was an additional sixth option to state 'don't know' if needed. The survey aimed to assess attitudes to digital pathology and AI for use in liver disease, respectively. The full text of the questionnaire is in online supplemental document 1.

Survey distribution

The survey was circulated to all full members of the UKLPG via email (CM and JW) with a link to the SurveyMonkey online tool (www.surveymonkey.com) and included accompanying guidance for completion. It was sent to members in a routine mailing list email and was not mandatory to complete. Responses were collected from April 2021 to May 2021, and reminder emails and a deadline extension were given to encourage participation (CM and JW).

Data analysis

Analyses for each question were performed using Microsoft Excel software (CM). Given the range of knowledge levels across the community, with some members having no experience and others with extensive experience of these technologies, none of the questions were mandatory. Some questions allowed more than one response to gather a wider reflection of participant experience. Therefore, the summary data are given on a per question basis. Despite this, responses ranged from totals of 40–42 responses per question from the total of 42 participants, as only a maximum of two respondents skipped any individual question.

UKLPG membership involvement

Members of the UKLPG were involved throughout the conception of the work, study design, survey approval and distribution,

Table 1 Survey participant characteristics by case reporting numbers

No of cases reported per year	<20	20–49	50–199	200–500	>500
Total participant responses n (%)	2 (5)	10 (24)	15 (36)	10 (24)	5 (12)
Transplant centre n (%)	0 (0)	0 (0)	0 (0)	6 (14)	5 (12)
Tertiary non-transplant centre n (%)	0 (0)	4 (10)	13 (31)	3 (7)	0 (0)
Secondary care/District General Hospital (DGH) n (%)	2 (5)	6 (14)	2 (5)	1 (2)	0 (0)
Total response rate across all MCQs (%)	62	100	100	100	99

MCQs, multiple choice questions.

analysis and interpretation of the data. Results of the survey were presented at the UKLPG annual meeting in December 2021, providing opportunity for further discussion, involvement and questions from the entire membership.

RESULTS

Survey participants

A total of 42 of 116 consultant members of the UKLPG completed individual responses to the email survey, representing 36% of the full membership. Respondents were from a mixture of healthcare settings including transplant centres 26% (11/42), tertiary non-transplant centres 48% (20/42) and secondary care / district general hospitals 26% (11/42). **Table 1** demonstrates participant characteristics by cases reported. The participant reporting rates of liver specimens varied from <20 per year (5%) to >500 per year (12%). All pathologists at transplant centres declared reporting at least two hundred liver cases per year, those at tertiary non-transplant centres reported between 20 and 500 cases and pathologists in secondary care/ district general hospitals (DGHs) reported between 1 and 500 cases. Response rates for all possible MCQ responses are shown by number of cases reported per year. Completion rate was 99% or more for all those who report 20 or more liver cases per year.

Prior experience of digital pathology and AI

Respondents were asked about their experience to date of using of digital pathology and AI. This included use in primary diagnosis, second opinion, research, teaching and EQA. Respondents could select multiple options and results are shown in **figure 1**. Ninety per cent (38/42) of participants had used digital pathology for EQA, followed by 64% (27/42) for teaching and training. Use for primary diagnosis and second opinion was reported in 26% (11/42) and 21% (9/42) of cases, respectively, and only one participant (2%, 1/42) had no experience of using digital pathology. Other examples in the comments included acquiring a scanner to trial the technology to assess where it may be useful. Of those reporting 200 or more liver cases per year, all had prior experience of digital pathology.

As AI is not yet available for routine clinical use by UK histopathologists (apart from a few exceptions), questions on experience of AI focused on a research setting. **Figure 2** outlines participants descriptions of their levels of experience with AI research. The majority of participants (57%) had no prior experience but had a general interest in AI development for histopathology. Twelve per cent of respondents had no prior experience, knowledge or interest in AI research.

Digital pathology

Twelve uses and considerations for digital pathology were explored in this section of the survey, with responses shown in

Participants experience of digital pathology




Figure 1 Responses for participants experiences of digital pathology for a range of purposes. EQA, external quality assurance.

figure 3. For primary diagnosis of liver specimens, 83% (34/41) of respondents either agreed or strongly agreed that digital pathology could be useful. Ninety per cent (37/41) of respondents either agreed or strongly agreed that digital pathology could be useful for obtaining second opinions. Digital pathology was also identified as useful in both research and training/education, where 85% (35/41) and 95% (39/41) of respondents, respectively, either agreed or strongly agreed with its utility in these areas. There was less certainty when asked if digital pathology would improve accuracy of diagnosis, with the largest group of respondents at 37% (15/41) reporting to be undecided. However, flexible working was identified as a benefit of digital pathology with 90% of participants responding with 'agree' or 'strongly agree' when asked about this.

Artificial intelligence

Twenty-four topics were examined across four MCQs, exploring uses and potential of AI in pathology. Responses to many of these questions are shown across **figures 4–6**. It is not possible to include all questions within figures in the main paper. Therefore, questions with most agreement in the responses given are highlighted with the full responses are included in online supplemental document 2. For general concepts where AI may improve practice (**figure 4**), 90% (37/41) of respondents agreed or strongly agreed that AI could improve the range of tools available to the histopathologist in their clinical practice. Improving

the consistency of diagnosis and the potential for understanding tissue features not currently recognised by pathologists, 63% (26/41) and 65% (26/40) of participants either agreed or strongly agreed with these concepts. Again, there was less certainty around the ability of AI to improve the speed or accuracy of diagnosis with 27% (11/41) and 34% (14/41) undecided.

When exploring the utility of AI by liver disease groups (**figure 5**), 80% (33/41) of participants identified fatty liver diseases as an area where AI could be useful for their clinical practice. Fifty-nine per cent (24/41) of participants either agreed or strongly agreed that AI would be useful for diagnosis and assessment of neoplastic liver diseases. There was less certainty for inflammatory liver diseases and biliary diseases, but still 44% (18/41) and 41% (17/41), respectively, of participants agreed or strongly agreed that AI would be useful in these contexts. Forty-three per cent (17/40) of participants responded 'don't know' when asked about utility in transplantation.

Respondents were asked about 14 specific tasks where AI could potentially be applied in clinical practice. The list of tasks was derived from reviewing the literature and from the consultations prior to the design of the survey. The tasks where there was most recognised utility are shown in **figure 6**. Ninety-three per cent (38/41) and 88% (35/40) of participants agreed or strongly agreed that quantifying steatosis and quantifying collagen would be useful in their clinical practice, respectively. Other quantitative tasks (eg, counting bile duct or portal tract numbers),

Participants experiences of AI




Figure 2 Responses for participants descriptions of their AI experience, knowledge and interest. AI, artificial intelligence.




Figure 3 Responses for concepts around digital pathology.

identifying copper associated protein and identifying ballooning or Mallory bodies were also of interest to the liver pathologists with 78% (32/41), 78% (32/41) and 60% (24/40), respectively, either agreeing or strongly agreeing that AI for these tasks would be useful. There was less certainty with tasks such as predicting patient outcomes in medical liver diseases and neoplastic diseases with responses in both cases at 41% (29/41) indicating that they were undecided. Although, identifying lymphovascular invasion was highlighted as another potentially useful task with 59% (24/41) of participants agreeing or strongly agreeing with this concept.

Concerns

Eight areas of potential concern around the impact of AI on liver pathologists were explored. In this question, participants were most concerned about AI being developed for pathology laboratories without the involvement of a histopathologist, with 73% (30/41) responding ‘agree’ or ‘strongly agree’ when asked about this. This was followed by 63% (26/41) of respondents agreeing or strongly agreeing that AI may struggle with existing digital systems. However, when asked whether AI is likely to replace pathologists in the future, the majority were not concerned




Figure 4 Responses for areas where AI may improve practice generally. AI, artificial intelligence.




Figure 5 Responses for AI utility by disease group. AI, artificial intelligence.

about this with 63% (26/41) responding 'disagree' or 'strongly disagree'.

Comments

At the end of the survey, two free-text questions asked participants to first highlight priority areas of digital pathology or AI research and second to make any other comments. Common themes in research priorities included quantitative tasks, fatty liver diseases, neoplastic liver diseases, improving variation between pathologists and practical benefits of digital pathology (eg, accessing archive cases, sharing images at meetings, collaboration and seeking second opinions). Comments were in keeping with the findings of the quantitative questions. Example quotes from the first free text question included: 'help with grading of assessment', 'estimation of the amount of fatty change, fibrosis and inflammation', 'ALD/NAFLD (alcohol-related liver disease and non-alcoholic liver disease) when biopsies done', 'grading and typing of HCC' (hepatocellular carcinoma), 'reduction of inter/intraobserver variation', 'marking, measuring, photographing and use for teaching would all be improved', 'cross site collaboration' and 'national network so that digital slides can be viewed from anywhere'.

A range of points were raised in the final open question, with many expanding comments on the earlier questions and

discussing benefits and concerns around the technologies. Examples of comments from this question were: 'This is not only coming but has arrived. Pathologists have largely been resistant, indifferent or ignorant. We need to get ahead of the curve.', 'Should be done on a global basis and not piecemeal roll out', 'Requires pathologist input for development', 'The hype that advocates perpetuate in AI does everyone a disservice.', 'Digital pathology may not fulfil its potential for substantive pathologists, training and improving patient care' and 'If AI is widely adopted and proves to be useful, we need to rethink the specialty, training etc and be realistic about future'.

DISCUSSION

Key findings

Recent developments in digital pathology and AI have impacted many areas of pathology, including liver disease.^{10 19} A survey published in 2018 examined the access and usage of digital pathology across UK pathology institutions, and this showed that there was great interest in digital pathology at that time.²⁰ This survey also demonstrated that 41% of institutions were already using image analysis on digital slides in some capacity and there have been many developments in this field since the study was performed. While the 2018 survey was conducted at a




Figure 6 Responses for task specific uses for AI. AI, artificial intelligence.

more general and departmental level, and so direct comparison is difficult, it is notable that 60% of departments at this time had access to a digital pathology scanner but 58.8% of these reported that they did not produce any digital slides. Those who reported producing digital slides varied from 50 to 300 000 slides produced per year. This study of liver pathologists showed that only 2% (1/42) of participants had no experience of digital pathology and perhaps reflects wider general accessibility of digital slides compared with this earlier study. As well as uses of digital pathology for tasks such as primary diagnosis, second opinions and education; the availability of this technology facilitates the application of image analysis and AI.^{10 21 22} Image analysis techniques have been established in liver pathology research over several decades^{23–26}; however, more recent AI techniques such as deep learning have generated high levels of interest and hold promise for future clinical applications.^{13 19 27}

This study demonstrates the appeal and potential of digital pathology and AI to the clinical practice of liver pathologists. The majority of respondents had experience of using digital pathology and most were interested in the potential applications of AI. The usefulness of digital pathology for primary diagnosis, second opinions, research and teaching and training were all highlighted. As deployment of digital pathology is currently ongoing across the National Health Service (NHS), these findings emphasise current demand and recognition of utility within liver disease. Liver pathologists saw promise in AI expanding the range of diagnostic tools available to them and improving the consistency of diagnosis. However, there was uncertainty about its potential to improve the speed and accuracy of diagnosis. Although, it must be acknowledged that AI is not currently in use in the vast majority of laboratories, and so uncertainty around how it will function is not surprising. This was a brief snapshot survey of opinions around digital pathology and AI, intended to maximise participation from UK liver pathologists. It was therefore not possible to explore more extensive details such as precise numbers or proportions of digital cases reported, the digital pathology platforms used, the data storage solutions in place and the areas of AI research that some participants were working in, but these areas could be examined in future work. The comments illustrated some additional information about these topics, with one participant reporting that digital pathology was being ‘... implemented for primary rapid/urgent diagnosis and for second opinion’. Another reported trialling the use of digital pathology in principle. Several participants indicated issues with existing NHS information technology (IT) infrastructure as being potential barriers to using digital pathology for more cases. However, there were requests for digital pathology to be made available for use in a range of contexts included within the comments. There was also awareness of a variety of potential clinical applications of AI from many participants.

Fatty liver diseases and neoplastic liver diseases were determined as areas of demand for the application of AI, and many comments from participants discussed specific applications of AI for both of these disease groups. Transplantation was an area of uncertainty in terms of AI utility, however, this likely reflects that the majority of respondents (74%) do not practice routinely at transplant centres.

The task-specific questions and participant comments showed a focus on quantitative tools with the aim of removing some of these time consuming tasks from their practice. Identifying objects that may be difficult or laborious for a liver pathologist to find such as ballooning, Mallory bodies, copper associated protein and vascular invasion were highlighted as well. Participants were undecided regarding the potential of AI to predict

patient outcomes. This may reflect the increased complexity of this task when compared with simple quantitation or the existence of alternative tools in place to do this.

Key concerns to acknowledge were the potential lack of involvement of pathologists in AI development and the implementation of AI tools within existing digital systems and infrastructure within laboratories. These are concerns that will need to be addressed to achieve future successful implementation of AI products. Finally, it is of note that the majority of liver pathologists are not concerned (63%) that AI could replace them in the future.

Study limitations

While the response rate (36%) was good for an unsolicited email survey, there is a possible non-response bias when accounting for members who did not participate. To reflect the varying knowledge levels of the participants, not all questions were mandatory and so responses ranged from 40 to 42 responses per question from the 42 participants, and a non-response bias is possible here. It should be acknowledged that current variation in levels of experience and exposure to digital pathology and AI may have influenced responses both positively or negatively. This survey provides a useful snapshot of opinion at one point in time, and this may change with rapid advancement of these technologies. A formal Delphi exercise could be explored in the future to follow this initial piece of work and to expand on the themes identified here.

CONCLUSION

Given the current pace of change on a national scale in the UK, with increasing availability of digital pathology solutions and a growing body of AI research, this survey aimed to gather current attitudes and priorities around both of these technologies. Demand for digital pathology access and numerous applications of AI for tasks to aid the liver pathologist in clinical practice were identified. Findings highlighted within this study could be used to help inform the direction of future research within this field, to outline current areas of need and also potential concerns when implementing these tools.

Author affiliations

- ¹Pathology and Data Analytics, University of Leeds, Leeds, UK
- ²Department of Histopathology, Leeds Teaching Hospitals NHS Trust, Leeds, UK
- ³Faculty of Health Sciences, University of Bradford, Bradford, UK
- ⁴Wolfson Centre for Applied Health Research, Bradford, UK
- ⁵Division of Pathology, University of Edinburgh, Edinburgh, UK
- ⁶Translational and Clinical Research Institute, Newcastle University, Newcastle upon Tyne, UK
- ⁷Division of Digestive Diseases, Imperial College London, London, UK
- ⁸Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, UK
- ⁹Department of Cellular Pathology, Queen Elizabeth Hospital Birmingham, Birmingham, UK
- ¹⁰Department of Cellular Pathology, Royal Free Hospital, London, UK
- ¹¹Translational and Clinical Research Institute, Newcastle University, Newcastle, UK
- ¹²Department of Pathology, National and Kapodistrian University of Athens, Athens, Greece
- ¹³Department of Clinical Pathology and Department of Clinical and Experimental Medicine, Linköping University, Linköping, Sweden
- ¹⁴Centre for Medical Image Science and Visualization (CMIV), Linköping University, Linköping, Sweden

Handling editor Yoh Zen.

Contributors All authors contributed to the design of this study. CM gathered the survey responses with assistance from JW. CM analysed the data and wrote the initial draft of the manuscript. All authors reviewed the manuscript and contributed to editing and revising it. All authors have approved the manuscript for publication.

CM is the guarantor author, accepts full responsibility for the work and the conduct of the study, had access to the data, and controlled the decision to publish.

Funding The authors would like to thank the UKLPG membership for their kind participation. DTr and AC are funded by National Pathology Imaging Co-operative (NPIC). NPIC (project no. 104687) is supported by a £50m investment from the Data to Early Diagnosis and Precision Medicine strand of the Government's Industrial Strategy Challenge Fund, managed and delivered by UK Research and Innovation (UKRI). CM is funded by Leeds Hospitals Charity, the National Institute for Health Research (NIHR) and NPIC. AQ is supported by the National Institute For Health Research (NIHR) UCLH/UCL Biomedical Research Centre (BRC).

Disclaimer The funders had no role in the survey's design, implementation and analysis.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This was an anonymous online survey of NHS pathologists who report liver pathology specimens. The survey was voluntary and completion was not associated with any reward or disadvantage. No patient data was used in this study and it was performed in accordance with the principles of the Declaration of Helsinki.

Provenance and peer review Not commissioned; internally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Clare McGenity <http://orcid.org/0000-0002-0224-2340>

Desley A H Neil <http://orcid.org/0000-0001-9800-6811>

REFERENCES

- Asrani SK, Devarbhavi H, Eaton J, *et al*. Burden of liver diseases in the world. *J Hepatol* 2019;70:151–71.
- Pimpin L, Cortez-Pinto H, Negro F, *et al*. Burden of liver disease in Europe: epidemiology and analysis of risk factors to identify prevention policies. *J Hepatol* 2018;69:718–35.
- Public Health England. Liver disease profiles: short statistical commentary, 2018. Available: <https://www.gov.uk/government/statistics/liver-disease-profiles-november-2018-update/liver-disease-profiles-short-statistical-commentary-november-2018> [Accessed Aug 2022].
- Bedossa P. Liver biopsy. *Gastroenterol Clin Biol* 2008;32:4–7.
- Campbell MS, Reddy K. The evolving role of liver biopsy. *Aliment Pharmacol Ther* 2004;20:249–59.
- The Royal College of pathologists. *Meeting Pathology Demand. Histopathology Workforce Census* 2018 <https://www.rcpath.org/uploads/assets/952a934d-2ec3-48c9-a8e6e00fcdca700f/Meeting-Pathology-Demand-Histopathology-Workforce-Census-2018.pdf>
- Wyatt J, Hübscher S, Goldin R. *Dataset for histopathology reporting of liver resection specimens (including gall bladder) and liver biopsies for primary and metastatic carcinoma*. London, UK: The Royal College of Pathologists, 2012. <https://www.rcpath.org/uploads/assets/cbae24ce-220c-4d9f-95c5b83dd1f0ad2b/Dataset-for-histopathology-reporting-of-liver-resection-specimens-and-liver-biopsies-for-primary-and-metastatic-carcinoma-2nd-edition.pdf>
- Cross S, Furness P, Igalii L. *Best practice recommendations for implementing digital pathology*. London: The Royal College of pathologists, 2018. <https://www.rcpath.org/uploads/assets/f465d1b3-797b-4297-b7fedc00b4d77e51/Best-practice-recommendations-for-implementing-digital-pathology.pdf>
- Department of Health and Social Care. Press release: funding boost for artificial intelligence in NHS to speed up diagnosis of deadly diseases, 2020. Available: <https://www.gov.uk/government/news/funding-boost-for-artificial-intelligence-in-nhs-to-speed-up-diagnosis-of-deadly-diseases#:~:text=A%20%20C%20A350%20million%20funding,improve%20the%20diagnosis%20of%20disease> [Accessed August 2022].
- Niazi MKK, Parwani AV, Gurcan MN. Digital pathology and artificial intelligence. *Lancet Oncol* 2019;20:e253–61.
- Madabhushi A, Lee G. Image analysis and machine learning in digital pathology: challenges and opportunities. *Med Image Anal* 2016;33:170–5.
- Calderaro J, Kather JN. Artificial intelligence-based pathology for gastrointestinal and hepatobiliary cancers. *Gut* 2021;70:1183–93.
- Heinemann F, Birk G, Stierstorfer B. Deep learning enables pathologist-like scoring of NASH models. *Sci Rep* 2019;9:1–10.
- Forlano R, Mullish BH, Giannakeas N, *et al*. High-Throughput, machine Learning-Based quantification of steatosis, inflammation, ballooning, and fibrosis in biopsies from patients with nonalcoholic fatty liver disease. *Clin Gastroenterol Hepatol* 2020;18:e9:2081–90.
- Saillard C, Schmauch B, Laifa O, *et al*. Predicting survival after hepatocellular carcinoma resection using deep learning on histological slides. *Hepatology* 2020;72:2000–13.
- Brunt EM, Clouston AD, Goodman Z, *et al*. Complexity of ballooned hepatocyte feature recognition: defining a training atlas for artificial intelligence-based imaging in NAFLD. *J Hepatol* 2022;76:1030–41.
- Liu F, Goh GB-B, Tiniakos D, *et al*. qFIBS: an automated technique for quantitative evaluation of fibrosis, inflammation, ballooning, and steatosis in patients with nonalcoholic steatohepatitis. *Hepatology* 2020;71:1953–66.
- UKLPG. UKLPG - background and UKLPG committee meetings, 2017. Available: https://www.virtualpathology.leeds.ac.uk/eqa/specialist/liver/liver_group.php [Accessed Aug 2022].
- Nam D, Chapiro J, Paradis V, *et al*. Artificial intelligence in liver diseases: improving diagnostics, prognostics and response prediction. *JHEP Rep* 2022;4:100443.
- Williams BJ, Lee J, Oien KA, *et al*. Digital pathology access and usage in the UK: results from a national survey on behalf of the National cancer research Institute's CM-Path initiative. *J Clin Pathol* 2018;71:463–6.
- Paradis V, Quaglia A, pathology D. What is the future? *J Hepatol* 2019;70:1016–8.
- Williams BJ, Treanor D. Practical guide to training and validation for primary diagnosis with digital pathology. *J Clin Pathol* 2020;73:418–22.
- Tsiplakidou M, Tsipouras M, Giannakeas N, *et al*. Automated detection of liver histopathological findings based on biopsy image processing. *Information* 2017;8:36.
- Calvaruso V, Burroughs AK, Standish R, *et al*. Computer-Assisted image analysis of liver collagen: relationship to Ishak scoring and hepatic venous pressure gradient. *Hepatology* 2009;49:1236–44.
- Marsman H, Matsushita T, Dierkhising R, *et al*. Assessment of donor liver steatosis: pathologist or automated software? *Hum Pathol* 2004;35:430–5.
- Nativ NI, Chen AI, Yarmush G, *et al*. Automated image analysis method for detecting and quantifying macrovesicular steatosis in hematoxylin and eosin-stained histology images of human livers. *Liver Transpl* 2014;20:228–36.
- Arjmand A, Angelis CT, Christou V, *et al*. Training of deep convolutional neural networks to identify critical liver alterations in histopathology image samples. *Appl Sci* 2019;10:42.



Survey for members of UKLPG: Digital Pathology and AI in liver disease

Introduction

Thank you for participating in this survey. Your responses will build a picture of the UK liver pathology community's opinions on digital pathology and artificial intelligence (AI) and we anticipate the results will be published as a report. This report could influence future decisions about research priorities in this area. This survey aims to build on discussions from a focus group of interested pathologists working in liver research. Your individual answers will remain confidential throughout this process and will not be used for purposes other than those outlined above.

This first page contains some general introductory questions that we would like all participants to complete. The following pages are optional but we would be grateful for your insight. If you do not feel able to answer certain questions, please select "don't know" or you can skip to the next question.

* 1. Describe the hepatology services in your current place of work

- Transplant centre
- Tertiary non-transplant centre
- Secondary care / DGH
- Other (please specify):

* 2. Approximately how many liver specimens would you typically report per year

- <20
- 20-49
- 50-199
- 200-500
- >500

* 3. What is your experience of using digital pathology? (Please select all those that apply)

- Primary diagnosis
- Second opinion
- Research
- Teaching / training
- EQA
- No experience of digital pathology
- Other (please specify)

* 4. Do you have any experience, knowledge or interest in AI development? (Please select the best description)

- Yes, direct experience of AI research
- Yes, knowledge of AI research but no direct experience
- Generally interested but no prior knowledge or experience of AI research
- No prior experience, knowledge or interest in AI research



Survey for members of UKLPG: Digital Pathology and AI in liver disease

Digital Pathology

Digital pathology usually refers to images of histology generated by whole slide imaging and available to review by a pathologist on a computer.

Access and experience may vary across the group, but we would like to understand what considerations are important to liver pathologists in this area. With your current knowledge and future expectations, please rate how the questions relate to you.

Questions are rated from strongly disagree to strongly agree, with the option to say "don't know" or you may skip the question. Comments and suggestions may be added at the bottom of each question.

5. Digital pathology could be useful for:

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Primary diagnosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Second opinions	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Transplant frozen sections	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Training and education	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Flexible working	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Research	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Establishing national and international networks	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Creating large, multicentre image datasets	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Additional comments / suggestions

6. Digital pathology:

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Could improve speed of diagnosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Could improve accuracy of diagnosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Should have pathologists involved in its deployment for clinical use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Should be easy to use	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Additional comments / suggestions



Survey for members of UKLPG: Digital Pathology and AI in liver disease

Artificial intelligence (general concepts)


Artificial intelligence usually refers to machines demonstrating intelligent behaviour by evaluating their environment and taking resulting actions. In the context of histopathology, this is often referring to machines extracting information from an image.

Access and experience may vary across the group but we would like to understand what considerations in general are important to liver pathologists in this area. With your current knowledge and future expectations, please rate how the questions relate to you.

7. Artificial intelligence could improve:

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Speed of diagnosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Accuracy of diagnosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Consistency of diagnosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
The range of tools available to a pathologist	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Understanding of tissue features not currently recognised by pathologists	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Additional comments / suggestions



Survey for members of UKLPG: Digital Pathology and AI in liver disease


Artificial intelligence and liver diseases

We would like to establish where AI might be useful to assist with diagnostic work. Please rate how useful you think AI tools would be in assessment and diagnosis these diseases.

8. Artificial intelligence could be useful for the pathological assessment and diagnosis of

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Fatty liver diseases	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Inflammatory liver diseases	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Biliary diseases	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Neoplastic liver diseases	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Transplantation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Additional comments / suggestions



Survey for members of UKLPG: Digital Pathology and AI in liver disease

Artificial intelligence tools

We have suggested some potential tools that might be performed by a computer with an AI algorithm to assist with your diagnostic work. Please rate whether these would be useful.

Possible concerns are included at the bottom, please rate whether or not these are of concern to you.

9. Artificial intelligence tools performing the following for diagnostic work would be useful in **medical liver diseases**

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Quantifying steatosis	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Quantifying collagen	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Other quantitative tasks e.g. bile duct or portal tract numbers	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Identifying inflammation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Identifying ballooning or Mallory bodies	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Identifying copper associated protein	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Predicting patient outcomes	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Additional comments / suggestions

10. Artificial intelligence tools performing the following for diagnostic work would be useful in **neoplastic liver disease**


	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Quantify tumour grade	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Identifying lymphovascular invasion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Identifying perineural invasion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Classifying nodule type e.g. regenerative, dysplastic, HCC	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Providing morphomolecular classification	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Predicting patient outcomes	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Predicting genetics	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Additional comments / suggestions

11. I am concerned that **artificial Intelligence** may:

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Be developed without pathologist involvement	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Replace pathologists	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Make decisions that we cannot understand	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Create additional work for the pathologist	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Impair training	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
De-skill pathologists	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Be unsafe for patients	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Struggle with existing digital systems	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Additional comments / suggestions



Survey for members of UKLPG: Digital Pathology and AI in liver disease

Priorities and final comments

Please outline any concept(s) that you feel would be most important or useful to your practice in this area of research. Please feel free to add any final comments or suggestions.

Thank you for completing this survey.

12. If you were to highlight a priority / priorities for research and development in this area that would be most useful for your practice, what would you choose?

13. Please give any further comments or additional suggestions

Supplementary document 2 – Summary of responses

Responses tables are given as number of responses and percentage of responses for each question.

Q1 Describe the hepatology services in your current place of work

Transplant centre	Tertiary non-transplant centre	Secondary care / DGH	Other	Total skipped
11	20	11	0	0

Transplant centre	Tertiary non-transplant centre	Secondary care / DGH	Other
26%	48%	26%	0%

Q2 Approximately how many liver specimens would you typically report per year

<20	20-49	50-199	200-500	>500	Total skipped
2	10	15	10	5	0

<20	20-49	50-199	200-500	>500
5%	24%	36%	24%	12%

Q3 What is your experience of using digital pathology

Primary diagnosis	Second opinion	Research	Teaching / training	EQA	No experience of digital pathology	Other	Total skipped
11	9	16	27	38	1	2	0

Primary diagnosis	Second opinion	Research	Teaching / training	EQA	No experience of digital pathology	Other
26%	21%	38%	64%	90%	2%	5%

Q4 Do you have any experience, knowledge or interest in AI development?

Yes, direct experience of AI research	Yes, knowledge of AI research but no direct experience	Generally interested but no prior knowledge or experience of AI research	No prior experience, knowledge or interest in AI research
6	7	24	5

Yes, direct experience of AI research	Yes, knowledge of AI research but no direct experience	Generally interested but no prior knowledge or experience of AI research	No prior experience, knowledge or interest in AI research
14%	17%	57%	12%

Q5 Digital pathology could be useful for

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know	Total skipped
Primary diagnosis	2	0	5	16	18	0	1
Second opinions	2	0	1	13	24	1	1
Transplant frozen sections	2	2	7	3	8	19	1
Training and education	2	0	0	6	33	0	1
Flexible working	2	0	1	13	24	1	1
Research	2	0	1	13	22	3	1
Establishing national and international networks	2	0	0	14	25	0	1
Creating large, multicentre image datasets	2	0	0	11	28	0	1

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Primary diagnosis	5%	0%	12%	39%	44%	0%
Second opinions	5%	0%	2%	32%	59%	2%
Transplant frozen sections	5%	5%	17%	7%	20%	46%
Training and education	5%	0%	0%	15%	80%	0%
Flexible working	5%	0%	2%	32%	59%	2%
Research	5%	0%	2%	32%	54%	7%
Establishing national and international networks	5%	0%	0%	34%	61%	0%
Creating large, multicentre image datasets	5%	0%	0%	27%	68%	0%

Q6 Digital pathology:

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know	Total skipped
Could improve speed of diagnosis	1	5	14	12	9	0	1
Could improve accuracy of diagnosis	1	13	15	8	3	1	1
Should have pathologists involved in its deployment for clinical use	0	1	0	14	24	1	2
Should be easy to use	0	0	5	13	22	1	1

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Could improve speed of diagnosis	2%	12%	34%	29%	22%	0%
Could improve accuracy of diagnosis	2%	32%	37%	20%	7%	2%
Should have pathologists involved in its deployment for clinical use	0%	3%	0%	35%	60%	3%
Should be easy to use	0%	0%	12%	32%	54%	2%

Q7 Artificial intelligence could improve:

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know	Total skipped
Speed of diagnosis	1	2	11	11	8	8	1
Accuracy	1	2	14	12	7	5	1
Consistency of diagnosis	1	0	10	17	9	4	1
The range of tools available to a pathologist	1	1	1	23	14	1	1
Understanding of tissue features not currently recognised by pathologists	1	2	8	15	11	3	2

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Speed of diagnosis	2%	5%	27%	27%	20%	20%
Accuracy	2%	5%	34%	29%	17%	12%
Consistency of diagnosis	2%	0%	24%	41%	22%	10%
Range of tools available to a pathologist	2%	2%	2%	56%	34%	2%
Understanding of tissue features not currently recognised by pathologists	3%	5%	20%	38%	28%	8%

Q8 Artificial intelligence could be useful for the pathological assessment and diagnosis of

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know	Total skipped
Fatty liver diseases	1	0	4	25	8	3	1
Inflammatory liver diseases	1	3	13	13	5	6	1
Biliary diseases	1	3	14	11	6	6	1
Neoplastic liver diseases	1	3	9	18	6	4	1
Transplantation	1	3	9	7	3	17	2

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Fatty liver diseases	2%	0%	10%	61%	20%	7%
Inflammatory liver diseases	2%	7%	32%	32%	12%	15%
Biliary diseases	2%	7%	34%	27%	15%	15%
Neoplastic liver diseases	2%	7%	22%	44%	15%	10%
Transplantation	3%	8%	23%	18%	8%	43%

Q9 Artificial intelligence tools performing the following for diagnostic work would be useful in medical liver diseases

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know	Total skipped
Quantifying steatosis	0	0	2	22	16	1	1
Quantifying collagen	0	0	4	22	13	1	2
Other quantitative tasks e.g. bile duct or portal tract numbers	0	1	5	22	10	3	1
Identifying inflammation	0	3	11	16	4	7	1
Identifying ballooning or Mallory bodies	0	2	8	17	7	6	2
Identifying copper associated protein	0	1	5	23	9	3	1
Predicting patient outcomes	0	0	17	12	2	10	1

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Quantifying steatosis	0%	0%	5%	54%	39%	2%
Quantifying collagen	0%	0%	10%	55%	33%	3%
Other quantitative tasks	0%	2%	12%	54%	24%	7%
Identifying inflammation	0%	7%	27%	39%	10%	17%
Identifying ballooning or Mallory bodies	0%	5%	20%	43%	18%	15%
Identifying copper associated protein	0%	2%	12%	56%	22%	7%
Predicting patient outcomes	0%	0%	41%	29%	5%	24%

Q10 Artificial intelligence tools performing the following for diagnostic work would be useful in neoplastic liver disease

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know	Total skipped
Quantifying tumour grade	0	3	10	16	5	7	1
Identifying lymphovascular invasion	0	2	8	18	6	7	1
Identifying perineural invasion	0	1	11	15	8	6	1
Classifying nodule type e.g. regenerative, dysplastic, HCC	0	3	15	10	4	8	2
Providing morphomolecular classification	0	2	12	17	5	5	1
Predicting patient outcomes	0	0	17	12	3	9	1
Predicting genetics	0	0	16	12	2	10	2

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Quantifying tumour grade	0%	7%	24%	39%	12%	17%
Identifying lymphovascular invasion	0%	5%	20%	44%	15%	17%
Identifying perineural invasion	0%	2%	27%	37%	20%	15%
Classifying nodule type e.g. regenerative, dysplastic, HCC	0%	8%	38%	25%	10%	20%
Providing morphomolecular classification	0%	5%	29%	41%	12%	12%
Predicting patient outcomes	0%	0%	41%	29%	7%	22%
Predicting genetics	0%	0%	40%	30%	5%	25%

Q11 I am concerned that artificial intelligence may:

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know	Total skipped
Be developed without pathologist involvement	0	9	1	20	10	1	1
Replace pathologists	8	18	5	6	2	2	1
Make decisions that we cannot understand	0	8	11	11	5	6	1
Create additional work for the pathologist	0	9	9	16	4	3	1
Impair training	2	12	11	9	4	3	1
De-skill pathologists	3	13	11	11	3	0	1
Be unsafe for patients	2	13	12	9	2	3	1
Struggle with existing digital systems	1	3	6	18	8	5	1

	Strongly disagree	Disagree	Undecided	Agree	Strongly agree	Don't know
Be developed without pathologist involvement	0%	22%	2%	49%	24%	2%
Replace pathologists	20%	44%	12%	15%	5%	5%
Make decisions that we cannot understand	0%	20%	27%	27%	12%	15%
Create additional work for the pathologist	0%	22%	22%	39%	10%	7%
Impair training	5%	29%	27%	22%	10%	7%
De-skill pathologists	7%	32%	27%	27%	7%	0%
Be unsafe for patients	5%	32%	29%	22%	5%	7%
Struggle with existing digital systems	2%	7%	15%	44%	20%	12%