

Short Report

Consumption of nuts and seeds and pancreatic ductal adenocarcinoma risk in the European Prospective Investigation into Cancer and Nutrition

Mireia Obón-Santacana^{1,2,3}, Leila Luján-Barroso^{2,4,5}, Heinz Freisling⁶, Sabine Naudin⁶, Marie-Christine Boutron-Ruault^{7,8}, Francesca Romana Mancini^{7,8}, Vinciane Rebours^{9,10}, Tilman Kühn¹¹, Verena Katzke¹¹, Heiner Boeing¹², Anne Tjønneland^{13,14}, Anja Olsen¹³, Kim Overvad¹⁵, Cristina Lasheras¹⁶, Miguel Rodríguez-Barranco^{3,17}, Pilar Amiano^{3,18}, Carmen Santiuste^{3,19}, Eva Ardanaz^{3,20,21}, Kay-Thee Khaw²², Nicholas J. Wareham²³, Julie A. Schmidt²⁴, Dagfinn Aune^{25,26,27}, Antonia Trichopoulou²⁸, Paschalis Thriskos²⁸, Eleni Peppas²⁸, Giovanna Masala²⁹, Sara Grioni³⁰, Rosario Tumino³¹, Salvatore Panico³², Bas Bueno-de-Mesquita^{33,34,35,36}, Veronica Sciannameo³⁷, Roel Vermeulen³⁸, Emily Sonestedt³⁹, Malin Sund⁴⁰, Elisabete Weiderpass^{41,42}, Guri Skeie⁴³, Carlos A. González^{2,4}, Elio Riboli²⁵ and Eric J. Duell^{2,4}

¹Oncology Data Analytics Program (ODAP), Catalan Institute of Oncology (ICO), L'Hospitalet del Llobregat, Barcelona, Spain

²Bellvitge Biomedical Research Institute (IDIBELL), L'Hospitalet de Llobregat, Barcelona, Spain

³Consortium for Biomedical Research in Epidemiology and Public Health (CIBERESP), Madrid, Spain

⁴Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Catalan Institute of Oncology (ICO), L'Hospitalet del Llobregat, Barcelona, Spain

⁵Department of Nursing of Public Health, Mental Health and Maternity and Child Health School of Nursing, Universitat de Barcelona, Barcelona, Spain

⁶Nutritional Methodology and Biostatistics Group, International Agency for Research on Cancer, Lyon, France

⁷CESP, Fac. de médecine - Univ. Paris-Sud, Fac. de médecine - UVSQ, INSERM, Université Paris-Saclay, Villejuif, France

⁸Gustave Roussy, Villejuif, France

⁹Pancreatology Department, Beaujon Hospital, DHU Unity, AP-HP, Clichy, and Paris-Diderot University, Paris, France

¹⁰Inserm UMR1149, DHU Unity, and Paris-Diderot University, Paris, France

¹¹Division of Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg, Germany

¹²Department of Epidemiology, German Institute of Human Nutrition (Dife) Postdam-Rehbrücke, Nuthetal, Germany

¹³Danish Cancer Society Research Center, Copenhagen, Denmark

¹⁴Department of Public Health, University of Copenhagen, Copenhagen, Denmark

¹⁵Section for Epidemiology, Department of Public Health, Aarhus University, Aarhus, Denmark

¹⁶Department of Functional Biology, Faculty of Medicine, University of Oviedo, Oviedo, Spain

Key words: pancreatic cancer, nuts, seeds, diet, intake, EPIC, prospective cohort study

Abbreviations: 95% CI: 95% confidence intervals; BMI: body mass index; EPIC: European Prospective Investigation into Cancer and Nutrition; HR: hazards ratio; LRT: likelihood ratio test; NHS: Nurses' Health Study; PC: pancreatic cancer; PDAC: pancreatic ductal adenocarcinoma

Grant sponsor: LK Research Funds, Dutch Prevention Funds, Dutch ZON (Zorg Onderzoek Nederland); **Grant sponsor:** Associazione Italiana per la Ricerca sul Cancro (AIRC) and National Research Council (Italy); **Grant sponsor:** Cancer Research UK;

Grant numbers: C570/A16491, C8221/A19170; **Grant sponsor:** Danish Cancer Society (Denmark); **Grant sponsor:** Deutsche Krebshilfe, Deutsches Krebsforschungszentrum (DKFZ) and Federal Ministry of Education and Research (Germany); **Grant sponsor:** Dutch Ministry of Public Health, Welfare and Sports (VWS); **Grant sponsor:** Fundación Científica Asociación Española Contra el Cáncer; **Grant sponsor:** Ligue contre le Cancer, Institut Gustave Roussy, Mutuelle Générale de l'Éducation Nationale, Institut National de la Santé et de la Recherche Médicale (INSERM); **Grant sponsor:** Medical Research Council; **Grant number:** MR/M012190/1; **Grant sponsor:** Netherlands Cancer Registry (NKR); **Grant sponsor:** Swedish Cancer Society, Swedish Scientific Council and Regional Government of Skåne and Västerbotten (Sweden); **Grant sponsor:** The European Commission (DG-SANCO); **Grant sponsor:** the Health Research Fund (FIS) of the Spanish Ministry of Health; **Grant sponsor:** the Hellenic Health Foundation (Greece); **Grant sponsor:** the International Agency for Research on Cancer; **Grant sponsor:** World Cancer Research Fund (WCRF) and Statistics Netherlands (The Netherlands); **Grant sponsor:** Navarra and the Catalan Institute of Oncology, La Caixa; **Grant number:** BM 06-130; **Grant sponsor:** Regional Governments of Andalucía, Asturias, Basque Country, Murcia; **Grant number:** 6236

DOI: 10.1002/ijc.32415

History: Received 17 Jan 2019; Accepted 16 Apr 2019; Online 20 May 2019

Correspondence to: Eric J. Duell, Unit of Nutrition and Cancer, Cancer Epidemiology Research Program, Catalan Institute of Oncology (ICO), Bellvitge Biomedical Research Institute (IDIBELL), Avda Gran Via 199-203, 08907 L'Hospitalet del Llobregat, Barcelona, Spain, Tel.: +34-93-260-7401; Fax: +34-93-260-7787, E-mail: eduell@iconcologia.net

- ¹⁷Escuela Andaluza de Salud Pública, Instituto de Investigación Biosanitaria IBS GRANADA, Universidad de Granada, Granada, Spain
- ¹⁸Public Health Division of Gipuzkoa, BioDonostia Research Institute, San Sebastian, Spain
- ¹⁹Department of Epidemiology, Murcia Regional Health Council, IMIB-Arixaca, Murcia, Spain
- ²⁰Navarra Public Health Institute, Pamplona, Spain
- ²¹IdiSNA, Navarra Institute for Health Research, Pamplona, Spain
- ²²Department of Public Health and Primary Care, University of Cambridge, Cambridge, United Kingdom
- ²³MRC Epidemiology Unit, School of Clinical Medicine, University of Cambridge, Cambridge, United Kingdom
- ²⁴Cancer Epidemiology Unit, Nuffield Department of Population Health, University of Oxford, Oxford, United Kingdom
- ²⁵Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, United Kingdom
- ²⁶Department of Nutrition, Bjørknes University College, Oslo, Norway
- ²⁷Department of Endocrinology, Morbid Obesity and Preventive Medicine, Oslo University Hospital, Oslo, Norway
- ²⁸Hellenic Health Foundation, Athens, Greece
- ²⁹Cancer Risk Factors and Life-Style Epidemiology Unit, Institute for Cancer Research, Prevention and Clinical Network (ISPRO), Florence, Italy
- ³⁰Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Milan, Italy
- ³¹Cancer Registry and Histopathology Department, "Civic - M. P. Arezzo" Hospital, ASP Ragusa, Ragusa, Italy
- ³²Department of Clinical and Experimental Medicine, Federico II University, Naples, Italy
- ³³Department for Determinants of Chronic Diseases (DCD), National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands
- ³⁴Department of Gastroenterology and Hepatology, University Medical Centre, Utrecht, The Netherlands
- ³⁵Department of Epidemiology and Biostatistics, The School of Public Health, Imperial College London, London, United Kingdom
- ³⁶Department of Social & Preventive Medicine, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia
- ³⁷Unit of Epidemiology, Regional Health Service ASL TO3, Turin, Italy
- ³⁸Institute for Risk Assessment Sciences, Division of Environmental Epidemiology, Utrecht University, Utrecht, The Netherlands
- ³⁹Nutritional Epidemiology, Department of Clinical Sciences Malmö, Lund University, Malmö, Sweden
- ⁴⁰Department of Surgical and Preoperative Sciences, Umeå University, Umeå, Sweden
- ⁴¹Department of Community Medicine, University of Tromsø, The Arctic University of Norway, Tromsø, Norway
- ⁴²International Agency for Research on Cancer, Lyon, France
- ⁴³Department of Community Medicine, Faculty of Health Sciences, UiT-The Arctic University of Norway, Tromsø, Norway

Four epidemiologic studies have assessed the association between nut intake and pancreatic cancer risk with contradictory results. The present study aims to investigate the relation between nut intake (including seeds) and pancreatic ductal adenocarcinoma (PDAC) risk in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. Cox proportional hazards models were used to estimate hazards ratio (HR) and 95% confidence intervals (95% CI) for nut intake and PDAC risk. Information on intake of nuts was obtained from the EPIC country-specific dietary questionnaires. After a mean follow-up of 14 years, 476,160 participants were eligible for the present study and included 1,283 PDAC cases. No association was observed between consumption of nuts and PDAC risk (highest intake vs nonconsumers: HR, 0.89; 95% CI, 0.72–1.10; *p*-trend = 0.70). Furthermore, no evidence for effect-measure modification was observed when different subgroups were analyzed. Overall, in EPIC, the highest intake of nuts was not statistically significantly associated with PDAC risk.

What's new

Environmental and lifestyle factors probably play a major role in the development—or prevention—of pancreatic cancer (PC). For example, might the inclusion of nuts in one's diet reduce the risk of PC? Results of previous studies have been contradictory. In this analysis of data from the large, prospective EPIC study, the authors found no significant protective role for nuts or seeds against PC.

Introduction

Pancreatic cancer (PC) is one of the most aggressive human cancers, and it is projected to be the second leading cause of cancer mortality by 2030.¹ The most frequent histological type is pancreatic ductal adenocarcinoma (PDAC), and accounts for almost 95% of all exocrine pancreatic tumors.² PC incidence is increasing, and 5-year survival is the worst (<8%) of all

common cancers since it is usually diagnosed at late stages, and few treatment improvements have been achieved in recent years.³ Thus, scientific evidence for primary prevention is crucial.⁴

Chronic pancreatitis and long-standing diabetes are associated with higher PC risk, while family history and genetic syndromes account for <10% of all PC cases, suggesting that

environmental and lifestyle factors play a major role in PC development.^{5–7} Tobacco smoking, heavy alcohol consumption and body fatness are considered lifestyle risk factors.⁶ Red and processed meat intakes have also been associated with PC risk, but scientific evidence remains unclear. Likewise, inconsistent results have been reported for *Helicobacter pylori*, physical activity, adherence to the Mediterranean diet and dietary intakes of fruits, vegetables, magnesium and folate.^{6–8}

Nuts (comprising tree nuts and peanuts) are a food group that has largely been associated with beneficial health effects including reduced total and cause-specific mortality, cardiovascular disease, hypertension, diabetes, insulin resistance and cancer risk.^{9–12} The characteristic nutritional composition of nuts (rich in fiber, vitamins, minerals, mono- and polyunsaturated fatty acids and bioactive compounds) makes them an ideal food group to be studied as a preventive factor for PC.^{10,13}

One prospective epidemiologic study from the United States found evidence for an inverse association between nut intake and PC risk in women,¹⁴ whereas one case-control and one prospective cohort study, both from the Netherlands, observed no statistically significant associations.^{15,16} A third prospective cohort study from Iran also found no clear association.¹⁷ The purpose of the present study was to investigate the relation between the consumption of nuts and seeds and PDAC risk accounting for dietary and lifestyle factors in one of the largest prospective cohort studies of nutrition and chronic diseases.

Methods

Study population

The European Prospective Investigation into Cancer and Nutrition (EPIC) is a multicenter study that started between 1992 and 1998 and comprises 23 research centers in ten European countries. The study was approved by the International Agency for Research on Cancer ethical review boards and/or all local ethics committees. The design and methodology of the EPIC study has been published elsewhere.¹⁸

Of the 521,324 participants, a total of 45,164 were excluded because they had prevalent cancer other than nonmelanoma skin cancer at recruitment ($n = 25,184$), had incomplete follow-up data ($n = 4,128$), had missing data of diagnosis ($n = 20$), had no lifestyle or dietary information at recruitment ($n = 6,259$) or had an extreme ratio of energy intake to energy requirement (top or bottom 1%; $n = 9,573$); resulting in 476,160 participants (70% women) for the present analysis.

Identification of PC cases

PC incidence was ascertained through population-based cancer registries or active follow-up (Germany, Greece and France) and confirmed through a mixture of methods that included health insurance records, and cancer and pathology registries. Participants were followed until cancer diagnosis, death or last complete follow-up, whichever occurred first. Fifty-seven neuroendocrine PC cases were censored. After a mean follow-up of 14 years, 1,283 first incidence PDAC cases

were available for analysis and were classified according to International Classification of Diseases for Oncology third edition codes C25.0–C25.3 and C25.7–C25.9.

Information on lifestyle, dietary and nut intake

Anthropometric measures were assessed at baseline, and participants also answered a lifestyle questionnaire.¹⁸ Country-specific validated dietary questionnaires, with the timeframe referring to the preceding year, were used to assess dietary information at baseline.¹⁸ The determination of nut and seed intake in EPIC has been previously published.^{19,20} Briefly, the term “nut” denotes a combination of three terms: tree nuts (including almonds, Brazil nuts, cashews, hazelnuts, macadamia nuts, pecans, pine nuts, pistachios and walnuts), peanuts (including peanut butter) and nonspecific nuts (not specified by the participant). Generally, in the EPIC cohort, there was a low intake of specific seeds (i.e., sunflower, linseed and pumpkin), thus “seeds” were combined as a sum total variable. Finally, total intake of nuts and seeds was used as the main exposure variable (herein referred to as “total nut intake”). Consumers were defined as those who reported an intake >0 g per day on average.

Statistical analysis

Cox proportional hazards models were used to estimate hazards ratio (HR) and 95% confidence intervals (95% CIs) for total nut intake and overall PDAC risk. Total nut intake was analyzed both as a continuous variable (15 g/day; 15 g-increments correspond to half a standard serving),²¹ and as categorical variable with all nonconsumers as the reference category and consumers categorized in quartiles based on the distribution of total nut intake in the EPIC cohort. All statistical models had age as the primary time variable, were stratified by study center to control for center effects and by age at recruitment in 1-year categories. Covariates of gender, smoking status, diabetes, alcohol consumption, body mass index (BMI) and total energy intake were included in final models as they were known PC risk factors or potential confounders. Other variables such as physical activity using the Cambridge index, education level, magnesium (mg/day), red and processed meat, fiber, vegetable and fruit intake (all in g/day) were evaluated but not included as they did not change the HR estimates $\geq 10\%$. We also evaluated sex-specific and country-specific categorical variables for total nut intake; however, since HRs for total nut intake and PDAC risk did not vary from those of the main model, results were not shown.

Analyses for effect-measure modification were carried out by known PC risk factors: smoking (never, ever), diabetes (yes, no) and BMI (<25 , ≥ 25 kg/m²). Heavy alcohol consumption (>60 , 0.1–4.9 g/day) was evaluated for men and women combined.²² Stratified analyses by sex (male, female), by geographic region (northern: Norway, Denmark and Sweden; central: Germany, The Netherlands, the United Kingdom and northern of France; southern: southern of France, Italy, Spain and Greece) and by country intake (countries over vs

countries below the EPIC median nut intake) were also investigated.

Sensitivity analyses were performed: (i) exclusion of PDAC cases that were diagnosed during the first 2 years of follow-up to minimize the possible effect of preclinical disease on dietary intake; (ii) restriction to microscopically confirmed PDAC cases ($n = 910$) to reduce a possible disease misclassification; (iii) adjustment for the *adapted-relative* Mediterranean diet score (removing nuts from the score)²³; (iv) evaluation of nut intake in quartiles of frequency (never/almost never, 0.2–1 times/month, 0.25 to ≤ 1 times/week, >1 times/week), rather than absolute intake.²⁴ This analysis was performed excluding 47,171 participants from Cambridge and Malmö, as frequency data were not available; (v) removing BMI and diabetes from the multivariable model and (vi) modeling waist-to-hip ratio instead of BMI.

The proportional hazards assumption was evaluated using Schoenfeld residuals, which was satisfied in all models. The median value for each category was estimated and included in a score test to evaluate dose–response trends. The likelihood ratio test p -value was used to evaluate statistical significance of effect-measure modification based on the continuous intake variable. All analyses were performed using SAS v.9.3 and STATA v.14 was used to test the proportional hazards assumption. An α -level of 0.05 was used to set the cutoff for statistical significance.

Results

Basic information on cohort members

After a mean follow-up of 14 years, 1,283 PDAC cases (57% women) were observed. More than 90% of the populations from The Netherlands, Germany and Greece reported consuming nuts and seeds, whereas only 38.8% of the Spanish population reported nut/seed consumption. However, the highest median of intake among consumers was observed in Spain (5.9 g/day), followed by Greece (5.3 g/day) and the Netherlands (5.0 g/day) (Table 1). Even though the distributions of intake were skewed, means of intake by country are presented in Table 1 to compare to some previously published reports.

Participants classified at the highest levels of total nut intake were more likely to have higher energy, dietary fiber, vegetable and fruit intakes, while nonconsumers had higher intakes of processed meat. Furthermore, nonconsumers compared to high consumers tended to be nonalcohol drinkers, had higher BMI and a higher proportion of smokers and were more likely to report diabetes at baseline (Table 2).

Overall PDAC risk

No associations and no evidence for linear dose–response trends were observed between total nut intake and PDAC risk in EPIC (highest intake vs nonconsumers: HR, 0.89; 95% CI, 0.72–1.10; p -trend = 0.70) (Table 3). The continuous total nut intake variable, assessed in 15 g/day increments, was nonsignificantly inversely associated (HR_{15g/day}, 0.94; 95% CI, 0.84–1.07).

Table 1. Estimated total nut¹ intake and pancreatic ductal adenocarcinoma cases by country in the EPIC cohort

Country	Cohort sample N	Person-years	Total PDAC cases N (%) ²	Male PDAC cases N (%) ²	Female PDAC cases N (%) ²	Microscopically confirmed cases N (%)	Consumers of total nuts N (%)	Total nuts among consumers (g/day) Median (p25–p75)	Mean \pm SD
France	67,403	869,372	56 (4.4)	56 (7.7)	56 (7.7)	26 (2.9)	48,381 (71.8)	4.8 (2.1–8.6)	7.7 \pm 9.0
Italy	44,545	630,951	104 (8.1)	35 (6.3)	69 (9.5)	66 (7.3)	39,667 (89.0)	0.2 (0.2–1.0)	1.0 \pm 2.1
Spain	39,989	637,947	106 (8.3)	55 (9.9)	51 (7.0)	80 (8.8)	15,200 (38.8)	5.9 (2.1–13.7)	11.3 \pm 15.6
United Kingdom	75,416	1,122,765	188 (14.7)	69 (12.4)	119 (16.4)	19 (2.1)	63,134 (83.7)	2.3 (0.5–7.0)	7.1 \pm 12.3
The Netherlands	36,539	524,671	93 (7.3)	20 (3.6)	73 (10.0)	67 (87.4)	34,402 (94.2)	5.0 (1.8–11.9)	9.3 \pm 12.8
Greece	26,048	281,284	44 (3.4)	25 (4.5)	19 (2.6)	18 (2.0)	24,173 (92.8)	5.3 (0.2–8.0)	6.3 \pm 8.5
Germany	48,557	504,479	116 (9.0)	72 (13.0)	44 (6.1)	88 (9.7)	43,597 (89.9)	1.5 (0.6–3.7)	3.9 \pm 7.7
Sweden	48,674	801,130	204 (15.9)	93 (16.7)	111 (15.3)	197 (21.6)	30,489 (62.6)	0.3 (0.0–0.8)	1.5 \pm 4.2
Denmark	55,014	815,097	325 (25.3)	187(33.6)	138 (19.0)	303 (33.3)	40,815 (74.2)	0.8 (0.8–1.6)	2.4 \pm 4.5
Norway	33,975	452,171	47 (3.7)	47 (6.5)	47 (6.5)	46 (5.10)	16,105 (47.4)	3.0 (3.0–6.4)	4.6 \pm 3.9
Total	476,160	6,639,867	1,283 (100)	556 (100)	727 (100)	910 (100)	355,963 (74.8)	2.3 (0.7–5.7)	5.3 \pm 9.5

¹Total nut intake is the sum of the total intake of nuts and seeds.

²Percentages are calculated by country.

Abbreviations: EPIC, European Prospective Investigation into Cancer and Nutrition; PDAC, pancreatic ductal adenocarcinoma; SD, standard deviation.

Table 2. Estimated total nut¹ intake and covariates at baseline used in the analyses

	Total nut intake (g/day)				
	Nonconsumers n g/day	Q1 >0-0.7 g/day Median: 0.3 g/day	Q2 >0.7-2.3 g/day Median: 1.0 g/day	Q3 >2.3-5.7 g/day Median: 3.5 g/day	Q4 >5.7 g/day Median: 11.8 g/day
Participants (n)	120,197	89,014	86,213	91,788	88,948
PDAC (n)	400	223	303	196	161
Female (%)	72.0	65.4	65.7	75.0	71.7
Age at recruitment	53.6 (47.9-60.2) ²	50 (40.7-57.9)	53.7 (49.1-59)	49.8 (43.4-56.2)	49.2 (43.3-55.8)
Energy intake (Kcal/day)	1887 (1,536-2,315)	1908 (1,544-2,360)	2049 (1,695-2,467)	1964 (1,626-2,375)	2,216 (1,832-2,665)
Dietary fiber (g/day)	20.7 (16.4-25.8)	20.3 (16.1-25.2)	22.3 (17.9-27.6)	21.9 (17.8-26.8)	24.3 (19.8-29.7)
Magnesium (mg/day)	324.4 (266.6-394.1)	306.6 (251.3-372.8)	356.7 (297.7-426.2)	349.6 (293.2-416.5)	399.7 (333.5-479.7)
Vegetables (g/day)	169.3 (104.9-262.7)	143.7 (86.2-236.3)	163.3 (108.0-241.6)	203.7 (127.3-316.9)	207.3 (129.2-326.5)
Fruit (g/day)	185.2 (100.0-307.6)	206.8 (110.3-330.9)	170.7 (94.0-279.9)	200.9 (114.2-318.5)	210.6 (118.7-327.9)
Red meat (g/day)	35.1 (17.9-61.5)	26.2 (12.2-49.5)	49.5 (26.9-75.9)	31.4 (12.8-58.6)	35.6 (12.3-66.7)
Processed meat (g/day)	26.9 (13.4-46.4)	22.6 (9.7-41.0)	25.9 (14.2-45.0)	20.0 (5.0-40.6)	23.6 (6.5-45.4)
Body Mass Index (kg/m ²)	25.3 (22.8-28.4)	25.1 (22.7-27.9)	25.0 (22.6-27.8)	24.4 (22.1-27.3)	24.1 (21.9-27.1)
Time quitting smoking (year)	14.5 (6.5-23.0)	13.0 (6.5-21.0)	15.0 (7.0-23.5)	14.0 (6.5-22.0)	13.5 (96.5-21.5)
Smoking status (%)					
Never	48.4	53.0	44.8	50.3	48.4
Former	25.0	25.5	28.8	26.8	27.7
Smoker	24.0	20.6	25.1	20.4	21.4
Unknown	2.6	0.9	1.3	2.5	2.5
Diabetes (%)					
No	81.4	94.9	89.9	89.8	92.6
Yes	3.5	2.7	2.2	2.2	2.2
Do not know	0.5	0.0	1.6	0.2	0.1
Missing	14.5	2.4	6.3	7.7	5.1
Alcohol at recruitment					
Nondrinker (%)	26.7	9.6	8.1	9.5	8.6
Consumers among women (g/day)	4.4 (1.4-10.9)	2.9 (0.8-9.8)	6.9 (2.2-14.7)	5.3 (1.6-11.8)	7.0 (2.3-15.3)
Consumers among men (g/day)	14.1 (5.4-33.1)	8.0 (2.8-23.7)	17.2 (8.1-35.9)	15.1 (6.5-30.6)	16.6 (6.8-33.1)

¹Total nut intake is the sum of the total intake of nuts and seeds.

²Median (p25-p75).

Abbreviation: PDAC, pancreatic ductal adenocarcinoma.

Table 3. Hazard ratios and 95% confidence intervals for estimated total nut intake¹ and PDAC risk in EPIC

	Total nut intake										Trend test p-value ²	LRT p-value ³	
	Nonconsumers 0 g/day					Consumers							
	-15 g/day increments	Q1 >0-0.7 g/day Median: 0.3 g/day	Q2 >0.7-2.3 g/day Median: 1.0 g/day	Q3 >2.3-5.7 g/day Median: 3.5 g/day	Q4 >5.7 g/day Median: 11.8 g/day	HR (95% CI)	N cases	HR (95% CI)	N cases	HR (95% CI)			N cases
Final model													
N cases	1,283	223	303	196	161								
HR (95% CI) ⁴	0.94 (0.84-1.07)	0.92 (0.74-1.14)	0.92 (0.77-1.09)	0.99 (0.82-1.20)	0.89 (0.72-1.10)	1.00 (ref)							0.70
Sensitivity analysis													
Cases diagnosed ≥2 years after recruitment													
N cases	1,195	208	284	181	148	374							
HR (95% CI)	0.95 (0.84-1.08)	0.93 (0.74-1.17)	0.92 (0.77-1.10)	0.99 (0.81-1.20)	0.89 (0.72-1.10)	1.00 (ref)							0.75
Microscopically confirmed cases													
N cases	910	140	242	125	104	229							
HR (95% CI)	0.95 (0.66-1.38)	0.72 (0.46-1.13)	0.87 (0.65-1.15)	0.99 (0.71-1.37)	0.68 (0.44-1.04)	1.00 (ref)							0.28
Mediterranean diet score													
N cases	1,283	223	303	196	161	400							
HR (95% CI) ⁵	0.94 (0.84-1.07)	0.92 (0.74-1.14)	0.92 (0.77-1.09)	0.99 (0.82-1.20)	0.89 (0.72-1.10)	1.00 (ref)							0.70
Analyses of effect-measure modification													
Alcohol consumption													
Light drinkers (0.1-4.9 g/day)													
N cases	373	99	75	53	35	111							
HR (95% CI) ⁶	0.72 (0.52-1.01)	0.91 (0.62-1.32)	1.10 (0.79-1.54)	0.82 (0.57-1.19)	0.65 (0.42-0.99)	1.00 (ref)							0.74
Heavy drinkers (≥60 g/day)													
N cases	71	4	24	7	10	26							
HR (95% CI) ⁶	0.79 (0.43-1.47)	0.53 (0.14-1.99)	0.76 (0.40-1.46)	0.45 (0.17-1.19)	0.61 (0.25-1.52)	1.00 (ref)							0.57
Diabetes													
No													
N cases	1,075	195	260	159	145	316							
HR (95% CI) ⁷	0.94 (0.82-1.07)	0.82 (0.65-1.04)	0.85 (0.71-1.03)	0.91 (0.73-1.13)	0.89 (0.71-1.12)	1.00 (ref)							0.38
Yes													
N cases	74	19	14	12	6	23							
HR (95% CI) ⁷	0.93 (0.58-1.51)	1.96 (0.83-4.62)	1.28 (0.57-2.86)	1.35 (0.60-3.07)	0.72 (0.26-1.99)	1.00 (ref)							0.40

(Continues)

Table 3. Hazard ratios and 95% confidence intervals for estimated total nut intake and PDAC risk in EPIC (Continued)

	Total nut intake						Trend test p-value ²	LRT p-value ³
	Nonconsumers 0 g/day		Consumers					
	-15 g/day increments	Q1 >0-0.7 g/day Median: 0.3 g/day	Q2 >0.7-2.3 g/day Median: 1.0 g/day	Q3 >2.3-5.7 g/day Median: 3.5 g/day	Q4 >5.7 g/day Median: 11.8 g/day			
Body Mass Index								
<25 kg/m ²	535	88	118	82	78			
N cases								
HR (95% CI) ⁸	0.98 (0.82-1.16)	1.00 (ref)	0.84 (0.65-1.10)	0.92 (0.69-1.24)	0.90 (0.66-1.22)		0.66	
≥25 kg/m ²	748	135	185	114	83			0.34
N cases								
HR (95% CI) ⁸	0.91 (0.77-1.09)	1.00 (ref)	0.97 (0.77-1.21)	1.03 (0.80-1.33)	0.87 (0.66-1.15)		0.81	
Smoking Status								
Never smokers	465	90	99	73	61			
N cases								
HR (95% CI) ⁹	0.91 (0.73-1.14)	1.00 (ref)	0.97 (0.72-1.30)	0.99 (0.72-1.37)	0.94 (0.66-1.32)		0.98	
Ever smokers ¹⁰	783	128	197	120	94			0.88
N cases								
HR (95% CI) ⁹	0.89 (0.75-1.05)	1.00 (ref)	0.88 (0.71-1.09)	1.00 (0.78-1.27)	0.81 (0.62-1.06)		0.47	

¹Total nut intake is the sum of the total intake of nuts and seeds.

²All p-values for trend are based on the quintile medians.

³All LRT p-values for effect measure modification are based on the continuous total nut intake variable.

⁴Stratified by age at recruitment and center. Adjusted for gender, total energy intake (1,000 kcal/day), BMI (kg/m²), smoking status (never smokers, current pipe or cigar or occasional smokers, current cigarette smokers: 1-15, 16-25, or ≥26 cigarettes/day, former cigarette smokers who quit >20 years, 11-20 years or ≤10 years before recruitment), diabetes (no, yes, do not known, missing) and alcohol consumption (nondrinkers, drinkers of 0-6 g/day, >6-12 g/day, >12-24 g/day, ≥24-60 g/day, female drinkers with >60 g/day, male drinkers >60-96 g/day, male drinkers >96 g/day).

⁵Final model further adjusted by the adapted relative Mediterranean diet score.

⁶Stratified by age at recruitment and center. Adjusted for gender, total energy intake (1,000 kcal/day), BMI (kg/m²), smoking status (never smokers, current pipe or cigar or occasional smokers, current cigarette smokers: 1-15, 16-25 or ≥26 cigarettes/day, former cigarette smokers who quit >20 years, 11-20 years or ≤10 years before recruitment) and diabetes (no, yes, do not known, missing).

⁷Stratified by age at recruitment and center. Adjusted for gender, total energy intake (1,000 kcal/day), BMI (kg/m²), smoking status (never smokers, current pipe or cigar or occasional smokers, current cigarette smokers: 1-15, 16-25 or ≥26 cigarettes/day, former cigarette smokers who quit >20 years, 11-20 years or ≤10 years before recruitment) and alcohol consumption (nondrinkers, drinkers of 0-6 g/day, >6-12 g/day, >12-24 g/day, ≥24-60 g/day, female drinkers with >60 g/day, male drinkers >60-96 g/day, male drinkers >96 g/day).

⁸Stratified by age at recruitment and center. Adjusted for gender, total energy intake (1,000 kcal/day), smoking status (never smokers, current pipe or cigar or occasional smokers, current cigarette smokers: 1-15, 16-25 or ≥26 cigarettes/day, former cigarette smokers who quit >20 years, 11-20 years or ≤10 years before recruitment), diabetes (no, yes, do not known, missing) and alcohol consumption (nondrinkers, drinkers of 0-6 g/day, >6-12 g/day, >12-24 g/day, ≥24-60 g/day, female drinkers with >60 g/day, male drinkers >60-96 g/day, male drinkers >96 g/day).

⁹Stratified by age at recruitment and center. Adjusted for gender, total energy intake (1,000 kcal/day), BMI (kg/m²), diabetes (no, yes, do not known, missing) and alcohol consumption (nondrinkers, drinkers of 0-6 g/day, >6-12 g/day, >12-24 g/day, ≥24-60 g/day, female drinkers with >60 g/day, male drinkers >60-96 g/day, male drinkers >96 g/day).

¹⁰Ever smokers: former and current smokers.

Abbreviations: EPIC, European Prospective Investigation into Cancer and Nutrition; HR, hazards ratio; LRT, likelihood ratio test; PDAC, pancreatic ductal adenocarcinoma.

The three sensitivity analyses performed excluding cases with follow-up of <2 years, restricting to microscopically confirmed cases and adjusting for adapted relative Mediterranean diet score showed similar results (Table 3). Likewise, when total nut intake was analyzed using frequency of consumption, no statistically significant inverse association was observed (p -trend = 0.23; data not shown). Results remained unchanged when diabetes and BMI were not included in final models, and when BMI was replaced by waist-to-hip ratio (data not shown).

Effect-measure modification

No effect-measure modification was observed for any of the stratified analyses according to heavy alcohol consumption, diabetes, smoking status or BMI (Table 3). There was also no evidence for modification of HRs for total nut intake and PDAC by sex, geographic region or country intake (likelihood ratio test p -value = 0.31, 0.42 and 0.50 respectively; data not shown).

Discussion

The present study prospectively assessed the association between total nut intake and PDAC risk in the EPIC cohort. Although all relative risk estimates were below the null value, our study failed to detect any statistically significant inverse associations for men and women. Likewise, relative risks were somewhat lower when we restricted the analysis to microscopically confirmed PDAC cases, but no statistically significant associations were observed. Results for total nut intake and PDAC remained unchanged when we evaluated effect-measure modification by various subgroups.

Regular nut consumption has been associated with health benefits in both epidemiological and clinical studies. Regular nut consumption may play a role in reducing insulin resistance, inflammation, hyperglycemia and oxidative stress among others.^{12,24} Despite differences in nutritional composition by nut subtypes (i.e., walnuts have the highest content in linoleic acid and α -linolenic acid, hazelnuts in fiber, peanuts in protein and folate, pine nuts in polyunsaturated fatty acids), they are considered highly nutritious.²⁵ Therefore, nuts have been postulated as a food group that might have potential in cancer prevention and in lowering cancer mortality; however, the epidemiologic evidence remains limited, particularly for specific cancers.^{9–11}

To our knowledge, only four published studies have evaluated the potential preventive role of nut intake on PC risk, with inconclusive results. The first study was conducted in the Netherlands and encompassed 164 PC cases and 480 controls from both genders. The authors concluded that there was no association between the intake of “nuts and tasty snacks” (including peanuts and other nuts, peanut butter and chips among others) and PC risk.¹⁵ The second study was performed in the prospective Nurses’ Health Study of 75,680 women, where the frequency of nut consumption (defined by the sum of peanuts and other nuts) was statistically significantly inversely associated with PC risk showing an HR of 0.65 (95% CI, 0.47–0.92; p -trend = 0.007).¹⁴ The third study,

from the Golestan Cohort Study (Iran), included 50,045 participants and 54 PC cases, and found no association between nut intake and PC risk.¹⁷ The most recent study was conducted in the Netherlands Cohort Study and evaluated the association between consumption of nuts (sum of peanuts and tree nut), tree nuts, and peanut butter and the risk of PC overall and by sex. Despite observing lower relative risks for higher consumers compared to nonconsumers (HR, 0.84; 95% CI, 0.63–1.11; p -trend = 0.17), none of the associations or trend tests were statistically significant.¹⁶

Our results are consistent with those from the case-control study, the Netherlands Cohort Study and the Golestan Cohort Study studies, but not with the Nurses’ Health Study. Nonetheless, we advise caution when comparing results across studies since nut consumption was assessed differently, and the types of nuts consumed differed between studies. In EPIC, as discussed by Jenab *et al.*, the exposure variable was a combination of tree nuts and seeds (\approx 90% nuts, of which walnuts, almonds and hazelnuts were the most regularly consumed),^{19,20} whereas in both the Nurses’ Health Study and the NCLS, peanuts were more frequently consumed than tree nuts. Similarly, in EPIC-Netherlands, peanuts composed more than half of total nut and seed intake, but no associations with PDAC were observed in our investigation when country-specific analyses were performed, including EPIC-Netherlands. Nuts and peanuts have different nutritional composition, and thus, they may play different roles in human health. In the present study, we could not analyze them as a unique variable, but additional studies should try to evaluate these foods items separately.

Dietary guidelines recommend a minimum portion of 30 g/day of nuts, seeds and legumes as they may have beneficial effects on human health.²¹ In the present study, only 2% of nut consumers reported an intake >30 g/day. The general population may have a misconception about nut consumption, that they are thought to increase weight due to their high caloric value; however, in our study, as well as in other published prospective and clinical studies, it has been observed that high nut consumers have lower BMI and less weight gain compared to nonconsumers.^{24,26} Some studies have suggested a lower risk of Type-2 diabetes among nut consumers as well, although data have not been consistent.¹² Both excess weight and Type-2 diabetes are established risk factors for PC; however, removing BMI or diabetes from the multivariable model, or analyzing waist-to-hip ratio as a measure of abdominal fatness instead of BMI, did not materially alter our results.

The present study had the following limitations: although we tried to control for confounding effects, we could not adjust our models for all known PC risk factors (i.e., family history, ABO blood group, chronic pancreatitis) because this information was not collected in EPIC. Some but not all EPIC dietary questionnaires were designed to capture nut consumption, thus misclassification of the exposure and the possibility that some foods that contribute to total nut intake in EPIC were not assessed (i.e., *turrón* in Spain) could have also influenced

our results. Furthermore, as mentioned before, specific analyses by type of nut could not be performed, including for peanut butter. The range of total nut intake in EPIC was narrower than reported by the Netherlands Cohort Study and the Nurses' Health Study.^{14,16} Lastly, there is only one dietary assessment on all EPIC participants (which was conducted at baseline), thus we were not able to evaluate changes in diet over time.

One of the strengths of our study is its prospective design, in which recall bias is less likely than in case-control studies, and EPIC is a multicountry cohort with heterogeneity in diets and lifestyle factors. We performed a sensitivity analysis excluding cases diagnosed within the first 2 years of follow-up to avoid any influence of prediagnostic PC on dietary intakes, which showed similar results to the overall model. Moreover, this is the largest study evaluating the association between total nut intake and PDAC risk to date (including men and women and over a thousand of PC cases), which allowed us to evaluate effect measure modification by several parameters.

In conclusion, the results of the present study indicate that there were no statistically significant inverse associations between total nut intake and PDAC risk within a large European cohort.

Acknowledgements

The authors thank the CERCA Program/Generalitat de Catalunya for their institutional support. M.O.-S. received a post-doctoral fellow from

References

- Rahib L, Smith BD, Aizenberg R, et al. Projecting cancer incidence and deaths to 2030: the unexpected burden of thyroid, liver, and pancreas cancers in the United States. *Cancer Res* 2014;74:2913–21.
- Kamisawa T, Wood LD, Itoi T, et al. Pancreatic cancer. *Lancet Lond Engl* 2016;388:73–85.
- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. *CA Cancer J Clin* 2019;69:7–34.
- Cooperman AM, Iskandar ME, Wayne MG, et al. Prevention and early detection of pancreatic cancer. *Surg Clin North Am* 2018;98:1–12.
- Landi S. Genetic predisposition and environmental risk factors to pancreatic cancer: a review of the literature. *Mutat Res* 2009;681:299–307.
- Maisonneuve P, Lowenfels AB. Risk factors for pancreatic cancer: a summary review of meta-analytical studies. *Int J Epidemiol* 2015;44:186–98.
- Barone E, Corrado A, Gemignani F, et al. Environmental risk factors for pancreatic cancer: an update. *Arch Toxicol* 2016;90:2617–42.
- Koushik A, Spiegelman D, Albanes D, et al. Intake of fruits and vegetables and risk of pancreatic cancer in a pooled analysis of 14 cohort studies. *Am J Epidemiol* 2012;176:373–86.
- Aune D, Keum N, Giovannucci E, et al. Nut consumption and risk of cardiovascular disease, total cancer, all-cause and cause-specific mortality: a systematic review and dose-response meta-analysis of prospective studies. *BMC Med* 2016;14:207.
- Gonzalez CA, Salas-Salvado J. The potential of nuts in the prevention of cancer. *Br J Nutr* 2006;96(Suppl 2):S87–94.
- Bao Y, Han J, Hu FB, et al. Association of nut consumption with total and cause-specific mortality. *N Engl J Med* 2013;369:2001–11.
- de Souza RGM, Schincaglia RM, Pimentel GD, et al. Nuts and human health outcomes: a systematic review. *Nutrients* 2017;9:1311.
- Casari I, Falasca M. Diet and pancreatic cancer prevention. *Cancer* 2015;7:2309–17.
- Bao Y, Hu FB, Giovannucci EL, et al. Nut consumption and risk of pancreatic cancer in women. *Br J Cancer* 2013;109:2911–6.
- Bueno de Mesquita HB, Maisonneuve P, Runia S, et al. Intake of foods and nutrients and cancer of the exocrine pancreas: a population-based case-control study in The Netherlands. *Int J Cancer* 1991;48:540–9.
- Nieuwenhuis L, van den Brandt PA. Total nut, tree nut, peanut, and peanut butter consumption and the risk of pancreatic cancer in The Netherlands cohort study. *Cancer Epidemiol Biomarkers Prev* 2018;27:274–84.
- Ghorbani Z, Poursahms A, Fazeltabar Malekshah A, et al. Major dietary protein sources in relation to pancreatic cancer: a large prospective study. *Arch Iran Med* 2016;19:248–56.
- Riboli E, Hunt KJ, Slimani N, et al. European Prospective Investigation into Cancer and Nutrition (EPIC): study populations and data collection. *Public Health Nutr* 2002;5:1113–24.
- Jenab M, Ferrari P, Slimani N, et al. Association of nut and seed intake with colorectal cancer risk in the European Prospective Investigation into Cancer and Nutrition. *Cancer Epidemiol Biomarkers Prev* 2004;13:1595–603.
- Jenab M, Sabate J, Slimani N, et al. Consumption and portion sizes of tree nuts, peanuts and seeds in the European Prospective Investigation into Cancer and Nutrition (EPIC) cohorts from 10 European countries. *Br J Nutr* 2006;96(Suppl 2):S12–23.
- World Cancer Research Fund/American Institute for Cancer Research. Continuous Update Project Expert Report 2018. Recommendations and public health and policy implications. [Internet]. 2018. Available from: <https://www.wcrf.org/dietandcancer>.
- Naudin S, Li K, Jaouen T, et al. Lifetime and baseline alcohol intakes and risk of pancreatic cancer in the European Prospective Investigation into Cancer and Nutrition study. *Int J Cancer* 2018;143:801–12.
- Buckland G, Travier N, Cottet V, et al. Adherence to the Mediterranean diet and risk of breast cancer in the European Prospective Investigation into Cancer and Nutrition cohort study. *Int J Cancer* 2013;132:2918–27.
- Freisling H, Noh H, Slimani N, et al. Nut intake and 5-year changes in body weight and obesity risk in adults: results from the EPIC-PANACEA study. *Eur J Nutr* 2017;57:2399–408.
- Ros E. Health benefits of nut consumption. *Nutrients* 2010;2:652–82.
- Martínez-González MA, Bes-Rastrollo M. Nut consumption, weight gain and obesity: epidemiological evidence. *Nutr Metab Cardiovasc Dis* 2011;21:S40–5.

24 days of stem cells

Shape the future of stem cell innovation
October 1- November 1, 2019

Join us for 24 Days of Stem Cells; a premiere virtual event featuring the latest advances in stem cell research.

This year's format will feature a new hour of cutting edge content every week day starting October 1st. Attend the sessions that are most relevant to your work - at your convenience and at your pace.

During the 24-day long event, you can:

- Access leading scientific presentations from thought leaders around the world
- Watch live training demonstrations from our stem cell experts
- Download key stem cell tools and resources
- Complete weekly challenges to earn points towards certification and prizes

Register today at
www.24daysofstemcells.com

ThermoFisher
SCIENTIFIC

WILEY