

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Hidden paleosols at high elevation in the Alps (Stolenberg Plateau - NW Italy): evidence for a Lateglacial Nunatak?**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1888625> since 2023-01-30T15:47:56Z

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.


(Article begins on next page)



# Hidden paleosols at high elevation in the Alps (Stolenberg Plateau - NW Italy): evidence for a Lateglacial Nunatak?






Pintaldi E.<sup>1</sup>, D'Amico M.<sup>1</sup>, Colombo N.<sup>1,2,3</sup>, Giardino M.<sup>2,3</sup>, Freppaz M.<sup>1,2</sup>  
<sup>1</sup>University of Turin - DISAFA, <sup>2</sup>University of Turin, NATRISK, <sup>3</sup>University of Turin – DST.



The study was performed in the periglacial environment of the Stolenberg Plateau (LTER site Istituto Mosso), located at the foot of the southern slope of the Monte Rosa Massif (Western Italian Alps, elevation: ca. 3030 m a.s.l.). The plateau is covered by thick periglacial blockfields and blockstreams, with a plant cover that reaches no more than 3-5% of the surface.

Under the periglacial landforms well-developed soils were observed, characterized by thick (between 30 and 65 cm) umbric horizons, under which discontinuous cambic Bw ones were developed. The organic carbon (C) stocks were surprisingly high (above 5 kg\*m<sup>-2</sup>). Geophysical investigations showed that these soils are widespread under the stony cover, with a thickness ranging between 20 and 90 cm.



Radiocarbon dating ranged between 20.5 and 4.4 ka cal. BP, indicating that these soils are paleosols, probably originated during the main warming phases/interstadials occurred between the end of Last Glacial Maximum and the beginning of the Neoglacial.

The results suggest that the Stolenberg Plateau could represent a Lateglacial Alpine Nunatak on which specific pedoclimatic conditions were probably suitable for alpine plant life and pedogenesis already since 21-20 ka BP.