

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**EFFECT OF THERANOSTIC PHOSPHOLIPIDIC SRCL2/SRI2AND  $\beta$ -D FRUCTOPYRANOSE BASED NANOPARTICLES ON CELL VIABILITY AND UPTAKE**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1891490> since 2023-02-09T11:38:59Z

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

## Introduction

Nanoparticles can be engineered with precise sizes, shapes, composition and with surface targeting ligands that allow a specific interaction with target cells.

The aim of this work was to analyze the effect of non-toxic and low-cost Sr/I-based nanoparticles, in order to develop a new  $^{89}\text{Sr}$ - and/or  $^{131}\text{I}$ -based radiotherapy method on cell lines. In particular we analyzed the effect on cellular uptake and cell viability of compound 1 ( $^{89}\text{Sr}(\text{fructose})_2\text{Cl}_2 \cdot \text{H}_2\text{O}$ ) and compound 2 ( $^{89}\text{Sr}(\text{fructose})_2\text{I}_2$  and  $\text{Sr}(\text{fructose})_2^{131}\text{I}_2$ ), on a non-tumour cell line, hTERT-HME1, and a tumour cell line, HT-29. The compounds studied exhibit some features that could be interesting for biomedicine applications, such as the biocompatibility due to their non-toxic components and the significant SH emission, that can permit exploitation for *in vitro* bio-imaging.

## Materials and methods

We tested the effect of nanoparticles encapsulated in a hydrophilic shell of mPEG-DSPE that increases their biocompatibility on non-tumour and tumour cell lines, hTERT-HME1 and HT-29, respectively. In particular, we analyzed the cellular uptake of nanoparticles and the effect on cell viability. To evaluate the effect of nanoparticles on cell viability, we treated hTERT-HME1 and HT-29 cell lines with five different nanoparticles concentrations, and cell viability was assessed using ATP content and the CellTiter-Glo® Luminescent Cell Viability Assay (Promega).

## Results

hTERT-HME1 cell line showed sensitivity to compound 1 already at the lowest concentration, while compound 2 reduced hTERT-HME1 cell viability in a dose-dependent manner; HT-29 viability was reduced of about 70% and 50% only at the highest concentration of nanoparticles of compounds 1 and 2, respectively. The nanoparticles covered with mPEG-DSPE can be detected for few days in a biological medium and can be internalized into hTERT-HME1 and HT-29 cells.

## Discussion and Conclusions

In this work we have analyzed the effect of Sr/I-based nanoparticles, on cell uptake and viability. The nanoparticles formulation could be useful to concentrate the effect of  $^{89}\text{Sr}$  and  $^{131}\text{I}$  radio-API in a specific tissue, allowing a selective activity of the activated compounds. The nanoparticles are biocompatible and are internalized into the cell; the viability of cell lines utilized is influenced by the presence of the nanoparticles themselves rather than their chemical composition.