



ABSTRACT

Aeromedical retrievals as a measure of potentially preventable hospitalisations and cost comparison with provision of GP-led primary health care in a remote community

Part of Special Series: WONCA World Rural Health Conference Abstracts 2022 

AUTHORS



Lara Wieland¹ lwieland@vtown.com.au, Medical Student *



Gail Abernethy² Research Assistant



CORRESPONDENCE

*Dr Lara Wieland

AFFILIATIONS

¹ James Cook University, Tarzali, Qld, Australia; and Torres and Cape Hospital and Health Service, Weipa, Qld, Australia

² James Cook University, Tarzali, Qld, Australia

PUBLISHED

10 January 2023 Volume 23 Issue 1

HISTORY

RECEIVED: 20 September 2022

ACCEPTED: 20 September 2022

CITATION

Wieland L, Abernethy G. Aeromedical retrievals as a measure of potentially preventable hospitalisations and cost comparison with provision of GP-led primary health care in a remote community. *Rural and Remote Health* 2023; 23: 8150.

<https://doi.org/10.22605/RRH8150>

This work is licensed under a [Creative Commons Attribution 4.0 International Licence](https://creativecommons.org/licenses/by/4.0/)

ABSTRACT:

Introduction: Kowanyama is a very remote Aboriginal community. It is ranked amongst the top five most disadvantaged communities in Australia and has a very high burden of disease. Currently, the community has access to GP-led Primary Health Care (PHC) 2.5 days a week for a population of 1200 people. This audit aims to assess whether GP access correlates with retrievals and/or hospital admissions for potentially preventable conditions and whether it is cost effective and improves outcomes to provide the benchmarked staffing of GPs.

Methods: A clinical audit was undertaken of aeromedical retrievals for the year 2019 to assess whether access to a rural GP would have prevented the need for retrieval and whether the condition for

retrieval was 'preventable' or 'not preventable'. A cost-analysis was undertaken to compare the cost of providing accepted benchmark levels of GPs in community with the cost of potentially preventable retrievals.

Results: There were 89 retrievals of 73 patients in 2019. 61% of all retrievals were potentially preventable. Most (67%) of the preventable retrievals occurred with no doctor on site. For preventable condition retrievals, the mean number of visits to the clinic compared with non-preventable condition retrievals was higher for registered nurse or health worker visits (1.24 vs 0.93) and lower for GP visits (0.22 vs 0.37). The conservatively calculated costs

of retrievals for 2019 matched the maximum cost of providing benchmark numbers (2.6 FTE) of rural generalist (RG) GPs in a rotating model for the audited community.

Conclusion: Greater access to GP-led PHC appears to lead to fewer retrievals and hospital admissions for potentially preventable conditions. It is likely that some preventable condition retrievals would be avoided if a GP was always on site. Providing benchmarked numbers of RG GPs in a rotating model in remote communities is cost-effective and would improve patient outcomes.

This PDF has been produced for your convenience. Always refer to the live site <https://www.rrh.org.au/journal/article/8150> for the Version of Record.