

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

One gene, two proteins: alternative splicing in human CD157/Bst1 unmasked

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1797255> since 2021-08-17T17:23:31Z

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

One gene, two proteins: alternative splicing in human CD157/Bst1 unmasked

Ferrero E¹, Lo Buono N², Morone S¹, Parrotta R¹, Giacomino A¹, Augeri S¹, Mancini C³, Rosal-Vela A⁴, Garcia-Rodriguez S⁴, Zubiaur M⁴, Sancho J⁴, Ortolan E¹ & Funaro A¹.

¹*Immunogenetics Lab, Dept. of Medical Sciences, University of Torino, Torino, Italy.*

²*San Raffaele Diabetes Research Institute, San Raffaele Hospital, Milan, Italy.*

³*Medical Genetics Lab, Dept. of Medical Sciences, University of Torino, Torino, Italy.*

⁴*Dept. of Cell Biology and Immunology, IPBLN-CSIC, Granada, Spain.*

Until now, CD157 (alias Bst1) has lived a double life: one as a receptor, expressed in human neutrophils and monocytes, that regulates leukocyte diapedesis through interactions with the extracellular matrix, and one as an ectoenzyme of the ADP-ribosyl cyclase family of NAD-cleaving enzymes that generates nucleotide derivatives that somehow trigger endocellular Ca²⁺ mobilization. For decades, we have known but one form of human CD157: a 318 amino acid protein generated from one transcript (BST1-001) encoded by the 9-exon BST1 gene on chromosome 4(p15). However, RT-PCR experiments revealed the existence of a second BST1 transcript. Database mining, molecular cloning and sequence analysis of this transcript, designated BST1-002, revealed that it contained a previously undescribed exon. Located between canonical exons 1 and 2, this exon was named exon 1b. Exclusion of exon 1b by alternative splicing yields canonical CD157, whereas inclusion of the exon 1b-encoded sequence in frame yields CD157-002, a novel proteoform of 333 amino acids. Neutrophils express both canonical and CD157-002 proteins, although the canonical transcript is the dominant form. Both forms are also co-expressed in other tissues. By comparative functional assays, the proteoforms show overlapping properties in mAb (anti-CD157 SY/11B5) binding, subcellular localization and migration. However, NAD glycohydrolase activity was found in canonical CD157 alone, CD157-002 appearing to be catalytically inert in our experimental conditions. As the 10-exon organization of human BST1 is a departure from the 9-exon organization found in other mammals, we performed phylogenetic analyses to trace the origins of the novel exon, identifying the exon 1b sequence as dynamically evolving and conserved in BST1 in primate evolution. We believe that it is important to bear in mind its quadruple life forms when dealing with human CD157 in basic molecular and biomedical studies.